Curvature from unlabelled two-dimensional causal orders

Alexander Kolpakov
Center for Science, Technology, Engineering, and Mathematics
University of Austin, Austin, Texas, United States
akolpakov@uaustin.org
The Wolfram Institute for Computational Foundations of Science
akolpakov@wolframinstitute.org

Research draft, 21 October 2026

Abstract

Consider independent events in a two-dimensional causal rectangle with a positive smooth volume density, and retain only their unlabelled causal order. We prove that every two-order representation recovers the empirical null ranks to within $O(\log n/n)$, up to exchanging the two directions, outside an event of probability $O(\log^2 n/n)$. The proof combines classical modular decomposition with an empty-rectangle argument for densities bounded above and below. For a bounded C^4 class with fixed volume and uniform null marginals, this gives an estimator of scalar curvature at the marginal centre with mean squared error $O(n^{-2/5})$. A matching lower bound holds even when all event coordinates are observed. The permutation restrictions control the ambiguity of the latent coordinates; the remaining two quotient frames form a genuine two-element torsor. The statistical rate and the modular machinery have established antecedents. The contribution addressed here is their quantitative transfer to the partial-order observation model.

1 The observation and the question

Let c be a positive density on $[0,1]^2$ with uniform marginals:

$$\int_0^1 c(u,v)\,dv = \int_0^1 c(u,v)\,du = 1. \tag{1.1}$$

The metric on the open square is

$$g_c = -2c(u,v)\,du\,dv. \tag{1.2}$$

Its volume is one and its chronology is the strict product order. Sample $Z_i = (U_i, V_i)$ independently with density c, then observe only the unlabelled poset

$$i <_P j \quad\Longleftrightarrow\quad U_i < U_j,\ V_i < V_j.$$

A causal Hasse diagram supplies the same information after transitive closure. Sampling labels, coordinates, an embedding and a null frame are not observed.

Uniform marginals calibrate the null coordinates by marginal volume. They do not assert that the two coordinates are independent. For a physical rectangle of known volume V, the normalized model has metric $-2Vc\,du\,dv$ and all curvatures below are divided by V. The rectangle and its

volume are part of the model. There is no claim of selecting a frame equivariantly in an unbounded Lorentz-invariant sprinkling.

The target point is $z_* = (1/2, 1/2)$, invariant under exchange of the two null directions. The scalar curvature, with the convention used throughout these notes, is

$$R_c(z_*) = \left(\frac{2c_{uv}}{c^2} - \frac{2c_u c_v}{c^3} \right) \Bigg|_{z_*} . \tag{1.3}$$

Indeed $\Gamma^u_{uu} = \partial_u \log c$ and $\Gamma^v_{vv} = \partial_v \log c$ are the only nonzero Christoffel symbols; contracting the Ricci tensor gives $R = 2c^{-1}\partial_u \partial_v \log c$.

Order-based embedding and curvature recovery are established subjects. Two-order models of causal rectangles already underlie Brightwell–Henson–Surya [7], including their discussion of correlated null coordinates for a nonconstant conformal factor. Henson [6] reconstructs approximate flat embeddings. Benincasa–Dowker [9] construct order-based curvature operators. The copula–permutation correspondence is standard; see Grübel [4]. Tian [12] studies curvature from paired null ranks, explicitly distinguishing those ranks from their partial-order intersection. Our question is whether that intersection suffices at the same statistical exponent on a full-support smooth class.

2 The restrictions and their ambiguity

For a finite set V, the bijections $\mathrm{Lab}(V) = \mathrm{Bij}(V, [n])$ form an S_n-torsor. Pairs of rank bijections form an $S_n \times S_n$-torsor. The admissible subset for P is

$$\mathcal{R}(P) = \{(r_1, r_2) : x <_P y \iff r_1(x) < r_1(y),\ r_2(x) < r_2(y)\}.$$

This subset is generally not itself a torsor. Its members are the ordered two-realizers of P. The history presentation in [11] also uses restrictions of permutation torsors, there to describe token wirings and schedules. Here the restrictions specify latent coordinate orders with a given intersection; a causal sampling law is an additional assumption.

Fix one such pair. A common interval is a vertex set consecutive in both orders. Let b be the largest size of a proper common interval, counting singletons. Relabel the first order increasingly to obtain the usual permutation representation.

Lemma 2.1. *Suppose $b < n/2$. For any other ordered two-realizer (r'_1, r'_2), either directly or after exchanging its two orders,*

$$\max_{x \in V} \max_{j=1,2} |r'_j(x) - r_j(x)| \le b - 1. \tag{2.1}$$

After contracting the maximal proper strong modules, exactly two ordered quotient realizers remain. Exchange of the two orders acts freely and transitively on them.

Proof. We use the classical modular representation of a permutation graph: strong modules are precisely strong common intervals in every realizer, and all realizers are obtained by composing realizers of the quotient graphs in its modular decomposition. Prime quotients have four ordered graph realizers, obtained by exchanging the orders and reversing both. These facts, including the strong-module assertion, are stated in Crespelle–Paul [1, Section 2.2, Proposition 1].

Apply them to the incomparability graph of P. The root of its modular decomposition is prime. Otherwise its quotient is complete or empty, so the children occur monotonically in the two orders. Each child has size at most b. Remove an end child: the union of the remaining children is a proper common interval with size at least $n - b > b$, a contradiction.

Let $B_1, \dots, B_k$ be the root's children. Each is consecutive in both orders of every realizer, and $|B_i| \le b$. The prime quotient has four graph realizers. Reversing both orders reverses the nonempty comparable relation of the quotient poset. It therefore cannot preserve P. The two remaining quotient realizers exchange the orders. After making this global exchange if necessary, the blocks have exactly the same order and the same rank ranges in both representations. Only ranks within a block can change, by at most $|B_i| - 1$. This proves (2.1) and the final assertion. □

The decomposition, recognition and prime-realizer statements are known. The lemma records their quantitative consequence for latent sampling ranks. It does not claim a new modular decomposition theorem. The residual two-element set is a $\mathbb{Z}/2$-torsor of null frames. Fine-scale choices inside modules are additional ambiguities. Uniqueness of two-realizers for prime posets is also classical; see Pouzet–Zaguia [8, Section 7.2] for the finite result of El-Zahar–Sauer and its extension. The two-element frame ambiguity itself is not a novelty claim.

3 Localization for positive sampling densities

Assume for this section that

$$0 < a \le c \le B < \infty \tag{3.1}$$

and that (1.1) holds. No differentiability is needed yet.

Theorem 3.1. *There are constants K, C depending only on a, B such that, for all sufficiently large n, the following event has probability at least $1 - C\log^2 n/n$: every proper common interval of the sampled ranks has at most $K \log n$ elements. Consequently every ordered two-realizer of the observed poset agrees with the empirical null ranks within $K \log n/n$ after normalization, up to one global exchange of axes.*

Proof. Set $\tau = A \log n/n$, with a sufficiently large constant A depending on a. We establish three events.

Empty rectangles. With failure probability $O(n^{-6})$, every open axis-aligned rectangle of area greater than τ contains a sampled point. To prove this, use the uniform grid with $\lceil 8/\tau \rceil$ subdivisions. An empty rectangle of area greater than τ contains an empty grid rectangle of area at least $\tau/2$: moving each of its four sides inward by at most one mesh loses area at most $\tau/2$. There are $O(\tau^{-4})$ grid rectangles. Any specified one of area at least $\tau/2$ is empty with probability at most $\exp(-an\tau/2)$. A union bound proves the claim when A is large enough.

Occupancy of short intervals. With failure probability $O(n^{-6})$, every interval of the u-axis of length at most 4τ contains at most $12n\tau$ sampled U_i's. Such an interval is contained in one of the $O(\tau^{-1})$ intervals $[j\tau, (j+6)\tau] \cap [0,1]$. Uniform marginals bound the corresponding binomial mean by $6n\tau$. The exponential Markov bound for a binomial variable gives $\mathbb{P}(X \ge 2\mu_0) \le \exp(-\mu_0/3)$ when its mean is at most μ_0. Take $\mu_0 = 6n\tau$ and use another union bound.

Corner exclusion. With failure probability at most $64Bn\tau^2$, no sample lies in one of the four corner squares of side 4τ. This follows from the density upper bound and a union bound over the n points.

On these three events, let S be a proper common interval with at least two elements. Write its coordinate bounding rectangle as $I \times J$, with widths s, t. There is no sample in $I \times ([0,1] \setminus J)$ or in $([0,1] \setminus I) \times J$. Each set is a union of two rectangles. The empty-rectangle event gives

$$s(1-t) \le 2\tau, \qquad t(1-s) \le 2\tau.$$

If one width is at most $1/2$, the other is at most 4τ, and substituting back gives both widths at most 4τ, for $\tau \le 1/8$. If both widths exceed $1/2$, both are at least $1-4\tau$. Any sample outside S must then lie outside both I and J, and therefore in a corner square of side 4τ. Corner exclusion contradicts properness of S. Thus both widths are small. The occupancy event gives $|S| \le 12n\tau = 12A\log n$. Singletons satisfy the same bound. The total failure probability is $O(\log^2 n/n)$. For large n, $12A\log n < n/2$; apply Lemma 2.1. □

For uniform random permutations, substantially sharper common-interval results are classical: typically only size-two proper blocks remain. See Bouvel–Chauve–Mishna–Rossin [3, Theorem 3] and the earlier work cited there. Theorem 3.1 uses a weaker logarithmic bound uniformly over the positive copula class. It does not assume that the sample permutation is uniform.

4 An estimator from the order alone

Fix $0 < a < 1 < B$ and $M > 0$. Let $\mathcal{C}$ be the densities satisfying (1.1)–(3.1), extending as C^4 functions to the closed square, with

$$\max_{1\le|\nu|\le 4} \|\partial^\nu c\|_\infty \le M.$$

The model is genuinely nonparametric. For example $1 + \theta\sin(2\pi u)\sin(2\pi v)$ belongs to a suitable fixed class for every $|\theta| \le \theta_0 < 1$.

Uniformly label the observed abstract poset, using its labelling torsor, and apply any fixed rule that selects an ordered two-realizer. Such a rule can use transitive orientation of the incomparability graph, a standard permutation-graph algorithm [1]. The ranks $(\widehat{U}_i, \widehat{V}_i) = (r_1(i)/n, r_2(i)/n)$ are the only coordinates used by the estimator. Uniform labelling makes its law independent of any presentation of the unlabelled input. It is not an assertion that all two-realizers are equiprobable under the unknown sampling law.

Choose the even, compactly supported kernel

$$k(t) = \frac{315}{256}(1-t^2)^4 \mathbf{1}_{\{|t|\le 1\}}.$$

It is C^3, integrates to one and has zero first moment. For $\nu = (r,s) \in \{(0,0),(1,0),(0,1),(1,1)\}$ put

$$T_\nu = \frac{1}{nh^{2+|\nu|}} \sum_{i=1}^n k^{(r)}\left(\frac{1/2-\widehat{U}_i}{h}\right) k^{(s)}\left(\frac{1/2-\widehat{V}_i}{h}\right). \tag{4.1}$$

Project T_{00} onto $[a,B]$ and each other T_ν onto $[-M,M]$; denote the results by $\widetilde{T}_\nu$. Define

$$\widehat{R} = \frac{2\widetilde{T}_{11}}{\widetilde{T}_{00}^2} - \frac{2\widetilde{T}_{10}\widetilde{T}_{01}}{\widetilde{T}_{00}^3}. \tag{4.2}$$

Exchange of axes leaves (4.2) unchanged. It is bounded in absolute value by $R_{\max} = 2M/a^2 + 2M^2/a^3$.

Theorem 4.1. *With $h = n^{-1/10}$, there is a constant $C = C(a,B,M)$ such that*

$$\sup_{c\in\mathcal{C}} \mathbb{E}_c|\widehat{R} - R_c(z_*)|^2 \le Cn^{-2/5}$$

for sufficiently large n. The bound holds for every realizer-selection rule described above.

Proof. Let E_n be the event of Theorem 3.1 and $\epsilon_n = K \log n/n$. Since (4.2) is invariant under exchanging axes, we may align the selected ranks with the true empirical ranks on E_n. Writing $F_{U,n}, F_{V,n}$ for their empirical distribution functions, set

$$\Delta_n = \max(\|F_{U,n} - \mathrm{id}\,\|_\infty, \|F_{V,n} - \mathrm{id}\,\|_\infty).$$

The uniform marginals and the Dvoretzky–Kiefer–Wolfowitz inequality [5] give

$$\mathbb{P}(\Delta_n > t) \le 4e^{-2nt^2}, \qquad \mathbb{E}\Delta_n^4 \le C/n^2. \tag{4.3}$$

Dependence of U_i and V_i within a point does not affect either one-dimensional empirical distribution bound. On E_n, the aligned estimated coordinates satisfy

$$\max_i \|(\widehat{U}_i, \widehat{V}_i) - (U_i, V_i)\|_\infty \le \Delta_n + \epsilon_n. \tag{4.4}$$

Let T_ν^0 be (4.1) evaluated at the true sample coordinates. Integration by parts, a symmetric-kernel Taylor expansion, and $|\nu| \le 2$ give

$$|\mathbb{E}T_\nu^0 - \partial^\nu c(z_*)| \le Ch^2, \qquad \mathrm{Var}(T_\nu^0) \le C/(nh^{2+2|\nu|}). \tag{4.5}$$

The first bound uses derivatives of c up to order $|\nu| + 2 \le 4$. For the second, the kernel is bounded by $Ch^{-2-|\nu|}$ and supported on a square of probability at most $4Bh^2$.

For large n, $\epsilon_n \le h/2$. Restrict first to $\Delta_n \le h/2$. Let N_h count the true points in the square of radius $2h$ about z_*. A summand can change in (4.1) only if its true point lies in this square. The bounded derivative of the scaled kernel and (4.4) imply

$$|T_\nu - T_\nu^0| \le Ch^{-3-|\nu|}(\Delta_n + \epsilon_n)N_h/n \quad \text{on } E_n \cap \{\Delta_n \le h/2\}. \tag{4.6}$$

Here N_h is binomial with success probability at most $16Bh^2$. When $nh^2 \ge 1$, expanding its second and fourth moments yields

$$\mathbb{E}(N_h/n)^2 \le Ch^4, \qquad \mathbb{E}(N_h/n)^4 \le Ch^8.$$

Cauchy–Schwarz and (4.3) give

$$\mathbb{E}\{\Delta_n^2(N_h/n)^2\} \le Ch^4/n.$$

These bounds do not assume independence of N_h and Δ_n. Squaring (4.6), with $|\nu| \le 2$, gives

$$\mathbb{E}\big[|T_\nu - T_\nu^0|^2; E_n, \Delta_n \le h/2\big] \le Ch^{-6}(n^{-1} + \epsilon_n^2).$$

The true c and its derivatives belong to the projection intervals, so projecting cannot increase their estimation errors. The map in (4.2) is Lipschitz on the product of these intervals. On the complementary event its squared error is at most $4R_{\max}^2$. Combining the preceding bounds gives

$$\mathbb{E}|\widehat{R} - R_c(z_*)|^2 \le C\left\{h^4 + \frac{1}{nh^6} + \frac{\log^2 n}{n^2h^6} + \frac{\log^2 n}{n} + e^{-nh^2/2}\right\}.$$

For $h = n^{-1/10}$ the first two terms are $n^{-2/5}$; all the others are smaller for sufficiently large n. □

Corollary 4.2. *Let $\widehat{R}, \widehat{R}'$ use the same kernel, bandwidth $h = n^{-1/10}$ and projection intervals, but any two possibly randomized rules for selecting the two-realizer of the same input. Under any coupling of these selections,*

$$\sup_{c \in \mathcal{C}} \mathbb{E}_c|\widehat{R} - \widehat{R}'|^2 \le C \log^2 n/n.$$

Thus the choice between admissible realizers contributes a smaller order than the minimax estimation error.

Proof. On E_n, align each pair with the true empirical ranks by its global axis exchange. The resulting pairs differ by at most $2\epsilon_n$ pointwise. On $\{\Delta_n \le h/2\}$ the argument of (4.6) therefore gives

$$|T_\nu - T'_\nu| \le Ch^{-3-|\nu|}\epsilon_n N_h/n.$$

The binomial second-moment bound gives squared expectation at most $C\epsilon_n^2 h^{-6}$. Projection is nonexpansive and the curvature map is Lipschitz on the projection intervals. On the complement both scalar estimators are bounded by $R_{\max}$. Hence their squared difference has expectation at most

$$C\{\epsilon_n^2 h^{-6} + \log^2 n/n + e^{-nh^2/2}\}.$$

Substitution proves the claim. Every step is uniform in the selection rules and their auxiliary randomness. □

5 A matching information bound

Theorem 5.1. *There is $c_0 > 0$ such that*

$$\inf_{\widehat{S}} \sup_{c\in\mathcal{C}} \mathbb{E}_c|\widehat{S} - R_c(z_*)|^2 \ge c_0 n^{-2/5}$$

for sufficiently large n, even if $\widehat{S}$ observes all point coordinates. Thus the unlabelled-order experiment attains the full-coordinate minimax exponent.

Proof. Let q be a smooth odd function supported in $(-1,1)$ with $q'(0) \ne 0$. For a small fixed $\gamma > 0$ set

$$c_h(u,v) = 1 + \gamma h^4 q((u-1/2)/h)q((v-1/2)/h).$$

The integral of q is zero, so both marginals remain exactly one. For $h < 1/2$ and sufficiently small γ, c_h and the flat density belong to $\mathcal{C}$. Derivatives of order $j \le 4$ are bounded by a constant times γh^{4-j}. At z_*, the density equals one, both first derivatives vanish, and $(c_h)_{uv} = \gamma h^2 q'(0)^2$. Thus the curvature difference is $2\gamma h^2 q'(0)^2$.

For one full coordinate observation,

$$\mathrm{KL}(c_h\,du\,dv\|du\,dv) = \int (c_h \log c_h - c_h + 1)\,du\,dv \le C\gamma^2 h^{10}.$$

For n independent observations, the bound is $Cn\gamma^2 h^{10}$. Take $h = n^{-1/10}$ and γ small enough that Pinsker's inequality bounds total variation by $1/2$. The nearest-curvature test associated with any estimator has sum of testing errors at least $1/2$. If the curvature difference is d_h, a classification error forces squared estimation error at least $d_h^2/4$. The larger of the two risks is therefore at least $d_h^2/16$, of order $n^{-2/5}$. Forgetting coordinates or labels cannot improve this testing bound. □

The exponent is the classical two-dimensional second-derivative estimation exponent. The content here is its attainment from the partial order rather than from supplied coordinates or paired ranks. It is not a new density-estimation rate.

Corollary 5.2. *Replace the C^4 assumption by bounded derivatives through order two, with each second derivative Hölder continuous of exponent $0 < \alpha \le 1$ and seminorm at most M. For this class, curvature at z_* can be estimated from the unlabelled order with minimax mean squared error of order*

$$n^{-2\alpha/(6+2\alpha)}.$$

The corresponding expected absolute error has minimax order $n^{-\alpha/(6+2\alpha)}$.

Proof. The localization theorem is unchanged. In (4.5) the bias is bounded by Ch^α: for the mixed derivative use its Hölder continuity directly, and for the lower derivatives use the same symmetric Taylor expansion. The perturbation and variance terms are unchanged. Choose $h = n^{-1/(6+2\alpha)}$. The term $\log^2 n/n$ remains smaller than the resulting squared risk. For the lower bound replace h^4 in $c_h - 1$ by $h^{2+\alpha}$. This preserves the Hölder bound and both marginals. The curvature separation is of order h^α and the n-sample relative entropy is at most $Cnh^{6+2\alpha}$. The same two-point test proves the squared-risk lower bound. Cauchy–Schwarz gives the absolute-error upper bound; the nearest-curvature test gives its matching lower bound. □

This is the exponent reported for the paired-rank experiment in Tian's abstract [12]. Corollary 5.2 observes only the intersection order. This comparison concerns the estimation rate in the stated positive class, not finite-sample equivalence of the two observation models.

6 Example, torsor role, and remaining work

For the Farlie–Gumbel–Morgenstern density

$$c_\theta(u,v) = 1 + \theta(1-2u)(1-2v), \qquad |\theta| \le \theta_0 < 1,$$

the central curvature is 8θ. This classical dependence family [10] has comparable-pair probability $1/2 + \theta/9$. Indeed its lower-left cumulative distribution is $uv\{1+\theta(1-u)(1-v)\}$, and comparability has probability twice the integral of this function against c_θ. Expanding the integral gives the stated value. Uniformly label an unlabelled sample of $2m$ events and pair the labels. The m comparability indicators Y_j are then independent Bernoulli variables with this parameter. Consequently

$$\widehat{R}_{\mathrm{pair}} = 72\left(m^{-1}\sum_{j=1}^{m} Y_j - \tfrac{1}{2}\right)$$

is unbiased for central curvature, with mean squared error at most $1296/m$. This is a parametric worked example, not a new copula or a general reconstruction method. Theorem 4.1 also applies to nonparametric perturbations and to asymmetric positive copulas. There is no requirement that a finite list of pair frequencies identify the density.

For an asymmetric worked example, put

$$f(x) = 1 - 2x, \qquad p(x) = 6x^2 - 6x + 1,$$

and take

$$c(u,v) = 1 + \theta\{f(u)f(v) + \lambda f(u)p(v) + \mu p(u)f(v)\}.$$

Both f and p have zero integral and absolute value at most one. Thus the marginals are uniform, and $|\theta|(1 + |\lambda| + |\mu|) < 1$ ensures positivity. At the marginal centre the density is one, its first derivatives are $\theta\lambda, \theta\mu$, and its mixed derivative is 4θ. Consequently

$$R_c(z_*) = 8\theta - 2\theta^2\lambda\mu.$$

For $(\theta, \lambda, \mu) = (2/5, 3/10, 1/2)$ the density lies between $7/25$ and $43/25$, and the curvature is $394/125$. The order-only estimator applies without knowing these parameters or assuming this polynomial family. Exchanging the axes interchanges λ, μ and preserves the target.

The torsor has two precise uses. Uniform labelling gives an invariant randomized observation procedure. Restricting the product of the two labelling torsors defines the admissible coordinate

pairs. Their modular quotient leaves a genuine two-element torsor; curvature descends under its axis exchange. The proof also controls the fine ambiguity left inside modules. Calling the whole realizer fibre a torsor would be false.

Positivity throughout the rectangle is substantive. With disconnected or vanishing-density support, large common blocks can persist, and coordinate choices inside those blocks need not agree up to one global axis exchange. The theorem does not cover that case, noisy causal comparisons, higher order dimension, or arbitrary rewrite ensembles.

Wolfram's discussion of causal coordinates [14, Section 5.14] motivates the question of extracting geometry from an event order. The present result supplies a rigorous reconstruction protocol under specified sampling hypotheses. It does not derive those hypotheses from a rewriting rule or derive an equation of motion. Two-dimensional Einstein dynamics is not claimed.

The strongest candidate contribution is the transfer from partial orders to curvature estimation through localization of all realizers. The relevant combinatorial structure is classical [1, 2, 3], as are copula ranks [4] and statistical derivative estimation. Tian's recent preprint [12] is especially close; its abstract was consulted, but the full text was unavailable during this audit. Ghosh [13] also reports finite causal-set sampling and identifiability results; only the repository description was accessible. These comparisons must be completed before claiming publication priority.

Reproduction

The exact checker `check_rank_ambiguity.py` enumerates all ordered two-realizers for the 1 840 base permutations of sizes four through seven that satisfy the deterministic interval hypothesis. All 7 456 realizers obey the bound in Lemma 2.1. The separate checker `check_curvature.py` checks the kernel's normalization and boundary jets, the asymmetric example's marginals, and its curvature directly from the metric. These finite checks supplement the proofs; they do not test an asymptotic rate by simulation. Run `make causal-order-paper` and `make causal-order-check` from the repository root. The exact outputs are included beside the source.

References

[1] C. Crespelle and C. Paul, *Fully dynamic algorithm for recognition and modular decomposition of permutation graphs*, Algorithmica **58** (2010), 405–432. Author's full version.

[2] F. Bassino, M. Bouvel, V. Féray, L. Gerin and A. Pierrot, *Dense and nondense limits for uniform random intersection graphs*, arXiv:2402.06394.

[3] M. Bouvel, C. Chauve, M. Mishna and D. Rossin, *Average-case analysis of perfect sorting by reversals*, Author's full version, Theorem 3.

[4] R. Grübel, *Ranks, copulas, and permutons*, Metrika **87** (2024), 155–182, doi:10.1007/s00184-023-00908-2.

[5] P. Massart, *The tight constant in the Dvoretzky–Kiefer–Wolfowitz inequality*, Annals of Probability **18** (1990), 1269–1283, doi:10.1214/aop/1176990746.

[6] J. Henson, *Constructing an interval of Minkowski space from a causal set*, Classical and Quantum Gravity **23** (2006), L29–L35, arXiv:gr-qc/0601069.

[7] G. Brightwell, J. Henson and S. Surya, *A 2D model of causal set quantum gravity: the emergence of the continuum*, Classical and Quantum Gravity **25** (2008), 105025, arXiv:0706.0375.

[8] M. Pouzet and I. Zaguia, *Hereditary classes of ordered sets of width at most two*, arXiv:2112.02633, Section 7.2.

[9] D. M. T. Benincasa and F. Dowker, *The scalar curvature of a causal set*, Physical Review Letters **104** (2010), 181301, arXiv:1001.2725.

[10] D. J. G. Farlie, *The performance of some correlation coefficients for a general bivariate distribution*, Biometrika **47** (1960), 307–323, doi:10.1093/biomet/47.3-4.307.

[11] A. Kolpakov, *Causal histories, observable states, and cubical geometry*, accompanying research manuscript, 2026, repository source and PDF.

[12] C. Tian, *Recovering Lorentzian curvature from finite volumes and paired ranks*, preprint posted 16 September 2026, SSRN 7472426. Abstract-only comparison.

[13] A. Ghosh, *Sample-complexity bounds and an exact bulk-sector channel obstruction for curvature estimation on finite causal sets*, 11 August 2026, Zenodo, version 4. Repository-description-only comparison; the earlier version is doi:10.5281/zenodo.21894198.

[14] S. Wolfram, *A Class of Models with the Potential to Represent Fundamental Physics*, 2020, Section 5.14.