

UMGS: UAV Multispectral Gaussian Splatting for Spectral-Index Synthesis – Supplementary Material

ANONYMOUS AUTHOR(S)
SUBMISSION ID: TBD

CCS Concepts: • **Computing methodologies** → **Image-based rendering**; **Reconstruction**.

Additional Key Words and Phrases: multispectral imaging, gaussian splatting, 3D reconstruction, UAV

ACM Reference Format:

Anonymous Author(s). 2026. UMGS: UAV Multispectral Gaussian Splatting for Spectral-Index Synthesis – Supplementary Material. 1, 1 (May 2026), 3 pages. <https://doi.org/10.1145/nnnnnnn.nnnnnnn>

1 Protocol Notes and Scope

This supplement aligns with the current main-paper draft and records additional tables that are useful for reproducibility and reviewer inspection.

The main paper focuses on 10 UAV/UAV-style scenes. This supplement separately reports four additional official aligned scenes whose viewpoints are not part of the UAV-focused main scope. The retained-subset diagnostic for raw unaligned input is reported as an applicability check and should not be interpreted as the main raw full-scene comparison protocol.

Depth diagnostics follow the relative reference-depth protocol. Since raw UAV reconstructions are not in a globally calibrated metric coordinate system, all thresholds are relative-depth ratios rather than metric distances.

Comparison shims. UMGS-I-shim and UMGS-J-shim denote UMGS renders bilinearly resized to the external adapter resolution solely for common-mask comparison. They are not distinct training variants and are not used to define the UMGS method.

Grouped split sanitization. For official aligned scenes with inconsistent split and transform metadata, we sanitize splits at the capture-group level against the available transforms. A D/G/R/RE/NIR group is removed only when one or more required members are absent from the transform file, making it unusable as a complete multispectral comparison unit. This is a protocol repair for data completeness, not a performance-based selection.

Depth-reference status. Official aligned depth diagnostics use sparse point-splat references derived from available transforms and point support. They are secondary reference-depth evidence, not dense MVS or LiDAR ground truth. The main geometry claims therefore

use relative-depth consistency and do not claim absolute dense geometry accuracy.

2 Additional Non-UAV Official Scene-Wise External Comparison

Table 1 reports the four additional official aligned scenes that are intentionally kept out of the main UAV-focused tables. Table 2 provides their scene-wise common-mask comparison between UMGS-J-shim and MS-Splatting. We also include the full seven-scene official aligned average for reproducibility.

3 Additional Relative-Depth Threshold Sweep

The clean raw-self depth-agreement sweep across six thresholds is reported in the next table block. It complements the main paper’s compact geometry indicators and makes the trend from strict to relaxed thresholds explicit.

4 Why Unfrozen-Support Is Not a Product Model

The Unfrozen-support branch is included as a single-band upper-bound ablation. It starts from the same RGB anchor as UMGS, but allows band-specific geometry and opacity updates in Stage 2. This can improve selected per-band render metrics, but it breaks the shared-support contract required by model-level spectral products.

The failure is concrete in our audit logs. On `ms_lake`, Unfrozen-support improves several band PSNR values, but product construction fails with an xyz mismatch between bands:

$$\max |\Delta xyz| = 6.84 \times 10^{-2}.$$

Scratch-MS training diverges further and triggers Gaussian-count mismatch across bands. These variants can still produce independent band images, and one could compute post-hoc 2D ratios after rendering; however, those ratios are not defined over a common 3D Gaussian support and therefore are not used as valid model-level multispectral products in our protocol.

5 UMGS-J Initialization Ablation (UMGS-J vs UMGS-JR)

The main paper keeps UMGS-JR out of the main tables and reports UMGS-J as the representative joint branch. This section records the initialization ablation that motivated that decision. Both variants use the same geometry-locked joint multispectral training design and export per-band evaluation views from a unified checkpoint. They differ only in appearance-bank initialization.

UMGS-J initializes each spectral bank from the reset-appearance state. UMGS-JR initializes each spectral bank from the RGB-tied carrier.

Table 5 compares these variants on the two joint-branch pilot scenes used during method screening. UMGS-JR does not improve the branch: it degrades image/index quality on both pilot scenes.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than the author(s) must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.
© 2026 Copyright held by the owner/author(s). Publication rights licensed to ACM.
ACM XXXX-XXXX/2026/5-ART
<https://doi.org/10.1145/nnnnnnn.nnnnnnn>

Table 1. Additional official aligned non-UAV scenes, reported separately from the UAV-focused main-paper suite. RGB metrics for UMGS-J use the shared UMGS-I RGB anchor. Spectral metrics average G/R/RE/NIR.

Method	RGB PSNR	RGB SSIM	MS PSNR	MS SSIM	MS LPIPS	Idx. RMSE	Idx. SSIM	AbsRel	Ag@5%	Ag@10%
UMGS-I	21.67	0.690	25.75	0.763	0.262	0.108	0.710	0.164	0.616	0.688
UMGS-J	21.67	0.690	25.75	0.763	0.262	0.108	0.710	0.164	0.616	0.688

Table 2. Additional non-UAV official aligned scene-level common-mask comparison between UMGS-J-shim and MS-Splatting. PSNR/SSIM/LPIPS average G/R/RE/NIR; RMSE and SAM are spectral 4-band metrics.

Scene	UMGS-J PSNR	MMS PSNR	UMGS-J SSIM	MMS SSIM	UMGS-J LPIPS	MMS LPIPS	UMGS-J RMSE	MMS RMSE	UMGS-J SAM	MMS SAM
ms_bud	24.77	25.09	0.797	0.781	0.225	0.257	0.0630	0.0601	6.46	7.34
ms_fruit	24.18	23.81	0.698	0.701	0.318	0.306	0.0642	0.0670	8.12	8.66
ms_garden	29.44	27.36	0.874	0.820	0.179	0.283	0.0378	0.0460	3.60	4.40
ms_tree	26.45	24.83	0.812	0.757	0.251	0.325	0.0525	0.0644	5.90	7.07

Table 3. Full seven-scene official aligned common-mask average, included only for reproducibility. Lower is better for LPIPS, 4-band RMSE, and SAM.

Method	PSNR	SSIM	LPIPS	4-band RMSE	SAM ^o
UMGS-J-shim	26.55	0.820	0.231	0.0544	5.75
MS-Splatting	26.03	0.803	0.272	0.0571	6.23

Table 4. Clean raw-self relative-depth agreement sweep (mean over G/R/RE/NIR). Thresholds are relative-depth ratios.

Method	0.5%	1%	2.5%	5%	10%	20%
UMGS-I	0.288	0.395	0.607	0.893	0.991	0.997
UMGS-J	0.288	0.395	0.607	0.893	0.991	0.997
Scratch-MS	0.189	0.304	0.469	0.601	0.635	0.638
Unfrozen-support	0.277	0.388	0.599	0.889	0.991	0.997

Table 6 reports the formal reference-depth geometry block on self_m3m. Identical geometry numbers are expected because both joint variants share the same frozen RGB support.

6 Takeaway

The supplementary tables support three practical points used by the draft: (1) additional official aligned scenes remain available for reproducibility but are outside the main UAV-focused scope, (2) relative-depth trends are stable under threshold sweeps when support is preserved, and (3) UMGS-JR is an initialization ablation rather than a promoted branch.

Table 5. UMGS-J and UMGS-JR quality/cost comparison from pilot runs. Band metrics are G/R/RE/NIR means. Index metrics are NDVI/GNDVI/NDRE means.

Scene	Variant	Band SSIM $\uparrow$	Band PSNR $\uparrow$	Band LPIPS $\downarrow$	Index MAE $\downarrow$	Index RMSE $\downarrow$	Index PSNR $\uparrow$	Index SSIM $\uparrow$	Time (s) $\downarrow$
self_m3m	UMGS-J	0.7657	25.330	0.2113	0.0607	0.0863	27.701	0.7943	8129
self_m3m	UMGS-JR	0.7434	24.966	0.2061	0.0623	0.0880	27.519	0.7854	7339
Vineyard A	UMGS-J	0.8804	28.690	0.1092	0.0339	0.0513	32.169	0.8948	5805
Vineyard A	UMGS-JR	0.8590	27.847	0.1266	0.0375	0.0574	31.326	0.8800	6730

Table 6. UMGS-J and UMGS-JR reference-depth geometry on self_m3m. Values are averaged over G/R/RE/NIR. Abs., Med., and Missing are threshold-independent.

Variant	τ	Abs. $\downarrow$	Med. $\downarrow$	Missing $\downarrow$	Agree $\uparrow$	Front $\downarrow$	Too-deep $\downarrow$
UMGS-J	0.05	0.0925	0.0419	0.0002	0.5490	0.2560	0.1949
	0.10	0.0925	0.0419	0.0002	0.7255	0.1697	0.1047
	0.25	0.0925	0.0419	0.0002	0.9150	0.0472	0.0377
	0.50	0.0925	0.0419	0.0002	0.9805	0.0050	0.0143
	1.00	0.0925	0.0419	0.0002	0.9957	0.0003	0.0038
UMGS-JR	0.05	0.0925	0.0419	0.0002	0.5490	0.2560	0.1949
	0.10	0.0925	0.0419	0.0002	0.7255	0.1697	0.1047
	0.25	0.0925	0.0419	0.0002	0.9150	0.0472	0.0377
	0.50	0.0925	0.0419	0.0002	0.9805	0.0050	0.0143
	1.00	0.0925	0.0419	0.0002	0.9957	0.0003	0.0038