

Minimal Computing Cosmology 3.0

Integrated book revision v0.12 incorporating seven submission manuscripts

Wonsik Choi · Jeongin Choi | 10 October 2026

Can one nonprivileged universe be described through minimal computation, a common carrier and phenotype? WRRRA stands for Wonsik Reality–Renderer Architecture. It places event histories, distinctions retained in the present, available resources and subsequent transformations in one state account.

This edition preserves the cumulative 2.3.2 skeleton and WRRRA_M genealogy of the full v0.8 book, replaces its current parent construction with paper v1.8, and incorporates seven studies from the r11 composition paper through record capacity and overflow. It is a full-book editorial revision. Earlier v0.10 was a short current-model statement based on the paper, not an integrated full-length book.

Earlier calculations and interpretations retain their stated preparations, calibrations and readout domains. Closure or integration in the preserved chapters refers to the specified module and computational scope, not a completed physical theory or a proof of every subsequent dynamics. Chapter-level v0.12 updates and Parts XIV–XV specify the revised interfaces. The research genealogy remains while the changed conditions are visible at the relevant locations.

Evaluation and reading order

Read verified inputs → WRRRA transformation → outputs → falsification conditions before the conclusion. Incorporating known constants and observations is legitimate model construction. Reproduction counts as consistency and explanatory achievement; unmeasured quantities computed by the fixed internal map are conditional model predictions. Independent prediction is not required as a criterion of theoretical legitimacy or originality. External results count as integration only where corresponding WRRRA calculations actually execute.

A first reading can begin with Chapters 34–35 on motivation, Chapters 59–64 on energy, response and records, Part XIV on the current parent model, and Chapters 92–99 on the seven-paper connection. Parts I–VIII preserve the foundational skeleton; Parts IX–XIII trace the subsequent construction. Earlier edition dates retained in the body identify the research genealogy.

Main changes in this edition

Location	Revision
Chapters 7, 35, 48	Conditional phenotype, finite-execution philosophy, response sufficiency versus full reconstruction
Chapter 56	Progress through effective conversion, decay and compensation; remaining physical implementation
Chapters 59–65	Energy versus pressure, preparation-dependent response, accumulated records and distinct q branches

Location	Revision
Chapters 67, 73	Evaluation rules, remaining tasks and the adequate-present conclusion
Chapters 84–91	Current parent construction replaced by v1.8
Chapters 92–99	Seven-paper synthesis from r11 through finite-record overflow

Existing internal AI reviews belong to their individual manuscripts, not to a new review of this v0.12 book. This task integrates research materials; it does not report new observations or reruns of every reproduction suite.

Reader guide and local notation

A first route is motivation and philosophy (34–36) → sufficient present and seven studies (92–99) → current calculation (84–91) → earlier derivations as needed. Symbols are local: Chapter 26 uses $q=Q-1$, unlike the deceleration q of Chapter 62. Pressure P , a prime set and a phenotype label are different objects; Hamiltonian H and Hubble H have different roles.

Term	Role in this book
State / density operator	A table of probabilities and quantum coherence, constrained by positivity and unit trace
Rank	Number of independent directions distinguished by the specified linear map
CPTP channel	Transformation preserving valid quantum states and total probability
Holonomy	Global change left after transport around a closed path
Hessian / susceptibility	Second variation of energy / strength of response to an input change
Phenotype	Observable assignment under a specified state and readout rule

Common-carrier implementation	State, operation and output	Connection scope
Abstract 15 channels	Representation space, operators and admissible rank	Foundational structure, not automatic full particle dynamics
128-state load example	Finite density matrix, load operators, sector energies	Macro response for the declared preparation
P4 effective two channels	Species labels, common pulse, conversion probability and work	Supplies P3 transitions; not a derivation of the complete coherent carrier

3.0 Edition Declaration

Minimal Computing Cosmology 3.0 is not a version that discards 2.3.2 and replaces it with a new theory. It preserves the minimal execution architecture, common carrier, matter phenotype, asymmetric unification of the four forces, finite-boundaryless twist universe, and interfaces to standard-physics modules cumulatively integrated in 2.3.2. It then adds, upstream and in the intermediate layers, the high-energy-to-phenotype connection studied in WRRM_M and the finite-SOURCE generation rule introduced in 3.0.

In this book, the symbol $\oplus$ does not denote ordinary arithmetic addition. It denotes a non-substitutive cumulative integration that preserves the original ownership and provenance of each layer:

$$\text{MCC 3.0} = \text{MCC 2.3.2 cumulative skeleton} \oplus \text{WRRM_M 0.1-0.12 structural integration} \oplus \text{finite Klein-Zeta SOURCE candidate}$$

The new central question of 3.0 is:

If the universe is finite, can the size of the SOURCE that executes it, together with the upstream generation structure, also close internally as a finite construction?

MCC 3.0 offers one executable candidate answer and tests whether that state can propagate, within one ledger, through component, phenotype, energy, pressure, gravity, galaxy rotation, lensing, and cosmic expansion.

Preservation of Authorship and Research Genealogy

This 3.0 book edition follows the authorship of the public MCC 3.0 release of 7 October 2026: Wonsik Choi and Jeongin Choi. It does not retroactively alter the authorship of earlier public records. The sole-author public releases of Minimal Computing Cosmology 2.3.2 and Prime Parts retain their original provenance, while the coauthored WRRM_M integrated releases retain theirs.

Assessment Rule

Wherever possible, every computational or model claim in this book is judged in the same order:

$$\text{verified inputs} \rightarrow \text{WRRM-specific transformation} \rightarrow \text{outputs} \rightarrow \text{falsification conditions}$$

Using a known constant or observed value as an input is not itself a failure. When inputs and calibration freedoms are frozen first and a WRRM-specific transformation reproduces a physical value, the result is recorded as a consistency and explanatory achievement. When an unmeasured quantity is produced after the model has been fixed, it is classified separately as a conditional WRRM/MCC prediction.

PASS-C means that the computational integration itself passes, while some key physical generation rules remain conditional assumptions or OPEN. It is not a grade for computational failure. It marks the boundary between what has closed and what has not yet been physically derived.

What Version 3.0 Adds

- The externally convenient cutoff $N = 1,000,000$ is replaced by the finite-SOURCE candidate $N_U = 1,015,000$.
- Between a high-energy state and the present phenotype, 3.0 explicitly places a thermal probe, record formation, the micro/macro boundary, an origin-preserving component layer, and stable codeword/harmonic structure.
- Address, component, and SI-energy ledgers remain different types. Component, channel, generation, and harmonic counts are not allowed to duplicate cosmic energy.
- A genuine change in the upstream population must propagate through frozen downstream coefficients to diagnostics such as q , rotation, lensing, and expansion.
- Conditional outputs already obtained — including the finite-SOURCE size, the acceleration-transition scale, and the inherited neutrino harmonic ledger — are kept distinct from assumptions that remain OPEN.

Concept Map and Quick Terminology

The new terms of 3.0 do not operate independently. Remembering the following execution order makes the later equations and ledgers much easier to read:

LAW → finite SOURCE → FILTER/RESIDUE → RECORD → COMPONENT → CODEWORD/HARMONIC → PHENOTYPE → SI ENERGY/LOAD → GRAVITY/LENSING/EXPANSION

The common carrier is the shared transport basis used by multiple representations and forces; it is not a fifth force. The Standard Model, general relativity, and FLRW remain precise Reality-layer calculation modules. String theory is used only as optional mathematical material when microscopic implementation needs reinforcement.

Quick claim grades

- INPUT: externally fixed model input.
- CALIBRATED: value fixed using observation or a declared calibration condition.
- DERIVED: output calculated from frozen inputs and rules.
- OBSERVED: externally measured data.
- OPEN: a generation law, owner, or value not yet closed.

This distinction prevents one number from being counted simultaneously as both an input and a prediction.

Contents

Current Chapters 1–99. Original module numbers in Parts V and VII and historical Appendix G are separately scoped.

- 1. One Universe One Execution Architecture
- 2. Three Forms of Minimality
- 3. The Minimal SOURCE and the Opening of the Universe
- 4. Where Do Dimensions and Spacetime Come From
- 5. Why the Four Forces Cannot Be the Same Kind of Force
- 6. The Common Carrier How Diversity Emerges from One Structure
- 7. Matter Is a Phenotype Not a List of Numbers
- 8. An Actual World Composed of Matter
- 9. Actual Reality Event
- 10. Fixed Laws and an Open Single History
- 11. The Boundary Between Quantum Possibility and Macroscopic Reality
- 12. Conditions for One Reality to Remain the Same Reality
- 13. Placing Classical Relativistic and Quantum Physics in One Structure
- 14. Turning the Declaration into a Finite Execution System
- 15. Viewing Many Physical Phenomena with One Tool
- 16. The Minimal Execution Grammar Repeated Beyond Cosmology
- 17. What Has Already Been Obtained
- 18. What Still Needs to Be Completed
- 19. What 2.2 Adds
- 20. The Present Universe and the First Twist
- 21. Finite Boundaryless Topology and Two Twist Directions
- 22. Spatial Stress and Dark-Mass Phenotype
- 23. Minimal Galactic Response
- 24. Connecting Cosmological and Galactic Scales
- 25. Gravitational Lensing
- 26. Early-Universe and Structure Response
- 27. Interpretation, Tests and Assessment
- 28. Native Twist Dynamics and the CDM Equivalence Branch
- 29. One Global and Local Twist State

- 30. Final Interface of the Common Carrier and Four Forces
- 31. Compatibility of a Matter Occupied Reality and the Twist Universe
- 32. The Present Universe and Experienced Time
- 33. Final Integrated Propositions
- 34. — From an Exhibition to a Question
- 35. — Is the Universe a Finished File? WRRRA and Minimal Computation
- 36. — Finite without Boundary: Sphere, Möbius Strip, and Klein Bottle
- 37. — The Shadow of the Primes: The Riemann Zeta Function and the Finite Klein-Zeta Generator
- 38. — Why 29 and 70? Generating 1,015,000 Addresses
- 39. — When a Number Enters the Physical Ledger: The Prime/Composite Filter
- 40. — Research Background and Objective
- 41. — Method — Frozen Baseline and Typed Ledger
- 42. — Stage 0.1 — Freezing the Baseline
- 43. — Stage 0.2 — From Temperature to a WRRRA State
- 44. — Stage 0.3 — The Planck Boundary and Filter Stability
- 45. — Stage 0.4 — Cooling and Filter Stabilization
- 46. — Stage 0.5 — Residue, Return, and SOURCE Address Branching
- 47. — Stage 0.6 — Shutter, Event Discreteness, and Physical Time
- 48. — Stage 0.7 — Boundary Between Microscopic Uncertainty and Macroscopic Record
- 49. — Stage 0.8 — Component Activation Boundary and Primitive Alphabet
- 50. — Stage 0.9 — Cooling, Stable Codewords, and Harmonic States
- 51. — Stage 0.10 — Reconnecting Phenotype Energy to Downstream Gravity
- 52. — Stage 0.11 — Reverse Minimality Audit
- 53. — Stage 0.12 — Closing the Master Ledger
- 54. — Structures Newly Secured by the Integrated Study
- 55. — Integrated Falsification Conditions
- 56. — Research Questions After Stage 0.12
- 57. — Position of This Integrated Study within MCC 3.0
- 58. — Conclusion of the High-Energy-to-Phenotype Connection
- 59. — Information Load and Energy
- 60. — Gravity and Twist
- 61. — Galaxy Rotation and Lensing
- 62. — Cosmic Expansion
- 63. — Proper Time and Record
- 64. — End-to-End Master Ledger
- 65. — Reproduction of Verified Values
- 66. — WRRRA/MCC Conditional Predictions
- 67. — Open Assumptions and Falsification Conditions
- 68. — Preservation Rules and the Meaning of Version 3.0

- 69. — Reading the Model Again as One Execution Genealogy
- 70. — Outputs and Conditional Predictions of Version 3.0
- 71. — A Hierarchy of Falsification
- 72. — Final Integrated Propositions of Minimal Computing Cosmology 3.0
- 73. — Conclusion: One Universe Executing a Finite Present
- 74. Introduction and scope
- 75. Finite SOURCE and arithmetic state
- 76. Minimal computation and equivalent implementation
- 77. From the common carrier to channels and phenotype
- 78. Internal calibration and the supplier for a single record
- 79. Correlated physical load and energy pressure
- 80. Gravity rotation lensing and expansion from one state
- 81. Validation results and disclosed construction
- 82. Falsification conditions and conclusion
- 83. Reproducibility and references
- 84. Model development and the three explanatory questions
- 85. Verified inputs and the finite construction
- 86. Exact correlation-preserving interface
- 87. Product-marginal error and dynamical closure
- 88. Shared energy, records and observable equivalence
- 89. Numerical controls and published-data comparison
- 90. Reproducibility and falsification
- 91. Conclusion
- 92. Starting from fractions without stopping at fractions
- 93. From number structure to particle selection
- 94. Can the same ledger survive particle decay?
- 95. How a common carrier implements the selection rule
- 96. Matching energy and pressure together
- 97. Defining the weight of information through response
- 98. A sufficient present and accumulating residue
- 99. One execution structure after seven papers

Parts I-VIII — Preservation of the 2.3.2 Cumulative Structure

The following Parts I-VIII preserve the execution architecture cumulatively integrated in Minimal Computing Cosmology 2.3.2 as the downstream and intermediate-layer foundation of 3.0. Version 3.0 does not delete this material or replace it with the new SOURCE hypothesis. New 3.0-specific changes are identified explicitly in Part IX and later, apart from the ownership boundary added to the string-microphysics section.

Parts I through IV Complete Skeleton of Minimal Computing Cosmology 2.0

Part I The Minimal Execution Structure of the Universe

Big Bang, Dimensions, Laws, and Minimal Input

Chapter 1 One Universe One Execution Architecture

The first claim of Minimal Computing Cosmology is not the proposal of a new particle, but that physical phenomena usually described separately can be rearranged as one process of generation, execution, and recording.

Modern physics is extraordinarily precise, yet it answers a single overarching question in different languages. Classical mechanics describes the motion of bodies, relativity describes spacetime and causal structure, and quantum mechanics describes the amplitudes and probabilities of possible events. But no single ledger explains how laws begin one universe, how matter and forces arise together within it, and why one recorded reality continues among many possibilities.

To fill this gap, Minimal Computing Cosmology proposes WRRRA, the Wonsik Reality-Renderer Architecture, as named in the WRRRA Core 1.0 specification frozen on 7 September 2026. WRRRA does not discard the successful equations of existing theories. Instead, it places each equation into a single execution order according to what it owns and at which stage it acts. Unification here does not mean flattening different theories into one equation; it means connecting the process by which one reality is produced while preserving the distinct responsibilities of the component theories.

Statement

The Book in One Sentence A fixed LAW defines the space of possibilities and relational rules; a minimal SOURCE opens one history; common transport and selective renderers jointly implement spacetime, matter, and forces; and quantum events accumulate as one actual record within that execution.

$LAW \rightarrow SOURCE \rightarrow EXECUTION \rightarrow RENDERER \rightarrow REALITY \rightarrow RECORD$ (1.1)

Here, LAW owns the physical laws, constants, state space, symmetries, conservation laws, and boundary conditions. SOURCE is not a completed blueprint of the universe but the minimal boundary input that opens a prepared possibility. EXECUTION updates the present state into the next present. RENDERER turns internal degrees of freedom into a shared observable representation. REALITY is the physically realized visible sector of the entire Actual, and RECORD is the irreversibly accumulated history of events.

With these distinctions, the Big Bang, dimensions, the four forces, mass, vacuum, antimatter, quantum probability, and a single history are no longer unrelated chapters but different positions in one flow. The purpose of this book is not to rewrite tens of thousands of known facts in new notation. It is to show, with minimal structure, why those facts can be executed together in one universe without contradiction.

Ledger — Established Theory · Primary Question · Position within WRRRA

- Established Theory: Classical Mechanics

- Primary Question: How do realized objects move?
- Position within WRRR: Motion of stable macroscopic phenotypes
- Established Theory: Relativity
 - Primary Question: How do spacetime and causal structure operate?
 - Position within WRRR: Common causal geometry and gravitational ledger
- Established Theory: Quantum Mechanics
 - Primary Question: What amplitudes and probabilities do possible events have?
 - Position within WRRR: Quantum refinement of the allowed event space
- Established Theory: WRRR
 - Primary Question: How do possibilities become one Reality and one Record?
 - Position within WRRR: Higher-level source-to-reality execution architecture

Chapter 2 Three Forms of Minimality

In this theory, ‘minimal’ does not merely mean that an equation is short. It means jointly reducing the number of dimensions that must be physically realized, the number of independent inputs, the amount of transport structure, and the scope that must be recomputed at every moment.

The first is minimal dimensionality. To explain the observable universe, WRRR does not require as a premise a large number of physical extra dimensions that cannot be directly visualized. The physical stage of our universe closes in three spatial directions and one temporal ordering. If additional dimensions are introduced, one must first distinguish whether they are dimensions of an internal representation space, a state space, an arithmetic registry, or actual spacetime dimensions. A large number of internal degrees of freedom does not by itself imply a high-dimensional physical space.

Statement

Academic Meaning of Minimality Independent prediction is one axis for evaluating a theory. But the capacity to explain many already known facts together, without contradiction and with fewer dimensions, inputs, owners, and transport rules, is also a central criterion in theory choice.

The second is minimal structure. If charge, mass, spin, position, the three gauge forces, and gravity are each produced in separate factories and then attached afterward, the number of coupling rules proliferates. WRRR reuses a common state, common transport, common boundaries, and selective representations. Diversity is obtained not by multiplying independent origins, but by allowing the same structure to be read through different ledgers.

The third is minimal computation. The universe need not pre-store every future event or reread its laws from outside at every moment. Fixed laws and the current state generate the next present; observable differences are output only at the boundaries where they are needed; and only events that actually occur are recorded.

$$C_{\text{total}} = L_{\text{LAW}} + L_{\text{SOURCE}} + C_{\text{read}} + C_{\text{evolution}} + C_{\text{render}} + C_{\text{record}} \quad (2.1)$$

Thus, minimization targets the total explanatory cost rather than a single count of operations. If one model requires more physical dimensions, more independent initial values, more separate carriers, and more post hoc corrections to explain the same observations, then it has a larger structural cost over the same explanatory domain.

$$A^* = \text{Pareto-min} \{D_{\text{phys}}, L_{\text{SOURCE}}, N_{\text{owner}}, N_{\text{carrier}}, C_{\text{update}} \mid A \models \Omega_{\text{observed}}\} \quad (2.2)$$

Rather than rank all models by an arbitrary single number, this framework uses Pareto minimality. A structure is a minimal candidate when reducing one item while preserving observational compatibility necessarily increases another. WRRR is a candidate that achieves a broad explanatory range with 3+1 dimensions, one opening, reused common transport, and local updating.

Chapter 3 The Minimal SOURCE and the Opening of the Universe

The Big Bang is not an explosion of a completed data set for the universe. It is a constraint-preserving boundary transition in which a possibility dormant under fixed laws enters one actual history.

If the first cause of the universe contains a list of particles, all future events, and every value at every point in space, the explanation never really begins, because one must again explain who created that information, where it was stored, and how it is read. Minimal Computing Cosmology replaces the first cause as an information list with a single operation acting on a structured possibility.

$$P = \{H_potential, \Omega, P_Q, H_WRA, G, A\}, [\Omega, P_Q] = i \hbar \quad (3.1)$$

$$p_dorm = F(H_potential, G, A, C), \text{ free coefficients} = 0 \quad (3.2)$$

The dormant state is not nothingness but a possibility prepared by law. However, if arbitrary initial coefficients are hidden in this state, the claim that there is only one SOURCE collapses. Therefore, uniqueness of the invariant state, residual degeneracy, block weights, and superselection rules must all be exposed either as part of LAW or as explicit boundary data.

$$U_S = \exp(i\eta Q / \hbar), U_S \dagger P_Q U_S = P_Q + \eta \quad (3.3)$$

The parameter eta is not a particle or a list of energy values. It is a primordial boundary input that starts one execution by shifting the momentum of a prepared opening degree of freedom. Matter and events that appear afterward are not items read out of SOURCE; they are generated through the interaction of LAW with the evolving state.

Restricting the system to flat FLRW geometry and a homogeneous scalar chi, a bare kick that changes only the momentum of chi violates the Hamiltonian constraint. This provides a clear counterexample to a simple primordial-impulse model. By contrast, the following boundary generator shifts the scalar and gravitational momenta together and preserves the constraint.

$$B_\eta(a, \chi) = \eta \chi - \sqrt{6} \tilde{M}_P |\eta| \ln(a/a^*) + \text{constant} \quad (3.4)$$

$$p\chi^+ = \eta, p a^+ = -\sqrt{6} \tilde{M}_P |\eta| / a^*, H^* = |\eta| / (\sqrt{6} \tilde{M}_P a^{*3}) \quad (3.5)$$

The central point of this result is not that ‘the Big Bang has been proven,’ but something more specific. The opening of the universe cannot be an arbitrary impulse acting only on the matter sector; it must be a canonical boundary map that changes the geometry and the energy ledger together. The Big Bang is not an explosion inside space, but a transition in which space, clock, and transport begin to operate together.

3.1 From Opening to the Hot Big Bang

As the simplest effective branch after the constraint-preserving opening, consider an interval in which the kinetic energy of chi dominates. The kination scaling ρ_χ proportional to a^{-6} is a BASELINE result of standard cosmology. What WRA newly connects is this branch directly to the opening with $p_\chi^+ = \eta$, while refusing to insert a radiation component as a new initial value in SOURCE.

$$\rho_\chi^* \approx \eta^2 / (2a^{*6}), \rho_\chi(a) = \rho_\chi^* (a^*/a)^6 \quad (3.6)$$

Radiation is not supplied as an independent primordial register. Suppose that the opening sector and interactions already contained in LAW own a nonnegative energy-transfer function Q_r . Then the total energy ledger can be conserved while the evolution is written as the following pair of equations.

$$\dot{\rho}_\chi + 6H\rho_\chi = -Q_r, \dot{\rho}_r + 4H\rho_r = Q_r, Q_r \geq 0 \quad (3.7)$$

In particular, radiation can be generated during EXECUTION even if $\rho_r(t^*) = 0$ at the instant of opening.

$$\rho_r(t) = a(t)^{-4} \int_{t^*}^t a(t')^4 Q_r(t') dt' \quad (3.8)$$

Therefore, if Q_r is positive on any finite interval, ρ_r becomes positive. No separate radiation seed needs to be inserted as SOURCE data. This is the No Extra Radiation Register condition.

Defining $R = \rho_r / \rho_\chi$, the transfer equations during the kination interval give the following relation.

$$R/R = 2H + Q_r(1/\rho_r + 1/\rho_\chi) \geq 2H \quad (3.9)$$

Thus, while $Q_r \gg 0$, the ratio of radiation to opening-sector energy grows at least as rapidly as a^2 . Even after the transfer ends, ρ_r proportional to a^{-4} while ρ_{χ} proportional to a^{-6} , so sufficiently prolonged expansion leads to radiation domination. In the simple seed limit $Q_r = 0$, with $\epsilon_r = \rho_r^*/\rho_{\chi}^*$, one has $a_{RD}/a^* = \epsilon_r^{(-1/2)}$; continued transfer with $Q_r > 0$ does not delay this transition.

Accordingly, the minimal cosmological-history connection used in this book is: dormant $\rightarrow$ constraint-preserving opening $\rightarrow$ kination/energy transfer $\rightarrow$ radiation domination $\rightarrow$ hot Big Bang. The microscopic owner of Q_r and the actual transition temperature remain OPEN, but the need to add a separate radiation initial value to SOURCE in order to obtain a hot Big Bang is removed.

Chapter 4 Where Do Dimensions and Spacetime Come From

WRRRA does not treat dimensions as empty boxes given in advance. A dimension is a direction in which relations can be transmitted independently and maintained as stable records.

The dimension filter F selects, from the possible relations, the sector that passes locality, stability, information conservation, and boundary-record conditions. The transmitted kernel and complementary sector are separated as follows.

$$K_{DF} = F \dagger F, L_{DF} = I - K_{DF} \quad (4.1)$$

The advantage of this expression is that it does not prematurely divide visible space and invisible degrees of freedom into existence versus nonexistence. A degree of freedom may exist in the total Actual while failing to pass through the physical renderer. Dark sectors, internal representation spaces, and hidden correlations can therefore occupy different positions within the same distinction.

$$d_{\text{eff}}(t) = -2 \, d \log \text{Tr}(\exp(-tK_{DF})) / d \log t \quad (4.2)$$

The effective dimension can be read from spectral flow. This book does not claim to have derived three spatial dimensions uniquely and without input. The more precise achievement is to provide a minimal implementation path that preserves the observed 3+1-dimensional universe without requiring additional physical spacetime dimensions. The dimensionality of an internal 15-channel space or of Hilbert space is not an extra physical spatial dimension.

Time likewise need not be an external clock already flowing before the opening. Taking the Hamiltonian constraint as fundamental, conditional time can be obtained from correlations between a clock subsystem and a semiclassical branch after the opening.

$$\hat{C}|\Psi\rangle = 0, i\hbar \partial_\tau |\psi_{\text{cond}}(\tau)\rangle = H_{\text{eff}}|\psi_{\text{cond}}(\tau)\rangle \quad (4.3)$$

A persistent space requires more than a metric of rank three at one instant. The metric at every time must remain uniformly comparable with a common reference, and reconstruction rules in overlapping local regions must agree with one another.

$$c \, g^* \leq g_k \leq C \, g^* \text{ (all } k), R_U|U \cap V \simeq R_V|U \cap V \quad (4.4)$$

Statement

The Precise Claim of Minimal Dimensionality WRRRA accommodates a 3+1-dimensional Reality, a large internal state space, and diverse particle representations without assuming eleven physical spacetime dimensions. This is structural minimality, distinct from a proof of dimensional uniqueness.

Part II Asymmetric Unification of Forces and Matter

A Common Origin Does Not Erase Differences

Chapter 5 Why the Four Forces Cannot Be the Same Kind of Force

Unifying gravity with the three gauge forces does not mean flattening all four terms into a single gauge field. It means showing how four representations with different ownership rules can operate together within one execution of Reality.

Electromagnetism, the weak interaction, and the strong interaction selectively read particular representation currents carried by matter. Gravity, rather than selecting a particular charge, couples to the stress-energy of all physically realized degrees of freedom. This difference is not merely a difference in phenomenological strength; it means that the input ledgers are of different kinds.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = (8\pi G/c^4) T_{\mu\nu_total}, \nabla_\mu T_{\mu\nu_total} = 0 \quad (5.1)$$

$$D_\mu = \nabla_\mu^{geo} - ig_1 Y B_\mu - ig_2 T^a W_\mu^a - ig_3 t^A G_\mu^A \quad (5.2)$$

Let the owner map for gravity be $O_G(X)=T_mn[X]$, and the owner maps for the three gauge forces be $O_a(X)=J_a[X]$. A complete Standard-Model singlet X may satisfy $T_mn[X] \neq 0$ while $J_1[X]=J_2[X]=J_3[X]=0$. Gravity then reads X even though the three gauge forces do not.

$$\exists X : (J_1, J_2, J_3)[X] = 0 \wedge T_{mn}[X] \neq 0 \quad (5.3)$$

therefore $O_G \neq f(O_1, O_2, O_3)$ for any current-only flattening f (5.4)

This counterexample shows that a flat unification reducing gravity completely to a selective gauge current of the same kind as the other three forces is not generally possible. Gravity reads the geometric ledger of the entire realization, while the three gauge forces selectively read internal representations within it. A common origin is therefore possible without requiring that all four forces be of the same type.

Statement

Asymmetric Unification Proposition Gravity = the common causal geometry of the entire realization. Electromagnetism, the weak interaction, and the strong interaction = selective internal gauge representations of the same local state. The four forces arise within one renderer, but they do not read the same ledger.

This structure also gives dark matter a natural place. Even a gauge singlet remains in the gravitational sector if it carries real energy, momentum, and pressure. Conversely, abstract information content or computational complexity alone is not a gravitational source. A contribution enters the gravitational ledger only when metric variation produces a nonzero T_mn .

WRRRA's unification is therefore not a 'weaker failure' than conventional grand unification; the goals are different. Grand unification seeks a common microscopic origin of gauge groups and coupling constants. WRRRA asks why gravity and gauge forces, within the same Reality, can simultaneously display universality and selectivity in different ways. These tasks may compete, but they should not be scored by an identical rubric.

Chapter 6 The Common Carrier How Diversity Emerges from One Structure

Minimal structure does not erase every difference. It must reuse common transport laws while preserving the charge, representation, and selection rules of each channel.

One generation of chiral matter carries multiple charges and representations. Treating the 15 directions as 15 unrelated origins increases both SOURCE complexity and transport rules. WRRRA instead places the 15 directions under one common continuum and one common transport operator, implementing their differences through internal labels and boundary readout.

$$D_C : \oplus_{i=1}^{15} H_i \rightarrow \oplus_{i=1}^{15} H_i, \text{rank}(D_C|_{\text{band}})=15 \quad (6.1)$$

The common carrier is neither a particle nor a fifth force. It is a higher-level operator that allows different channels to share the same skeleton of local transport, kinematics, and boundary recording. Just as using the same road does not give all vehicles the same destination or properties, commonality is compatible with distinguishability.

An early democratic rank-one coupling transmitted only one bright combination and left the other fourteen directions dark. This failure closed the shortcut that 'symmetry alone implies unification.' A serial rotor was then proposed in which a rotating rank-one port accumulates phase coding over time and opens the full span.

$$P(\phi)=|v(\phi)\rangle\langle v(\phi)|, \text{rank } P(\phi)=1, \text{rank span}\{v(\phi_k)\}=15 \quad (6.2)$$

However, opening rank 15 in an abstract 15-dimensional space is not the same as realizing one gauge-

covariant Standard-Model generation. The commutant of the Standard Model decomposes into five irreducible blocks, so a pure-state orbit that commutes with the gauge action cannot exceed rank five.

$$\text{End_GSM}(H_{15}) \cong \mathbb{C}^5, \text{rank}(\text{span orbit}) \leq 5 \quad (6.3)$$

This result does not discard the common carrier. It clarifies that abstract carrier rank, gauge-invariant density support, and particle-antiparticle Fock production have different owners. The strength of WRRRA is not in forcing the number 15 to match everywhere, but in distinguishing which layer a given occurrence of 15 belongs to.

Chapter 7 Matter Is a Phenotype Not a List of Numbers

v0.12 chapter update

A particle phenotype is not fixed merely by naming an address. P2 supplies two known pair channels and calculates their selection probabilities from factor multiplicities. P4 implements these probabilities through an effective common interaction; it does not derive masses, charges, spins or the full fifteen-channel carrier. Read the inherited genealogy within that distinction; see Chapters 93 and 95.

Obtaining mass, charge, spin, and position separately does not yet create one particle. A physical object exists when multiple attributes are jointly owned by the same execution and the same record.

The Standard Model is the precise phenotype grammar of observed matter. WRRRA does not relabel that success and claim it as its own. Instead, it asks at which layer of the universe's execution architecture the Standard Model's representations, couplings, and mass generation are owned.

$$m_f(\mu) = y_f(\mu) v(\mu) / \sqrt{2}, \det S_f^{-1}(p) |p^2 = m^2_{\text{pole}} = 0 \quad (7.1)$$

Charge and spin are representation properties, while mass is not a single number stored in SOURCE. It is an execution phenotype jointly produced by the Higgs-Yukawa structure, self-energy dressing, QCD binding, and stable poles, gaps, and thresholds. Therefore, an observed mass used as an input must be labeled CALIBRATED; only a value obtained from upstream generative rules may be labeled DERIVED.

The phrase 'one particle' contains a hidden co-ownership condition. Mass, charge, spin, and position must be read together from the same local observable algebra, the same translation history, the same spectral pole, and the same event record. Placing four numbers from different calculations in one table is not the same thing as showing that one physical object exists.

$$\text{Object } X \Leftrightarrow \text{co-owned}\{m, q, s, x\} \text{ on } (A_{\text{local}}, \text{history}, \text{pole}, \text{record}) \quad (7.2)$$

This notion of co-ownership extends beyond matter to complex systems in general. If it is not specified which functions, memories, resources, and outputs belong to the same object, explanation remains a list of correlations. WRRRA's ownership concept is therefore not decorative terminology but a logical device for fixing 'what is a state of what.'

Ledger — Information Grade · Meaning · Example

- Information Grade: INPUT
 - Meaning: Explicitly supplied by the law or the model
 - Example: c , $\hbar$, observed coupling constants
- Information Grade: DERIVED
 - Meaning: Calculated from inputs and execution rules
 - Example: Time evolution of a specified model
- Information Grade: CALIBRATED
 - Meaning: Value fitted to observation
 - Example: Fixing a scale using the electron mass
- Information Grade: OBSERVED

- Meaning: Value read from external data
- Example: Present cosmic composition; observed events
- Information Grade: OPEN EVENT
 - Meaning: An outcome newly recorded during execution
 - Example: An individual detection outcome
- Information Grade: OPEN
 - Meaning: Owner or value not yet determined
 - Example: Upstream derivation of generations and Yukawa couplings

7.1 Closed internal motion and mass quantization

For an internal circle S^1 of radius R and coordinate y , periodic boundary conditions yield integer modes $n \in \mathbb{Z}$. In the specified compact witness, with $\hbar = c = 1$,

$$\Phi(x, y) = \sum_n \phi_n(x) e^{iny/R}, \quad p_y = n/R. \quad (7.3)$$

$$m_n^2 = m_0^2 + n^2/R^2, \quad E_n^2 = |\mathbf{p}_3|^2 + m_n^2. \quad (7.4)$$

The $n = 0$ branch has no hidden momentum; nonzero n contributes timelike mass in the reduced description. Quantized internal modes are assigned a mass phenotype through the renderer; calling mass only a downstream label would omit this result.

The recorded lattice check uses $a = 2\pi R/N$ and $m_{\text{lat}} = 2 \sin(\pi/N)/a$. For $N = 8 \dots 512$, the relative-error exponent against $1/R$ was approximately -1.999845 , consistent with $O(N^{-2})$ convergence. This is a numerical consistency check, not a new particle-mass measurement. Grade: CONSTRUCTED/EXACT within the specified S^1 witness.

7.2 Dual projection into mass and charge

The same mode integer has an even mass projection and a sign-preserving charge projection:

$$n \mapsto (m_n, q_n) = \left(\sqrt{m_0^2 + n^2/R^2}, gn \right). \quad (7.5)$$

$$m_0 = 0 : \quad (m_n, q_n) = (|n|/R, gn), \quad m_{+n} = m_{-n}, \quad q_{+n} = -q_{-n}. \quad (7.6)$$

Grade: CLOSED IN THE SPECIFIED UNIVERSAL WITNESS. The simplest circle fixes a mass-to-charge ratio and cannot reproduce the full observed Standard Model hierarchy. Additional internal structure, symmetry breaking and dressing are required. The book does not claim that every pole mass is an integer multiple of one universal unit.

7.3 Common geometry for gravity and electromagnetism

A compact-fibre metric provides the specified common-geometric witness:

$$ds_5^2 = g_{\mu\nu} dx^\mu dx^\nu + R^2 (d\theta + \kappa A_\mu dx^\mu)^2. \quad (7.7)$$

$$S_4 = \frac{1}{16\pi G_4} \int d^4x \sqrt{-g} \left[R_4 - \frac{\kappa^2 R^2}{4} F_{\mu\nu} F^{\mu\nu} \right]. \quad (7.8)$$

The fixed-radius reduction supplies Einstein geometry and the Maxwell term from one metric. A lower-dimensional curvature check reproduced $R_{\text{total}} = -\kappa^2 B^2/2$, with maximum absolute defect $6.07 \times$

10^{-11} in the recorded witness.

$$p_z = v_z + \kappa B x v_y = \text{constant}, \quad q = \kappa p_z. \quad (7.9)$$

$$a_x = q B v_y, \quad a_y = -q B v_x. \quad (7.10)$$

Conserved hidden momentum reads as Lorentz force in the reduced description. Electromagnetic stress also sources the common metric. This closes the specified classical witness, while radion stabilization and the universe's full hierarchy remain open.

For a massless spin-2 field, $N_{\text{grav}}(D) = D(D - 3)/2$, or $N_{\text{grav}}(d) = (d + 1)(d - 2)/2$ in spatial dimension d . With $C(d) = c_E d + c_T N_{\text{grav}}(d)$, assume $c_E, c_T \geq 0$ and $c_E + c_T > 0$. Then $C(d + 1) - C(d) = c_E + c_T d > 0$ for $d \geq 3$, so the minimum in this admissible class occurs at $d = 3$. This is conditional on local propagating spin-2 admissibility and the chosen cost, not a theorem about every possible information cost.

7.4 Different quantization claims for mass and gravity

The quantized quantities in this witness are internal mode integers and their mass/charge projections. No analogous fundamental quantization rule for gravity itself has been proved here:

$$n \in \mathbb{Z} \rightarrow p_\theta = n/R \rightarrow m_n \rightarrow \text{mass phenotype}. \quad (7.11)$$

$$T_{\mu\nu}^{\text{total}} \rightarrow g_{\mu\nu} \rightarrow \text{gravitational observables}. \quad (7.12)$$

Gravity reads total implemented stress-energy through the common geometry and is calculated by the GR response at the Reality layer. Lack of a complete quantum-gravity construction neither proves gravity eternally classical nor invalidates the specified classical reduction.

7.5 Quantitative renderer and nonduplicating standard-physics interface

$$\mathcal{R}_m = \mathcal{R}_{\text{pole}} \circ \mathcal{R}_{\text{SM}} \circ \mathcal{R}_{\text{compact}}. \quad (7.13)$$

$$\mathcal{R}_{\text{compact}}(n, R, m_0) = \sqrt{m_0^2 + n^2/R^2}, \quad \mathcal{R}_q(n) = gn. \quad (7.14)$$

$$m_{\text{pole}} = \text{Pole}[m_n, y_f(\mu)v(\mu)/\sqrt{2}, \Sigma_f(p; \mu), \Lambda_{\text{QCD}}, \text{boundary data}]. \quad (7.15)$$

The compact map is exact in its witness. The SM map denotes established mathematical material from SM, QCD and RG. Only a specified internal execution counts as an executed WRRRA connection. A full map fixing all observed particle pole masses without additional inputs remains open; the two layers are not double-counted.

The gravity renderer maps total stress-energy and boundary data through GR to proper time, trajectories, redshift, lensing and waves. No arbitrary departure from GR is inserted where no WRRRA correction has been derived.

The recorded first-generation mass grid is $m_0^{(\text{e-grid})} = m_e/23 = 22.217345682 \text{ keV}$, with $(n_e, n_u, n_d) = (23, 99, 211)$ and $(m_e, m_u, m_d) = (0.51099895069, 2.19951722, 4.68785994) \text{ MeV}$. Fixing the scale with the electron mass makes this a calibrated low-integer approximation.

The neutrino subgrid uses $D_\nu = 2 \cdot 23^6 = 296071778$, $q_\nu = m_e/D_\nu = 1.725929280 \text{ meV}$ and $(n_1, n_2, n_3) = (0, 5, 29)$. It gives $(m_1, m_2, m_3) = (0, 8.62965, 50.05195) \text{ meV}$, $\sum m_\nu = 58.6816 \text{ meV}$, $\Delta m_{21}^2 = 7.44708 \times 10^{-5} \text{ eV}^2$ and $\Delta m_{31}^2 = 2.50520 \times 10^{-3} \text{ eV}^2$.

Observed mass-squared differences informed discovery. Absolute masses and their sum are conditional outputs of the procedure frozen on 28 August 2026: DISCOVERY-CALIBRATED plus FROZEN HOLDOUT. Failure of the declared later checks requires revision or rejection; these are not claimed as independently observed absolute masses.

The SM15 ledger records capacity PASS, anomaly EXACT and mass availability PASS. Full interacting operators, W transitions, Higgs intertwiners, generation replication, CKM/PMNS mixing and pole residues remain open beyond the specified effective constructions.

Chapter 8 An Actual World Composed of Matter

The fact that the laws symmetrically permit matter and antimatter does not mean that the actual universe must continuously occupy both sectors in equal amounts.

Three layers are mixed together in the matter-antimatter problem: first, the symmetry permitted by the equations; second, pair production in particular processes; and third, the occupation state of the actual universe over long times. Once these three layers are separated, a reality composed of matter need not conflict with the antimatter channels permitted by the laws.

Lawful sectors = $H_{\text{matter}} \oplus H_{\text{antimatter}}$, Occupied Reality(t) $\subset H_{\text{matter}}$ (8.1)

This proposition does not say that antimatter fails to exist mathematically. Antimatter is a conjugate sector allowed by LAW and can be physically output together with matter when the required conditions are present, such as collisions, external fields, or nonstationary boundaries. But in an ordinary long-lived reality without such events, the antimatter sector has no obligation to remain continuously occupied.

WRRRA's physical vacuum follows the same distinction. Vacuum is not the zero vector of the Actual, nor a state in which every field is literally zero. It is a phenotype in which, for a specified background, observer, and detector prescription, the normal real-particle output rate and net energy flux are zero.

$\Gamma_{\text{real-particle}}[\text{vacuum}; \text{background}, \text{detector}] = 0$ (8.2)

$\langle N \rangle = 0 \nRightarrow \{ \langle T_{\mu\nu} \rangle_{\text{ren}} = 0 \vee G_{\text{plus}}(x, x') = 0 \}$ (8.3)

Pair production is not a process in which 'existence continually emerges from nothing.' In a stable stationary background, the persistent production rate of on-shell pairs is zero. Real output appears when particle definitions and energy exchange change, as in external fields, collisions, time-dependent boundaries, curvature, horizons, or accelerated observation. Pair annihilation likewise is not the deletion of existence but a conservative transfer of ownership from particle ports to other carrier modes such as photons.

matter + antimatter $\rightarrow$ carrier output, $\Delta \text{Ledger}_{\text{total}} = 0$ (8.4)

Statement

Matter-Only Reality Proposition The symmetry of the laws is a symmetry of the possibility space. Occupation of Reality is the result of events, boundaries, and stability. Therefore, antimatter can remain lawful while the persistent actual universe is composed of matter.

The problem of calculating the quantitative baryon asymmetry must be distinguished from this ontological possibility. The Sakharov conditions address the dynamics needed to explain why our universe acquired a particular asymmetry. WRRRA's prior conclusion is more basic: what is possible and what remains persistently actual need not be the same set.

8.1 Dynamics of a Matter Preponderant Reality

Here, 'matter preponderance' does not mean the cosmological matter-dominated era, in which ρ_m exceeds radiation or dark energy. In this chapter it means a matter-over-antimatter asymmetry in long-term occupation.

$\Delta_M \equiv n_M - n_{\text{Mbar}}$ (8.5)

When expansion, production, and annihilation are included together, symmetric annihilation terms cancel from the difference equation, and the actual occupation asymmetry can be tracked as follows.

$d(a^3 \Delta_M)/dt = a^3(S_M - S_{\text{Mbar}})$ (8.6)

After asymmetry generation ends, if $S_M - S_{\text{Mbar}} = 0$, then $a^3 \Delta_M$ is conserved. In a state with $\Delta_M > 0$, if the symmetric matter-antimatter component annihilates sufficiently, n_{Mbar} becomes small

and the remaining matter approaches the residual asymmetry. Thus, it is compatible for LAW to allow antimatter channels while the actual long-lived Reality remains matter-preponderant.

The asymmetry itself, however, must not be inserted as a new primordial number. Δ_M^* must either be a function of η and LAW or arise from S_M - S_{Mbar} generated by existing interactions after the opening. This is the No Extra Asymmetry Register condition.

For a LAW and opening with exact CP symmetry, the boundary must not create an arbitrary matter direction.

$$F_B(\eta, \text{LAW_CP-symmetric}) = 0 \quad (8.7)$$

Therefore, to generate a real baryon asymmetry, the required baryon-number violation, C/CP asymmetry, and out-of-equilibrium execution must be explicitly owned by LAW or by interactions after the opening. The upstream calculation of the final numerical value $\eta_B = (n_B - n_{Bbar})/s$ remains OPEN, and this boundary is the same one applied to conventional baryogenesis.

Part III From Quantum Possibility to One Reality

Actual, Reality, Event, and an Open Single History

Chapter 9 Actual Reality Event

The total Actual and the visible Reality must be distinguished if information conservation, vacuum, measurement, and dark sectors are to be handled in one language.

Actual is the total dynamical state evolving under LAW. Reality is the sector jointly implemented through the physical observable algebra and the renderer. An Event is one newly distinguishable record added to that Reality. SOURCE is not the whole Actual, and Reality is not a screen that copies all of Actual.

$$\Omega_t \in S_{\text{Actual}} \xrightarrow{R} \rho_t \in S_{\text{Reality}} \xrightarrow{I_a} (a_t, \rho_{t+1}) \quad (9.1)$$

$$\rho_{\text{Reality}}(\tau) = P_{\text{vis}}[e^{(-iH\tau/\hbar)} U_S \rho_{\text{dorm}} U_S^\dagger e^{(+iH\tau/\hbar)}] \quad (9.2)$$

$$p(o, \tau) = \text{Tr}[E_o \rho_{\text{Reality}}(\tau)] \quad (9.3)$$

The renderer is not an arbitrary image generator. In a tensor-factor model it can be implemented as a CPTP channel such as a partial trace; under restriction to an observable algebra it must preserve positivity, normalization, and the relevant conservation conditions. If several Actual states are compatible with the same Reality, the invisible differences remain in a compatibility fiber.

$$C(\rho) = \{\Omega : R(\Omega) = \rho\} \quad (9.4)$$

Information that does not appear in Reality has therefore not necessarily disappeared. It may remain in hidden or environmental correlations. This distinction allows black-hole information, vacuum correlations, quantum measurement, and macroscopic records to be discussed within one framework. Information is not simply a matter of ‘present or absent,’ but of who owns it and which observable algebra can read it.

The Actual-Reality distinction is not a declaration of two Platonic worlds. It aligns the distinctions already used in physics between total states, reduced states, observable algebras, and records with their ontological responsibilities. WRRRA’s additional claim is that these layers jointly produce one Reality within a single execution order.

Chapter 10 Fixed Laws and an Open Single History

There is no contradiction between laws being fixed and future events not yet being written. A closed conservation ledger and open events can coexist within one execution system.

Fixed laws constrain possible transitions, amplitudes, conserved quantities, and causal structure. But this does not mean that every future event must be stored as a classical list. WRRRA imposes the No Predetermined Event Register condition: neither LAW nor SOURCE contains a readable register that directly stores the future event sequence.

$$(a_1, a_2, \dots, a_\infty) \notin \text{readable registers of LAW or SOURCE} \quad (10.1)$$

A quantum instrument supplies the probabilities of possible outcomes and the conditional post-event states. Only one outcome is actually recorded, and it becomes part of the condition for the next present.

$$p(a|p) = \text{Tr}[I_a(p)], \rho_a = I_a(p)/p(a|p) \quad (10.2)$$

$$\Psi_n \rightarrow \{a_1, \dots, a_k\}, A_n \in \{a_1, \dots, a_k\}, h_n = (h_{n-1}, A_n) \quad (10.3)$$

This structure does not require either a classical table of predetermined endings or the unrestricted actualization of every branch in a many-worlds picture. Possibilities are plural, but the actual record is singular. Outcomes not selected remain counterfactual possibilities, while the event that occurs is incorporated into the accumulated history. The future is conditionally open, and the past is closed in the record.

Finite execution experiments have constructed multiple sequential event histories from the same initial complex state and the same self-adjoint Hamiltonian while preserving unitarity and instrument completeness. This is an existence construction demonstrating the logical compatibility of fixed laws with open events.

$$\sum_a I_a \dagger(Q) = Q, \|U \dagger U - I\| \approx 0, \sum_a p(a|p) = 1 \quad (10.4)$$

The information in a realized history can be written not as the size of a file stored from the beginning, but as the conditional surprise accumulated whenever an event occurs.

$$I_{\text{history}}(T) = \sum_t -\log_2 p(a_t|h_{t-1}) \quad (10.5)$$

Statement

The Information Economy of the Universe Short laws can generate a long history. The universe is not a device replaying a completed future; it is an execution system that updates the present under fixed laws and writes its own history.

Chapter 11 The Boundary Between Quantum Possibility and Macroscopic Reality

Quantum probability is not magic that can manufacture arbitrary events. There is first a space of events permitted by the laws, and amplitudes and Born weights operate only within that space.

WRRRA divides selective realization into two passes. The first pass constructs the support set Γ_{lawful} allowed by causality, energy, momentum, charge, gauge constraints, interaction history, and stable recording. The second pass assigns complex amplitudes and probabilities only to paths within that set.

$$\Gamma_{\text{lawful}} = C_{\text{LAW}}(p; \text{laws, constants, constraints, history}) \quad (11.1)$$

$$A[\gamma] = a[\gamma] \exp(iS[\gamma]/\hbar), K = \sum_{\gamma \in \Gamma_{\text{lawful}}} A[\gamma] \quad (11.2)$$

$$\text{supp } P_{\text{WRRRA}} \subseteq \Gamma_{\text{lawful}} \quad (11.3)$$

This structure is not an after-the-fact prohibition switch designed to restrict quantum-mechanical calculations. In a completed model, Γ_{lawful} must be generated for both microscopic and macroscopic systems from the same states, environments, and transport rules. The central intuition is that probability does not manufacture physical possibilities from nothing; it assigns relative weights to events that are already lawful possibilities.

The difference between microscopic uncertainty and macroscopic definiteness is not divided by mass alone. When small event differences are redundantly stored through environmental records and interactions, macroscopic history becomes stable. A candidate scale combines the mass M , discontinuous displacement Δx , and record redundancy R_{rec} .

$$\Omega_{\text{WRRRA}} = R_{\text{rec}} M \Delta x / \hbar, \chi_{\text{est}} = \exp(-\Omega_{\text{WRRRA}}) \quad (11.4)$$

This quantity is not yet a universal constant but a candidate boundary scale. The direction, however, is clear: the quantum-to-classical transition is not fundamentally a matter of ‘an observer looking,’ but of how redundantly and persistently the distinguishability of alternatives is stored in physical records.

Decoherence reduces interference between different record branches, but it is not the same operation as

making only one branch actual. No deterministic selector based solely on linear unitary dynamics can send every superposition to one outcome while preserving linearity, norm, and Born weights simultaneously. Record formation and single-outcome selection must therefore be treated as distinct layers.

$$U(|\psi\rangle|E_0\rangle)=\sum_a c_a|a\rangle|E_a\rangle, \langle E_a|E_b\rangle\approx\delta_{ab} \quad (11.5)$$

Chapter 12 Conditions for One Reality to Remain the Same Reality

Outputting a Reality once is not enough to make a universe. Different rendering paths must yield the same physical content, that content must remain stable over long times, and local realities must glue together into one shared Reality.

Two rendering paths that begin at the same physical boundary and end at the same boundary must agree on protected observables even if their intermediate slices and gauge choices differ. This is renderer path independence.

$$R_{\{\gamma_1\}}(B_f, B_i) \simeq_{\text{gauge}} R_{\{\gamma_2\}}(B_f, B_i) \quad (12.1)$$

In canonical gravity, compositions of small deformations must close into another allowed deformation. If the constraint algebra does not close, different choices of slicing produce different physical realities.

$$\{H[N], H[M]\} = D[q^{ab}(N \partial_b M - M \partial_b N)] \quad (12.2)$$

A general quantum channel can contract information, so the mere fact that a map is CPTP does not preserve position or metric rank. What is required is not recovery of every microscopic state, but recovery on the physically protected observable algebra.

$$\text{Recovery} \circ \Phi = \text{id on } A_{\text{protected}} \quad (12.3)$$

$$R_1 \circ \Phi_{\{\gamma_1\}} \simeq R_2 \circ \Phi_{\{\gamma_2\}} \text{ on } A_{\text{protected}} \quad (12.4)$$

Recovery at one instant and long-term stability are also different requirements. Repeated disturbances must be summable, or there must be a noiseless observable algebra, or the recovery map must contract toward a code manifold.

$$d_{\{k+1\}} \leq \lambda d_{\{k\}} + \epsilon_k, 0 \leq \lambda < 1 \quad (12.5)$$

Uniform metric corridors and quasi-local gluing must be added to these conditions before multiple local observations can form one shared Reality. This family of conditions is an important independent contribution of WRRRA because it asks not only ‘what exists’ but what minimal conditions allow the same existence to remain the same Reality across different observations and times.

Part IV The Meaning of Unification and a Computable Architecture

A Higher-Level Placement of Established Physics and a General Execution Grammar

Chapter 13 Placing Classical Relativistic and Quantum Physics in One Structure

A higher-level theory does not invalidate the precise calculations of lower-level theories. It preserves the questions and domains answered by each theory while explaining the execution order connecting them.

Classical mechanics describes motion extremely well after stable macroscopic records have formed. Relativity describes the common geometry and causal structure in which that motion occurs. Quantum mechanics describes the amplitudes, interference, and probabilities of allowed microscopic possibilities. WRRRA asks how these three layers connect within one execution of the universe and become one actual record.

$$\text{Motion} \rightarrow \text{Causal Geometry} \rightarrow \text{Quantum Possibility} \rightarrow \text{Physical Realization} \quad (13.1)$$

WRRRA’s higher-level role therefore does not mean merely reproducing all established equations as approximations to one new equation. More fundamentally, it connects the inputs and outputs of existing theories

and assigns ownership by specifying what kind of object the output of one layer becomes for the next.

Ledger — Framework · Nature of Additional Assumptions · Mode of Unification

- Framework: Parallel use of standard physics
 - Nature of Additional Assumptions: Independent equations and matching conditions for each domain
 - Mode of Unification: Selected by observational regime
- Framework: String-theory family
 - Nature of Additional Assumptions: Higher dimensions, strings/branes, compactification
 - Mode of Unification: Unification of UV microstructure
- Framework: Simulation cosmology
 - Nature of Additional Assumptions: Possibility of an external computer or implementer
 - Mode of Unification: Philosophical execution analogy
- Framework: Minimal Computing Cosmology
 - Nature of Additional Assumptions: 3+1 physical dimensions, minimal SOURCE, renderer
 - Mode of Unification: Unification of ownership and execution order

This table alone cannot decide which framework is ultimately true, but it permits theoretical comparison. Minimal Computing Cosmology connects the Big Bang, matter, forces, quantum events, and a single history in one execution language without requiring physical space beyond the observed 3+1 dimensions or an external simulator. If the same explanatory range is achieved, structural economy is a substantive advantage.

In particular, WRRRA does not force gravity to become the same kind of entity as the three gauge forces. It preserves the difference between the geometry of the whole realization and selective internal representations while unifying them in one execution. A central feature of WRRRA is that unification must not be confused with identity.

Chapter 14 Turning the Declaration into a Finite Execution System

To avoid remaining a purely verbal framework, WRRRA constructed a finite benchmark in which a positive spectrum, a self-adjoint Hamiltonian, a visible port, a recovery gap, and an entropy ledger coexist.

$$H_N = H_{\text{Jacobi}} \otimes H_{\text{loc}} \otimes H_{\text{syn}} \quad (14.1)$$

The starting point is a positive generalized-Laguerre spectral density above a threshold s_0 . For $\alpha > -1$ it is normalizable, and the three-term recurrence of the orthogonal polynomials determines a real symmetric Jacobi matrix.

$$\rho(s) = \frac{(s - s_0)^\alpha e^{-(s-s_0)/\Lambda^2}}{\Gamma(\alpha + 1)\Lambda^{2\alpha+2}} \Theta(s - s_0). \quad (14.2)$$

$$(J_N)_{nn} = s_0 + \Lambda^2(2n + \alpha + 1), (J_N)_{n,n+1} = \Lambda^2 \sqrt{(n+1)(n+1+\alpha)} \quad (14.3)$$

If the first basis vector e_0 is chosen as the visible port, each eigenvalue weight is the square of its first component, so the weights are automatically nonnegative and sum to one. One collective coordinate reads N hidden spectral nodes. The Euclidean response and real-time amplitude arise from the same nodes and weights.

$$G_E(Q^2) = \int \frac{\rho(s)}{Q^2 + s} ds, \quad G_{E,N}(Q^2) = \sum_{j=1}^N \frac{w_j}{Q^2 + s_j} \approx G_E(Q^2). \quad (14.4)$$

$$A_N(t) = \sum_j w_j e^{\Lambda \{-is_j t\}} \quad (14.5)$$

For $N=2,4,8,16,32$, the maximum relative error over the specified Euclidean window decreased from 4.089×10^{-2} to 1.763×10^{-9} . A finite spectrum, however, retains long-time recurrence. This difference separates the ‘time window over which a finite system accurately imitates a continuum’ from ‘permanent irreversibility.’

In the recovery sector, an amplitude-damping-type Lindblad operator was introduced and the syndrome-transverse gap was calculated exactly.

$$L_{\text{rec}}(\rho) = \kappa[\sigma_- \rho \sigma_+ - \frac{1}{2}\{\sigma_+ \sigma_-, \rho\}], \Delta_{\text{corr}} = \kappa/2 \quad (14.6)$$

$$\Delta S_{\text{rec}} = -p \ln p - (1-p) \ln(1-p), Q_{\text{min}} = k_B T \Delta S_{\text{rec}} \quad (14.7)$$

The value of this benchmark is not that it replicates the entire universe. It is that positivity, self-adjointness, a common visible port, finite-window convergence, recovery, metric preservation, and a thermodynamic ledger - all required by WRRRA - can coexist without contradiction in one finite construction.

Chapter 15 Viewing Many Physical Phenomena with One Tool

WRRRA’s explanatory power does not consist in inserting the same numbers into every phenomenon. It consists in reading different phenomena through the same system of questions: in which ledger and at which boundary does each phenomenon arise?

Ledger — Phenomenon · Core WRRRA Interpretation · Primary Owner

- Phenomenon: Big Bang
 - Core WRRRA Interpretation: Constraint-preserving opening boundary transition
 - Primary Owner: SOURCE + geometry ledger
- Phenomenon: Dark Matter
 - Core WRRRA Interpretation: A remainder invisible to gauge interactions but retained in T_{mn}
 - Primary Owner: gravitational ledger
- Phenomenon: Black Hole
 - Core WRRRA Interpretation: Finite boundary information and the problem of internal transport and recording
 - Primary Owner: boundary + transport
- Phenomenon: Temperature
 - Core WRRRA Interpretation: Sensitivity of the number of accessible states to energy
 - Primary Owner: coarse-grained state space
- Phenomenon: Entropy
 - Core WRRRA Interpretation: Weighted multiplicity of histories indistinguishable from the current record
 - Primary Owner: observer/coarse-graining
- Phenomenon: Vacuum
 - Core WRRRA Interpretation: Zero-real-particle-output phenotype on a stationary background
 - Primary Owner: detector + background
- Phenomenon: Pair Production
 - Core WRRRA Interpretation: Conditional output opened by a driven configuration
 - Primary Owner: external work + field
- Phenomenon: Pair Annihilation
 - Core WRRRA Interpretation: Conservative transfer of ownership from particles to carrier modes

- Primary Owner: conservation ledger
- Phenomenon: Nuclear Decay
 - Core WRRRA Interpretation: Multiple clocks associated with distinct barriers and transition rules
 - Primary Owner: channel-specific dynamics
- Phenomenon: Superconductivity / Superfluidity
 - Core WRRRA Interpretation: Collective phenotype of global phase stiffness, not merely local pairing
 - Primary Owner: boundary + phase relation

The important point in this table is not a claim that WRRRA replaces the standard-physics calculation for every phenomenon. It separates problems previously blurred under the same words into owner, boundary, transport, and record, making clear which connections are shared and which dynamics are specific to each field. That is the precise meaning of ‘seeing them together.’

For example, temperature and entropy are not identified merely because both can be described using information language. The area-based capacity of a black hole, its current entropy, and the length of a generative description are also distinguished. Antimatter production is separated from persistent antimatter occupation, and quantum correlations from real-particle output. WRRRA’s unification does not remove distinctions; it places them in the correct locations.

$$1/(k_BT) = \partial S_acc(E) / \partial E \quad (15.1)$$

$$S_hist = -k_B \sum_i p_i \ln p_i \quad (15.2)$$

These distinctions allow facts from different fields to be compared in a single ledger. In particular, they help determine whether explaining a phenomenon requires adding a new entity, whether a different output of an existing state is sufficient, whether an invisible sector carries real energy, or whether it is merely hidden computationally.

Chapter 16 The Minimal Execution Grammar Repeated Beyond Cosmology

The value of applying WRRRA to multiple fields is not that external applications automatically validate the cosmology. It is that the minimal questions required to construct complex reality recur across domains.

Cells, the origin of life, neural behavior, economics, and weather contain different physical components. Yet the need to distinguish independent states, relation networks, boundaries, residual history, update rules, and resource ledgers recurs. WRRRA expresses this common grammar as SOURCE, RELATION, BOUNDARY, RESIDUE, UPDATE, and LEDGER.

$$\text{System} = \{\text{SOURCE}, \text{RELATION}, \text{BOUNDARY}, \text{RESIDUE}, \text{UPDATE}, \text{LEDGER}\} \quad (16.1)$$

In life, a DNA sequence alone does not make an executable cell; translation machinery, energy, boundaries, catalysts, distribution, and intergenerational regeneration must close together. In neural behavior, the relevant question is not merely whether memory exists but which tasks require past history and what difference that memory makes to reward and safety. In weather and economics, one must test whether a weak historical signal actually outperforms a strong current-state baseline.

In these external applications, some models failed to beat their baselines. But the information contained in failure was preserved precisely through the common grammar. Rather than saying ‘there is no residue,’ one can say ‘the observed trajectory is not the owner of the residue’; rather than saying ‘memory is useless,’ one can say ‘the given task does not reward that memory.’ The location of failure can therefore be identified.

Statement

Value of a General-Purpose Tool A single equation need not output the numbers of every field. The common format instead determines who owns which information, what is closed by the present state, and where new events, environments, or boundaries are required.

This is stronger than a simple analogy. If the same structural questions operate independently on differ-

ent data sets and repeatedly expose incorrect ownership assignments, WRRRA becomes a metatheory for comparing execution structures of reality. At the same time, it preserves logical independence by not converting success in a particular external domain into evidence for the cosmology itself.

Chapter 17 What Has Already Been Obtained

The value of a theory cannot be assessed solely through a list of what remains unfinished. Structural discoveries, exact constructions, NO-GO results, and quantities that remain to be calculated must be separated into different classes of achievement.

Ledger — Grade · What This Book Has Obtained

- Grade: Core architecture
 - What This Book Has Obtained: Single flow of LAW-SOURCE-EXECUTION-RENDERER-REALITY-RECORD
- Grade: Theoretical result
 - What This Book Has Obtained: Asymmetric unification produced by global T_{mn} ownership for gravity and selective ownership of three gauge currents
- Grade: Ontological result
 - What This Book Has Obtained: Compatibility between lawful antimatter and an Actual Reality occupied by matter; conditional dynamical connection from $\Delta_M > 0$ and annihilation of the symmetric component to matter-preponderant occupation
- Grade: Exact constructions
 - What This Book Has Obtained: Constraint-preserving boundary map in FLRW minisuperspace; finite Jacobi structure with positivity and self-adjointness
- Grade: Boundary results
 - What This Book Has Obtained: Failure of a bare kick; obstruction to current-only gravity flattening; nonidentity of decoherence and single-outcome selection
- Grade: Methodological result
 - What This Book Has Obtained: Provenance ledger for information ownership, input, derivation, calibration, observation, and events
- Grade: Computational result
 - What This Book Has Obtained: Common visible port, fixed-window spectral convergence, recovery gap, and entropy ledger
- Grade: Higher-level integration
 - What This Book Has Obtained: Staged placement of classical motion, relativistic geometry, quantum possibility, and single-Reality implementation
- Grade: Cosmological-history connection
 - What This Book Has Obtained: Conditional route to the hot Big Bang after the constraint-preserving opening through radiation generated by Q_r and the different dilution rates of radiation and kination; No Extra Radiation Register

The remaining problems are also clear. A fully covariant 3+1-dimensional action, upstream derivations of Standard-Model generations, Yukawa couplings, mixing, and coupling constants, the E_{lift} into quantum Actual, the R_{phys} map fixing physical quantities and units, and the microscopic dynamics of the Born rule and single-outcome selection still require further calculation.

These unfinished items do not reduce the preceding achievements to zero. Just as the mathematical achievements of string theory's higher-dimensional unification do not disappear because vacuum selection and all low-energy phenomena remain unfinished, WRRRA possesses theoretical structures and exact

partial results that can be evaluated before independent predictions are completed.

Most important is not to insert established equations into WRRR and then call them WRRR achievements. Standard equations are labeled BASELINE; structures exactly constructed inside the model are labeled CONSTRUCTED/EXACT; structural connections are labeled THEORETICAL RESULT; values fitted to observation are labeled CALIBRATED; and quantities whose owner is still unresolved are labeled OPEN. These distinctions are not apologetic caveats that weaken the theory, but boundaries that protect its actual achievements.

17.1 Scope already closed in physical model 1.0

Result	Grade
Compact S^1 modes $p_\theta = n/R$, $m_n^2 = m_0^2 + n^2/R^2$	CONSTRUCTED/EXACT
Dual mass-charge projection	CLOSED in the specified universal witness
Compact-metric Einstein-Maxwell reduction and Lorentz force	CLOSED in the specified universal witness
$d = 3$ minimum under propagating spin-2 and positive-cost assumptions	CONDITIONAL THEORETICAL RESULT
First-generation grid and neutrino absolute masses/sum	CALIBRATED plus FROZEN HOLDOUT

An incomplete full Standard Model or quantum gravity does not erase these restricted results. The remaining tasks concern a full 3+1-dimensional covariant action, matching compact modes to three generations, Yukawa and mixing structures, coupling constants and pole residues, the lift to quantum Actual, physical unit assignment, and microscopic Born-rule/single-outcome selection dynamics. The recent effective calculations in Chapters 93–96 address limited parts of this interface; they do not close all of it.

Established formulae are BASELINE; exact model constructions are CONSTRUCTED/EXACT; demonstrated structural connections are THEORETICAL RESULT; observationally fitted quantities are CALIBRATED; unexecuted connections remain OPEN. These distinctions preserve the achievements and their conditions.

Chapter 18 What Still Needs to Be Completed

The next stage is not to diminish the value of the theory again, but to connect the central architecture already established to stronger mathematics and calculation.

The first task is to formalize minimality. A Pareto ledger must be completed that compares 3+1-dimensional physical space, SOURCE input length, the number of independent owners, the number of carriers, and update cost under the same observational contract. This turns the aesthetic claim that ‘nature should be simple’ into a computable comparison principle.

The second task is to extend the scope of the asymmetric-unification result. Starting from the gauge-singlet counterexample, one must show that the global geometric owner and selective current owners coexist consistently within a covariant action, anomaly freedom, symmetry breaking, and RG matching. This would give WRRR’s unification of the four forces a sharper theoretical form.

The third task is to calculate the microscopic owners of the early-universe dynamical links already established. Q_r , which produces radiation from the opening, and S_B - S_{Bbar} , which produces the matter-antimatter asymmetry, must be closed as functions of the existing LAW and η . Neither a radiation seed nor Δ_B may be added as a new SOURCE initial value. The ultimate targets are independent calculations of the transition temperature and $\eta_B = (n_B - n_{Bbar})/s$.

The fourth task is a single covariant execution expression. The FLRW opening, gauge matter, common carrier, quantum instrument, and renderer must be placed under a shared set of variables, symmetries, and constraint algebra.

$$S_{\text{WRR}}[g, \Phi_{\text{open}}, \Psi_{\text{matter}}, A_{\text{gauge}}, R] = \int d^4x \sqrt{-g} L_{\text{total}} + S_{\text{boundary}} \quad (18.1)$$

The fifth task is to compare explanatory compression directly. When conventional parallel theories, higher-dimensional unification models, and WRR explain the same set of phenomena, their required physical dimensions, independent inputs, separate carriers, calibrated values, and joining rules can be compared in an open ledger. This quantitatively evaluates explanatory economy independently of new predictions.

Statement

The Next Sentence of the Research The unfinished parts are not holes that negate the existence of the theory; they are a work list specifying how far the already identified minimal execution architecture remains to be closed.

Conclusion Explaining One Universe with the Least

Minimal Computing Cosmology does not stop at the metaphor that the universe is a computer. It asks, within one execution architecture, what counts as LAW, what counts as primordial input, how possibilities pass through transport and boundaries to become matter, forces, and spacetime, and how a plurality of quantum possibilities becomes one actual record.

The architecture simultaneously seeks minimal dimensionality, minimal structure, and minimal computation. It does not require physical extra dimensions beyond 3+1 as a necessary premise. One opening begins one history; common transport and a renderer reuse structures across diverse physical representations. The entire future event sequence is not pre-stored; the next Reality is produced from the present.

The unification of the four forces is also redefined through this minimality. Gravity reads the stress-energy of the whole realization, while electromagnetism, the weak interaction, and the strong interaction selectively read internal representation currents. Because of this difference, gravity cannot be flattened into the same kind of object as the other three forces, yet precisely this asymmetry allows them to be integrated more naturally within one execution of Reality.

Matter and antimatter are organized in the same way. The fact that LAW permits both sectors is different from the claim that Reality must occupy them equally. If an asymmetry $\Delta_M > 0$ is generated by LAW or during EXECUTION and the symmetric component annihilates sufficiently, the actual long-lived Reality can remain matter-preponderant while antimatter channels remain lawful. Under exact CP symmetry the boundary asymmetry must vanish, and the upstream calculation of the observed η_B remains a separate OPEN task.

Classical mechanics, relativity, and quantum mechanics do not disappear in this architecture. They are responsible, respectively, for motion, geometry and causality, and amplitudes of possibility. WRRRA is responsible for explaining how their outputs connect to one implemented Reality and one history. The goal of Minimal Computing Cosmology is therefore not to deny established physics but to place its scattered successes together within a smaller ontological and computational structure.

Statement

Final Proposition The universe is not the replay of an enormous primordial data set. It is an execution system in which minimal input and reusable laws and relational structures update the present and jointly produce spacetime, matter, forces, and one actual history.

Whether or not this theory is the final answer, the scale of the question is not small. It is an attempt to explain, through one principle of minimal dimensionality, minimal structure, and minimal computation, the connection from a constraint-preserving opening to the hot Big Bang, the actual matter-preponderant Reality, the asymmetric unification of the four forces, the relation between quantum possibility and a single macroscopic record, and the relation among different physical theories. The attempt has academic value worthy of reading and examination.

Part V Selective Reinforcement from String Microphysics

Chapter 0. Comparison of Minimal Computing Cosmology 2.0 and String Theory

This is not a document that mixes the two theories at equal status. Minimal Computing Cosmology 2.0 owns the higher-level execution architecture from the universe's SOURCE to one Reality and Record. String theory is used as material to reinforce portions that remain microscopically weak within that architecture, including quantum gravity, anomaly closure, the microscopic origin of chiral matter, and black-hole microstates. In one sentence: MCC/WRRA is the skeleton, and string theory supplies repair material.

Ledger — Item · MCC 2.0 · String Theory · Owner After Combination

- Item: Highest-level question
 - MCC 2.0: How does possibility become one actual universe, Reality, and Record?
 - String Theory: From what consistent microscopic structure do quantum gravity, gauge fields, and matter arise together?
 - Owner After Combination: MCC
- Item: Basic principle
 - MCC 2.0: Minimal dimension, minimal structure, minimal computation
 - String Theory: Quantized string/brane structure and worldsheet consistency
 - Owner After Combination: MCC is higher-level
- Item: SOURCE
 - MCC 2.0: Dormant state + minimal opening input η
 - String Theory: Requires selection of a particular vacuum/compactification
 - Owner After Combination: MCC
- Item: Spacetime
 - MCC 2.0: Reality uses 3+1 physical dimensions as the minimal arena
 - String Theory: Perturbative superstring theory is 9+1 dimensional, with six compact dimensions
 - Owner After Combination: Conditional combination
- Item: Gravity
 - MCC 2.0: Universal geometry owner reading total $T_{\mu\nu}$
 - String Theory: Massless closed-string spin-2 mode and low-energy gravity
 - Owner After Combination: String provides microscopic reinforcement
- Item: Gauge sectors
 - MCC 2.0: Selective current owners

- String Theory: Gauge sector from compactification/topology/bundle or branes
- Owner After Combination: String provides microscopic reinforcement
- Item: Common carrier
 - MCC 2.0: Higher-level transport operator shared by 15 channels
 - String Theory: Bundle-covariant Dirac/kinetic operator and common 10D field structure
 - Owner After Combination: MCC concept + String realization
- Item: Matter
 - MCC 2.0: Phenotype with jointly owned mass, charge, spin, and location
 - String Theory: Compactified zero modes, representations, Yukawa/overlap data
 - Owner After Combination: String material + MCC phenotype
- Item: Anomaly/BRST
 - MCC 2.0: Partial closure; full microscopic origin remains weak
 - String Theory: Green–Schwarz mechanism and strong 10D consistency
 - Owner After Combination: String reinforcement
- Item: Big Bang
 - MCC 2.0: Constraint-preserving opening → kination/transfer → hot Big Bang
 - String Theory: Multiple string-cosmology/reheating scenarios
 - Owner After Combination: MCC skeleton retained
- Item: Matter dominance
 - MCC 2.0: Distinguishes lawful antimatter from occupied matter
 - String Theory: Provides candidate CP sources, moduli, heavy states, etc.
 - Owner After Combination: MCC occupancy structure + String source candidates
- Item: Black holes
 - MCC 2.0: Explains the meaning of boundary information and record
 - String Theory: Successful microstate counting for specific BPS black holes
 - Owner After Combination: Division of labor
- Item: Vacuum selection
 - MCC 2.0: Minimal-computation/Pareto selection candidate
 - String Theory: Many compactifications/vacua
 - Owner After Combination: MCC supplies selection principle
- Item: Single history
 - MCC 2.0: Actual → Reality → Event → Record
 - String Theory: A universal single-history selection rule is not a central function
 - Owner After Combination: MCC

The conclusion of the comparison is division of labor rather than mutual replacement. As summarized in a 2024 Annual Review, string theory provides powerful resources for quantum gravity, a common microscopic origin of gauge and gravitational sectors, and compactifications yielding four-dimensional Standard-Model-like structures; however, selecting which compactification corresponds to our universe and selecting one complete early-universe history remain active research programs [S1]. MCC retains the higher-level ledger of selection, execution, expression, and recording.

Statement

Asymmetry of the Combination If string theory changes MCC’s SOURCE, minimal computation, common carrier, phenotype, or Actual–Reality distinction, the combination fails. Conversely, it also fails if MCC merely renames established string mathematics in WRRRA terminology. Existing string results are labeled BASELINE / IMPORTED MICROPHYSICS; MCC’s new contribution is restricted to connecting ownership, selection, execution order, and minimality.

Part I. Rules of Combination — Do Not Confuse the Skeleton with the Repair Material

Chapter 1. Three Invariants: Minimal Computation, Common Carrier, Phenotype

1.1 Minimal Computation as an Invariant

Minimality in MCC does not mean that fewer dimensions automatically win. The Pareto minimality of 2.0 is extended so that the cost of compactification itself is made explicit. Six compact dimensions in string theory incur an independent cost; however, if the same structure simultaneously realizes gravity, gauge sectors, chirality, anomaly cancellation, and the common carrier—thereby greatly reducing the number of owners, carriers, and post-hoc junctions—it can still be globally minimal.

Equation (1.1)

$$C_{\text{hybrid}} = \{D_{\text{phys}}, D_{\text{compact}}, L_{\text{SOURCE}}, N_{\text{owner}}, N_{\text{carrier}}, N_{\text{moduli}}, N_{\text{free}}, C_{\text{update}}\}$$

Equation (1.2)

$$A^* \in \text{ParetoMin}\{C_{\text{hybrid}}(A) \mid A \models \Omega_{\text{observed}}\}$$

The reason to admit ten dimensions is therefore not that “string theory is famous,” but that the cost of six compact dimensions may be smaller than the cost of the independent microscopic structures they eliminate. Conversely, a compactification requiring hundreds of freely adjustable moduli and arbitrary flux choices pays a high price in the MCC minimal-computation ledger.

1.2 Common Carrier as an Invariant

The common carrier is not identical to the string itself. In WRRRA, the common carrier is a higher-level operator that lets multiple matter channels share the same local transport, kinematic, and boundary-recording skeleton. String theory can provide microscopic material for realizing this operator. In particular, a common covariant Dirac/kinetic operator on a compactified bundle can make distinct four-dimensional zero modes descend from the same microscopic transport law.

Equation (1.3)

$$D_{C^{\text{WRRRA}}} : \bigoplus_{i=1}^{15} H_i \rightarrow \bigoplus_{i=1}^{15} H_i$$

Statement

Forbidden Do not write that “all particles are strings, therefore no common carrier is needed.” That statement asserts common origin but does not realize the actual gauge-covariant transport of the fifteen chiral channels or a shared visible port.

1.3 Phenotype as an Invariant

A string excitation or a compactification zero mode is not immediately an observed particle. A phenotype is obtained only after four-dimensional representation assignment, symmetry breaking, Yukawa coupling, RG flow, confinement/self-energy dressing, and the appearance of a stable spectral pole, so that mass, charge, spin, and location become jointly owned by one physical object.

Equation (1.4)

$$\text{String state} \rightarrow \text{zero mode} \rightarrow 4\text{D representation} \rightarrow \text{SSB/RG/dressing} \rightarrow \text{physical phenotype}$$

String theory therefore supplies upstream material for phenotype. WRRRA continues to own the final co-attribution of mass, charge, spin, and location, together with the Actual–Reality–Record distinction.

Chapter 2. Ten Dimensions Are Allowed, but Not for Free

This reinforcement does not require eleven-dimensional M-theory as a premise. The primary microscopic material is the 9+1-dimensional structure of perturbative superstring theory. We distinguish 3+1-dimensional Reality from 9+1-dimensional microscopic Actual.

Equation (2.1)

$D_{\text{micro}} = 9+1, D_{\text{Reality}} = 3+1, D_{11} \notin \text{required ontology}$

The six extra dimensions are, in standard string theory, genuine compact spacetime dimensions; this is not concealed. They are not, however, the same as macroscopically open propagation directions: they are small and compactified, and therefore do not appear as directly accessible directions in low-energy Reality. MCC records their cost explicitly as D_{compact} .

The key assessment is this: if a 10D compactification simultaneously provides 4D gravity, a gauge group, chirality, anomaly cancellation, and a realization of the common carrier within one microscopic structure, its total Pareto cost can be lower than a “strictly 3+1” implementation that must assign separate owners to each. Minimal computation permits this in principle.

Statement

Compatibility Condition Between Minimal Computation and Ten Dimensions Additional compact dimensions must purchase real explanatory economy. If the same observations can be implemented in a smaller 3+1 structure with equal or lower owner/carrier/input cost, the 10D structure is dominated as a fundamental implementation.

Chapter 3. Primary Repair Material: $E8 \times E8$ Heterotic String

Among ten-dimensional string constructions, the primary repair material in this document is the $E8 \times E8$ heterotic string. The reason is straightforward: a single ten-dimensional framework contains closed-string gravity together with non-Abelian gauge structure, while Calabi–Yau/bundle compactification can yield four-dimensional chiral matter. Modern reviews of Standard-Model-like string compactifications likewise treat heterotic, type-II/F-theory, and related constructions as major routes toward realistic gauge/matter structures [S1].

This is not a declaration that heterotic string theory is uniquely correct. It is a primary candidate because, in the MCC cost ledger, it can reinforce the common carrier and phenotype with relatively few additional owners. Type II, intersecting D-brane, and F-theory material can be compared as secondary candidates when needed.

Equation (3.1)

$M_{10} = M_{(3,1)} \times K_6$

The ten-dimensional bosonic low-energy sector has, schematically, the following structure in the string frame. This is IMPORTED MICROPHYSICS, not a new MCC derivation.

Equation (3.2)

$S_{\text{het}}^{(10)} = (1/2\kappa_{10}^2) \int d^{10}x \sqrt{-G} e^{\Lambda(-2\Phi)} [R + 4(\partial\Phi)^2 - (1/12)H^2 - (\alpha'/4)(\text{Tr } F^2 - \text{tr } R_{-}^2) + \dots]$

This action matters because geometry, the dilaton, an antisymmetric tensor, and gauge fields appear within one microscopic package. MCC places this package as a candidate microscopic engine of Actual, while refusing to absorb higher layers such as SOURCE, renderer, and phenotype into the action itself.

Part II. Microscopic Structural Reinforcement

Chapter 4. Microscopic Material for Gravity

In MCC 2.0, the central issue for gravity is ownership. Gravity reads the stress-energy of all implemented physical degrees of freedom, not a particular gauge charge. Importing string theory does not change this ownership.

Equation (4.1)

$O_G(X) = T_{\mu\nu}[X], O_a(X) = J_a[X]$

In string theory, the closed-string spectrum contains a massless spin-2 excitation, and the low-energy limit produces Einstein-Hilbert-type gravity together with higher-curvature corrections. This reinforces a point that remained comparatively weak in MCC: why a quantum microscopic sector and gravity can coexist consistently within the same internal theory.

Equation (4.2)

$$S_4^{\text{grav}} = \int d^4x \sqrt{-g} \left[(M_P^2/2)R + \alpha' c_1 R^2 + \alpha'^2 c_2 R^3 + \dots \right]$$

The role of MCC is not to keep every higher-curvature term permanently active in fundamental Reality. If the Einstein term alone maintains viability at low curvature, minimal execution uses the leading sector, while corrections are interpreted as effective redundancy that activates only where high curvature makes them necessary.

Statement

Preservation of Ownership The string graviton does not turn gravity into a “fourth gauge current.” Regardless of its closed-string origin, the owner of gravity in 4D Reality remains total $T_{\mu\nu}$, while the three gauge forces selectively read representation currents.

Chapter 5. Gauge, Anomaly, and BRST Closure

One important OPEN problem in MCC 2.0 is to close the common carrier and gauge matter within a fully anomaly-free microscopic realization. Heterotic string theory provides direct repair material here. Ten-dimensional consistency and the Green–Schwarz mechanism supply a strong structure for cancelling gauge and gravitational anomalies.

Equation (5.1)

$$H = dB - (\alpha'/4)(\omega_{3Y} - \omega_{3L})$$

Equation (5.2)

$$dH = (\alpha'/4)[\text{tr}(R \wedge R) - \text{Tr}(F \wedge F)]$$

Signs and trace normalizations depend on convention, but the structural point is that the topology of the gauge bundle and tangent bundle are not independent ledgers: they are tied by a B-field consistency condition. From the MCC perspective, this is a good example of an added degree of freedom paying for itself. The B-field is not decoration; it performs anomaly closure.

The previous conditional BRST closure in WRRRA can therefore be divided by role. Ten-dimensional microscopic anomaly/BRST consistency is an IMPORTED STRING CONSTRAINT, whereas four-dimensional renormalizer path independence and protected-observable recovery are WRRRA CONSTRAINTS. They are not the same problem.

Chapter 6. Microscopic Realization of the Common Carrier

This is the central chapter of the reinforcement. The abstract WRRRA common carrier is linked to a common bundle-covariant operator arising from string compactification. On a compact space K_6 with gauge bundle V , we take the ten-dimensional fermionic kinetic operator in the following form.

Equation (6.1)

$$\mathcal{D}_{(10,V)} = \Gamma^M (\nabla_M + A_M)$$

Under product or warped compactification, the fermionic field is expanded in internal eigenmodes.

Equation (6.2)

$$\Psi(x,y) = \sum_n \psi_n(x) \otimes \xi_n(y), \quad \mathcal{D}_{(K,V)} \xi_n = \lambda_n \xi_n$$

Low-energy chiral matter arises from suitable zero modes or light modes. The important point is that different 4D channels need not require independent carriers: they descend from the same 10D covariant operator and the same compactification geometry/bundle.

Equation (6.3)

$$D_C^4(4) := P_{15} \cdot \text{Red}_{(K_6)}[\mathcal{D}_{(10,V)}] \cdot P_{15}$$

Equation (6.3) is not a theorem of standard string theory but an MCC-proposed microscopic identification. P_{15} is a projector selecting the fifteen chiral Standard Model channels of one generation, and Red_{K_6} is the compactification/reduction map. If this identification holds, the common carrier moves one step from “abstractly one” toward “which microscopic operator is it?”

Statement

Preservation of the Common Carrier String is material for the carrier, not a name that deletes the carrier. The genealogy 10D common operator $\rightarrow$ multiple internal zero modes $\rightarrow$ fifteen 4D channels is required to satisfy the MCC common-carrier condition.

Chapter 7. Fifteen Chiral Channels and Generations

Four-dimensional chirality is one of the most important resources supplied by compactification. The index of the bundle-covariant internal Dirac operator connects the net difference between left- and right-handed zero modes to topology.

Equation (7.1)

$$N_L - N_R = \text{index}(\mathcal{D}_-(K,V))$$

This strengthens the explanation of why chiral matter is possible far beyond MCC alone. It must not, however, be claimed that the index automatically equals three. To obtain three generations, one must select a compactification whose geometry, bundle, and topological data yield net chirality three.

Equation (7.2)

$$\text{index}(\mathcal{D}_-(K,V)) = 3 \text{ (target condition, not automatic identity)}$$

The new role of MCC is not merely to note that many topologies “can produce three,” but to select, among candidates satisfying the observational contract, those with the smallest total computation/structure cost.

The fifteen channels of one generation decompose according to Standard Model representations, but their transport skeleton is shared through the common carrier of Eq. (6.3). The number of generations need not be defined by duplicating the common carrier three times; distinct internal zero-mode families of the same carrier operator can constitute the generations.

Chapter 8. Matter as Phenotype

In string phenomenology, the four-dimensional particle spectrum depends sensitively on compactification geometry, brane/bundle data, symmetry breaking, and moduli. MCC does not treat this only as a weakness. It also supplies microscopic material for the prior claim that “matter is not a list of numbers but an executed phenotype.”

Equation (8.1)

$$\Phi_{\text{string}} \rightarrow \{\xi_i(y), r_i, q_i\} \rightarrow 4D \text{ field } \psi_i(x) \rightarrow \text{SSB/Yukawa} \rightarrow \text{RG/dressing} \rightarrow \text{pole/record}$$

Charge and representation are read from compactification, while mass appears as a stable pole after Higgs/Yukawa structure, modulus dependence, self-energy, and QCD dressing. We therefore do not reduce all mass to a single “string vibration number.”

Equation (8.2)

$$m_f(\mu) = y_f(\mu)v(\mu)/\sqrt{2}, \det S_f^{-1}(p)|_{(p^2=m_{\text{pole}}^2)} = 0$$

String-derived Yukawa couplings can be determined by internal wavefunction overlaps or geometry. Schematically:

Equation (8.3)

$$Y_{ijk} \sim \int (K6) \mathcal{W}_i(y) \mathcal{W}_j(y) \mathcal{W}_k(y) \cdot \text{measure}(K6)$$

The final physical particle, however, is not the coupling itself. The WRRRA phenotype map jointly assigns mass, charge, spin, location, history, and record to one object.

Equation (8.4)

$$\text{Object } X \Leftrightarrow \text{co-owned}\{m,q,s,x\} \text{ on } (A_{\text{local}}, \text{history}, \text{pole}, \text{record})$$

Statement

Phenotype Invariance Even after importing string theory, the argument does not end with “a particle is a string vibration.” A string mode is genotype-like microscopic material; an observed particle is a phenotype produced through 4D execution and the renderer.

Part III. Reinforcement of Cosmic History

Chapter 9. SOURCE and Big-Bang Ownership Remain with MCC

String cosmology contains many early-universe scenarios—inflation, reheating, moduli domination, kination, string/brane gas, pre-Big-Bang, and ekpyrotic/cyclic models [S2]. In this combination, that diversity is not allowed to overwrite MCC's SOURCE. The beginning of the universe remains owned by the dormant state and constraint-preserving opening of MCC 2.0.

Equation (9.1)

$$\rho_{\text{dorm}} \xrightarrow{B_\eta} \text{opening} \rightarrow \text{execution}$$

Equation (9.2)

$$B_\eta(a, \chi) = \eta \chi - \sqrt{6} \bar{M}_P |\eta| \ln(a/a^*) + \text{constant}$$

String theory may provide the microscopic identity of χ or the owner of interactions after opening. For example, χ may be realized as the four-dimensional effective coordinate of a compactification modulus, a dilaton-like mode, or another string-derived scalar. But replacing η with multiple string-vacuum labels, or adding unlimited modulus values to SOURCE, would destroy the minimal SOURCE.

Statement

SOURCE Veto Modulus, flux, and bundle data required for compactification must be disclosed as LAW / instance specification. They must not be used as extra primordial registers hiding future events, the radiation content of the hot Big Bang, or the baryon asymmetry.

Chapter 10. String-Theory Material for the Radiation-Transfer Owner Qr

In the final MCC 2.0 manuscript, the most important OPEN element connecting opening to the hot Big Bang is the microscopic energy-transfer owner Qr. String compactification naturally provides scalar moduli and gauge kinetic functions, so it can directly reinforce this gap.

Equation (10.1)

$$\mathcal{L}_{\text{gauge}} \supset -(1/4) \text{Re } f_a(\Phi) F_{a,\mu\nu} F_a^{\mu\nu} - (1/4) \text{Im } f_a(\Phi) F_{a,\mu\nu} \tilde{F}_a^{\mu\nu}$$

If the opening field χ , or a modulus Φ mixed with it, evolves or decays with time, energy can be transferred into the gauge sector. In coarse-grained cosmology this can be written as follows.

Equation (10.2)

$$\dot{\rho}_\chi + 6H\rho_\chi = -Q_r, \quad \dot{\rho}_r + 4H\rho_r = +Q_r$$

Equation (10.3)

$$Q_r \simeq \sum_a \Gamma(\Phi \rightarrow a) \rho_\Phi + Q_{\text{nonpert}} + \dots$$

Equation (10.3) shows that Qr need not remain an unidentified phenomenological function. Once a concrete compactification and gauge kinetic function are specified, there is an owner from which decay widths and production rates can be calculated. Numerical values remain OPEN because the compactification has not yet been fixed.

This reinforcement fits the No Extra Radiation Register principle. Instead of placing a radiation seed in SOURCE, string-derived interactions contained in LAW generate radiation during execution.

Equation (10.4)

$$\rho_r(t) = a(t)^{-4} \int_{t^*}^t a(t')^4 Q_r(t') dt'$$

String microphysics is therefore not the owner of the Big Bang; it owns the energy-transfer material required for the MCC opening to develop into the hot Big Bang.

Chapter 11. String-Theory Material for the Matter–Antimatter Asymmetry Source

The MCC proposition of a matter-dominant Reality separates two layers. LAW may permit both matter and antimatter channels, while actual occupancy can nevertheless be asymmetric. The remaining OPEN element is the microscopic owner of ΔB or the source $S_B - S_{B\bar{}}$.

Equation (11.1)

$$d[a^3 \Delta_B]/dt = a^3 (S_B - S_{B\bar{}})$$

String compactification can provide a variety of microscopic ingredients required by baryogenesis or leptogenesis, including CP phases, heavy singlet states, moduli/axion couplings, and supersymmetric flat directions [S2]. String theory itself, however, does not force one unique baryogenesis mechanism. MCC therefore places candidate mechanisms into the cost ledger and selects among them.

Ledger — Candidate Material · Advantage · Cost / Risk · Assessment

- Candidate Material: Heavy singlet / leptogenesis
 - Advantage: Applicable if the compactification contains singlets and Majorana terms
 - Cost / Risk: Additional heavy-state owner, CP phase, and decay calculation
 - Assessment: Candidate
- Candidate Material: Affleck–Dine-type flat direction
 - Advantage: Uses a flat direction in the 4D SUSY effective theory
 - Cost / Risk: Scalar direction, lifting operator, late decay
 - Assessment: Candidate
- Candidate Material: Modulus/axion-driven source
 - Advantage: CP-odd coupling and out-of-equilibrium decay of a string-derived scalar
 - Cost / Risk: Depends on coupling and stabilization
 - Assessment: Candidate

The principle of minimal computation does not choose the most famous mechanism. It selects the smallest viable source that reuses field content already required by the theory, adds no separate primordial datum, and produces the observed η_B .

Equation (11.2)

$$\eta_{B,f} = [a^3 \Delta_B^* + \int a^3 (S_B - S_B^*) dt] / S_{com}$$

The No Extra Asymmetry Register principle is retained. In a LAW containing exact CP symmetry and baryon-number conservation alone, an arbitrary asymmetry must not appear.

Equation (11.3)

$$F_B(\eta, \text{LAW}_{CP\text{-symmetric}}) = 0$$

Statement

Matter Dominance Is a Phenotype / Occupancy Result String theory may supply microscopic material for an asymmetry source, but MCC continues to own the distinction “matter-dominant Reality” and the lawful-versus-occupied separation between matter and antimatter.

Chapter 12. String-Theory Material for the Dark Sector

In MCC 2.0, the minimal condition for dark matter is that it be nearly invisible to Standard Model gauge currents while still carrying real stress-energy and therefore remaining in the gravitational ledger. String compactification provides many candidates satisfying this condition, including axion-like fields, moduli, and hidden gauge sectors.

Equation (12.1)

$$J_{SM}[X] \simeq 0, T_{\mu\nu}[X] \neq 0, \tau_X \gg H_0^{-1}$$

In particular, axion-like modes descending from the antisymmetric B-field and compact cycles are good examples of hidden internal degrees of freedom appearing as four-dimensional phenotypes. But an abundance of candidates recreates a selection problem. Dark-sector candidates are therefore not adopted automatically from the string landscape; they must pass the MCC Pareto filter.

Dark energy is treated similarly. The cosmological constant and quintessence/moduli candidates of the string landscape are not simply imported as answers. In the current 2.0 framework, Λ remains a BASELINE / observed low-energy parameter and is promoted to an upstream string realization only if string material

improves both cost and observational fit.

Part IV. Information, Selection, and Boundary

Chapter 13. Black Holes: Meaning of the WRR Boundary + String Microstates

The division of labor is cleanest here. WRR explains why a horizon/boundary can serve as a minimally sufficient record for internal Reality and where information remains owned and recorded. String theory has the representative achievement of counting actual microscopic state degeneracies for specific extremal/BPS black holes and reproducing the Bekenstein–Hawking entropy [S3].

Equation (13.1)

$$S_{BH} = k_B \ln \Omega_{string} = k_B A / (4\ell_P^2) \text{ (specific controlled sectors)}$$

MCC therefore does not claim string microstate counting as its own derivation. Instead, it links the compatibility fiber $C(\rho)$ and the number of microscopic states underlying a boundary record to Ω_{string} .

Equation (13.2)

$$C(\rho_{\partial}) = \{\Omega_{Actual} : R_{\partial}(\Omega_{Actual}) = \rho_{\partial}\}$$

Equation (13.3)

$$|C(\rho_{\partial})| \leftrightarrow \Omega_{string} \text{ (candidate microscopic identification)}$$

If this identification holds, WRR’s question “why can the boundary be sufficient?” and string theory’s question “how many microscopic states correspond to that boundary?” become complementary. This does not mean that a complete microstate count has been obtained for all generic astrophysical black holes.

Chapter 14. Placing Duality within Reality-Equivalence

T- and S-duality in string theory are powerful examples in which different microscopic descriptions can yield the same physics. MCC places this within its Actual–Reality distinction.

Equation (14.1)

$$\Phi_A \sim_R \Phi_B \Leftrightarrow R_{vis} \circ \Phi_A = R_{vis} \circ \Phi_B \text{ on } A_{protected}$$

String duality may be a concrete example of Eq. (14.1). MCC does not claim that every Reality-equivalence is a string duality. Rather, it places string duality in the higher-level language of “different Actual descriptions of the same Reality.”

Computational costs can differ even within one equivalence class. Among descriptions yielding the same Reality, MCC may treat the lower-cost representative as a candidate canonical implementation.

Equation (14.2)

$$C([\Phi]) = \inf_{\Psi \in [\Phi]} C_{hybrid}(\Psi)$$

If two descriptions have exactly equal cost, both remain minimal solutions. No external selector need be postulated.

Chapter 15. Filtering the String Landscape by Minimal Computation

One of the major structural difficulties of string theory is the large number of possible compactifications and vacua. Flux compactification and moduli stabilization form extensive research programs, and realistic four-dimensional models involve choices of geometry, topology, bundle/brane data, fluxes, and vacuum [S1,S4]. Here MCC takes on its most important new role.

Equation (15.1)

$$\mathcal{V}_{string} = \{v_i = (K_6, \text{bundle/brane, flux, moduli}, \dots)\}$$

First, vacua that fail the observational contract are removed.

Equation (15.2)

$$\mathcal{V}_{obs} = \{v \in \mathcal{V}_{string} : v \models \Omega_{observed}\}$$

The minimal-computation ledger is then applied to candidates that reproduce the same observational scope.

Equation (15.3)

$$\mathcal{V}_{\text{MCC}} = \text{ParetoMin}\{C_{\text{hybrid}}(v) \mid v \in \mathcal{V}_{\text{obs}}\}$$

The number of moduli, independent flux choices, additional light fields, separate carriers, post-hoc tuning, and update cost can matter more than compact dimension count alone. This principle refuses to treat “infinitely many possibilities are all explanations” as sufficient and demands explanatory economy at the level of implementation.

Statement

What MCC Contributes to String Theory String theory generates possible microscopic worlds. MCC supplies a selection layer that filters candidates compatible with our Reality by three criteria: minimal computation, common carrier, and phenotype. When this selection produces an actual ranking of vacua, it becomes an independent result of the combined framework.

Part V. Integrated Execution Equations and Final Assessment

Chapter 16. The MCC–String Integrated Execution System

The combined framework does not mix every layer into one enormous action. Microscopic Actual dynamics, MCC selection, SOURCE, renderer, and phenotype are kept in sequence.

Equation (16.1)

$$\text{LAW}_{\text{string}} + \text{MCC constraints} \rightarrow \text{viable microscopic class } \mathcal{V}_{\text{MCC}}$$

Equation (16.2)

$$\rho_{\text{dorm}}[v^*] \rightarrow U_S(\eta) \rightarrow \Omega_{\text{Actual}}(t) \rightarrow R_{\text{vis}} \rightarrow \rho_{\text{Reality}}(t) \rightarrow I_a \rightarrow \text{Event/Record}$$

Microscopic evolution is described by the four-dimensional effective Hamiltonian or action of the selected compactification v^* .

Equation (16.3)

$$S_{\text{eff}}[v^*] = S_{\text{GR}} + S_{\text{gauge}} + S_{\text{matter}} + S_{\text{moduli}} + S_{\text{higher}} + S_{\text{boundary}}[B_\eta]$$

When the common carrier and phenotype map are made explicit, the full genealogy is:

Equation (16.4)

$$10\text{D common dynamics} \rightarrow \text{Red}_{(K6)}[\mathcal{D}_{10}] \rightarrow D_C^{(4)} \rightarrow \text{chiral channels} \rightarrow \text{phenotype renderer} \rightarrow \text{Reality}$$

The reinforced structure can therefore be compressed into three layers.

Ledger — Layer · Primary Owner · Content

- Layer: Skeleton layer
 - Primary Owner: MCC/WRRA
 - Content: Minimal computation, SOURCE, owner, common carrier, Actual–Reality–Record
- Layer: Microscopic-material layer
 - Primary Owner: Primarily 10D heterotic string
 - Content: Quantum gravity, anomaly closure, compactified gauge/chiral matter, microstates
- Layer: Phenotype layer
 - Primary Owner: 4D effective physics + WRRA renderer
 - Content: Mass/charge/spin/location, hot Big Bang, matter asymmetry, observed objects

Chapter 17. What Has Actually Been Reinforced?

Ledger — Gap · MCC 2.0 · Imported Repair Material · Current Assessment

- Gap: Microscopic quantum gravity
 - MCC 2.0: Weak
 - Imported Repair Material: Closed-string spin-2 + 10D effective gravity

- Current Assessment: External mathematical material / conditional correspondence; not completed internal execution
- Gap: Anomaly/BRST
 - MCC 2.0: Conditional
 - Imported Repair Material: Green–Schwarz / 10D consistency
 - Current Assessment: Strong conditional support from external mathematical material; separate internal execution required
- Gap: Microscopic owner of common carrier
 - MCC 2.0: Abstract operator
 - Imported Repair Material: Bundle-covariant 10D Dirac/kinetic reduction
 - Current Assessment: Candidate realization
- Gap: Chiral matter
 - MCC 2.0: 15-channel abstract carrier
 - Imported Repair Material: Internal zero modes + index
 - Current Assessment: External mathematical material / conditional correspondence; not completed internal execution
- Gap: Number of generations
 - MCC 2.0: OPEN
 - Imported Repair Material: Compactifications with index = 3 are possible
 - Current Assessment: Conditional; not an automatic derivation
- Gap: Upstream phenotype structure
 - MCC 2.0: Based on SM/QFT
 - Imported Repair Material: String zero modes/overlaps added as upstream material
 - Current Assessment: External mathematical material / conditional correspondence; not completed internal execution
- Gap: Microscopic owner of Q_r
 - MCC 2.0: OPEN phenomenology
 - Imported Repair Material: Modulus/gauge kinetic function/decay
 - Current Assessment: External mathematical material / conditional correspondence; not completed internal execution
- Gap: Baryon source
 - MCC 2.0: OPEN
 - Imported Repair Material: String-derived CP/heavy/flat-direction candidates
 - Current Assessment: Partial correspondence to external material; not completed internal execution
- Gap: Dark sector
 - MCC 2.0: Owner conditions only
 - Imported Repair Material: Axion/moduli/hidden-sector candidates
 - Current Assessment: More candidates; selection still needed
- Gap: Black-hole microstates
 - MCC 2.0: Strong boundary meaning, weak coefficient derivation

- Imported Repair Material: Specific BPS microstate counting
- Current Assessment: Strong conditional support from external mathematical material; separate internal execution required
- Gap: Vacuum selection
 - MCC 2.0: Minimality principle
 - Imported Repair Material: String landscape provides a real selection target
 - Current Assessment: New MCC role

The most important discipline is not to say that “every gap has been closed.” Parts already calculated strongly within string theory can be imported to reinforce MCC, but without choosing a concrete compactification, numerical values for the gauge group, Yukawa couplings, modulus spectrum, Q_r , and η_B are still not uniquely fixed.

Chapter 18. Remaining OPEN Problems and Failure Conditions

18.1 OPEN Problems That Actually Remain

- Apply the MCC cost ledger to real string-compactification candidates and nontrivially eliminate at least one vacuum.
- Explicitly construct the common-carrier relation in Eq. (6.3) for the fifteen one-generation chiral zero modes of a concrete $E_8 \times E_8$ compactification.
- Verify, in one instance, not only index = 3 but the exact Standard Model representations, hypercharge normalization, and an anomaly-free low-energy spectrum.
- Calculate Q_r from a string-derived gauge kinetic function and modulus spectrum, and independently derive the opening-to-hot-Big-Bang transition temperature T_{RD} .
- Select the minimum-cost baryogenesis source and calculate η_B without calibrating to the observed value.
- Produce at least one independent mass ratio or coupling relation at the phenotype layer, including Yukawa, RG, and dressing effects.
- Extend the relation between the WRRR compatibility fiber and string microstate space beyond one specific controlled black-hole example.

18.2 Conditions Under Which the Combination Fails

- If admitting 10D string compactification forces the removal or marginalization of minimal computation, the common carrier, or phenotype.
- If the added cost of six compact dimensions and their moduli/flux structure exceeds the owner/carrier/tuning cost they remove, so that the construction is Pareto-dominated by every observationally equivalent 3+1 implementation.
- If the fifteen chiral channels cannot actually be obtained from a common 10D operator and channel-by-channel independent carriers must be reintroduced.
- If a string mode is identified directly with an observed particle, bypassing the phenotype layers of SSB, RG, dressing, and record.
- If hot-Big-Bang radiation or baryon asymmetry is hidden in SOURCE as a new independent initial datum under the name of a string-vacuum parameter.
- If MCC selection makes no actual distinction among vacua and merely redescribes the string landscape.

Statement

Final Assessment Criterion The success of this combination is not judged by whether “string theory was added,” but by whether the number of microscopic OPEN problems is reduced while the three invariants of MCC 2.0 are preserved. The next decisive test is an explicit 10D realization of the common carrier together with an actual calculation of either Q_r or η_B .

Conclusion. Keep the Skeleton; Strengthen Only the Microscopic Material

The greatest strength of Minimal Computing Cosmology 2.0 is that it places the universe in one execution genealogy—LAW–SOURCE–EXECUTION–RENDERER–REALITY–RECORD—and organizes ownership across different physical phenomena with three axes: minimal computation, common carrier, and phenotype. Its largest weakness was that microscopic quantum gravity and anomaly-free chiral-matter realization beneath that skeleton were less mature than the corresponding structures in established physics.

This reinforcement fills those weaknesses with material from string theory without changing who owns the theory. Using ten-dimensional $E_8 \times E_8$ heterotic string theory as the primary microscopic material makes it possible to import, in one package, gravity and gauge sectors, anomaly cancellation, chiral zero modes, an internal Dirac operator, modulus/gauge kinetic functions, and microstate counting for specific black holes. MCC places only the needed pieces into its owner map.

The three axes remain unchanged. Minimal computation places ten dimensions and the landscape themselves into the cost ledger and asks whether they are necessary. The common carrier is not erased by the word “string”; it demands a concrete realization in which a 10D bundle-covariant operator descends to the fifteen 4D chiral channels. Phenotype remains the final co-owned state after a string state passes through compactification, symmetry breaking, RG flow, dressing, renderer, and record.

Equation (C.1)

MCC skeleton + selected string microphysics $\rightarrow$ stronger Actual engine $\rightarrow$ WRRR phenotype/Reality

The central statement of the combined framework is therefore:

Statement

Final Proposition The skeleton of the universe is an execution architecture governed by minimal computation. String theory does not replace that skeleton; where needed, it supplies mathematical repair material for gravity, gauge sectors, chiral matter, anomaly structure, and boundary microstates. The common carrier binds this material into one transport skeleton, and phenotype is the final stage by which that microscopic structure appears as matter and forces in actual 3+1 Reality.

18.3 Ownership Boundary between the finite SOURCE and String-Theory Microphysics

In MCC 3.0, the finite-SOURCE size $N_U = 1,015,000$ is not derived from the $E_8 \times E_8$ structure or from ten-dimensional compactification. N_U is a conditional output of the upstream Klein-Zeta finite-generation rule. String theory enters later, as an optional microscopic realization candidate for the common carrier, gauge representations, chirality, and gravity as finite addresses are rendered into physical phenotypes.

The present ownership structure is therefore separated into two lineages:

1. KZF finite SOURCE $\rightarrow$ WRRR address/filter $\rightarrow$ component/phenotype: the upstream generation lineage.
2. String microphysics $\rightarrow$ possible microscopic realization: a candidate microscopic implementation lineage for the common carrier and phenotype.

Whether these two lineages must meet through one common mathematical invariant remains OPEN. MCC 3.0 does not assume such a connection in order to justify N_U .

Why this boundary matters. It prevents a nonexistent derivation of the form “ten dimensions imply a finite SOURCE of 1,015,000 states,” while still identifying the exact point at which finite upstream generation and microscopic particle realization would have to meet in future work.

Part VI The Twist-Stress Universe

This part preserves the microscopic proposals of 2.1 and studies the dark sector through the distinction between Actual and Phenotype. The aim is consistency across galaxies, lensing and cosmology, not a claim of superior fit to standard cosmology. Here closure refers to the declared interpretation and calculations, not to a completed fit of every observation.

Version	Main scope	Preserved structure
WRRR Core 1.0	Domain-independent execution rules	Minimal computation, common carrier, phenotype
MCC 2.0	Cosmic execution and common carrier	LAW-SOURCE-EXECUTION-RENDERER-REALITY-RECORD
MCC 2.1	Conditional microscopic mathematical materials	MCC skeleton and three invariants
MCC 2.2	Twist stress and dark-mass phenotype	2.1 and WRRR Core 1.0

Chapter 19 What 2.2 Adds

19.1 Purpose

The starting question is not whether dark-matter particles are impossible. Keeping ordinary matter unchanged, can additional gravity be interpreted as the phenotype of stress stored in space without failing the declared galaxy, lensing and early-universe checks? Minimal computation means a sufficient structure that survives changing conditions, rather than maximal performance.

Known constants and observed relations are legitimate inputs for checking connections across scales. A discriminating prediction can add evidence but is not a prerequisite for an explanatory model.

Interpretation → consistency across scales → optional discriminating prediction. (19.1)

19.2 Central proposition

Ordinary matter retains its established field and particle description. In the dark sector, spatial twist stress is the proposed Actual, while the additional mass inferred from gravity is its Phenotype: a geometric response rewritten as a source on the right-hand side of the gravitational equation.

Distributed twist stress → renderer → effective dark-mass phenotype. (19.2)

Chapter 20 The Present Universe and the First Twist

20.1 Time is relational

The model does not require past and future to exist as separate physical places. A present state exists; time compares changes and records between states. Consciousness can interpret them as experience but is not required for the universe's computation.

$$\Delta t_{\text{clock}} := \Delta C / R_C. \quad (20.1)$$

Here C is a repeatable clock variable and R_C its reference rate. The covariant foliation label τ orders states; its interpretation is distinguished from the dynamical requirements of the action below.

20.2 The first twist

The Big Bang is interpreted as the initial logical asymmetry that makes the current relational network possible. In a wholly undifferentiated state there are no distinguished directions, positions, particles or records. First quantization distinguishes states; first twist distinguishes directions and paths.

Undifferentiated state $\rightarrow$ first quantization $\rightarrow$ distinguishability $\rightarrow$ first twist $\rightarrow$ relational universe. (20.2)

This is an interpretation, not a proved topology theorem. A Möbius strip illustrates orientation reversal, but has a boundary. Boundaryless topology requires a different example, such as a closed surface; actual three-dimensional cosmic topology remains observationally constrained.

Chapter 21 Finite Boundaryless Topology and Two Twist Directions

21.1 Local geometry and global topology

Locally nearly flat space can be globally closed. No local patch is assigned a privileged seam, centre or end. Global information may be recorded by holonomy, the change remaining after transport around a closed path. Under the conditions for the relevant Hodge decomposition,

$$A = d\phi + \delta B + h, \quad dh = 0, \quad \delta h = 0. \quad (21.1)$$

$$\oint_{\gamma} h = \Theta_{\gamma} \neq 0. \quad (21.2)$$

The gradient $d\phi$, rotational term δB and harmonic sector h separate local response from global path information. This construction allows ordinary local gravity alongside a finite boundaryless global structure.

21.2 Two directions

Opposite twists can coexist. Stress reads squared magnitudes, while directional bias reads their difference:

$$\kappa^2 = \kappa_+^2 + \kappa_-^2, \quad \chi_T = \frac{\kappa_+^2 - \kappa_-^2}{\kappa_+^2 + \kappa_-^2}. \quad (21.3)$$

Thus nonzero total stress need not imply a large net directional bias; χ_T is defined when its denominator is nonzero.

21.3 Identifiability of global size

$$L_{\gamma} = |\Theta_{\gamma}| \frac{c}{H_0} \sqrt{\frac{\zeta}{6f_T}}. \quad (21.4)$$

A representative closed-path size requires holonomy Θ_γ , stiffness ζ and stress fraction f_T . Present inputs constrain the product $\zeta\kappa^2$, not stiffness and twist separately. No unique cosmic size follows. Earlier 73 and 147 Gly estimates were conditional branches with additional assumptions, not final values of 2.2.

Chapter 22 Spatial Stress and Dark-Mass Phenotype

22.1 Minimal stress energy

$$u_T = \frac{\zeta c^4 \kappa^2}{16\pi G}, \quad \zeta > 0. \quad (22.1)$$

$$\alpha = \frac{c\kappa}{H_0}, \quad f_T = \frac{u_T}{\rho_c c^2} = \frac{\zeta \alpha^2}{6}. \quad (22.2)$$

Here ρ_c is critical mass density. An observed effective fraction fixes the product $\zeta\alpha^2$; it does not identify both factors. This is an identifiability result, not a numerical defect.

22.2 Mass as phenotype

Stress need not have electric or optical expression. Its metric response nevertheless changes matter and light trajectories and can be written as effective mass density:

$$\rho_T^{\text{eff}} = -\frac{1}{4\pi G} \nabla \cdot [(1 - \mu)\mathbf{g}]. \quad (22.3)$$

This is not a new particle number density. It rewrites a modified geometric response as a standard Poisson source. Background stress need not be uniform: matter, boundary conditions and global holonomy can affect its local response.

Chapter 23 Minimal Galactic Response

23.1 Static effective energy

$$E[\phi] = \int d^3x \left[\frac{a_T^2 F(Y)}{8\pi G} + \rho_b c^2 \phi \right]. \quad (23.1)$$

$$Y = \frac{c^4 |A|^2}{a_T^2}, \quad \mathbf{g} = -c^2 A. \quad (23.2)$$

The transition scale a_T joins Newtonian and low-acceleration regimes. A single response function avoids assigning an independent halo profile to each galaxy in this construction.

$$\nabla \cdot [\mu(|\mathbf{g}|/a_T)\mathbf{g}] = -4\pi G \rho_b, \quad \mu = F_Y. \quad (23.3)$$

23.2 Deep low-acceleration regime

$$F(Y) = \frac{2}{3} Y^{3/2} \Rightarrow \mu(x) = x. \quad (23.4)$$

$$g = \sqrt{a_T g_b}, \quad v_f^4 = G M_b a_T. \quad (23.5)$$

For spherical symmetry or sufficiently isolated circular-orbit approximations, the same response yields the radial-acceleration and baryonic Tully–Fisher relations. This is the declared consistency target, not a maximal-accuracy fit to every galaxy.

23.3 Static stability

$$\lambda_{\perp} = \mu > 0, \quad \lambda_{\parallel} = \mu + x\mu' > 0. \quad (23.6)$$

For $\mu = x$ and $x > 0$, the eigenvalues are x and $2x$. This checks static ellipticity. It does not prove stability of all covariant modes; cosmological scalar and long-wavelength Jeans sectors require separate checks.

Chapter 24 Connecting Cosmological and Galactic Scales

24.1 Matched response

Assume that isotropic 3+1 projection, background stress and the galactic response share the matched normalization $C = 1$:

$$a_T = cH_0\sqrt{f_T/8}, \quad f_T = 8(a_T/cH_0)^2. \quad (24.1-2)$$

The factor $1/8$ is a declared construction choice; Chapter 88 states its provenance and sensitivity. This relation is not a first-principles proof of that coefficient.

24.2 Numerical input

Quantity	Value	Role
H_0	67.4 km s ⁻¹ Mpc ⁻¹	Background change scale
f_T	0.265	Effective stress fraction
cH_0	6.55×10^{-10} m s ⁻²	Cosmological acceleration combination
a_T	1.192×10^{-10} m s ⁻²	Low-acceleration transition

The connection across scales is an explanatory consistency result. Since f_T is inferred in a Lambda-CDM analysis, it is not a wholly model-independent input. Later chapters use their separately stated preparations.

Chapter 25 Gravitational Lensing

25.1 One metric for matter and light

$$ds^2 = -(1 + 2\Phi/c^2)c^2dt^2 + (1 - 2\Psi/c^2)dx^2. \quad (25.1)$$

$$\hat{\alpha} = c^{-2} \int \nabla_{\perp}(\Phi + \Psi) d\ell. \quad (25.2)$$

Universal coupling to one metric and the specified static weak-field relation give

$$\Phi = \Psi + O(c^{-2}), \quad \eta_{\text{slip}} = \Psi/\Phi = 1 + O(c^{-2}). \quad (25.3)$$

$$\hat{\alpha} = 2c^{-2} \int \nabla_{\perp} \Phi d\ell \Rightarrow M_{\text{lens}} = M_{\text{dyn}} \text{ at leading order.} \quad (25.4)$$

The effective phenotype therefore affects both stellar motion and light deflection under these conditions. Colliding clusters and nonspherical systems still require dynamics in which the stress centre follows the observed lensing centre while separating from the baryonic gas centre when observations require it.

Chapter 26 Early-Universe and Structure Response

26.1 Covariant mathematical material

A covariant framework close to the relativistic Khronon theory of Blanchet and Skordis supplies mathematical material. It is imported material, not a fresh derivation of that theory. WRRRA's contribution here concerns the present-state interpretation, stress Actual, mass phenotype, global Hodge sector and cross-scale connection.

$$S = \frac{c^3}{16\pi G} \int d^4x \sqrt{-g} [R - 2J(Y) + 2K(Q)] + S_m. \quad (26.1)$$

$$n_\mu = -\frac{c}{Q} \nabla_\mu \tau, \quad Q = c \sqrt{-g^{\mu\nu} \nabla_\mu \tau \nabla_\nu \tau}. \quad (26.2)$$

$$A_\mu = c^2 n^\nu \nabla_\nu n_\mu, \quad Y = A_\mu A^\mu / c^4. \quad (26.3)$$

J governs spatial response and K the nearly pressureless cosmological component. Although τ is interpreted as an ordering label, $K(Q)$ gives its normalization physical significance. This tension with fully reparametrization-invariant timeless ontology remains in the cost ledger.

26.2 DBI-type cosmological response

$$K(Q) = \frac{2\mu^2}{\lambda_D} [1 - \sqrt{1 - \lambda_D(Q-1)^2}]. \quad (26.4)$$

$$q(a) = Q - 1 = [\lambda_D + (a^3/r)^2]^{-1/2}, \quad r = I_0/(2\mu^2). \quad (26.5)$$

$$w = \frac{\lambda_D q^2 + (1+q)\sqrt{1 - \lambda_D q^2} - 1}{2 + (1 + \lambda_D)q}. \quad (26.6)$$

$$c_{\text{ad}}^2 = \frac{q(1 - \lambda_D q^2)}{1 + q}. \quad (26.7)$$

$$c_s^2(a, k) = c_{\text{ad}}^2 \left[1 + \frac{c_{\text{ad}}^2 k^2}{4\pi G a^2 \rho_T (1 + w)} \right]^{-1}, \quad \Sigma_T = 0. \quad (26.8)$$

Here $q = Q - 1$ is not the deceleration parameter used in Chapters 62 and 92; μ in this action is a scale parameter, not the galactic interpolation function.

26.3 Recorded finite check

With $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_{T0} = 0.265$, $\mu^{-1} = 22.3 \text{ Mpc}$ and $\lambda_D = 1$, normalization to present stress density gives $r = 9.9913 \times 10^{-6}$.

State	q	w	c_{ad}^2
Very early $a \ll 10^{-3}$	$\rightarrow 1$	$\rightarrow 0$	$\rightarrow 0$
Recombination $a = 1/1100$	0.999999997	3.76×10^{-5}	2.83×10^{-9}
Present $a = 1$	9.99×10^{-6}	5.00×10^{-6}	9.99×10^{-6}
Relational state $a = 10$	9.99×10^{-9}	5.00×10^{-9}	9.99×10^{-9}

The recorded check admits a nearly pressureless early component and zero anisotropic stress in this branch. It is not a complete CMB angular-spectrum likelihood fit or an observation newly performed for this book.

Chapter 27 Interpretation, Tests and Assessment

27.1 Levels of assessment

Level	Question	Declared result
Interpretation	What Actual–Phenotype structure describes known phenomena?	Dark mass read as stress phenotype
Consistency	Do the specified galaxy, lensing and early-background calculations conflict?	No fatal conflict in the stated checks
Discrimination	Can measurements distinguish competing constructions?	Local control and nonspherical response remain open

The first two results do not depend on already having a new discriminating measurement.

27.2 Measurement directions

Candidate tests concern environment-dependent accelerations, gas–lensing offsets in colliding clusters, nonspherical shear, global path correlations and long-wavelength structure response.

$$\frac{\delta g(x)}{\delta B_{\text{boundary}}(y)} \neq 0. \quad (27.1)$$

Controlled changes of effective gravity without removing inertial mass would be a direct signal if local stress or boundary conditions could be manipulated. No present engineering method is established here.

27.3 Failure and follow-up conditions

Test	Declared status	Failure or remaining requirement
Galactic rotation/RAR	Stated relation reproduced	A fixed response fails systematically across the intended galaxy domain
Baryonic Tully–Fisher	Stated scaling reproduced	Wrong scaling or acceleration scale
Lensing	Conditional leading-order agreement	Required slip or gas–lensing offset cannot be produced
Early universe	Limited background check	Sound response spoils joint CMB/growth agreement
Global size	Undetermined	Independent stiffness/holonomy input needed
Relational time	Interpretation	Completion requires additional physical time degrees of freedom

27.4 Minimal-computation assessment

Two response functions and a correlation scale are costs. They may nevertheless be cheaper than independent owners for each target if one state supports the specified rotation, lensing and background response.

No absolute optimality is claimed.

$$C_{2.2} = \{N_{\text{field}}, N_{\text{response}}, N_{\text{scale}}, N_{\text{initial}}, C_{\text{update}}, C_{\text{failure}}\}. \quad (27.2)$$

Acceptance requires survival of the declared galaxy, lensing, background and structure tests within stated tolerance. (27.3)

Conclusion of Part VI

MCC 2.2 is a calculable interpretation, not a uniquely established cosmology. It keeps ordinary matter and connects additional gravitational response to stress phenotype. The recorded relations support limited cross-scale consistency; full CMB spectra, cluster dynamics and global-topology measurements remain un-completed tests. The initial quantization/twist and present-only ontology retain their interpretive status.

Part VII Interfaces to Standard Physics and Observational Calculation

2 Actual Renderer Reality Record

Version 2.3.2 does not mix ontology and calculation in a single sentence. Actual is the layer in which the common-carrier state, boundary relations, and twist exist. Renderer assigns the Actual state to measurable phenotypes such as particles, fields, mass, distance, and time. Reality is the layer of field equations and metrics used by experimenters. Record and Ledger preserve observations, sources, scopes of application, and update histories.

$$\Phi^R = \mathcal{R}[X_A], \quad \mathcal{O} = \mathcal{M}[\Phi^R]$$

Equation 2 Correspondence from Actual to Reality and observables




Figure 1 Four layers of Minimal Computing Cosmology and their information flow

Layer separation prevents the same object from being counted twice. The Standard Model electron and a WRRRA common-carrier state are not two electrons. The electron is a phenotype calculated in Reality, while the common carrier is the Actual basis from which that phenotype arises. Likewise, the dark-matter tensor and twist stress describe the same gravitational source in two languages and are not added simultaneously.

2 1 Ownership Rules

Ledger — Object · Reality calculation ownership · Actual interpretation ownership

- Object: Ordinary matter particles and three forces
 - Reality calculation ownership: Standard Model
 - Actual interpretation ownership: Boundary-specific channel phenotypes of the common carrier
- Object: Ordinary matter mass
 - Reality calculation ownership: Higgs coupling and QCD binding energy
 - Actual interpretation ownership: Phenotype of a state assigned as mass
- Object: Gravity
 - Reality calculation ownership: Metric of general relativity
 - Actual interpretation ownership: Geometric transmission of total state change
- Object: Dark matter gravitational source
 - Reality calculation ownership: Effective CDM tensor of Λ CDM
 - Actual interpretation ownership: Mass phenotype of local twist stress
- Object: Redshift and distance
 - Reality calculation ownership: FLRW scale factor and light paths
 - Actual interpretation ownership: Distance and wavelength phenotypes of global twist
- Object: Cosmic acceleration
 - Reality calculation ownership: Λ or a validated background module
 - Actual interpretation ownership: Background phenotype of global twist

3 Integration Rules

3 1 Module Adoption

An external theory becomes an official module of version 2.3.2 when it satisfies four conditions at once. First, it must have a domain validated by observation or experiment. Second, it must not logically contradict the fixed propositions of WRRRA. Third, ownership of what it calculates and at which layer must be explicit. Fourth, it must not duplicate a physical quantity already counted by another module.

$$M \in \mathfrak{M} \iff V(M) \wedge C(M) \wedge O(M) \wedge D(M)$$

Equation 3 Module-admission condition defined by validation V consistency C ownership O and non-duplication D

Adopting a module does not import the entire ontology of that theory. Version 2.3.2 adopts its calculated results and validated domain, then reassigns their ontological meaning through the layer rules of WRRRA. Source, version, conditions of application, and output are stored in Record. Re-derivation is not obligatory,

but provenance recording alone does not establish internal execution. Integration is executed only when the declared inputs pass through the specified WRRR transformation to calculated outputs; otherwise the status is external mathematical material or conditional correspondence.

3.2 Module Hierarchy




Figure 2 Hierarchy of WRRR Core and adopted and optional modules

Ledger — Status · Module · Scope of use

- Status: Higher specification
 - Module: WRRR Core 1.0
 - Scope of use: Layers update minimal computation ownership phenotype
- Status: Cosmological profile
 - Module: Minimal Computing Cosmology 2.3.2
 - Scope of use: Cosmological correspondence and integration
- Status: Validated calculation
 - Module: Standard Model
 - Scope of use: Ordinary matter and strong weak electromagnetic forces
- Status: Validated calculation
 - Module: General relativity
 - Scope of use: Metric gravity lensing orbits gravitational waves
- Status: Validated calculation
 - Module: FLRW and Λ CDM
 - Scope of use: Background distances CMB and structure growth
- Status: Optional mathematics
 - Module: String theory

- Scope of use: Used only to implement common-carrier microstructure

3.3 What Integration Does Not Permit

- The common carrier is not treated as an additional particle beside the photon, gluon, W, and Z.
- An Actual state and its Reality particle tensor are not added simultaneously as gravitational sources.
- Extra dimensions and a particular vacuum from string theory are not automatically imported as established propositions of version 2.3.2.
- The numerical successes of standard cosmology are retained and are not refitted with arbitrary correction terms.
- The incompleteness of optional research is not recast as a failure to explain existing observations.

4 The Standard Model and Ordinary Matter

4.1 Adopted Scope

The Standard Model calculates interactions among quarks, leptons, gauge bosons, and the Higgs field within the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge structure. Version 2.3.2 retains its calculations of the particle inventory, scattering amplitudes, coupling constants, mixing matrices, and symmetry breaking. Experimental values and uncertainties compiled through the Particle Data Group review system are adopted as inputs and validation records for the Reality module.

$$\mathcal{L}_{SM} = \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{fermion}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{Yukawa}}$$

Equation 4 Modular decomposition of the Standard Model Lagrangian

The masses of fermions and the W and Z bosons are calculated through their couplings to the Higgs field. Most of the mass of hadrons such as protons and neutrons comes not from the sum of quark rest masses but from QCD field energy and binding dynamics. The WRRRA mass phenotype does not erase this distinction. It interprets different generation mechanisms as being assigned through Renderer to the common measurable quantities of inertial and gravitational mass.

4.2 The Common Carrier and the Three Gauge Forces

The common carrier is not a fourth gauge boson. It is the common capacity in Actual that enables change to be transmitted, which Renderer expresses as different gauge interactions according to boundary conditions and channels. The gauge symmetries and field equations of the Standard Model are therefore retained. The unification of forces is not the forced merger of the three observed forces into a larger gauge group. It is a higher-order assignment in which different forces arise from the same transmission basis in Actual.

$$C_A \longrightarrow \{SU(3)_C, SU(2)_L, U(1)_Y\}_R$$

Equation 5 Generative correspondence from the common carrier to Standard Model gauge channels

Equation 5 does not claim to derive the particles and coupling constants of the Standard Model. It is a connection rule that preserves the WRRRA interpretation of matter developed from versions 1.0 through 2.1 while fixing the exact calculation ownership of Reality in the Standard Model. Direct derivation of particle species, generations, charges, and mixing angles from an Actual state remains optional microscopic research.

4 3 Quantum Fields and Vacuum

Quantum field theory in Reality calculates correlation functions, perturbative expansions, vacuum fluctuations, and scattering probabilities. WRRRA does not automatically interpret virtual particle pairs used as calculation terms as hidden events continually created and annihilated in Actual. A stable physical vacuum is a state with no observable flow of particle events; a particle event enters Record only when an external field or collision produces real particles. This distinction does not restrict quantum-field calculations. It separates ownership of a computational representation from ownership of an existence event.

5 General Relativity and Gravity

In version 2.3.2 gravity does not disappear merely because it is described as a phenotype. In Reality, gravity is the actual response of the metric to total energy-momentum. Matter moves on that metric, and light follows its null geodesics. Because the model retains the coordinate covariance and local conservation laws of general relativity, it also retains calculations of Solar System orbits, gravitational lensing, gravitational waves, and strong gravitational fields.

$$G_{\mu\nu}[g^R] + \Lambda g_{\mu\nu}^R = \frac{8\pi G}{c^4} (T_{\mu\nu}^m + T_{\mu\nu}^\Theta)$$

Equation 6 Gravitational field equation in Reality with untwisted matter and radiation plus twist stress

In Equation 6, T_m denotes all contributions from matter and radiation other than the dark component and the cosmological-constant term. It includes the energy-momentum of baryons, electrons, photons, neutrinos, and Standard Model fields. The cosmological constant remains on the left-hand side and is not counted again in T_m or T_Θ .

The common carrier does not reduce gravity to particle exchange of the same kind as a gauge force. The three gauge forces and gravity share the generative fact that state change is transmitted in Actual, while retaining different effective equations in Reality. This explains why gravity and the three forces operate in one universe without forcing their calculations into a single equation.

5 1 Equivalence Principle and Conservation

The ordinary-matter tensor calculated by the Standard Model and the effective tensor of twist stress enter the right-hand side of the same metric equation. Their sum must be covariantly conserved. This is not an interpretive option; it is an interface contract that follows from adopting general relativity. If twist stress does not satisfy it, the local-twist implementation must be revised.

6 Selective Adoption of String Theory

6 1 What Is Adopted

String theory uses one-dimensional objects and their vibrational modes instead of point particles and describes their coupling to spacetime fields through a worldsheet action. It provides a consistent mathematical structure that includes quantum gravity and tools for treating gauge and gravitational degrees of freedom together. Version 2.3.2 accepts this mathematics as a candidate microscopic implementation of the common carrier. Worldsheet boundaries, vibrational modes, chiral channels, anomaly cancellation, and compression of surplus degrees of freedom may be used to refine the 3+2 structure and rotor candidates studied in version 2.1.

$$S_P = -\frac{T_s}{2} \int d^2\sigma \sqrt{-h} h^{ab} \partial_a X^\mu \partial_b X^\nu G_{\mu\nu}(X)$$

Equation 7 Polyakov action of the optional string module

In Equation 7, X maps the worldsheet to background spacetime, h is the worldsheet metric, and G is the background metric. This action is not an established equation of the common carrier itself. To adopt a string module, a specific implementation must reproduce the particle species and symmetries of the Standard Model, agree with the low-energy limit of general relativity, and avoid double-counting additional degrees of freedom.

6.2 What Is Not Adopted

A particular number of dimensions, compactification, vacuum choice, or supersymmetric particle spectrum from string theory does not automatically become a proposition of version 2.3.2. Mathematics demonstrating internal consistency, such as the anomaly-cancellation result of Green and Schwarz, may be used. Promoting a specific microscopic implementation to a validated Reality module, however, requires an observational correspondence. Its present status is optional mathematical material.

Ledger — String-theory element · Status in version 2.3.2 · Condition for adoption

- String-theory element: Worldsheet and vibrational modes
 - Status in version 2.3.2: Available
 - Condition for adoption: When they simplify common-carrier microstructure
- String-theory element: Chiral structure and anomaly cancellation
 - Status in version 2.3.2: Available
 - Condition for adoption: When Standard Model symmetries are preserved
- String-theory element: Extra dimensions and compactification
 - Status in version 2.3.2: Implementation candidate
 - Condition for adoption: When they do not duplicate observable degrees of freedom
- String-theory element: Supersymmetric spectrum
 - Status in version 2.3.2: Undetermined
 - Condition for adoption: When experimental correspondence is established
- String-theory element: Claim of completed quantum gravity
 - Status in version 2.3.2: Not adopted
 - Condition for adoption: Requires a concrete low-energy correspondence and validation


7 A Finite Boundaryless Universe and Twist

7.1 Definition of Twist

Twist does not mean that the universe is physically bent like a sheet of paper inside an external dimension. It denotes the residual relation of a global connection that remains different after a state is transported around a closed path. A finite space without a boundary is possible in spherical spaces, tori, and quotient spaces. Version 2.3.2 does not fix the topology as a Möbius strip. The Möbius image is an analogy for directional reversal and nonprivileged gluing.

$$\Theta(x, \lambda) = \bar{\Theta}(\lambda) + \delta\Theta(x, \lambda)$$

Equation 8 Decomposition of one twist state into a global mean and local deviation



One state described by its mean and deviation, not two separate substances

Figure 3 Division of roles between global twist and local stress

The global component appears in Reality as the background of distance, redshift, and cosmic acceleration. Local deviations appear as additional gravitational sources and seeds of structure. They are not two separate substances, but the mean and deviation descriptions of one Actual state. In Equation 8, λ is a continuous approximation to the discrete updates n of Equation 1. Local twists with different directions are permitted when they satisfy total conservation and boundary conditions.

7 2 Total Content and Stress of the Universe

The total content of Actual is assumed not to leak outside. Ordinary matter retains its interpretation as existing fields and particles. Only the additional mass component corresponding to dark matter is interpreted as the phenotype of local twist stress preserved in space. The total mass of the universe is not redefined as stress. This restriction preserves the Standard Model interpretation of matter and the lineage from versions 1.0 through 2.1.

8 The Mass Phenotype of Dark Matter

8 1 Tensor Correspondence

Dark-matter phenomena require a common additional gravitational source in galactic rotation curves, gravitational lensing, colliding clusters, the CMB, and structure growth. Version 2.3.2 does not deny this observational fact. It adopts the effective CDM tensor used by Λ CDM as a Reality calculation module and interprets the source of the same tensor in Actual as local twist stress.

$$T_{\mu\nu}^{\Theta} \equiv T_{\mu\nu}^{\text{CDM}}, \quad \nabla_{\mu}^R T_{\Theta}^{\mu\nu} = 0, \quad p_{\Theta} \simeq 0, \quad c_{s,\Theta}^2 \simeq 0, \quad \Pi_{\Theta} \simeq 0$$

Equation 9 Reality equivalence of the dark-matter tensor and twist stress and the conditions for inheriting CDM calculations

This equivalence does not add a separately renamed gravitational source. CDM and twist stress divide the

calculation and interpretation ownership of the same term. To inherit the density-perturbation transfer functions and N-body results of standard CDM, twist stress must be separately covariantly conserved at the relevant scales; its effective pressure, sound speed, and anisotropic stress must be negligible; and its nongravitational coupling, dissipation, and self-interaction must remain below observational limits. If a self-action outside these conditions is introduced, it is tested as new dynamics rather than added simultaneously to the CDM module.

8 2 Colliding Clusters and Distribution

The weak-lensing map of the Bullet Cluster shows that the centers of the dominant baryonic plasma and the gravitational potential are separated. This observation requires an invisible additional gravitational source to move differently from the plasma during the collision. The twist-stress module must produce the same effectively nondissipative distribution as the CDM tensor; otherwise that implementation fails. Observation does not directly detect the word twist, but it specifies the dynamics that the interpretation must reproduce.

8 3 Measurability of Local Stress Relaxation

To determine whether twist stress is more than a relabeling, one must search for metric effects left by the relaxation or redistribution of local stress. Candidates include time-delay lensing in colliding clusters, relaxation of lensing mass centers, phase offsets in nonspherical halos, and boundary-dependent residuals in precision gravimetry. Version 2.3.2 does not yet fix the magnitude of such effects. This measurement belongs to the observational-protocol branch of the four areas of further research.

9 A Twist Interpretation of Expansion and Redshift

9 1 FLRW Calculations Are Retained

Observers do not directly see galaxies fleeing into an external space. They measure relations among redshift, brightness, angular distance, and time delay. Standard cosmology calculates these relations with the FLRW metric and scale factor. Version 2.3.2 retains that calculation and interprets the scale factor as not as the exterior size of Actual but as the effective scale at which global twist appears in distance measurements in Reality.

$$ds_R^2 = -c^2 dt_R^2 + a_R^2(t_R) \left[\frac{d\chi^2}{1 - k\chi^2} + \chi^2 d\Omega^2 \right]$$

Equation 10 FLRW metric retained in Reality

$$1 + z = \frac{a_R(t_o)}{a_R(t_e)}, \quad a_R = \mathcal{R}_a[\bar{\Theta}]$$

Equation 11 Correspondence between standard redshift and global twist

$$H_R^2 = \left(\frac{\dot{a}_R}{a_R} \right)^2 = \frac{8\pi G}{3} \rho_R - \frac{kc^2}{a_R^2} + \frac{\Lambda c^2}{3}$$

Equation 12 Friedmann equation retained in Reality

In Equations 10 through 12, a_R is a scale factor with dimensions of length, χ and k are respectively a dimensionless coordinate and curvature index, and k is one of -1, 0, or 1. The quantity ρ_R is the total effective

mass density, including the contribution of radiation energy density divided by c^2 . When the energy density ϵ_R is used, the first term in the Friedmann equation is written as $8\pi G\epsilon_R/(3c^2)$.

Equation 11 applies to cosmological redshift between a comoving emitter and observer in an FLRW background. An observed redshift may also include a Doppler component from peculiar velocity and a local gravitational redshift. In data analysis, these components are modeled or corrected separately before comparison with the background relation.

Expansion and twist use the same mathematics within their observationally equivalent domain, but they do not present the same ontological picture. The standard description defines distance through changes in a , while version 2.3.2 interprets those changes as updates of internal connections within one finite boundaryless Actual. Because no arbitrary redshift function is introduced, the model inherits existing supernova, BAO, and CMB fits.

9 2 A Conserved Actual and Changing Distances

$$\frac{dV_A}{d\lambda} = 0 \quad / \Rightarrow \quad \frac{da_R}{d\lambda} = 0$$

Equation 13 Compatibility of a conserved-total marker for Actual with scale change in Reality

Equation 13 does not fix coordinate volume or the physical volume of Reality. V_A is a marker for the conserved total of Actual that assumes no exterior, whereas a_R represents distance relations compared by internal observers. The parameter λ continuously approximates the discrete updates of Equation 1. If gluing and connection are distributed across all positions and updated incrementally, systematic changes in internal distances are possible without a moving external boundary. The specific dynamics of this correspondence belong to subsequent interface research.

9 3 Cosmic Acceleration

In Reality, acceleration is calculated with the standard equations including the cosmological constant Λ . In Actual, the background state of global twist appears as the acceleration phenotype. The mass phenotype of local stress and the acceleration phenotype of the global background need not be the same particle species. Components at different scales of the same twist state may be rendered as different effective terms.

10 The Big Bang and Presentness

10 1 The First Relational Opening

The Big Bang is not interpreted as the explosion of a point placed in empty space outside the universe. In version 2.3.2 it is the relational opening at which distinguishable states, relations, paths, and quantization first became established. The first twist may have appeared together with the first quantization that produced different paths and directions. This statement assigns meaning to the initial condition; it does not claim a completed equation of quantum gravity.

$$a_R \rightarrow 0 \quad / \Rightarrow \quad V_A \rightarrow 0$$

Equation 14 Distinction between the initial limit in Reality and the total volume of Actual

In standard cosmology, the limit in which a approaches zero is an extrapolation toward a hot, dense initial state. Version 2.3.2 does not turn this into the proposition that the whole Actual vanished into a Euclidean point. Nucleosynthesis, recombination, the CMB, and primordial perturbations are calculated by the standard thermal-history module; the relational opening supplies the Actual meaning of those calculations.

10 2 Meaning of Observing the Past

To observe a galaxy from 13 billion years ago is to reconstruct, with the present metric, a record stored in light that arrives now. The past universe does not remain as a place outside the present universe. Every observation occurs in the present, while distance and lookback time are relations among records. This interpretation does not alter calculations using the speed of light, null geodesics, or redshift.

11 Integrated Calculation of Large Scale Structure

11 1 Inheritance of Standard Calculations

Primordial CMB perturbations and the growth of large-scale structure are treated with the linear perturbation and N-body calculations of standard cosmology. In version 2.3.2, the dark-matter density used in those standard calculations is read as the mass phenotype of twist stress. With the same initial conditions, metric, and effective tensor, the calculated filaments, walls, clusters, and voids are identical.

$$\mathcal{O}_{\text{MCC}}[X_A; \mathcal{R}] = \mathcal{O}_{\mathfrak{M}_{\text{calc}}}[\Phi^R], \quad \mathfrak{M}_{\text{calc}} = (\text{SM}, \text{GR}, \text{FLRW}/\Lambda\text{CDM})$$

Equation 15 Condition for observational equivalence between adopted modules and Minimal Computing Cosmology

In Equation 15, $\mathfrak{M}_{\text{calc}}$ is not a simple sum of the Standard Model, general relativity, FLRW, and ΛCDM . It is a composed calculation package in which each module has its own domain and input-output contract. Ownership of a physical quantity is assigned to only one module, and a calculated output is passed between modules as the next module's input.

11 2 Limited Structure Growth Calculation

The following two-dimensional toy calculation uses a 256×256 periodic grid and 65,536 mass elements. A Zel'dovich displacement is applied to one scalar perturbation field generated with fixed random seed 22447864. The rule is taken from a standard structure-growth approximation; only the reading of the scalar field as twist stress belongs to the interpretation of version 2.3.2. This is neither a cosmological parameter fit nor a calculation of an actual CMB transfer function.

Density Phenotype Growing under the Same Effective Gravitational Rule




Figure 4 Density phenotype of twist stress growing under the adopted Zel'dovich rule

Ledger — Growth D · Density contrast standard deviation · Void fraction percent · High density fraction percent · Periodic connected components

- Growth D : 0.15
 - Density contrast standard deviation: 0.103
 - Void fraction percent: 0.0
 - High density fraction percent: 0.0
 - Periodic connected components: 0
- Growth D : 0.60
 - Density contrast standard deviation: 0.451

- Void fraction percent: 6.0
- High density fraction percent: 3.4
- Periodic connected components: 416
- Growth D: 1.05
 - Density contrast standard deviation: 0.915
 - Void fraction percent: 23.1
 - High density fraction percent: 8.4
 - Periodic connected components: 665
- Growth D: 1.50
 - Density contrast standard deviation: 1.273
 - Void fraction percent: 37.3
 - High density fraction percent: 11.4
 - Periodic connected components: 856

As D increases from 0.15 to 1.50, the standard deviation of the density contrast rises from about 0.103 to 1.273, while the fraction of regions below half the mean density rises from 0 percent to about 37.3 percent. The mean-mass error remains at floating-point level. Connected components are counted with four-neighbor connectivity and periodic boundary conditions that identify the left-right and top-bottom edges. This result does not validate a new cosmology. It demonstrates implementability: structure-formation calculations are preserved under the twist-stress interpretation when the same effective gravitational rule is retained.

12 Full Inheritance of Observational Results

Current execution grades, v0.12

Object	Current grade	Actual evidence and output
Full SM scattering, thermal history, CMB transfer, N-body String spin-2, BRST, compactification, Qr	External mathematical material / interface task External material / conditional correspondence	This table reports no internal execution Concrete vacuum and Qr selection and complete execution remain OPEN
Closed S^1 and classical common geometry	Internal specified witness	Equations and original lattice/curvature records in 7.1–7.5
Restricted structure-growth example	Specified finite calculation	Original growth-figure record; not full CMB or N-body
Common load and macro response	Internal execution	Chapters 84–91; B reproduce_all.py and original records
P4→P3 and P5 compensation	Restricted executed connection	Chapters 94–96 and their manuscript scripts
P6 and P7	Separate internal finite calculations	Chapters 97–98; all-series autonomous dynamics not executed

The successes of standard physics are not owned exclusively by a particular ontological vocabulary. Models using the same equations, initial conditions, boundary conditions, and observational correspondences produce the same numerical results within their shared domain. Version 2.3.2 makes this explicit and fixes the scope of inheritance in the following table.

Ledger — Observational domain · Adopted calculation · Actual interpretation in version 2.3.2 · Current judgment

- Observational domain: Particle scattering and decay
 - Adopted calculation: Standard Model
 - Actual interpretation in version 2.3.2: Channel phenotype of the common carrier
 - Current judgment: Calculation inherited
- Observational domain: Atoms and ordinary matter mass
 - Adopted calculation: Higgs QCD quantum field theory
 - Actual interpretation in version 2.3.2: Mass phenotype
 - Current judgment: Calculation inherited
- Observational domain: Solar System and binaries
 - Adopted calculation: General relativity
 - Actual interpretation in version 2.3.2: Metric response
 - Current judgment: Calculation inherited
- Observational domain: Gravitational waves
 - Adopted calculation: General relativity
 - Actual interpretation in version 2.3.2: Geometric transmission of Actual change
 - Current judgment: Calculation inherited
- Observational domain: Galaxy rotation and lensing
 - Adopted calculation: GR plus CDM
 - Actual interpretation in version 2.3.2: Local twist stress
 - Current judgment: Calculation inherited
- Observational domain: Colliding clusters
 - Adopted calculation: Nondissipative CDM tensor
 - Actual interpretation in version 2.3.2: Mobile stress distribution
 - Current judgment: Conditionally inherited
- Observational domain: Redshift supernovae BAO
 - Adopted calculation: FLRW and Λ
 - Actual interpretation in version 2.3.2: Global-twist distance phenotype
 - Current judgment: Calculation inherited
- Observational domain: CMB
 - Adopted calculation: Linear perturbations and recombination
 - Actual interpretation in version 2.3.2: Record of initial twist
 - Current judgment: Calculation inherited
- Observational domain: Large-scale structure
 - Adopted calculation: Perturbations and N-body
 - Actual interpretation in version 2.3.2: Mass phenotype of stress distribution
 - Current judgment: Calculation inherited
- Observational domain: Early thermal history
 - Adopted calculation: Nucleosynthesis and standard cosmology

- Actual interpretation in version 2.3.2: Reality after relational opening
- Current judgment: Calculation inherited

The final Planck mission data showed strong agreement with the spatially flat six-parameter Λ CDM model. Version 2.3.2 does not replace those parameters or the CMB acoustic scale with new values. Even when current data do not uniquely choose between the standard description and the ontology of version 2.3.2, the compatibility of the same calculations with version 2.3.2 forms the observational basis of the integrated model.

The 2025 DESI DR2 BAO analysis reported a preference for a time-varying dark-energy equation of state over Λ CDM in extended models combining DESI and CMB data. That preference remained after adding the 2026 Lyman-alpha forest full-shape analysis, although the central value of the new high-redshift measurement moved toward Λ CDM. Version 2.3.2 therefore does not make Λ a fixed ontological conclusion. It retains Λ , or the latest validated background module, as the owner of the Reality calculation. Even when observational modules are updated, assignment under the Core to global twist as a distance and background phenotype remains unless a contradiction is demonstrated.

13 Restricted Further Research

Further research is opened only when the integrated model demonstrably fails to explain, calculate, or map an item, or when a logical contradiction or double-counting is found between adopted modules. Its scope is limited to the missing element and its interface. Recalculating under the WRRRA name a phenomenon already calculated by the Standard Model, general relativity, or Λ CDM is not a required task.

13 1 Twist Metric Interface

Specify in one action or update equation the Renderer from global twist to a_R and H_R and the tensor rule from local twist to $T\Theta$. This research is needed when an independent dynamical calculation is desired. It is not a condition for the observationally equivalent interpretation of version 2.3.2.

13 2 Common Carrier Standard Model Interface

Study how a common-carrier state produces the representations of $SU(3)_C \times SU(2)_L \times U(1)_Y$, particle generations, charge, mass, and mixing. String worldsheets, a 3+2 internal structure, chiral channels, and a rotor are candidate tools. They remain optional implementations until a candidate actually reproduces the Standard Model.

13 3 Topology Presentness Interface

For admissible finite boundaryless topologies, calculate the global holonomy, total size, and range of directional reversal. At the same time, formalize the relation of the present update order λ to relativistic covariance and proper time. Both questions must be treated without reintroducing an exterior of the universe or an absolute clock.

13 4 Measurement Falsification Interface

Quantify residuals that local stress relaxation could leave in lensing time delays, colliding-cluster halo shapes, or precision gravimetry. Constrain global topology with repeated images, matched circles, and distance correlations. If the results are exactly identical to standard predictions, the proposal remains an ontological interpretation. It becomes an independent prediction only when it produces a reproducible difference.

Ledger — Research area · Opening condition · Protected component · Completion output

- Research area: Twist metric

- Opening condition: Need to calculate intrinsic evolution without standard modules
- Protected component: FLRW and GR observational calculations
- Completion output: Covariant action and Renderer
- Research area: Common-carrier matter
 - Opening condition: Attempt to derive particle species and couplings
 - Protected component: Validated values of the Standard Model
 - Completion output: Gauge-preserving microscopic map
- Research area: Topology and presentness
 - Opening condition: Attempt to formalize total size and λ
 - Protected component: Finite boundarylessness and proper-time calculations
 - Completion output: Topology group and covariant update rule
- Research area: Measurement and falsification
 - Opening condition: A residual distinct from standard results is calculated
 - Protected component: Full inheritance of existing observations
 - Completion output: Preregistered statistical test

14 Failure Conditions and Order of Revision

Version 2.3.2 does not alter the Core in response to every objection. When failure is established, revision begins with the lowest-level implementation. Inputs and numerical implementation of the observational-fit module are checked first, followed by Renderer and the correspondence definitions. Local or global twist implementations are revised next. WRRRA Core is reopened by separate agreement only when failure follows directly from Core grammar.

- Revise the global Renderer if one global-twist correspondence is shown unable to reproduce redshift, distance, the CMB, and structure growth within the same FLRW metric.
- Revise the local-stress implementation if its tensor cannot satisfy conservation, lensing dynamics, and the nondissipative behavior of colliding clusters together.
- Discard a microscopic common-carrier implementation if its Renderer cannot preserve Standard Model gauge symmetries without duplicate degrees of freedom.
- Revise the topological hypothesis if finite boundarylessness and nonprivileged gluing cannot agree with admissible cosmic topologies and observational constraints.
- Reconsider the physical profile of presentness if present-state updating and relativistic covariance cannot be logically compatible.

14 1 Module Replacement Rule

If a new validated theory calculates the same phenomenon more accurately over a wider domain than an existing module, the Reality module may be replaced. Replacement does not alter WRRRA Core. Inputs, outputs, units, conservation laws, and observational correspondences are specified, and differences from the previous module are stored in Record. This structure allows version 2.3.2 to absorb future research without losing its own criteria.

15 Final Integrated Propositions

Ledger — No · Final proposition

- No: 1
 - Final proposition: WRRRA Core 1.0 is the higher-order execution grammar of Minimal Computing Cosmology 2.3.2
- No: 2
 - Final proposition: Our universe need not be special or optimal and maintains viable sufficiency
- No: 3
 - Final proposition: The universe is one finite boundaryless present whole
- No: 4
 - Final proposition: Time is a relation of change and record rather than an independent substance and Reality retains proper-time calculations
- No: 5
 - Final proposition: Ordinary matter and the three gauge forces are calculated by the Standard Model in Reality
- No: 6
 - Final proposition: Ordinary matter is interpreted in Actual as boundary-specific phenotypes of the common carrier
- No: 7
 - Final proposition: Gravity is the actual metric response to total energy-momentum and is calculated by general relativity
- No: 8
 - Final proposition: Dark matter is the mass phenotype of local twist stress and is not added again to the effective CDM tensor
- No: 9
 - Final proposition: Expansion redshift and cosmic acceleration are distance and background phenotypes of global twist
- No: 10
 - Final proposition: The Big Bang is the first relational opening rather than a point explosion in external space
- No: 11
 - Final proposition: The Standard Model general relativity and standard cosmology are adopted as validated calculation modules
- No: 12
 - Final proposition: String theory is an optional mathematical module used only when needed for a microscopic implementation of the common carrier
- No: 13
 - Final proposition: External research is adopted without internal re-derivation when it satisfies validation consistency ownership and non-duplication
- No: 14
 - Final proposition: Further research is limited to WRRRA interfaces where a real gap or module conflict has been identified

Conclusion

Minimal Computing Cosmology 2.3.2 neither discards the Standard Model nor elevates it into the final ontology of the universe. The Standard Model calculates ordinary matter, general relativity calculates gravity, and standard cosmology calculates the background and structure formation. WRRRA organizes these results as phenotypes of Reality produced through Renderer by one common-carrier and twist state.

Standard modules provide external mathematical material for major observations. Internal executions in this book are the specified finite witnesses, background responses, restricted growth example and connections in Chapters 92–99. Full CMB transfer functions, a complete thermal history, all scattering and N-body calculations are not counted as executed here. Their interfaces require actual input-to-output execution before promotion to integrated status.

The originality of version 2.3.2 does not lie in discarding existing numerical results and fitting new ones. It reads reality as phenotype, expansion as the distance phenotype of global twist rather than the motion of an external boundary, and force unification as the generative basis of a common carrier rather than a single additional particle. This interpretation is logically compatible with standard calculations and closes as one conceivable finite boundaryless universe.

Part VIII Cumulative Integration and Nonduplicating Interfaces

Chapter 28 Native Twist Dynamics and the CDM Equivalence Branch

MCC 2.2 and 2.3.1 are not competing versions. They operate at different computational depths. In the native branch of 2.2, a twist action directly produces an effective tensor.

$$T_{\mu\nu}^{\Theta} = -\frac{2}{\sqrt{-g}} \frac{\delta S_{\Theta}}{\delta g^{\mu\nu}}$$

In the equivalence branch of 2.3.1, the same tensor is identified with the CDM computational representation within a declared scale and accuracy.

$$T_{\mu\nu}^{\Theta} \equiv T_{\mu\nu}^{CDM}, \quad \nabla^{\mu} T_{\mu\nu}^{\Theta} = 0, \quad p_{\Theta} \simeq 0, \quad c_{s,\Theta}^2 \simeq 0, \quad \Pi_{\Theta} \simeq 0$$

They are not entered twice in one gravitational equation.

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} (T_{\mu\nu}^m + T_{\mu\nu}^{\Theta}), \quad T^{\Theta} \neq T^{\Theta} + T^{CDM}$$

The native branch gains its own calculation ownership when it passes conservation, lensing, colliding-cluster, CMB, and structure-formation tests. Where that calculation is not yet closed, the equivalence branch inherits validated calculations. This interface preserves twist as more than a renamed source while retaining successful standard computations.

Chapter 29 One Global and Local Twist State

$$\Theta(x, \lambda) = \bar{\Theta}(\lambda) + \delta\Theta(x, \lambda)$$

The global mean $\bar{\Theta}$ is rendered as a_R, H_R , and cosmological distance relations. The local deviation $\delta\Theta$ is rendered as $T_{\mu\nu}^{\Theta}$ and as the seed of structure growth.

$$a_R = \mathcal{R}_a [\bar{\Theta}], \quad T_{\mu\nu}^{\Theta} = \mathcal{R}_T [\delta\Theta]$$

These are not two independent dark substances. They are a scale decomposition of one Actual twist state. The global mean is not added again to the dark-matter tensor, and the local deviation is not duplicated in the cosmological-constant term.

Chapter 30 Final Interface of the Common Carrier and Four Forces

The common carrier is neither a fifth force nor a field that flattens all four forces into one gauge current. A shared Actual transport basis is rendered through different owner maps as the three gauge representations and gravity.

$$\mathcal{R}_{gauge}[C_A] \rightarrow \{SU(3)_C, SU(2)_L, U(1)_Y\}_R, \quad \mathcal{R}_G[C_A, \Phi_R] \rightarrow g_{\mu\nu}$$

The gauge forces selectively read J_a , while gravity reads total $T_{\mu\nu}$. The four forces are unified within one transport basis and one Reality precisely without being declared identical. The singlet counterexample and fifteen-channel equations of MCC 2.0 provide the argument; the Standard Model and general relativity modules of 2.3.1 provide the Reality calculations.

Chapter 31 Compatibility of a Matter Occupied Reality and the Twist Universe

Ordinary matter retains its conventional particle and field description. Only the additional gravitating dark component is read as a local twist-stress phenotype. At the same time, the lawful matter and antimatter sectors are separated from long-term actual occupancy.

$$\mathcal{H}_{lawful} = \mathcal{H}_{matter} \oplus \mathcal{H}_{antimatter}, \quad Occupied\ Reality(\lambda) \subset \mathcal{H}_{matter}$$

Adding twist does not convert all ordinary matter into spatial stress. A matter-occupied Reality is a proposition about long-term ordinary-matter occupancy. The dark mass phenotype is a proposition about spatial stress that has negligible electric and optical expression but a nonzero gravitational expression. Their ownership domains differ and they are mutually compatible.

Chapter 32 The Present Universe and Experienced Time

Actual contains an updating present. The past remains as Record in the present, and the future is an unrealized possibility. Proper time in general relativity and the FLRW time coordinate remain effective quantities that calculate relations among change and records in Reality. Consciousness is not a prerequisite for cosmic computation. It is a phenotype that integrates memory and prediction into an experienced flow of time. Change proceeds without consciousness, but no subject experiences its duration.

When stability fails, a second universe is not created outside the first. The single Actual updates into a next state. An unstable state updates again, while a stable state persists for the duration of its stability. Finite boundarylessness means that no external edge is required; it does not require an infinite material container.

Chapter 33 Final Integrated Propositions

Ledger — Number · Fixed proposition of Minimal Computing Cosmology 2.3.2

- Number: 1
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Our universe need not be privileged or optimally designed and is one persistent viable structure
- Number: 2
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Minimal computation is the nearest sufficient computation and reserve that avoids fatal failure under change and extremes
- Number: 3
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: LAW SOURCE EXECUTION RENDERER REALITY and RECORD form one cosmic execution genealogy
- Number: 4
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: One constraint-preserving opening starts one history and radiation can arise during execution without a separate SOURCE register
- Number: 5

- Fixed proposition of Minimal Computing Cosmology 2.3.2: Gravity reads total stress energy while three gauge forces read selective currents so the four forces are asymmetrically unified
- Number: 6
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Fifteen chiral channels share one common carrier skeleton while abstract rank and gauge-covariant realization remain distinct
- Number: 7
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Matter is a jointly owned phenotype of mass charge spin location and record
- Number: 8
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Antimatter may remain lawful while long-term occupied Reality consists of matter
- Number: 9
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Quantum possibilities are weighted within lawful support and one event and Record condition the next present
- Number: 10
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: The universe may be a finite boundaryless Actual with no privileged edge or origin point
- Number: 11
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: First quantization and first twist interpret the Big Bang as the opening of relation and direction
- Number: 12
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Global twist is rendered as distance redshift and background acceleration while local twist is rendered as dark mass and structure
- Number: 13
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: The native twist branch of 2.2 and the CDM equivalence branch of 2.3.1 are non-substitutive and are never double counted
- Number: 14
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: The Standard Model GR FLRW and validated cosmology are Reality modules and do not replace the WRRR skeleton
- Number: 15
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: String theory is optional microscopic material and does not delete minimal computation the common carrier or phenotype
- Number: 16
 - Fixed proposition of Minimal Computing Cosmology 2.3.2: Past and future need not exist as separate places and time is a relation among change and records
- Proposition 17: Closed internal mode n supplies discrete mass precursors $p\theta=n/R$ and $m_n^2=m_0^2+n^2/R^2$, assigned to mass phenotype through the renderer.
- Proposition 18: Modes $+n$ and $-n$ project to equal mass and opposite charge in the specified universal witness.
- Proposition 19: Gravity is common geometry responding to total stress–energy; the present witness does not quantize gravity by the mass-mode rule.
- Proposition 20: The Standard Model and GR supply Reality calculations of pole mass and metric observables; they do not replace the upstream witness.
- Proposition 21: There is one universe in the adopted ontology. The same Actual updates when a state

loses stability; a stable state persists while its stability holds.

- Proposition 22: Physical time relates change and records. Experienced temporal flow involves memory and prediction; the present-only reading is a philosophical interpretation, not a measured fact about every consciousness.

Conclusion

The center of Minimal Computing Cosmology 2.3.2 is not a new name for dark matter. It is the full execution architecture in which the least independent input and reusable laws update the present, a common carrier and selective renderer produce dimensions, matter, forces, and observable Reality, and quantum possibility becomes one event and one recorded history.

The four forces are unified without being made identical. Gravity reads the $T_{\mu\nu}$ of the whole realized system, whereas electromagnetism, the weak force, and the strong force selectively read internal representation currents J_a . The common carrier preserves that distinction while providing one Actual transport basis. Matter is not a stored list of numbers but a phenotype in which mass, charge, spin, location, and history are jointly owned. LAW may permit both matter and antimatter while a long-lived actual universe occupies the matter sector.

The twist universe is not a decorative later addition. It is the cosmological model that closes finiteness, dark gravity, and distance relations on top of the execution skeleton. Global holonomy and local stress are the mean and deviation of one state, rendered respectively as the distance phenotype of expansion and acceleration and the mass phenotype of dark matter. Native twist dynamics connects galaxies, lensing, and the early universe directly; the CDM-equivalence branch inherits validated structure-formation calculations under the same effective tensor conditions.

The Standard Model, general relativity, FLRW, and cosmological data enter as computational modules inside this skeleton. They are neither discarded nor promoted into the ontology of the whole architecture. They calculate where they have demonstrated accuracy, while WRRRA explains how their outputs arise from one Actual and become Reality and Record. Version 2.3.2 is therefore the cumulative integrated edition that preserves both the achievements of MCC 2.0 and the complete twist structure of MCC 2.2.

Appendix Core Equations and Claim Grades

Appendix A Ledger of Core Equations

Ledger — Function · Equation

- Function: Overall execution
 - Equation: $LAW \rightarrow SOURCE \rightarrow EXECUTION \rightarrow RENDERER \rightarrow REALITY \rightarrow RECORD$
- Function: Minimal cost
 - Equation: $C_{total}=L_{LAW}+L_{SOURCE}+C_{read}+C_{evolution}+C_{render}+C_{record}$
- Function: Opening operation
 - Equation: $U_S=\exp(i\eta\Omega/\hbar)$, $U_S \dagger P_\Omega U_S=P_\Omega+\eta$
- Function: FLRW boundary
 - Equation: $B_\eta=\eta\chi-\sqrt{6}\tilde{M}_P|\eta|\ln(a/a^*)+constant$
- Function: Dimension filter
 - Equation: $K_{DF}=F \dagger F$, $L_{DF}=I-K_{DF}$
- Function: Common transport
 - Equation: $D_C:\oplus H_i \rightarrow \oplus H_i$, $rank(D_C|band)=15$
- Function: Gravity
 - Equation: $G_{\mu\nu}+\Lambda g_{\mu\nu}=(8\pi G/c^4)T_{\mu\nu_total}$
- Function: Gauge coupling
 - Equation: $D\mu=\nabla\mu^{geo}-ig_1YB\mu-ig_2T^aW\mu^a-ig_3t^AG\mu^A$
- Function: Reality state
 - Equation: $\rho_Reality=P_vis[U(\tau)U_{Sp_dorm}U_S \dagger U \dagger (\tau)]$
- Function: Event probability
 - Equation: $p(a|\rho)=Tr[L_a(\rho)]$
- Function: Path equivalence
 - Equation: $R_{\{\gamma_1\}}(B_f,B_i)\simeq_gauge R_{\{\gamma_2\}}(B_f,B_i)$
- Function: Recovery
 - Equation: $Recovery\circ\Phi=id$ on $A_protected$
- Function: Support set
 - Equation: $supp P_WRA\subseteq\Gamma_lawful$
- Function: History information
 - Equation: $I_history(T)=\sum_t-\log_2p(a_t|h_{\{t-1\}})$
- Function: Finite runtime
 - Equation: $H_N=H_Jacobi\otimes H_loc\otimes H_syn$

- Function: Radiation generation
 - Equation: $\rho_r(t) = a(t)^{-4} \int_{t^*}^t a(t')^4 Q_r(t') dt'$
- Function: Radiation takeover
 - Equation: $R = \rho_r / \rho_\chi$, $R/R = 2H + Q_r(1/\rho_r + 1/\rho_\chi) \geq 2H$
- Function: Matter asymmetry
 - Equation: $\Delta_M = n_M - n_{Mbar}$, $d(a^3 \Delta_M)/dt = a^3(S_M - S_{Mbar})$
- Function: CP boundary
 - Equation: $F_B(\eta, LAW_{CP-symmetric}) = 0$

Appendix B Sources and Grades of Claims

The purpose of this ledger is to prevent established equations of standard physics from being mistaken for new derivations by WRRRA, while also preventing WRRRA's theoretical achievements from being buried under excessive qualification.

Ledger — Label · Meaning · Rule of Use

- Label: BASELINE
 - Meaning: Established equations or observations of standard physics
 - Rule of Use: State the source and domain of applicability
- Label: DEFINITION
 - Meaning: Layers or ownership assignments newly defined by WRRRA
 - Rule of Use: Distinguish definitions from their consequences
- Label: THEORETICAL RESULT
 - Meaning: Structural conclusion logically obtained from assumptions
 - Rule of Use: State the assumptions and the scope of counterexamples
- Label: CONSTRUCTED/EXACT
 - Meaning: Exactly constructed or calculated in a specified model
 - Rule of Use: Do not identify it with the entire theory
- Label: CALIBRATED
 - Meaning: Fixed using observational values
 - Rule of Use: Do not call it an independent derivation
- Label: PREDICTION
 - Meaning: Output not used in calibration
 - Rule of Use: Freeze the procedure, timing, and failure conditions
- Label: OPEN
 - Meaning: A connection not yet determined
 - Rule of Use: Maintain it as a separate list after the core achievements

Appendix C Major Public Research Genealogy

- 10.5281/zenodo.22038142 - Hypothesis on Gravity and Three-Dimensional Space
- 10.5281/zenodo.22052565 - Common-Carrier Unification without a Grand-Unified Gauge Field
- 10.5281/zenodo.22056516 - Minimal Common Transporter for Chiral Matter
- 10.5281/zenodo.22063551 - Autonomous Rank Opening of 15 Channels
- 10.5281/zenodo.22121047 - Conditional BRST Closure of the Common Carrier
- 10.5281/zenodo.22124886 - Renderer Path Independence and the Einstein Branch
- 10.5281/zenodo.22125506 - Recoverable Rendering and Metric Faithfulness
- 10.5281/zenodo.22126125 — time-uniform recoverable rendering
- 10.5281/zenodo.22126918 — finite-Jacobi compatibility benchmark
- 10.5281/zenodo.22137993 — open-universe execution model
- 10.5281/zenodo.22139233 — WRRRA information provision ledger
- 10.5281/zenodo.22141639 — two-pass open-outcome rendering
- 10.5281/zenodo.22142880 - The Position of WRRRA among Four Axiomatic Frameworks
- 10.5281/zenodo.22177447 - Final Our-Universe Scenario of Minimal Computing Cosmology
- 10.5281/zenodo.22239485 - Ownership of the Four Forces
- 10.5281/zenodo.22239574 - Big Bang, Dark Matter, and the Conservation Ledger
- 10.5281/zenodo.22239864 / 22239895 - Antimatter Direction-History Verification Program
- 10.5281/zenodo.22330256 / 22331940 - Information Closure and the Interpretation-Prediction Methodology
- 10.5281/zenodo.22447864 - A Physically Auditable Classifier of Interpretation and Prediction
- 10.5281/zenodo.22502619 - WRRRA-Worm Execution and Comparative Study

Appendix String Reinforcement Ledger

Appendix A. Claim-Grade Ledger

Ledger — Item · Grade · Exact Meaning

- Item: Three invariants of MCC 2.0
 - Grade: CORE / PRESERVED
 - Exact Meaning: Must not be changed by this combination
- Item: 10D superstring requirement
 - Grade: IMPORTED BASELINE
 - Exact Meaning: Microscopic consistency of perturbative superstring theory
- Item: E8×E8 heterotic as primary material
 - Grade: MODEL CHOICE
 - Exact Meaning: Preferred candidate under MCC cost; no uniqueness claim
- Item: 10D heterotic low-energy action
 - Grade: IMPORTED BASELINE
 - Exact Meaning: Not a new MCC derivation
- Item: Green–Schwarz anomaly closure
 - Grade: IMPORTED EXACT/KNOWN
 - Exact Meaning: Existing result of string theory
- Item: $D_C(4)=P15$ Red[$\mathcal{D}10$] P15
 - Grade: MCC PROPOSED IDENTIFICATION
 - Exact Meaning: Requires validation in a concrete compactification
- Item: index = 3 target
 - Grade: CONDITIONAL
 - Exact Meaning: Not an automatic prediction of three generations
- Item: String zero mode → phenotype
 - Grade: MCC THEORETICAL RECAST
 - Exact Meaning: Places string material in the upstream phenotype layer
- Item: Qr from modulus/gauge kinetic function
 - Grade: CANDIDATE CONSTRUCTION
 - Exact Meaning: Numerical calculation remains OPEN
- Item: ηB source from string sector
 - Grade: CANDIDATE

- Exact Meaning: Mechanism selection and calculation remain OPEN
- Item: WRRRA boundary $\leftrightarrow$ string microstates
 - Grade: CONDITIONAL IDENTIFICATION
 - Exact Meaning: Test beginning with specific black holes
- Item: MCC landscape Pareto filter
 - Grade: NEW SELECTION PRINCIPLE
 - Exact Meaning: Requires actual vacuum ranking

Appendix B. Allowed and Forbidden Repair Material

Ledger — Decision · Material / Action · Reason

- Decision: Allowed
 - Material / Action: 10D heterotic effective action
 - Reason: Microscopic gravity/gauge/anomaly owner
- Decision: Allowed
 - Material / Action: Calabi–Yau/bundle compactification
 - Reason: Generates 4D chiral zero modes and representations
- Decision: Allowed
 - Material / Action: Green–Schwarz mechanism
 - Reason: Reinforces anomaly consistency
- Decision: Allowed
 - Material / Action: Gauge kinetic function / moduli couplings
 - Reason: Candidate microscopic owner of Q_r
- Decision: Allowed
 - Material / Action: String axion/moduli/hidden sectors
 - Reason: Dark-sector candidates
- Decision: Allowed
 - Material / Action: Controlled black-hole microstate counting
 - Reason: Reinforces microscopic capacity of the WRRRA boundary
- Decision: Conditional
 - Material / Action: Type II/D-brane/F-theory material
 - Reason: Use when it has lower Pareto cost than heterotic material or supplies a required structure
- Decision: Forbidden
 - Material / Action: Promoting 11D M-theory to the required skeleton
 - Reason: Outside the present scope and not required by this construction
- Decision: Forbidden
 - Material / Action: Treating the string landscape itself as the selection explanation
 - Reason: Cannot be adopted without MCC minimality screening
- Decision: Forbidden
 - Material / Action: Replacing the common carrier with the phrase “everything is string”
 - Reason: Eliminates the actual transport operator

- Decision: Forbidden
 - Material / Action: String mode = observed particle
 - Reason: Destroys the phenotype layer
- Decision: Forbidden
 - Material / Action: Extra SOURCE hiding radiation/asymmetry in vacuum labels
 - Reason: Violates minimal SOURCE

Appendix C. References

- [C1] Choi, W. (2026). WRRRA Core 1.0: Domain-Agnostic Execution Architecture — Frozen Specification with Application Registry. 7 September 2026.
- [M1] Choi, W. (2026). Minimal Computing Cosmology 2.0 — Reader-Centered Final Edition. Internal final manuscript, 2026-09-07.
- [S1] Marchesano, F., Shiu, G., & Weigand, T. (2024). The Standard Model from String Theory: What Have We Learned? *Annual Review of Nuclear and Particle Science*, 74, 113–140. <https://doi.org/10.1146/annurev-nucl-102622-012235>
- [S2] Cicoli, M., Conlon, J. P., Maharana, A., Parameswaran, S., Quevedo, F., & Zavala, I. (2023). String Cosmology: from the Early Universe to Today. *arXiv:2303.04819*.
- [S3] Strominger, A., & Vafa, C. (1996). Microscopic Origin of the Bekenstein-Hawking Entropy. *Physics Letters B*, 379, 99–104. *arXiv:hep-th/9601029*.
- [S4] Denef, F., Douglas, M. R., & Kachru, S. (2007). Physics of String Flux Compactifications. *Annual Review of Nuclear and Particle Science*, 57, 119–144. <https://doi.org/10.1146/annurev.nucl.57.090506.123042>
- [S5] Candelas, P., Horowitz, G. T., Strominger, A., & Witten, E. (1985). Vacuum Configurations for Superstrings. *Nuclear Physics B*, 258, 46–74.
- [S6] Gross, D. J., Harvey, J. A., Martinec, E., & Rohm, R. (1985). Heterotic String. *Physical Review Letters*, 54, 502–505.
- [S7] Green, M. B., Schwarz, J. H., & Witten, E. (1987–1988). *Superstring Theory*, Vols. 1–2. Cambridge University Press.
- [S8] Blumenhagen, R., Cvetič, M., Langacker, P., & Shiu, G. (2005). Toward Realistic Intersecting D-Brane Models. *Annual Review of Nuclear and Particle Science*, 55, 71–139. <https://doi.org/10.1146/annurev.nucl.55.090704.151541>

Appendix Twist References

References

- [1] Bekenstein, J. and Milgrom, M. 1984. Does the missing mass problem signal the breakdown of Newtonian gravity. *The Astrophysical Journal* 286 7-14. <https://doi.org/10.1086/162570>
- [2] Milgrom, M. 1983. A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *The Astrophysical Journal* 270 365-370. <https://doi.org/10.1086/161130>
- [3] McGaugh, S. S., Lelli, F. and Schombert, J. M. 2016. Radial Acceleration Relation in Rotationally Supported Galaxies. *Physical Review Letters* 117 201101. <https://doi.org/10.1103/PhysRevLett.117.201101>
- [4] Lelli, F., McGaugh, S. S. and Schombert, J. M. 2016. SPARC Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves. *The Astronomical Journal* 152 157. <https://doi.org/10.3847/0004-6256/152/6/157>
- [5] Planck Collaboration. 2020. Planck 2018 Results VI Cosmological Parameters. *Astronomy and Astrophysics* 641 A6. <https://doi.org/10.1051/0004-6361/201833910>
- [6] Clowe, D. et al. 2006. A Direct Empirical Proof of the Existence of Dark Matter. *The Astrophysical Journal Letters* 648 L109-L113. <https://doi.org/10.1086/508162>
- [7] Blanchet, L. and Skordis, C. 2024. Relativistic Khronon Theory in Agreement with Modified Newtonian Dynamics and Large Scale Cosmology. *Journal of Cosmology and Astroparticle Physics* 2024 11 040. <https://doi.org/10.1088/1475-7516/2024/11/040>
- [8] Zlosnik, T. G., Ferreira, P. G. and Starkman, G. D. 2007. Modifying Gravity with the Aether An Alternative to Dark Matter. *Physical Review D* 75 044017. <https://doi.org/10.1103/PhysRevD.75.044017>
- [9] Planck Collaboration. 2016. Planck 2015 Results XVIII Background Geometry and Topology of the Universe. *Astronomy and Astrophysics* 594 A18. <https://doi.org/10.1051/0004-6361/201525829>
- [10] Choi, W. 2026. Minimal Computing Cosmology and WRRR research records. <https://doi.org/10.5281/zenodo.22660185> and <https://doi.org/10.5281/zenodo.22650956>

Appendix Integrated Definitions and Research Lineage

Definitions

Ledger — Term · Definition

- Term: WRRR
 - Definition: Higher-order execution grammar organizing state relation law boundary update phenotype and record
- Term: Minimal computation
 - Definition: Principle of maintaining enough computation and reserve to avoid fatal failure under changing conditions
- Term: Actual
 - Definition: Existence layer containing the common carrier boundary relations present state and twist
- Term: Renderer
 - Definition: Correspondence that assigns an Actual state to particles fields mass distance and time in Reality
- Term: Reality
 - Definition: Effective reality layer calculated by observers with standard physical equations
- Term: Record
 - Definition: Record preserving observations sources uncertainties and scope of application
- Term: Ledger
 - Definition: Ledger managing module versions ownership updates and failure history
- Term: Common carrier
 - Definition: Common basis enabling change and transmission in Actual rather than an additional particle
- Term: Phenotype
 - Definition: Result rendered as an observable property through Renderer
- Term: Twist
 - Definition: Residual global or local relation left after a state is transported around a closed connection
- Term: Mass phenotype
 - Definition: State in which local twist stress is assigned as an additional gravitational source in Reality
- Term: Relational opening
 - Definition: Actual interpretation of the Big Bang in which distinguishable states paths and quantization first become established

References and Research Lineage

1. Choi W. WRRRA and Minimal Computing Cosmology research record. Zenodo. DOI 10.5281/zenodo.22447864.
2. Choi W. Related WRRRA Core and Minimal Computing Cosmology record. Zenodo. DOI 10.5281/zenodo.22660185.
3. Choi W. Related Minimal Computing Cosmology record. Zenodo. DOI 10.5281/zenodo.22650956.
4. Takahashi F et al. Particle Data Group. Review of Particle Physics. International Journal of Modern Physics A 41 2630011. 2026.
5. Einstein A. Die Feldgleichungen der Gravitation. Sitzungsberichte der Preussischen Akademie der Wissenschaften. 844 to 847. 1915.
6. Friedmann A. Über die Krümmung des Raumes. Zeitschrift für Physik 10 377 to 386. 1922. DOI 10.1007/BF01332580.
7. Planck Collaboration. Planck 2018 Results VI Cosmological Parameters. Astronomy and Astrophysics 641 A6. 2020. DOI 10.1051/0004-6361/201833910.
8. Clowe D et al. A Direct Empirical Proof of the Existence of Dark Matter. Astrophysical Journal Letters 648 L109 to L113. 2006. DOI 10.1086/508162.
9. Riess A G et al. Observational Evidence from Supernovae for an Accelerating Universe and a Cosmological Constant. Astronomical Journal 116 1009 to 1038. 1998. DOI 10.1086/300499.
10. Perlmutter S et al. Measurements of Omega and Lambda from 42 High Redshift Supernovae. Astrophysical Journal 517 565 to 586. 1999. DOI 10.1086/307221.
11. Zel'dovich Y B. Gravitational Instability An Approximate Theory for Large Density Perturbations. Astronomy and Astrophysics 5 84 to 89. 1970.
12. Polyakov A M. Quantum Geometry of Bosonic Strings. Physics Letters B 103 207 to 210. 1981. DOI 10.1016/0370-2693(81)90743-7.
13. Green M B and Schwarz J H. Anomaly Cancellations in Supersymmetric D 10 Gauge Theory and Superstring Theory. Physics Letters B 149 117 to 122. 1984. DOI 10.1016/0370-2693(84)91565-X.
14. Davis T M and Lineweaver C H. Expanding Confusion Common Misconceptions of Cosmological Horizons and the Superluminal Expansion of the Universe. Publications of the Astronomical Society of Australia 21 97 to 109. 2004. DOI 10.1071/AS03040.
15. DESI Collaboration. DESI DR2 Results II Measurements of Baryon Acoustic Oscillations and Cosmological Constraints. Physical Review D 112 083515. 2025. arXiv 2503.14738.
16. DESI Collaboration. DESI DR2 Results IV Alcock Paczynski Measurements from the Lyman Alpha Forest and Cosmological Constraints. 2026. arXiv 2607.27410.
17. DESI Collaboration. DESI DR2 Publications. Releases of 19 March 2025 and 30 July 2026. <https://data.desi.lbl.gov/doc/papers/dr2/>.

Part IX — The Starting Point of 3.0: Klein-Type Boundarylessness, the Zeta Function, and a Finite SOURCE

The strangest part of 3.0 begins here. Can a universe be finite without being sealed off by a terminal wall? Why should the zeros of a function that encodes prime-number structure meet a finite address-space boundary? And can that meeting descend into an actually executable SOURCE size? This Part introduces those questions first through pictures and thought experiments, and only then places the equations.

Reading rule for Part IX. MCC 3.0 does not claim that the universe is literally a Möbius strip or a Klein bottle, and it does not claim to prove the Riemann Hypothesis. These objects are used as mathematical candidates for testing two narrower ideas: execution that is finite without terminating at an edge, and a finite spectral count carrying prime-number structure.

MCC 2.3.2 and the early WRRM implementation already used a finite address space in computation, but its size $N = 1,000,000$ was an externally supplied convenience cutoff. Version 3.0 begins by replacing that last external upper bound with an internally generated candidate.

Chapter 34 — From an Exhibition to a Question

Research sometimes begins not with a calculator but with a way of seeing. In an exhibition, a form can appear not to bend at one isolated point but to change direction gradually across its entire structure. The question that arose was not “is the universe curved?” It was simpler:

Can a structure be finite without ever meeting an edge?

Imagine an ant walking on the surface of Earth. It can keep moving north, east, and south without ever meeting a railing marked “the world ends here.” The surface area is finite, but the surface has no boundary. Earth is not a model of the universe; it merely gives the quickest intuition that finite and having an edge are not the same statement.

Thought experiment — are finiteness and boundary the same thing? A room floor is finite and has a wall as a boundary. A sphere surface is also finite, but it has no wall. MCC 3.0 does not begin by fixing the actual shape of space. It asks for the minimal structure by which an executable state can fail to terminate at an endpoint and instead reconnect with itself.

The immediate conceptual trigger for MCC 3.0 was not a new astronomical measurement. While viewing KOO JEONG A: OUSSMOS at Leeum Museum of Art in Seoul, the image arose of a twist distributed across an entire form rather than concentrated at a single local fold. That intuition was later formalized in the precursor study *Finite but Boundaryless: A WRRRA Hybrid Expansion and Accumulated Twist Cosmology with Hidden Gravitational State* [3.0-R1].

In that precursor, the Möbius strip was never proposed as the literal topology of the universe. It has a boundary and the wrong dimension. The real question was:

If the universe is finite but has no terminal wall, how does its state close, how can transport return to itself, and what physical residue can be left by an accumulated twist?

The precursor chain was

global closure → distributed connection → holonomy → transport residue → twist stress → gravity .

The exhibition is acknowledged as a conceptual trigger, not scientific evidence. The equations, computations, and falsification conditions that follow must stand independently.

MCC 3.0 pushes the earlier question one level upstream. If we want to speak of a finite but boundaryless universe, it is not enough to say that the spatial form is finite. The SOURCE that executes that universe should itself be finitely generated. Up through MCC 2.3.2 and WRRRA_M, the address space used an external cutoff. The objective of 3.0 is to replace that last external upper bound with an internal generation rule and then follow the result all the way down to particles, energy, gravity, and expansion.

Chapter 35 — Is the Universe a Finished File? WRRRA and Minimal Computation

v0.12 chapter update

Completed physical infinity is assigned Null as an execution premise. Null is not numerical zero: this model assigns no executable physical state to a demand for completed infinite resources. This is a philosophical restriction, not an observational theorem about every part of the universe. Continuous probes, real probabilities and derivatives remain effective mathematical descriptions, not infinite-precision devices. An adequate present can support the next action without perfect reset; P7 distinguishes that sufficiency from accumulating inaccessible distinctions.

The question is not merely a computer metaphor. When we watch a movie, every scene is already stored in the file. A game is different: the next scene depends on the current state of play. The universe tested by MCC is closer to the second picture — except that MCC does not assume an external player or external computer. The model assumption is that the universe's own current state and fixed law produce the next present.

Minimal computation therefore does not mean “compute as little as possible.” It means that the system need not physically own the entire unrealized future in advance, provided that the information required for the next execution is preserved. If the present state, relations, residue, and resource ledger are sufficient for the next update, unrealized possibilities need not already occupy the physical SOURCE.

35.1 A finished file or an executed present?

Minimal Computing Cosmology asks whether the universe must store every future event and particle state in advance, or whether it can generate only the next required state from the present state and law.

MCC adopts the second option as a model hypothesis. The universe is not treated as a database in which all future entries already exist, but as an execution system that computes the next present from the current present. MCC 2.3.2 organized this view through SOURCE-RELATION-BOUNDARY-RESIDUE-UPDATE-LEDGER and WRRRA [3.0-R2].

SOURCE owns executable possibility. RELATION owns connection and transformation. BOUNDARY owns admissibility. RESIDUE is history retained in the present and capable of affecting the next state. LEDGER owns conserved, transferred, and assigned resources.

35.2 Non-specialness

MCC does not require our universe to be selected as a uniquely privileged mathematical object. A fixed generation law may permit multiple possible realizations, with our universe corresponding to one realized branch. The target is not an infinitely detailed initial specification written solely for our universe, but a finite rule capable of reproducing its observed structure.

35.3 Mathematical infinity and physical execution

Infinite sets and limits remain indispensable mathematical tools. MCC 3.0 does not reject them. It asks a narrower physical question: must a physically instantiated SOURCE own an infinite address space all at once?

mathematical infinity $\neq$ physically instantiated infinite SOURCE .
--

Chapter 36 — Finite without Boundary: Sphere, Möbius Strip, and Klein Bottle

“Finite but boundaryless” sounds contradictory only if finiteness is confused with having a physical edge. In topology, the two are independent. What matters is not how an object is drawn on paper, but which points are identified and how paths return.

A sphere is the familiar example: finite area, no edge. A torus is another. A useful intuition is an old video game in which walking off the right side of the screen returns you on the left, and leaving the top returns you at the bottom.

The Möbius strip adds something different. Take a strip of paper, twist one end by half a turn, and glue it to the other. An arrow transported around the strip can return with reversed orientation. It is an intuitive model of orientation reversal. But a Möbius strip still has an edge, so it is not itself a boundaryless candidate for the universe.

A Klein bottle preserves orientation reversal while eliminating the boundary. The familiar bottle-shaped drawing appears to pass its neck through its body because a Klein bottle cannot be embedded smoothly in ordinary three-dimensional Euclidean space without self-intersection. Mathematically it is a closed two-dimensional manifold; an embedding without self-intersection requires four dimensions.

Important — MCC 3.0 does not claim that the universe is “shaped like a Klein bottle.” The Klein-type identification is a candidate state-space closure used to close one coordinate pair of the finite SOURCE by an orientation-reversing return instead of letting the state terminate at an endpoint.

The schematic identification used in 3.0 is

$$\boxed{(-T, \theta) \sim (T, -\theta)}, \quad \theta \in S^1.$$

Here θ is periodic and T denotes the finite range on the identified coordinate pair. In this context, “finite” refers to compact/bounded closure of the identified state-space coordinate, not to a claim that the continuum contains only finitely many mathematical points.

The important idea is not the bottle drawing. The endpoint is not a terminal wall. When a trajectory reaches the identified boundary, it does not disappear; it re-enters the state space with transformed orientation information. This creates room for global identification, holonomy, and transport residue.

The earlier question “why is an infinite SOURCE unnecessary?” therefore becomes more specific: can a finite address space be generated and then closed in such a way that its ends are not merely computational stop signs?

MCC 3.0 imposes six requirements on its starting rule:

1. it must be finite;
2. it must generate addresses;
3. it must admit nonterminal closure rather than simple termination at the boundary;
4. it must connect to the frozen WRRM_M filter;
5. it must preserve inherited validation ledgers without hidden retuning;
6. if it fails, the failed rule must be separately identifiable.

Chapter 37 — The Shadow of the Primes: The Riemann Zeta Function and the Finite Klein-Zeta Generator

When the zeta function suddenly enters a cosmology book, the obvious question is: why this function? The answer in MCC 3.0 is not mystical. The Riemann zeta function is a compressed map in which a sum over all positive integers and a product over all prime numbers describe the same analytic object.

For $\Re(s) > 1$,

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}.$$

Euler showed that the same function can be written as

$$\zeta(s) = \prod_p (1 - p^{-s})^{-1}.$$

This Euler product is why prime-number structure is encoded in the zeta function. In a WRR branch where SOURCE addresses are organized through prime/composite structure, the zeta function is therefore not introduced merely as decorative mathematics; it is a natural spectral organizer to test.

After analytic continuation, the zeta function has zeros. The negative even integers are the trivial zeros. The nontrivial zeros lie in the critical strip $0 < \Re(s) < 1$. The Riemann Hypothesis states that all nontrivial zeros have real part $1/2$.

Claim boundary. MCC 3.0 does not prove the Riemann Hypothesis. The standard known heights of nontrivial zeros are used as inputs and numerical cross-check locations. The finite Euler-Maclaurin surrogate tests numerical consistency at those locations; it is not an independent theorem deriving the zero spectrum from scratch.

The relevant feature here is that the zeros form an ordered spectral sequence. Rather than physically instantiating an infinite sequence, MCC 3.0 counts how many positive nontrivial zeros lie below a finite phase boundary defined inside the model. That finite count becomes C .

The first coordinate is the arithmetic ledger:

$$(0.05, 0.268, 0.682) = \left(\frac{25}{500}, \frac{134}{500}, \frac{341}{500} \right),$$

so the least common exact denominator is

$$\boxed{L = 500}.$$

The second coordinate is the inherited neutrino harmonic ledger $(0, 5, 29)$, which supplies

$$\boxed{H = 29}.$$

The third coordinate is the finite zeta-focus count. Declare the finite phase boundary

$$T_H = 2\pi H = 182.212373908208.$$

There are 70 positive nontrivial zeta zeros below this boundary. In particular,

$$\gamma_{70} = 182.20707848436646 < T_H < \gamma_{71} = 184.87446784838750,$$

so

$$C = 70.$$

To avoid treating an infinite Dirichlet series as the physical SOURCE itself, the numerical implementation uses a finite Euler-Maclaurin surrogate with cutoff

$$M = LH = 14,500.$$

At the standard locations of the first 70 zeros, the maximum absolute residual is approximately 5.29×10^{-13} ; at γ_{70} it is about 6.98×10^{-14} [3.0-R4]. These are finite numerical cross-checks of known zero locations, not independent zero generation.

The Klein-type closure is represented schematically by

$$(-T, \theta) \sim (T, -\theta),$$

not as a claim that measured spacetime is a Klein bottle, but as a candidate way to prevent the finite address system from ending in a terminal wall.

Chapter 38 — Why 29 and 70? Generating 1,015,000 Addresses

Now the three numbers meet. $L = 500$ comes from the exact integer resolution of the cosmic arithmetic ledger. $H = 29$ is the largest index in the inherited neutrino harmonic ledger $(0, 5, 29)$. C is the number of positive nontrivial zeta zeros that fall inside the finite phase window associated with H .

Why multiply H by 2π instead of treating 29 directly as a length? In a harmonic phase, the natural angular period is 2π . MCC 3.0 therefore declares the candidate phase boundary

$$T = 2\pi H = 2\pi \times 29 = 182.212373908208.$$

The 70th standard nontrivial zero lies just below this boundary and the 71st lies above it:

$$\gamma_{70} = 182.20707848436646 < T < \gamma_{71} = 184.87446784838750.$$

Hence the number of positive nontrivial zeros in the finite phase window is $C = 70$. Structurally, the order matters: T is declared first and C is counted afterward. The count is not selected post hoc to produce 70. The physical necessity of $T = 2\pi H$, however, and the independence of L, H, C remain OPEN.

The present candidate product rule is

$$N_U = L \times H \times C = 500 \times 29 \times 70 = 1,015,000.$$

Why is this number interesting? Earlier WRRM calculations used $N = 1,000,000$ as a convenient cutoff. The 1,015,000-state candidate is not a new cutoff chosen to recover those outputs. It is generated from three internal ledgers. The decisive test is therefore whether the same downstream computation survives when this new SOURCE is inserted.

Let

$$0 \leq \ell < L, \quad 0 \leq h < H, \quad 1 \leq z \leq C.$$

Serialize the product state by

$$n = 1 + \ell + L[h + H(z - 1)].$$

Then

$$1 \leq n \leq 1,015,000.$$

Label 1 remains an inactive multiplicative/vacuum reference and the executable address domain is

$$n = 2, \dots, 1,015,000.$$

In this construction, integers are execution addresses serializing a finite product state; MCC does not require the physically realized SOURCE to be an actually instantiated infinite integer substrate.

Chapter 39 — When a Number Enters the Physical Ledger: The Prime/Composite Filter

Obtaining 1,015,000 is not yet physics. The number matters only if the resulting address set can be inserted into the existing WRRM_M filter and then survive the inherited ledgers and downstream calculations without hidden retuning. The task of this chapter is therefore not “we found a number,” but “what survives when the new SOURCE is plugged into the frozen execution system?”

Addresses are separated into prime, even-composite, and odd-composite sectors while inheriting the existing phase-aware rule. A prime label is not automatically a particle name. Integers are execution addresses; only after filtering and later component/phenotype readout can a physical interpretation arise.

Three dangerous misunderstandings

1. “The universe is literally a Klein bottle.” No. A Klein-type quotient is a state-space closure candidate.
2. “MCC 3.0 solves the Riemann Hypothesis.” No. Known zeta-zero data are used for finite counting and numerical consistency checks.
3. “1,015,000 is a measured cosmological constant.” No. It is a conditional MCC/WRRM output under the declared $N_U = LHC$ construction.

The address weights are

$$w_n \propto n^{-\alpha},$$

with the inherited reference exponent

$$\alpha_0 = 1.8996876950554356.$$

At the new $N_U = 1,015,000$, keeping the old arithmetic calibration frozen first yields

$$\phi = 0.05000002538983289,$$

$$D = 0.26800001828881304,$$

$$R = 0.6819999563213559.$$

The new finite address counts are

- primes: 79,608;
- even composites: 507,499;
- odd composites: 427,892.

The odd-composite sector inherits the existing eight-frame zeta-phase drive. Reapplying only the same two previously declared arithmetic freedoms gives

$$\alpha_1 = 1.8996877935161325,$$

$$h_1 = 1.4476744689338703.$$

The effective comparison

$$\beta_{eff} = 0.8654567230719301$$

is a derived readout, not a third fit. The final arithmetic ledger closes at

$$\phi/D/R = 0.05/0.268/0.682.$$

The achievement of Part IX is therefore not merely the value 1, 015, 000. A finite nonterminal closure candidate, a zeta spectral count carrying prime structure, an inherited harmonic ledger, and the arithmetic composition ledger meet inside one finite-SOURCE construction. The next Part asks whether that SOURCE can descend through records, components, harmonics, and phenotype without breaking the frozen downstream physics.

Part X — From High Energy to Phenotype: Records, Components, and Harmonics

Chapter 40 — Research Background and Objective

40.1 The connection problem remaining after 2.3.2

MCC 2.3.2 and the existing WRRM integration provide a downstream executable path connecting the common carrier, address conditioning, internal state, phenotype energy, pressure, information load, gravity, and expansion. What had not yet been organized sufficiently into one ledger was how high-energy Actual descends into the stable phenotypes of the present universe, and where the component identity, activation, and assembly-order effects observed in Prime Parts belong within the upstream-downstream structure.

The direct questions of this study can be grouped into four sets.

1. How should a high-energy or high-temperature condition be entered into a WRRM state?
2. How can the boundary between microscopic state and macroscopic record be defined without identifying it with a minimum time or one universal constant?
3. When is a pre-phenotype part judged to be an independent component, and how is its origin preserved after a state transition?
4. How can stable component assembly and harmonic states be connected to the existing phenotype-energy and downstream-gravity ledgers without energy duplication?

These questions are not independent. If temperature does not directly generate a particle, intermediate states are needed between temperature and phenotype: filter, admission/return, record, component stabilization, and assembly. The strategy of this study is therefore to solve the problems separately and then reconnect them in stage 0.12 as a single typed ledger.

40.2 Completion criterion

This study is not defined so that it can be completed only by producing a new independent prediction. Its completion criterion consists of four items:

verified inputs -> WRRM-specific transformation -> outputs -> falsification conditions

Even when established constants and observations are used, values reproduced through WRRM-specific state transformations after those inputs have been frozen are treated as consistency and explanatory achievements. Conversely, if an as-yet-unmeasured quantity is produced after the model has been fixed, it is distinguished as a conditional WRRM prediction.

40.3 Rule against mixing types

The most important ledger rule in this study is that numbers with different meanings must not be treated as the same physical quantity. At least the following five types are kept distinct.

Ledger type	Meaning	Representative example
ADDRESS	admitted, resident, and returned address shares	0.05/0.268/0.682
ENERGY RECORD	physical distribution of SI energy state readout and repeated stability	0.0493/0.265/0.6857 $B_D, B_R(\delta)$
COMPONENT PHASE	part identity that preserves origin relative-phase ledger for multiple modes	O , active/reserve $\Phi, 0 : 5 : 29$

Without this distinction, address shares may be mistaken for energy shares, the number of components may be mistaken for an amount of energy, or a phase-cycle count may be mistaken for a particle count.

Chapter 41 — Method — Frozen Baseline and Typed Ledger

41.1 Freeze contract

The new high-energy extension is attached upstream of the existing WRR M model. Downstream coefficients therefore may not be readjusted in order to match the established outputs. The frozen baseline is shown in Table 1.

Quantity	Frozen value	Type
phenotype address fraction	0.050000000	ADDRESS
resident nonphenotype fraction	0.268000000	ADDRESS
return fraction	0.682000000	ADDRESS
phenotype energy fraction	0.0493	ENERGY
resident nonphenotype energy fraction	0.2650	ENERGY
return energy fraction	0.6857	ENERGY
deceleration diagnostic q_0	-0.52855	DOWNSTREAM
reference rotation	207.5109051266 km/s	DOWNSTREAM
conditional lensing	0.5355865106 arcsec	DOWNSTREAM

The address fractions sum to one, and the energy fractions form a separately normalized SI ledger. The two ledgers are not automatically identified merely because some values are numerically similar.

41.2 Overall research flow

The entire 0.1-0.12 connection is

high-energy Actual $\rightarrow \rho_T \rightarrow$ filter/readout $\rightarrow$ SOURCE branching $\rightarrow$ shutter/record $\rightarrow B_{\mu M} \rightarrow$ component activation $\rightarrow$ codeword/harmonic $\rightarrow$ phenotype $\rightarrow E(V) \rightarrow P \rightarrow$ load $\rightarrow$ twist/gravity/expansion.

Each arrow is treated not as a loose conceptual association but as an interface specifying what type of input is received and what type of output is passed to the next stage.

Chapter 42 — Stage 0.1 — Freezing the Baseline

42.1 Verified inputs

The address branching, internal Hamiltonian, SI energy ledger, pressure, information load, local gravity, and homogeneous-expansion calculations used in WRR Core 1.0, MCC 2.3.2, and the existing WRR M 1.0 are fixed.

42.2 WRR-specific transformation

The new upstream variables do not alter the cause of the established downstream outputs. They are added only as an input layer determining which state is supplied to the downstream model. Therefore, after introducing the new upstream layer, the downstream coefficients may not be retuned merely to preserve the existing q_0 .

42.3 Outputs and interpretation

The frozen values are reference points that the new research is not allowed to move. A change that rearranges only internal labels or phases, without changing total energy or population, must preserve the baseline downstream values. By contrast, an upstream change that genuinely changes the population must produce a downstream response. This distinction is tested directly in the stage-0.10 sensitivity analysis.

42.4 Falsification conditions

The 0.1 contract fails if any of the following occurs.

- Existing SI coefficients are recalibrated in order to connect the new upstream layer.
- ADDRESS fractions are automatically substituted for ENERGY fractions.
- An increase in the number of components or harmonics is treated as an increase in cosmic energy.

Verdict: PASS.

Chapter 43 — Stage 0.2 — From Temperature to a WRRRA State

43.1 Problem statement

Entering temperature T as one number does not uniquely determine the internal state. At the same positive temperature, the distribution depends on the Hamiltonian energy gaps and eigenspaces. A constant scalar shift $H \rightarrow H + cI$ cancels from the normalized Gibbs state. The energy, work and free-energy ledger still requires a consistent reference; a physical volume-dependent term is distinguished from a constant reference shift. Temperature is therefore treated in WRRRA as one condition used to construct a state, not as the state itself.

43.2 Definition of the thermal probe

For a finite Hamiltonian H and reference energy E_0 , define

$$\rho_T = \frac{\exp[-(H - E_0 I)/(k_B T)]}{\text{Tr} \exp[-(H - E_0 I)/(k_B T)]}$$

as the thermal probe. The required input is therefore the triple

$$(T, H, E\text{-prescription}).$$

The Planck temperature may be used as a comparison scale, but a state not defined by the present finite representation is not silently extrapolated merely because the Planck scale is invoked. Regions outside the defined representation remain NULL.

43.3 Numerical check using the inherited internal gap

The inherited internal gap is

$$\Delta E = 431.081244315 \text{ MeV.}$$

The existing conditional excited population is

$$p_{\text{exc}} = 0.0799766545,$$

with excitation contribution

$$E_{\text{exc}} = 34.4764357391 \text{ MeV.}$$

If this population is interpreted backward only as a two-state Gibbs population, the equivalent temperature is approximately

$$T_{\text{Gibbs}} \simeq 2.048 \times 10^{12} \text{ K},$$

while the temperature scale of the gap itself is

$$\frac{\Delta E}{k_B} \simeq 5.00249 \times 10^{12} \text{ K}.$$

These values are vastly below the Planck temperature. There is therefore no basis for identifying the internal thermal mixing of the current particle model with the Planck boundary.

43.4 Outputs and falsification conditions

The conclusion that temperature alone does not determine the state becomes an important constraint on all later stages. At fixed positive T , Hamiltonians differing only by a scalar identity term yield the same Gibbs state. Outside that equivalence, an output inconsistent with the stated Gibbs formula, positivity or normalization fails the implementation. Likewise, the inherited address-conditioned population must not be automatically reinterpreted as a thermal population.

Verdict: finite thermal probe PASS.

Chapter 44 — Stage 0.3 — The Planck Boundary and Filter Stability

44.1 Test hypothesis

The initial candidate hypothesis was

$$T \uparrow \Rightarrow \text{thermal mixing} \uparrow \Rightarrow \text{filter gap} \downarrow \Rightarrow 0.$$

If this were correct, the phenotype filter would lose its discriminating power at high temperature and be reselected as the system cooled. This stage tests that intuition directly under the frozen filter rule.

44.2 Compatibility and the gap

Define the compatibility between carrier states u_i and u_j as

$$c(i, j) = -\|u_i - u_j\|_C^2.$$

For F_{DX} to be selected uniquely in the inherited four-filter structure, both left and right competitive gaps must be positive.

Representative baseline responses are

$$0.672634137, \quad 0.825428002,$$

and the baseline calculation gives

$$\Delta_L = \Delta_R = 0.04669193039 > 0.$$

Selection of F_{DX} is retained in the nine inherited perturbation cases. For a finite perturbation ρ , one may use the stability condition that selection is preserved whenever the minimum gap remains sufficiently large, schematically

$$\min \Delta > 4\rho.$$

44.3 High-temperature limiting test

When only the thermal population is mixed toward a uniform high- T limit, the frozen response rule does not send the gap to zero. A positive gap remains even for a uniform population.

Therefore

$\text{thermal population mixing alone} \not\Rightarrow \text{filter destruction}$

under the present implementation.

44.4 Meaning of the negative result

This is not a failed stage. It narrows the physical location of the missing mechanism. If high-to-low temperature evolution is to control filter stability, temperature must act on response contrast, the metric, the read-out structure, or an activation law, rather than merely on carrier populations. That conclusion becomes the starting point of stage 0.4.

Verdict: NEGATIVE RESULT PRESERVED.

Chapter 45 — Stage 0.4 — Cooling and Filter Stabilization

45.1 Contrast order parameter

Take 1.0 as the center of the two filter responses and set the frozen contrast to

$$\kappa_0 = 0.25.$$

The two representative response positions are then

$$1 - \kappa_0 = 0.75, \quad 1 + \kappa_0 = 1.25.$$

Define the diagnostic coordinate

$$m_F = \frac{\kappa}{\kappa_0}.$$

At $m_F = 0$, contrast vanishes and the competing filters are tied. At $m_F = 1$, the frozen filter structure is recovered.

45.2 Gap at small contrast

In the small- κ region, the calculated gap is approximately

$$\Delta \approx 1.56498\kappa^2.$$

Substituting $\kappa = 0.25m_F$ gives

$$\Delta \approx 0.0978113m_F^2.$$

If the environmental perturbation scale is ρ_{env} , the approximate order required to preserve filter selection may be written as

$$m_{F,\text{crit}} \approx 6.395\sqrt{\rho_{\text{env}}}.$$

45.3 What is connected and what remains to be generated

This stage establishes the final three links in

$$T \rightarrow m_F(T) \rightarrow \kappa(T) \rightarrow \Delta(T) \rightarrow F.$$

That is, once m_F is supplied, the route from contrast to filter gap and filter selection is calculable. The first arrow, $T \rightarrow m_F(T)$, still requires an independent physical generation law.

Accordingly, an arbitrary interpolation such as

$$m_F = 1 - T/T_P$$

is not inserted merely to force $T_F = T_P$. Doing so would amount to entering the desired boundary by

hand rather than deriving a new physical law.

Verdict: structural PASS. The absolute Kelvin law $T \rightarrow m_F$ is a follow-up research problem.

Chapter 46 — Stage 0.5 — Residue, Return, and SOURCE Address Branching

46.1 Separating two kinds of filters

Stages 0.3-0.4 concern carrier/channel selection. SOURCE admission/return, which generates the present-universe address composition, is a separate operation. The two operations are therefore kept distinct:

- Filter A: carrier/channel selection
- Filter B: SOURCE address admission/return

This distinction becomes important when the component layer is inserted. Selecting a channel does not automatically imply that a source address has been admitted into phenotype.

46.2 Address weighting and branching rule

Let addresses run over $n = 2, \dots, N$ with weights

$$w_n \propto n^{-\alpha}.$$

The baseline calculation uses

$$N = 10^6, \quad \alpha_0 = 1.8996876950554356, \quad \beta_0 = 0.8654570124136961.$$

Classifying the weighted sum gives the odd-composite weight

$$O = 0.05777294456,$$

the even-composite weight

$$E = 0.268,$$

and the prime weight

$$P = 0.6742270554.$$

If a fraction β of the odd-composite weight is admitted to phenotype, then

$$\phi = \beta O = 0.05,$$

while the rejected odd-composite weight is

$$O - \phi = 0.00777294456.$$

Return is therefore not a third independently fitted quantity. It follows from completeness:

$$R = P + (O - \phi) = 0.682.$$

The resident Actual share is

$$\phi + D = 0.05 + 0.268 = 0.318.$$

46.3 The 31.8% share is not an automatic recycling fixed point

A separate test asks whether 31.8% becomes an automatic equilibrium when the SOURCE repeatedly releases the remaining stock. Under a representative repeated-release control, a release fraction of 0.2 applied over 32 cycles moves the cumulative branching to about 87.79%. If release is allowed indefinitely in the mathematical iteration, the cumulative release approaches 100%.

Thus the present 31.8% Actual fraction is not an automatic fixed point of unlimited recycling. Maintaining a specific cosmic branching ratio therefore requires

a finite generation window

or another explicit balance closure.

46.4 Verdict

Address branching and completeness close at this stage. How α, β , the admission shutter, and the generation window are set by an actual cosmic thermal history is a separate generation-law problem.

Verdict: SOURCE address branching PASS.

Chapter 47 — Stage 0.6 — Shutter, Event Discreteness, and Physical Time

47.1 Meaning of a frame

A WRR frame is a finite execution unit ordered as

preparation $\rightarrow$ transformation $\rightarrow$ shutter $\rightarrow$ record.

Here, “minimum” means a readout event not further subdivided within the execution contract, not a universal minimum physical time shared by the whole universe.

Four objects are kept distinct:

1. SOURCE frame index k ,
2. readout shutter,
3. worldline proper time τ ,
4. Planck time t_P .

47.2 Inherited SI clock bridge and numerical scale

For the inherited electron mode,

$$23\mu_E = 22217.345682174 \text{ eV}$$

corresponds to the characteristic time

$$t_\mu = 2.962603932832 \times 10^{-20} \text{ s.}$$

Applying the conditional resolution $\epsilon = 0.05$ gives

$$\Delta\tau_* = 1.481301966416 \times 10^{-21} \text{ s.}$$

Relative to Planck time,

$$\frac{\Delta\tau_*}{t_P} \approx 2.75 \times 10^{22}.$$

Under the separate SOURCE condition $\xi = 0.1$, the time interval used is

$$\Delta\tau_{\text{source}} = 2.962603932832 \times 10^{-21} \text{ s.}$$

47.3 The same frame unitary with different physical times

If only the dimensionless frame transformation is fixed, one may increase the generator rate and shorten the frame duration, or decrease the rate and lengthen the duration, while preserving the same unitary transformation. The frame index therefore does not determine a unique SI time interval.

Hence

$$\text{event discreteness} \neq \text{physical-time discreteness}$$

in the current implementation.

The baseline choice $K = 8$ is an execution input, not a derivation that the universe possesses a universal shutter count of eight.

47.4 Verdict

Finite shutter/readout event: PASS.

Identifying a WRRRA frame with Planck time is not required and is not supported by the present calculations. This separation motivates stage 0.7, where the micro-macro boundary is reformulated as a record-structure problem rather than as a universal time constant.

Chapter 48 — Stage 0.7 — Boundary Between Microscopic Uncertainty and Macroscopic Record

v0.12 chapter update

P7 needs two states for its specified present response but sixteen for full history reconstruction. Record stability, response sufficiency, reconstruction and unique-outcome generation are different tasks. Finite saturation constrains subsequent policies without selecting an outcome or cosmic restart. The repeated-shutter example below is not such a selection law. See Chapter 98.

48.1 Two boundaries instead of one threshold

If the microscopic-macroscopic distinction is defined by one absolute number, it cannot naturally reflect the fact that the boundary depends on which observable is read, at what resolution, and with what tolerance. This study therefore uses two boundaries.

Distinguishability boundary

B_D : Can different states be distinguished by different record projectors?

Record-stability boundary

$B_R(\delta)$: Is instability under repeated readout below the tolerance δ ?

The micro-macro interface is then defined as

$$B_{\mu M} = B_D \cap B_R(\delta)$$

rather than by one universal constant.

48.2 Address 9/15 and coarse/fine readout

The inherited 9/15 state pair is distinguishable under fine readout, while a coarse bin can place both addresses in the same record class. The same physical state information may therefore remain as internal coherence or be separated into distinct records depending on the readout partition.

The readout scaling q is kept distinct from the projector itself. Numerical scaling of the displayed readout must not be confused with the partition structure that defines which states are distinguishable.

48.3 Repeated-shutter calculation

Let K be the number of fine shutter events over a fixed total evolution interval. The transition probability is

$$a_K = \frac{1 - \cos^K(2gs/K)}{2}.$$

For the baseline $g = 1$ and $s = 1.2$, the values are:

K	a_K
1	86.869686%
2	43.434843%
4	26.799767%
8	15.308671%
16	8.264840%
32	4.307302%
64	2.200630%
128	1.112503%
256	0.559356%
512	0.280461%
1024	0.140428%

For large K ,

$$a_K \sim \frac{1.44}{K}.$$

The transition probability therefore decreases continuously as repeated fine readout increases.

48.4 Record-stability boundary as a function of tolerance

If record instability is required to stay below a tolerance δ , the approximate minimum K shifts as follows.

tolerance δ	approximate minimum K
0.1	13
0.05	28
0.01	143
0.001	1,439
0.0001	14,399

The micro-macro boundary is therefore not best represented as one universal number. It is a family of boundaries determined by the distinction being demanded and the stability tolerance being imposed.

48.5 Separating decoherence from a unique record

Even when off-diagonal coherence disappears under a three-branch coarse shutter, the calculated purity remains

$$\mathcal{P} = 0.539448,$$

and the diagonal population remains

$$(0.05, 0.268, 0.682).$$

The disappearance of coherence therefore does not automatically imply that one unique record has been selected.

decoherence $\neq$ single selected record

This distinction leads directly to the stage-0.8 requirement that a component must preserve origin and ledger identity rather than merely correspond to a decohered branch.

Verdict: micro/macro record interface PASS.

Chapter 49 — Stage 0.8 — Component Activation Boundary and Primitive Alphabet

49.1 Why a component layer is needed before particle phenotype

If the upstream model moves directly into a final particle phenotype, there is no layer in which to represent the assembly order, origin identity, inactive reserve, and repeated-composition differences found in the Prime Parts studies. A component layer is therefore inserted before particle phenotype.

A component is defined here by four jointly required properties:

$$\text{Component} = \text{origin} + \text{activation} + \text{identity retention} + \text{ledger closure}$$

49.2 Component boundary

The structural component boundary can be written as

$$B_C = B_D \cap B_{\text{origin}} \cap B_{\text{ledger}}.$$

When the component must persist into a stable phenotype, repeated-record stability is also required:

$$B_C^{\text{persistent}} = B_C \cap B_R(\delta).$$

49.3 Origin-preserving transition

The inherited Prime Parts execution retains orthogonal origin tags through state transitions. For example,

$$j : n \rightarrow j : D$$

changes the current state label while preserving the origin j . In the control in which the origin tag is removed, the isometry test fails. Origin is therefore not merely a descriptive name. It is a degree of freedom required to preserve the history carried by the component.

49.4 Inactive reserve

A component that is not currently expressed in the phenotype is not treated as having disappeared.

$$\text{inactive reserve} = \text{origin preserved, phenotype inactive}$$

This separates the currently expressed phenotype from the larger pre-phenotype component inventory.

49.5 What Address 105 shows

Address 105 demonstrates that the same source address can produce different numbers of active component occurrences depending on assembly order.

assembly order	active component occurrences
small-first	11
big-first	4

One therefore cannot conclude directly that there are four fundamental component types merely because one assembly path produces four active occurrences. The quantity four in this case is an active occurrence count associated with one assembly path.

Thus

$$N_{\text{active}} \neq N_{\text{primitive}}$$

in general.

49.6 Primitive hierarchy

The minimum hierarchy is written as

$$A_{N_P} = \{b_1, \dots, b_{N_P}\},$$

for the primitive alphabet,

$$C_j = G(\text{sequence, order, phase, repetition}),$$

for the origin/code layer, and

$$P = \mathcal{R}(C_1, C_2, \dots)$$

for the final particle phenotype.

Within this hierarchy, prime labels may be used as address rules at the component/origin layer, but primality itself is not automatically asserted to be the fundamental primitive alphabet.

49.7 Exact status of the four-primitive candidate

Under the additional assumption of a code with two independent positions,

$$4^2 = 16$$

is an interesting candidate relation. However, the same 16-state capacity can also be generated by another grammar such as 2^4 , while a three-symbol alphabet with code length three has capacity $3^3 > 16$. The present data therefore do not uniquely fix $N_P = 4$.

Likewise, the inherited common-carrier inventory contains 15 base channels and permits a conditional 16th neutral extension. That statement is not equivalent to the claim that the universe fundamentally contains 16 primitive component types.

Verdict: Component Activation Boundary PASS. The size of the primitive alphabet remains a follow-up output target.

Chapter 50 — Stage 0.9 — Cooling, Stable Codewords, and Harmonic States

50.1 Cooling suppresses reverse transitions rather than creating components from nothing

After stage 0.8 defines the existence conditions for a component, stage 0.9 interprets the role of cooling not as “creating components from nothing” but as suppressing reverse transitions and stabilizing allowed assemblies.

If the reverse barrier of state j is ΔE_j , one may use the diagnostic relation

$$P_{\text{reverse},j} \sim \exp \left[-\frac{\Delta E_j}{k_B T} \right].$$

For a tolerated reverse probability δ , define the state-dependent stability scale

$$T_j(\delta) = \frac{\Delta E_j}{k_B \ln(1/\delta)}.$$

This is not a prediction of the actual cosmic temperature history. It is a diagnostic showing how much thermal suppression is required for a given barrier and tolerance.

50.2 Examples of state-dependent barriers

Inherited reference barriers already span substantially different scales, for example the deuteron scale 2.22588559 MeV and the $\alpha \rightarrow d + d$ breakup scale 23.92275894 MeV. This supports the use of state-dependent stability boundaries rather than the assumption that every component locks simultaneously at one universal T_C .

50.3 Distinguishing stable codewords from harmonic states

A stable codeword is defined as an assembly satisfying three conditions:

1. its origin is distinguishable,
2. reverse transitions are sufficiently suppressed,
3. its assembly identity is maintained under repeated record formation.

A harmonic state, by contrast, contains multiple eigenmodes while preserving relative-phase information. Such a state is more naturally represented by a multi-mode phase ledger than by one fixed component codeword.

50.4 Phase ledger of the neutrino 0 : 5 : 29 structure

The existing WRRRA neutrino candidate is retained:

$$(n_1, n_2, n_3) = (0, 5, 29).$$

The basic mass quantum is

$$q_\nu = 1.725929280 \text{ meV}.$$

Therefore

$$\begin{aligned} m_1 &= 0, \\ m_2 &= 5q_\nu = 8.6296464 \text{ meV}, \end{aligned}$$

and

$$m_3 = 29q_\nu = 50.0519491 \text{ meV}.$$

Their sum is

$$\sum m_\nu = 58.6815955 \text{ meV} \approx 58.6816 \text{ meV}.$$

The common rest-phase recurrence is

$$T_\Phi \approx 2.39619 \times 10^{-12} \text{ s},$$

and over this interval the three modes accumulate respectively

$$0, \quad 5, \quad 29$$

integer phase cycles. Thus 0 : 5 : 29 is not merely a list of three numbers; it is an integer phase ledger defined on a common recurrence interval.

50.5 What is claimed and what is not yet claimed

This result provides an internal basis for representing the neutrino state harmonically. It does not identify the rest-phase recurrence directly with an observable flavor-oscillation period. Nor does it claim that the microscopic origin of the PMNS mixing matrix has already been generated from primitive grammar.

A diagnostic ratio such as

$$\Lambda = \frac{\Gamma_R}{\Delta f_{ij}}$$

may be useful for characterizing the boundary between codeword locking and harmonic coherence, but it is not fixed here as a universal law.

Verdict: stable-codeword/harmonic interface PASS.

Chapter 51 — Stage 0.10 — Reconnecting Phenotype Energy to Downstream Gravity

51.1 The central conservation rule

After adding the component and harmonic layers, the first required test is energy nonduplication. Finer internal structure must not increase the total cosmic energy merely because more internal labels exist.

The conserved normalization rule is therefore

$$\sum_{g,j} E_{\phi,g,j} = E_{\phi}$$

where g may label generation and j a component/channel slot. The inherited 48-slot routing uses normalized weights, and its sum returns to the existing E_{ϕ} .

51.2 Reserve and harmonic phase are not new cosmic sectors

Inactive reserve is component inventory not currently expressed in phenotype. It is not a fourth cosmic fraction added to ordinary matter, resident nonphenotype, and return. Likewise, a harmonic phase register increases internal state information but does not automatically increase population or total energy.

Therefore

more components/channels/generations/phases $\nRightarrow$ more cosmic energy.

51.3 Separating internal rearrangement from real population change

If only labels or phases are rearranged within the same E_{ϕ} , the baseline q_0 , rotation, and lensing outputs should remain unchanged. If the SOURCE generation rule actually changes the phenotype population, however, the downstream pressure and load should change and q should respond.

To test this, the generation control is varied while the SI coefficients are kept frozen.

upstream control	phenotype fraction	downstream q
$K = 4$	3.704794619%	-0.547908522
$K = 8$	5.000000000%	-0.528550000
$K = 16$	5.682473858%	-0.518343025

These values are not obtained by refitting the coefficients to a target output. The upstream population moves, passes through the same frozen downstream mapping, and produces a corresponding change in q .

The sensitivity chain is therefore

upstream population change $\rightarrow$ same $E(V) \rightarrow P \rightarrow$ load $\rightarrow q$

and exists explicitly within the model.

51.4 Relation to inherited local and global outputs

At the baseline state $K = 8$, the frozen outputs remain

$$q_0 = -0.52855,$$
$$v_{\text{rot}} = 207.5109051266 \text{ km s}^{-1},$$

and

$$\theta_{\text{lens}} = 0.5355865106''.$$

Internal relabeling or phase rearrangement alone should not move these values. A genuine change in population or in the energy ledger, by contrast, should propagate through the same downstream rule.

Verdict: upstream-downstream structural reconnection PASS.

Chapter 52 — Stage 0.11 — Reverse Minimality Audit

52.1 Reversing the question

Stages 0.1-0.10 proceed from high-energy conditions toward phenotype and downstream physics. Stage 0.11 reverses the direction and asks which ingredients cannot be removed while preserving the current calculations.

The minimum interface required by the present executable structure is summarized as

$$G_{\min} = \{A_{N_P}, O, B_L, B_R, R, \Phi, L\}$$

with the following roles.

Symbol	Meaning	Failure when removed
A_{N_P}	primitive alphabet	minimum symbol space for assembly disappears
O	origin identity	component history/isometry breaks
B_L, B_R	independent filter distinctions	independent filter discrimination disappears
R	assembly/termination rule	the assembly result for a given address is not determined
Φ	phase register	harmonic multi-mode representation disappears
L	conservation/normalization ledger	energy and population may be duplicated

52.2 Origin is required

Without origin, different histories cannot be distinguished by the current state label alone, and the inherited isometry control fails. Preserving component identity therefore requires O .

52.3 Assembly order and termination rule are required

Address 105 produces different active component profiles under different assembly orders. An alphabet alone is therefore insufficient to determine the result. At minimum, order and a termination rule are also required.

52.4 Harmonic interpretation requires a phase register

To preserve relative-phase structure in a multi-mode state such as the neutrino $0 : 5 : 29$ configuration, Φ cannot be removed. The need for a phase register, however, does not by itself determine the primitive count.

52.5 What is not a required assumption

Removing the assumption $N_P = 4$ does not destroy the current component boundary, codeword/harmonic interface, energy normalization, or downstream sensitivity. Four primitive types are therefore not yet a necessary premise.

Likewise, removing the identification of the WRRR frame with Planck time leaves the executed results intact. Planck-time identification is not part of the minimum structure.

The three neutrino modes may imply a spectral capacity of at least three, but spectral-mode capacity is not identical to the number of primitive component types. The two counts remain distinct.

Verdict: reverse/minimality constraints PASS.

Chapter 53 — Stage 0.12 — Closing the Master Ledger

53.1 One typed chain

The results obtained in stages 0.1-0.11 are finally placed into the following chain:

high-energy Actual → finite thermal probe
 → filter/readout
 → SOURCE admission/return
 → shutter + conditional record
 → $B_D \cap B_R(\delta)$
 → origin-preserving component
 → stable codeword or harmonic state
 → phenotype
 → normalized SI energy
 → pressure/information load
 → twist/gravity/expansion.

53.2 Closed interfaces

The interfaces judged closed at stage 0.12 are listed below.

No.	Interface	What is closed
1	finite thermal probe	finite thermal state defined by T, H, E_0
2	SOURCE branching	phenotype/resident/return completeness
3	shutter/readout	finite conditional record event
4	micro/macro	record boundary $B_D \cap B_R(\delta)$
5	component	origin-preserving activation/ reserve accounting
6	codeword/harmonic	distinction between stable assembly and multi-mode phase
7	phenotype -> SI energy	normalized allocation without energy duplication

No.	Interface	What is closed
8	energy -> pressure	pressure calculated from the same $E(V)$
9	upstream -> q	downstream propagation of a genuine population perturbation
10	local gravity	retained conditional rotation/lensing outputs
11	homogeneous expansion	macroscopic output of the frozen energy-pressure ledger

53.3 What stage 0.12 closes is the research scope

Closure at stage 0.12 does not mean that every microscopic origin in the universe has already been derived. The question closed by this study is narrower and explicit:

Can high-energy states be connected through records, components, and phenotypes to the existing WRRAM energy-gravity ledger as one executable structure without mixing types, duplicating energy, or retuning the frozen downstream model?

The answer produced by stages 0.1-0.12 is yes.

The final verdict is therefore

WRRAM 0.1-0.12 Structural Integration 1.0 - CLOSED
--

Chapter 54 — Structures Newly Secured by the Integrated Study

54.1 A component layer now exists between upstream generation and downstream physics

The earlier simplified flow could be read as

carrier/filter $\rightarrow$ phenotype $\rightarrow$ downstream physics.

After the present study, the structure becomes

carrier/filter $\rightarrow$ record $\rightarrow$ component $\rightarrow$ codeword/harmonic $\rightarrow$ phenotype $\rightarrow$ downstream physics

The detailed Prime Parts assemblers have not all been absorbed into the WRRRA M core. Nevertheless, the origin-preserving component ontology identified in Prime Parts now functions as a formal bridge between upstream generation and downstream phenotype physics.

54.2 Negative results narrowed the structure

Two important hypotheses were not adopted.

First,

high $T \Rightarrow$ population mixing $\Rightarrow$ filter collapse

does not hold under the current frozen rule.

Second,

WRRRA frame $= t_P$

is not required by the current calculation.

These negative results do not weaken the research. They more precisely locate where filter dynamics and clock dynamics must enter the architecture.

54.3 The micro-macro boundary is recast as a record problem

Rather than assigning the microscopic-macroscopic transition to one length, time, or mass constant, the study places it at

$$B_D \cap B_R(\delta).$$

This incorporates directly into the structure the fact that the boundary can move with measurement resolution and tolerance.

54.4 Harmonic structure is a phase ledger rather than a metaphor

The neutrino $0 : 5 : 29$ structure is organized not merely as a musical analogy but as integer cycles on a common rest-phase recurrence. “Harmonic” therefore has a concrete phase-bookkeeping meaning inside the model.

54.5 Growth of internal structure is separated from growth of cosmic energy

Even when the numbers of components, channels, generations, or harmonic modes increase,

$$\sum E_{\phi,g,j} = E_{\phi}$$

is preserved. This rule prevents increasing internal state-space complexity from being misread as an automatic increase in the cosmic energy budget.

Chapter 55 — Integrated Falsification Conditions

This study closes a structural chain while also stating the conditions under which a connection must be revised or rejected.

Category	Falsification condition
probability/state conservation filter	failure of trace, positivity, or branch completeness failure of $Q/B/L/E$ or of declared normalization the same unique filter remains selected after contrast is removed
component	identical component history and isometry remain fully preserved after origin is removed
record	increasing K produces a trend opposite to the declared stability behavior
energy	total cosmic energy increases merely with component/channel/generation/harmonic count
bridge	a genuine upstream population change produces no response in frozen downstream output
clock	proper-time and phase units contradict one another within the same ledger
observation	future observations under matched preparation significantly disagree with a conditional output

A crucial distinction is maintained between an unresolved follow-up question and a falsification of a completed result. For example, the fact that a physical particle Hamiltonian has not yet been generated from primitive grammar is not a falsification of the stage-0.8 component boundary. Falsification occurs when the declared boundary violates its own input-output contract.

Chapter 56 — Research Questions After Stage 0.12

v0.12 chapter update

Several original tasks now have bounded executable examples: P2 conditional readout, P3 daughter-cohort dynamics, P4 a common effective pulse and P5 record-aware compensation. It is no longer correct to say that no interaction candidate or nonequilibrium extension exists. The full particle spectrum, a derivation of weak interactions, autonomous control and covariant spacetime remain separate tasks. The entries below preserve the original questions; Chapters 93–99 give their updated assessment.

Once stages 0.1-0.12 are closed, four research axes remain. They are not a failure list for the present work; they are independent next-stage problems demanded by the closed structure.

56.1 From temperature to generation controls

There is not yet a physical generation law

$$T \rightarrow \{\kappa, \alpha, \beta, K_*\}.$$

Deriving such a law would directly connect thermal history to filter contrast, address admission, and the generation window.

56.2 From primitive grammar to a particle Hamiltonian

The component and assembly grammar are structurally defined, but a separate study is still required for

$$\text{primitive grammar} \rightarrow H_{\text{particle}}.$$

Closing this adapter would complete, at executable-code level, the connection between Prime Parts assembly structure and the inherited internal Hamiltonian.

56.3 From 0 : 5 : 29 to flavor dynamics

An integer phase ledger currently exists. The next question is how that phase structure appears as an observable flavor-mixing dynamics through an interaction/readout grammar:

$$0 : 5 : 29 \rightarrow \text{mixing dynamics}.$$

56.4 Common SI geometry for homogeneous background and nonuniform local states

Homogeneous expansion and conditional local gravity are already connected in the same broad ledger, but an executable model that evolves background and nonuniform states simultaneously in one fully covariant SI geometry remains a separate research task.

Chapter 57 — Position of This Integrated Study within MCC 3.0

The 0.1-0.12 integration does more than add one new number to MCC 2.3.2. It fixes a new intermediate layer in the generation flow of the universe and therefore provides structural groundwork for version 3.0.

If the compact 2.3.2 flow is written as

common carrier → filter → phenotype → forces/cosmology,

the present architecture is

SOURCE/Actual → filter → record → component → codeword/harmonic → phenotype → energy/load → gravity/expansion

The next major MCC revision should therefore not attach this research merely as an appendix. It is more natural to rewrite the record/component layer as a formal part of the core architecture following the common carrier.

Chapter 58 — Conclusion of the High-Energy-to-Phenotype Connection

This study fills the missing intermediate structure between the high-energy upstream side of WRRM and the established phenotype-to-downstream physics through a staged 0.1-0.12 examination.

Stage 0.1 freezes the inherited cosmic ledgers and forbids retuning. Stage 0.2 converts temperature into a finite thermal probe, and stage 0.3 rejects the hypothesis that thermal population mixing alone removes the filter gap. Stage 0.4 constructs a filter-stabilization structure through a contrast order coordinate. Stage 0.5 separates channel selection from SOURCE branching, closes the 5%/26.8%/68.2% address ledger through completeness, and shows that a finite generation window is required rather than unlimited recycling.

Stage 0.6 retains the frame as a record event while separating it from a universal minimum physical time. Stage 0.7 defines the micro-macro boundary as $B_D \cap B_R(\delta)$ and quantifies the record-stability crossover through repeated-shutter calculations. Stage 0.8 introduces the origin-preserving component as a formal bridge, connecting Prime Parts to the upstream-downstream integration. Stage 0.9 distinguishes stable assemblies from harmonic states and organizes the neutrino $0 : 5 : 29$ structure as a concrete integer phase ledger.

Stage 0.10 preserves normalized phenotype energy so that richer internal structure does not generate extra cosmic energy, while confirming that a genuine upstream population change propagates into the frozen downstream q . Stage 0.11 identifies the minimum interface required by the current execution through a reverse audit. Stage 0.12 integrates all of these connections into one Master Ledger.

The conclusion of the study is therefore summarized by one statement:

WRRM 0.1-0.12 Structural Integration 1.0 - CLOSED

Here, CLOSED does not mean that there are no further research questions. It means that the starting points of the next questions are now explicit. The subsequent research axes are the law generating filter/admission/generation controls from temperature, the law generating a particle Hamiltonian from primitive grammar, the law connecting the $0 : 5 : 29$ phase ledger to flavor mixing, and a common covariant SI geometry for homogeneous and nonuniform states.

A central achievement of the integration is that these next-stage questions are not mixed into the completed 0.1-0.12 results. They are separated as the next interfaces of an already closed structural chain.

Appendix A. Stage-by-Stage Verdict Ledger

Stage	Core question	Main output	Verdict
0.1	What must be frozen?	address/energy/ downstream baseline	PASS
0.2	How is temperature entered into a state?	finite thermal probe	PASS
0.3	Does high- T mixing erase the filter?	gap remains positive	NEGATIVE RESULT
0.4	How are cooling and filter selection connected?	$m_F \rightarrow \kappa \rightarrow \Delta$	PASS
0.5	How do addresses branch?	$\phi/D/R =$ 0.05/0.268/0.682	PASS
0.6	Is a frame a minimum physical time?	event/physical-time separation	PASS
0.7	What is the micro/ macro boundary?	$B_D \cap B_R(\delta)$	PASS
0.8	What is a component?	origin-preserving component	PASS
0.9	How are stable states represented?	codeword/harmonic	PASS
0.10	How is the component layer reconnected downstream?	normalized energy + q sensitivity	PASS
0.11	What is minimally required?	$G_{\min}$	PASS
0.12	Does the whole chain close in one ledger?	Master Ledger	CLOSED

Appendix B. Core Numerical Ledger

Quantity	Value
uncalibrated phenotype residue	4.876893353%
address phenotype	5.0%
address resident nonphenotype	26.8%
address return	68.2%
inherited energy phenotype	4.93%
inherited energy resident	26.5%
inherited energy return	68.57%
internal gap ΔE	431.081244315 MeV
conditional excitation contribution	34.4764357391 MeV
filter gap $\Delta_L = \Delta_R$	0.04669193039
SOURCE α_0	1.8996876950554356
SOURCE β_0	0.8654570124136961
odd-composite weight O	0.05777294456
even-composite weight E	0.268
prime weight P	0.6742270554
proper-time step $\Delta\tau_*$	$1.481301966416 \times 10^{-21}$ s
neutrino quantum q_ν	1.725929280 meV
neutrino mass sum	58.6816 meV
common rest-phase recurrence	2.39619×10^{-12} s
baseline q_0	-0.52855
reference rotation	207.5109051266 km/s
conditional lensing	0.5355865106 arcsec

Appendix C. Reproducibility and Public Records

This study is written on the public provenance chain of the following WRRM records.

- WRRM M Integrated Upstream and Downstream Model 1.0 r1 - DOI: 10.5281/zenodo.23126800
- WRRM M Integrated Measurement Case 1.0 - DOI: 10.5281/zenodo.23149260
- WRRM Prime Parts 0.1-0.12 Reviewed Collection - DOI: 10.5281/zenodo.23134716
- WRRM Address 105 Prime Parts Case Study - DOI: 10.5281/zenodo.23137504
- WRRM M High-Energy-to-Phenotype Extension 1.0 - DOI: 10.5281/zenodo.23176529

GitHub integration path:

https://github.com/Wonsik-Choi-janefather/wrrm-m-0.1/tree/main/integrated/high_energy_extension_v1_0

This detailed integrated research manuscript expands the stage-by-stage calculations, negative results, intermediate ledgers, and connection logic of the existing 1.0 release. The original provenance of the sole-author Prime Parts public releases and the coauthored WRRM M public releases is preserved.

Part XI — Downstream Execution in 3.0: From Information Load to Cosmic Expansion

Part X closes the intermediate generation layer. Part XI checks how that result descends, in the finite-SOURCE branch of 3.0, into the inherited WRRM_M calculations of SI energy, pressure, gravity, galaxies, lensing, and cosmic expansion.

Chapter 59 — Information Load and Energy

v0.12 chapter update

Matching total energy and matching its volume derivative are separate conditions. P2 requires sectorwise constancy of energy and derivative for recovery across all preparations. A P5 compensator must contribute its energy and records to the total. The inherited SI numbers below belong to the original macro allocation and do not automatically apply to a new hot-particle readout. See Chapters 93 and 96.

The cosmic additive ledger contains only

$$\boxed{\phi \oplus D \oplus R}.$$

Address response is

$$g_s(n) = 1 + \lambda_s \frac{\log n}{\log N_U}, \quad (\lambda_\phi, \lambda_D, \lambda_R) = (0, 0.25, 0.1).$$

The finite-source address moments are

$$\boxed{(\mu_\phi, \mu_D, \mu_R) = (0.05000000000000001, 0.278925610976744, 0.687742539854239)}.$$

The frozen SI coefficients are

$$\eta_\phi = 7.561582499072265 \times 10^{-10}, \quad \eta_D = 7.285761328795971 \times 10^{-10}, \quad \eta_R = 7.646102818811323 \times 10^{-10} \text{ J m}^{-3}.$$

At $V_0 = 1 \text{ m}^3$,

$$E_\phi = 3.78079124953614 \times 10^{-11} \text{ J},$$

$$E_D = 2.03218543006515 \times 10^{-10} \text{ J},$$

$$E_R = 5.25855017259595 \times 10^{-10} \text{ J},$$

$$\boxed{E_{total} = 7.668814727614716 \times 10^{-10} \text{ J}}.$$

Prime-part and 48-channel descriptions are nested nonadditive views of the phenotype budget. Each view separately sums back to E_ϕ ; alternative views must never be added as new cosmic energy.

For dimensional clarity, $E_{s,0} = V_0 \eta_s \mu_s$, with $V_0 = 1 \text{ m}^3$. The quoted E_s values are initial energies in J. Matter-sector densities divide by $V_0 a^3$; the return energy is $E_R(a) = E_{R,0} a^3$, so its density is constant. Component energies obey $E_{\phi,j} = V_0 \eta_\phi \mu_{\phi,j}$.

Chapter 60 — Gravity and Twist

v0.12 chapter update

P6 calculates different susceptibilities for the same mean load and inherited snapshot when fluctuations or relaxation constraints differ. The local gravity law below is a frozen macro-response branch, not identical to the new finite equilibrium response. Combining them requires spatial mapping, coupling, bath and dimensional anchors. Chapter 97 defines the response meaning of information weight.

The precursor finite-boundaryless work interpreted accumulated spatial twist as a possible owner of hidden gravitational stress [1]. Later WRRRA twist-stress work connected that owner to a dark-mass phenotype and a low-acceleration gravitational response [5].

MCC 3.0 does not add a second fundamental force. The resident D-sector is mapped through the already frozen local response. With inherited reference $f_c = 0.265$,

$$a_{T,0} = cH_0 \sqrt{\frac{f_c}{8}},$$

and the candidate state uses

$$a_T = a_{T,0} \sqrt{\frac{\rho_D}{u_{crit} f_c}}.$$

At $a = 1$,

$$a_T = 1.1917875313971119 \times 10^{-10} \text{ m s}^{-2}.$$

The same a_T is used for both rotation and conditional lensing.

Chapter 61 — Galaxy Rotation and Lensing

The frozen test source has mass 6×10^{10} solar masses, Plummer scale 3 kpc, test radius 8.2 kpc, finite lens patch radius 200 kpc and impact parameter 10 kpc.

The direct baryonic reference rotation remains $161.4505105 \text{ km s}^{-1}$. The finite-source total reference rotation is

$$207.5102419 \text{ km s}^{-1}.$$

The same local gravity response yields conditional finite-patch lensing

$$0.5355822400''.$$

Relative shifts from the legacy one-million-address branch are approximately 3.20×10^{-6} for rotation and 7.97×10^{-6} for lensing. These are model continuity outputs, not new astronomical measurements.

Chapter 62 — Cosmic Expansion

v0.12 chapter update

The dust assignment and expansion table below follow the inherited macro energy law. Hot particles and decay daughters in P2–P3 can have positive pressure at the same energy. This table is therefore not an automatic result of microscopic refinement. Specify cold allocation, accept altered pressure, or supply the explicit P5 compensation. Conditional background closure is still not a complete cosmological perturbation theory.

The inherited volume exponents are $(n_\phi, n_D, n_R) = (0, 0, 3)$. Thus

$$\rho_\phi(a) = E_{\phi,0}/(V_0 a^3), \quad \rho_D(a) = E_{D,0}/(V_0 a^3), \quad \rho_R(a) = E_{R,0}/V_0.$$

Pressure is $P_\phi = P_D = 0, P_R = -\rho_R$.

The homogeneous diagnostics are

$$\frac{H(a)}{H_0} = \sqrt{\frac{\rho(a)}{u_{crit}}}, \quad q(a) = \frac{1}{2} \frac{\rho(a) + 3P(a)}{\rho(a)}.$$

a	density (J/m ³)	H/H0	q
0.5	2.4540666613e-9	1.7888556123	0.1785814590
1.0	7.6688147276e-10	0.9999913260	-0.5285585894
2.0	5.5598332420e-10	0.8514575347	-0.9187161585

The uniform-branch acceleration transition is

$$a_{trans} = 0.6119598067968817.$$

Chapter 63 — Proper Time and Record

v0.12 chapter update

P3 anchors one decay episode to a supplied muon lifetime; it does not derive the proper-time interval below. P7 calculates finite record accumulation and overflow, advancing beyond an entirely open record question. Autonomous measurement apparatus and a cosmic post-overflow law remain separate.

The inherited proper-time step is

$$\Delta\tau = 1.4813019664158691 \times 10^{-21} s.$$

MCC 3.0 does not identify this with Planck time or an absolute minimum physical time.

For the candidate finite carrier, noncommuting internal loads evolve while total energy remains conserved to a maximum relative drift of approximately

$$\boxed{9.99 \times 10^{-16}}.$$

State trace remains unity within numerical precision. The framework distinguishes state update and ledger conservation from a full autonomous theory of irreversible physical record formation. Physical apparatus, reset and persistent-record dynamics remain open where not explicitly executed.

Chapter 64 — End-to-End Master Ledger

v0.12 chapter update

Each arrow requires retained state information. Totals may fail to determine pressure, and current energy and pressure may fail to determine the next state. P4→P3 and P5 cohort-aware compensation are executed compositions. P6 equilibrium response and P7 record experiments have not thereby become one autonomous universe simulator. Read the connection table in Chapter 99.

The integrated chain is

finite rule → finite addresses → filter → residue → component/phenotype
→ information load → energy → pressure/gravity → rotation/lensing → expansion

Only the ϕ , D , R sector energies are additive. Component and channel descriptions are internal alternative decompositions of E_ϕ , not new resources. This ownership rule prevents explaining the same physical content twice and then counting both descriptions as additional energy.

Chapter 65 — Reproduction of Verified Values

v0.12 chapter update

The values below preserve the earlier SI-load branch. The r11 value -0.523 uses a different sector-energy allocation. Do not treat them as conflicting outputs of identical inputs or replace one by the other. Check counts are internal audits of each study, not additive counts of independent observations. Chapter 92 explains the distinction.

The frozen legacy branch reproduces $(0.05, 0.268, 0.682)$ and the reference outputs

$$q_0 = -0.52855, \quad v = 207.5109051266 \text{ km s}^{-1}, \quad \theta = 0.5355865106''.$$

The finite-source branch gives

$$q_0 = -0.5285585894, \quad v = 207.5102418659 \text{ km s}^{-1}, \quad \theta = 0.5355822400''.$$

The repository contains 54 declared Stage 1–5 checks and 11 Stage-6 cross-stage checks, for a total of 65 passing mathematical, implementation and provenance checks. These are not 65 independent physical experiments.

Chapter 66 — WRRRA/MCC Conditional Predictions

The finite source size

$$N_U = 1,015,000$$

is an unmeasured output generated after the declared KZF construction is fixed. Its status is therefore a conditional MCC/WRRRA prediction.

The homogeneous acceleration-transition scale

$$a_{trans} = 0.6119598068$$

is another derived output of the frozen energy/pressure branch.

MCC 2.3.2 also carries the earlier frozen neutrino mass pattern

$$m_1 = 0, \quad m_2 = 8.62965 \text{ meV}, \quad m_3 = 50.05195 \text{ meV},$$
$$\sum m_\nu = 58.6816 \text{ meV}.$$

MCC 3.0 inherits $H = 29$ from that prior ledger but does not claim to have independently rederived the neutrino masses in the present six-stage integration.

Chapter 67 — Open Assumptions and Falsification Conditions

The following remain open:

1. necessity of $T = 2\pi H$;
2. Cartesian independence of L, H, C ;
3. uniqueness of $N_U = LHC$;
4. physical necessity of the Klein-type quotient;
5. microscopic physical status of the zeta-focus rule;
6. unique Prime-Parts to particle-species mapping;
7. full spin/flavor/confinement/binding Hamiltonian;
8. baryon/lepton origin;
9. fully covariant inhomogeneous completion;
10. autonomous physical measurement and persistent-record dynamics.

The integrated model must be revised if the finite source fails its frozen checks, a hidden third arithmetic fit or new SI refit becomes necessary, internal component views are double-counted as energy, rotation and lensing require different tuned gravity responses, proper-time conservation fails, cross-stage handoffs disagree, or an open microscopic rule becomes necessary and cannot be consistently supplied.

Undefined calculations remain OPEN rather than being counted as successes.

67.1 Where MCC 3.0 Converges with Standard Physics

MCC 3.0 does not aim to produce a different number merely because the Standard Model, general relativity, or FLRW already calculate a domain precisely. When the same inputs, boundary conditions, and Reality-layer calculation modules are adopted, convergence on the same observed value is the expected result. In such domains, the achievement lies less in a new number than in connecting SOURCE, common carrier, component, phenotype, and gravity under one explicit ownership and execution order.

67.2 Where WRR/MCC May Differ

Independent discriminating power arises only where WRR-specific structure creates a different physical dependency from the standard interpretation. The clearest present candidates are boundary-condition-sensitive gravitational response, rotation and lensing owned by the same D-sector, nonspherical stress distributions, global-holonomy correlations, and the path by which a finite-SOURCE perturbation propagates through frozen downstream coefficients into macroscopic outputs.

Candidate discriminants include:

- changes in internal galactic acceleration correlated with external fields or boundary conditions;
- relative displacement between baryons and lensing centers in colliding clusters;
- nonspherical lensing shear tied to one stress state rather than an independently tuned halo;
- repeated correlations or topology-sensitive signatures associated with global holonomy;

- very-long-wavelength suppression or scale dependence in structure growth;
- upstream finite-SOURCE changes that produce a pre-registered response in multiple downstream domains without domain-specific refitting.

67.3 Outputs after input calibration and their assessment

Verified constants and observations may be disclosed construction inputs. Once relevant inputs, calibration freedoms and readouts are fixed, unmeasured quantities computed by the internal WRRRA transformation are conditional predictions. Reproduction of known values counts as consistency and explanatory achievement. Numerical agreement with another theory does not negate an internally executed WRRRA calculation; externally imported results are labelled separately. Effects below measurement uncertainty remain calculated outputs with limited current detectability. Significant disagreement after fixing the model requires revising the relevant assumption or connection. Independent prediction is an additional test route, not a prerequisite for theory or originality.

Part XII — Final Integration of Minimal Computing Cosmology 3.0


Chapter 68 — Preservation Rules and the Meaning of Version 3.0

The major-version change in 3.0 does not consist of deleting earlier achievements. The single execution architecture, common carrier, matter phenotype, twist universe, and standard-physics interfaces organized in 2.3.2 remain in place. What changes is that the finite SOURCE and the record/component layer are added, turning the genealogy from SOURCE to present-day downstream physics into a longer and more explicit typed chain.

Whenever 3.0 reuses an equation or result from an earlier version, it therefore distinguishes among inherited inputs, frozen reference values, and values newly calculated in 3.0. A known input does not invalidate the explanatory achievement of a WRRRA-specific connection; conversely, an input is not relabeled as a new prediction.

Chapter 69 — Reading the Model Again as One Execution Genealogy

The complete execution flow of 3.0 can be summarized as



No layer in this chain replaces the preceding layer. An address is not energy. A component is not automatically a particle species. The number of harmonic modes is not new cosmic energy. Even when one physical resource has several internal representations, the additive ledger owns and sums it only once. This is the nonduplication rule of MCC 3.0.

Chapter 70 — Outputs and Conditional Predictions of Version 3.0

Using $L = 500$, $H = 29$, and $C = 70$ in the finite Klein-Zeta SOURCE candidate gives

$$N_U = LHC = 1,015,000.$$

This is not a measured cosmological constant. It is a conditional MCC/WRRA output once the declared KZF construction is frozen.

With the new finite SOURCE inserted into the address filter, the unretuned ledger remains

$$5.000002539\%/26.800001829\%/68.199995632\%.$$

Re-freezing only the same two previously declared arithmetic freedoms closes the ledger again at $5\%/26.8\%/68.2\%$ to numerical precision.

Without refitting downstream SI response coefficients or the local gravity response, the reference finite-SOURCE branch records

$$\begin{aligned} q_0 &= -0.5285585894, \\ a_T &= 1.1917875314 \times 10^{-10} \text{ m s}^{-2}, \\ v_{ref} &= 207.5102419 \text{ km s}^{-1}, \\ \theta_{lens} &= 0.5355822400'', \end{aligned}$$

and, for the uniform constitutive branch,

$$a_{trans} = 0.6119598068.$$

The inherited neutrino absolute-mass candidate

$$\begin{aligned} m_1 &= 0, & m_2 &= 8.62965 \text{ meV}, & m_3 &= 50.05195 \text{ meV}, \\ \sum m_\nu &= 58.6816 \text{ meV} \end{aligned}$$

was not newly derived in 3.0. Version 3.0 inherits the previously frozen $0 : 5 : 29$ harmonic ledger and uses its largest index $H = 29$ inside the finite-SOURCE generator.

Chapter 71 — A Hierarchy of Falsification

Minimal Computing Cosmology 3.0 is not framed so that the entire program must be either wholly correct or wholly false in one step. Failure conditions are separated by interface so that a failed connection can be identified and revised without silently rewriting every layer.

- finite SOURCE: revise the SOURCE candidate if $T = 2\pi H$, the independence of L, H, C , the product rule $N_U = LHC$, or the physical need for Klein-type closure cannot survive a simpler alternative.
- filter/branching: reject the corresponding connection if probability, normalization, or completeness fails, or if closing the ledger requires new post-hoc freedoms beyond the frozen ones.
- record/component: revise the present interpretation if component history remains fully preserved after origin identity is removed, or if a unique filter remains selected even after contrast is removed.
- particle layer: reject the current decoder hypothesis if one fixed rule cannot reproduce known quantum numbers, or if uniqueness can only be claimed by ignoring persistent order sensitivity.
- energy/gravity: reject the integrated ledger if component, channel, generation, or harmonic counts duplicate energy, or if a genuine upstream population change produces no downstream response under frozen coefficients.
- geometry/cosmology: revise the downstream connection if the same D-sector and energy-pressure ledger cannot enter local and global calculations consistently, or if they cannot be embedded in a fully covariant SI geometry.

Chapter 72 — Final Integrated Propositions of Minimal Computing Cosmology 3.0

1. Our universe need not be assumed to be a privileged optimum; it is treated as one persistent viable execution.
2. Minimal computation is not a rule that makes every quantity as small as possible. It is the nearest sufficient structure that reduces independent inputs, owners, carriers, and update costs while preserving observational viability.
3. MCC does not assume that a physically realized SOURCE must instantiate an infinite address space.
4. The 3.0 finite-SOURCE size 1, 015, 000 is a conditional output under the KZF construction and is not established as a universal constant.
5. The common carrier is not a fifth force. It is a shared Actual transport basis selectively read by multiple renderers.
6. Gravity and the three gauge forces are asymmetrically unified because gravity reads total $T_{\mu\nu}$ while gauge sectors read selective currents J_a ; the forces are not flattened into one identical interaction.
7. Address, component, and SI-energy ledgers are different types and are not directly summed.
8. Before the present particle phenotype, a record/component/codeword-harmonic layer may exist, and component identity preserves origin.
9. Event discreteness is not identified with universal physical-time discreteness.
10. Components, channels, generations, and harmonics may decompose an existing phenotype energy budget, but they do not automatically generate additional cosmic energy.
11. A genuine upstream population change must be able to propagate through frozen downstream coefficients into q and other downstream outputs.
12. Version 3.0 judges the computational integration PASS while preserving the physical necessity of KZF, a unique particle decoder, full spin/flavor/confinement/binding closure, and a fully covariant SI geometry as OPEN.

Chapter 73 — Conclusion: One Universe Executing a Finite Present

v0.12 chapter update

After seven studies, the current conclusion is that the information sufficient for a declared query can be calculated, rather than that every detail is complete. Conditions for preserving totals, pressure, dynamics, equilibrium response and record capacity are connected. An adequate present without perfect reconstruction can coexist with accumulating residue. Part XV assesses the concrete achievements and remaining physical connections.

The center of Minimal Computing Cosmology 3.0 is not the single number 1,015,000. Its core achievement is to extend the single execution genealogy built in 2.3.2 by separating, and then reconnecting, finite SOURCE generation, record formation, component identity, phenotype energy, information load, gravity, and expansion under explicit ownership rules.

This book does not claim that every microscopic origin has already been solved. Calculations that close are marked closed; quantities that depend on construction assumptions are marked conditional; physical generation laws not yet executed remain OPEN. Maintaining these distinctions is what keeps MCC from becoming one untestable grand declaration and instead leaves it as a program that can be inspected, reproduced, modified, and rerun.

The current status of 3.0 is six-stage computational integration PASS / integrated scientific model PASS-C. The next task is not to hide the remaining interfaces, but to execute them with fewer freedoms: from primitive grammar to a physical particle Hamiltonian, from the harmonic ledger to flavor dynamics, and from local and global sectors to one covariant SI geometry.

Appendix A — Core Numerical Ledger of Version 3.0

Quantity	Value
address phenotype	5.0%
address resident nonphenotype	26.8%
address return	68.2%
uncalibrated phenotype residue	4.876893353%
inherited SI-energy phenotype	4.93%
inherited SI-energy resident	26.5%
inherited SI-energy return	68.57%
finite SOURCE L	500
harmonic H	29
zeta count C	70
finite SOURCE N_U	1,015,000
internal gap ΔE	431.081244315 MeV
filter gap $\Delta_L = \Delta_R$	0.04669193039
SOURCE α_0	1.8996876950554356
SOURCE β_0	0.8654570124136961
proper-time step $\Delta\tau_*$	$1.481301966416 \times 10^{-21}$ s
neutrino quantum q_ν	1.725929280 meV
neutrino mass sum	58.6816 meV
common rest-phase recurrence	2.39619×10^{-12} s
MCC 3.0 q_0	-0.5285585894
D-sector acceleration a_T	$1.1917875314 \times 10^{-10}$ m/s ²
reference rotation	207.5102419 km/s
conditional lensing	0.5355822400 arcsec
acceleration transition a_{trans}	0.6119598068

Appendix B — Verification Ledger

The public MCC 3.0 release reports 54 Stage 1-5 computational/implementation checks plus 11 Stage-6 cross-stage checks, for a total of 65 PASS items. These are not 65 independent physical experiments; they are a combined ledger of computation, implementation, provenance, conservation, and cross-stage consistency checks.

- Stage 1: finite SOURCE construction and zero-count consistency.
- Stage 2: finite addresses and prime/composite filter.
- Stage 3: component/phenotype replay and charge/color consistency.
- Stage 4: information load and SI energy-pressure ledger.
- Stage 5: gravity, galaxy rotation, lensing, and homogeneous expansion.
- Stage 6: end-to-end ownership and no-hidden-refit checks across stages.

Appendix C — Public Records and Reproducibility Paths

- Minimal Computing Cosmology 3.0 — DOI: <https://doi.org/10.5281/zenodo.23202732>
- WRRRA M High-Energy-to-Phenotype Extension 1.0 — DOI: <https://doi.org/10.5281/zenodo.23176529>
- WRRRA M Integrated Upstream and Downstream Model 1.0 r1 — DOI: <https://doi.org/10.5281/zenodo.23126800>
- WRRRA M Integrated Measurement Case 1.0 — DOI: <https://doi.org/10.5281/zenodo.23149260>
- WRRRA Prime Parts 0.1-0.12 Reviewed Collection — DOI: <https://doi.org/10.5281/zenodo.23134716>
- WRRRA Address 105 Prime Parts Case Study — DOI: <https://doi.org/10.5281/zenodo.23137504>
- GitHub: <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1>

Reproduction entry point from the repository root:

```
python -m pip install -r mcc_3_0/requirements.txt
python mcc_3_0/reproduce_mcc_3_0.py
```

Stage 1 is a provenance freeze. Stages 2-5 replay in dependency order, and Stage 6 audits cross-stage consistency. A separate clean-machine replay by an external verifier remains an operational reproducibility step.

Appendix D — 3.0 Terminology

- finite SOURCE: upstream candidate that generates a finite address window from a declared rule rather than assuming a physically instantiated infinite address space.
- common carrier: a shared transport basis in Actual used by multiple representations and forces; not a fifth force.
- Actual: the total state presently owned and updated.
- Reality: the layer in which the state is rendered into observables and calculated through validated physical modules.
- Record: information retained in the present and available to affect subsequent execution.
- Residue: a current ledger term retaining the effect of prior execution rather than a failed output to be discarded.
- Component: a pre-phenotype unit satisfying origin + activation + identity retention + ledger closure.
- Phenotype: rendered physical expression such as mass, charge, spin, location, or observable response.
- Harmonic state: a representation in which multiple modes preserve relative phase; the 0 : 5 : 29 neutrino ledger is the present concrete example.
- Typed ledger: a bookkeeping structure that keeps addresses, probabilities, components, energy, information load, and gravity in distinct types and ownership domains.

Appendix E — Edition Note

This is the English translation of the cumulative integrated book manuscript v0.12. The broad preservation of the 2.3.2 material is intentional: the purpose is not to erase the earlier research lineage but to place the new finite-SOURCE and record/component layers into a single continuous book. The English text preserves the scientific ownership and claim boundaries of the Korean v0.12 manuscript, with the shared historical appendix explicitly identified while using the already published English 2.3.2 and WRRM_M manuscripts where they are direct counterparts.

Appendix F — Additional References for Version 3.0

- [3.0-R1] Choi, W. (2026). Finite but Boundaryless: A WRRR Hybrid Expansion and Accumulated Twist Cosmology with Hidden Gravitational State, Version 1.0. GitHub repository: `Wonsik-Choi-janefather/wrra-finite-boundaryless-twist-cosmology`. The paper identifies KOO JEONG A: OUSSSMOS, Leeum Museum of Art, Seoul, as conceptual inspiration rather than scientific evidence.
- [3.0-R2] Choi, W. (2026). Minimal Computing Cosmology 2.3.2: Integrated Model. Zenodo. DOI: <https://doi.org/10.5281/zenodo.22733000>
- [3.0-R3] Choi, W.; Choi, J. (2026). WRRR M Integrated Upstream-Downstream Model 1.0 r1. Zenodo. DOI: <https://doi.org/10.5281/zenodo.23126800>
- [3.0-R4] Choi, W.; Choi, J. (2026). WRRR_M Finite Klein-Zeta Generation Window v0.1. GitHub: `Wonsik-Choi-janefather/wrra-m-0.1/upstream/finite_klein_zeta_window_v0_1`
- [3.0-R5] Choi, W. (2026). Twist Stress Universe and the Mass Phenotype of Dark Matter. Zenodo. DOI: <https://doi.org/10.5281/zenodo.22700557>
- [3.0-R6] Choi, W. (2026). WRRR Spatial Twist Selection of Galactic Disk Planes and the Dark Mass Phenotype. Zenodo. DOI: <https://doi.org/10.5281/zenodo.23003875>
- [3.0-R7] Choi, W. (2026). WRRR Prime-Parts Rendering: Finite Assembly, Neutrality, Composition and Decay. Zenodo. DOI: <https://doi.org/10.5281/zenodo.23128482>
- [3.0-R8] NIST Digital Library of Mathematical Functions. Riemann Zeta Function: Zeros and Euler-Maclaurin Representations.
- [3.0-R9] `mpmath` documentation, `zetazero`.

Author Contact

Wonsik Choi
Email: janefather@gmail.com
ORCID: 0009-0001-4263-9772

Part XIII Theory improvement and final review

Chapter 74 Introduction and scope

The theory improvement of Minimal Computing Cosmology 3.0 and WRRRA concerns an executable connection between a finite SOURCE, arithmetic addresses, a common carrier, phenotype, records, physical load, and macroscopic outputs. The cumulative edition described this genealogy, while the structural meaning of addresses, the scope of minimality, the supplier for jointly calibrated internal states, correlated loads, and gravitational responses required more explicit contracts. We assemble the six improvement stages and their final evidence audit into a mathematical construction and computational validation [1–8].

The evaluation order is fixed: verified inputs, WRRRA transformation, outputs, falsification conditions, and conclusion. Observed values and known constants may serve as calibration inputs, with their locations disclosed. Reproduction of calibrated physical values counts as consistency and explanatory success. Unmeasured quantities computed after freezing the model are conditional WRRRA predictions under the stated construction. Numerical novelty and additional independent predictions are not introduced as completion requirements for this improvement campaign.

The principles of cosmic nonexceptionality, minimal computation, and existence of the present remain starting premises. Computation verifies the model assembled from these premises and disclosed constitutive rules. The final stage consists of six cross-evidence checks of hashes and interfaces. It is neither six independent observations nor a new full rerun of every stage [8].

Chapter 75 Finite SOURCE and arithmetic state

The reference input fractions are (P,D,R)=(0.05,0.268,0.682), where P denotes phenotype, D resident non-phenotype, and R return. The least common multiple L=500 of reduced rational denominators, together with the disclosed construction factors 29 and 70, defines N. Address 1 is inactive, and normalization runs from 2 through N. This supplies one finite candidate; it neither requires nor derives mandatory temporal stability of cosmic capacity [2].

$$N = L \cdot 29 \cdot 70 = 1\,015\,000, \quad w_n = \frac{n^{-\alpha}}{\sum_{m=2}^N m^{-\alpha}}, \quad \alpha = 1.8996877935161325.$$

An address represents prime-factor assembly and readout structure rather than an arbitrary display number. Primes enter the return branch; even composites enter D. Admission of odd composites into P follows four-mode phase readout and an eight-frame remainder update. Prime labels alone are not claimed to derive individual particle identities uniquely [3].

$$d_k(n) = \sum_{j=1}^4 c_j \cos[\gamma_j(\log n + \xi k) + \varphi_j], \quad t_k(n) = \frac{1}{1 + \exp[-(d_k(n) - h)]}.$$

$$e_P(n) = 1 - \prod_{k=0}^7 [1 - t_k(n)], \quad e_R(n) = 1 - e_P(n), \quad e_D(n) = 0 \quad \text{for odd composites.}$$

For other addresses, primes receive (eP,eD,eR)=(0,0,1) and even composites receive (0,1,0). Effects are therefore nonnegative and sum to one at each address. The reference coefficients are cj=0.5, offsets zero, ξ=0.1, and h=1.4476744689338703. The adopted γ values are 14.134725141734695, 21.022039638771556, 25.01085758014569, and 30.424876125859512. These are disclosed spectral and calibration inputs.

The sums $\sum w_n e_s(n)$ reproduce the reference fractions. Return 0.682 separates into a prime contribution of 0.6742270361220081 and a nonadmitted odd-composite contribution of 0.007772963877991983. Passive relabeling preserves outputs when assembly structure is transported with labels; active reassignment can change them. Separate input probes at N=4,060,000 and 20,300,000 were propagated through frozen downstream responses and are not interpreted as observations of cosmic time evolution [2,3].

Chapter 76 Minimal computation and equivalent implementation

Minimal computation requires specifying what an alternative must preserve. Reductions that retain fractions, realizations that retain the full phase sequence, and implementations that execute the same operation more efficiently constitute different comparisons. Models with 1,2,4,8 frames and a scalar admission model can calibrate to the same reference fractions while differing in detailed readout and conditional outputs [4].

Proposition 1. An autonomous real linear time-invariant realization of the entire fixed four-mode phase continuation requires at least eight states. Preservation of the whole continuation and membership in the autonomous LTI class are assumptions of this statement.

$$\lambda_{j,\pm} = e^{\pm i\gamma_j\xi}, \quad H_8 = V \operatorname{diag}(a_1, \dots, a_8) V^T, \quad \operatorname{rank}(H_8) = 8.$$

Proof. The eight poles are distinct and the complex coefficients at the reference address are nonzero. The corresponding 8×8 Vandermonde matrix V is invertible, so H_8 has rank eight. A d -state autonomous LTI realization has Hankel rank at most d , implying $d \geq 8$. Four real 2×2 rotation blocks realize the same sequence and attain the bound. H_8 uses synthesized continuation values at indices 0 through 14. It is not constructed from the eight admission samples alone, and the result does not establish uniqueness among all nonlinear implementations [4].

$$\begin{pmatrix} C_{k+1} \\ S_{k+1} \end{pmatrix} = \begin{pmatrix} \cos \omega & -\sin \omega \\ \sin \omega & \cos \omega \end{pmatrix} \begin{pmatrix} C_k \\ S_k \end{pmatrix}, \quad \omega = \gamma_j \xi.$$

The rotation implementation preserves the generating sequence and its explanation while reducing trigonometric evaluations from 13,692,544 to 3,423,144. Additional multiplications and additions prevent interpreting this as a 75% reduction in total computation. Among the 15 nonempty mode subsets, the closest reduced model using modes 1 and 2 had a weighted admission L1 difference of 0.0005084783856417499. This is an internal model difference, not an observational fit loss [4].

Chapter 77 From the common carrier to channels and phenotype

The common carrier is defined on a 128-site load lattice, with a separate channel space. The 15 CCS transport directions and one neutral extension have transport ranks 15 and 16. Rank one of the default stochastic address-to-origin readout belongs to a different map. Three families and 16 channels provide 48 slots; these are not counted as 48 particles, masses, or independent record events. Their normalized weights partition one P-energy ledger [3,5].

Selection uses actual carrier responses and compatibility comparisons among four candidate filters. The F_DX placement was selected across the ten reviewed conditions, with a minimum score gap of approximately 0.0242867792. Zero selection contrast ties all four candidates and rejects selection. Optimization weights are not identified with Born probabilities. The selection result file is an actual dependency of phenotype construction [5].

$$K_s(n) = \sqrt{e_s(n)} P_{\text{sel}}, \quad \sum_s K_s(n)^\dagger K_s(n) = I_{16}, \quad A_s^{\text{joint}} = I_{16} \otimes A_s.$$

Under the classical address-block assumption, this map preserves positivity, normalization, and channel coherence before a separate label projection. Channel-blind load operators commute with the selected permutation, preventing passive placement from generating new energy. This conditional construction does not derive coherence between addresses or a collapse law. The anomaly cancellations of the disclosed charge assignment and the 48-slot ledger were computationally reproduced [5].

Chapter 78 Internal calibration and the supplier for a single record

Nucleon internal states connect the inherited finite spin-flavor construction to two-configuration mixing. Proton and neutron masses of 938.27208943 and 939.56542194 MeV and current-response coefficients are disclosed calibration inputs. The effective Hamiltonian uses $\Delta=(m_p+m_n)/6$ as a construction rule, not a measured QCD excitation spacing [5].

$$H = (m - e_{\min})I + \Delta \begin{pmatrix} 0 & -C\sqrt{r_3 r_5} \\ -C\sqrt{r_3 r_5} & 1 \end{pmatrix}.$$

The magnetic-response calibration gives $q_{\text{mix}}=0.07353193837872393$, $\theta=15.733765$ degrees, and $g_A=1.5686240821617012$, failing to reproduce the reference axial value 1.2753. The joint calibration uses $q_{\text{mix}}\approx 0.293525$ and an additional current coefficient $c_1\approx 0.439987793$ to reproduce masses, magnetic moments, $g_A=1.2753$, and the inherited lifetime calibration. The lifetime of 878.3 s and its coefficient belong to an already calibrated ledger and are not independent predictions. The mixing quantity q_{mix} differs from cosmic deceleration q [5].

A single effective record operation applies across the channels. Multiplying $W=P+\otimes \text{Irecord}+P-\otimes \text{Xrecord}$ by the channel identity commutes with the disclosed charges. Effective record occupancy distributes the P ledger across channels without duplicating record cost by the number of slots. The supplier must cover both record work and the energy required by possible outcome branches [5,6].

Branch	Effective gap MeV	P supply condition	20% supplier
Inherited effective record	431.081244315	Tested at 20%	Admitted
Magnetic internal calibration	366.935940492	15.9112205014%	Admitted
Joint internal calibration	757.895431800	31.8278353979%	Rejected

The joint branch requires work of $1.2033019830228727 \times 10^{-11}$ J and minimum supply, including outcome branches, of $1.2033440156388269 \times 10^{-11}$ J. The inherited 20% supply of $7.561582499072275 \times 10^{-12}$ J cannot execute it. An alternative branch is admitted after disclosing a minimum P-supply fraction of 31.82783539785394%. This is a supplier fraction within the record P ledger, not a cosmic matter fraction. A law for natural generation of the supplier is outside this computation [5].

Internal-state branches are alternative models and their energy ledgers are not added together. A composite nucleon is not identified with one Weyl slot. Freezing the current SOURCE in the magnetic branch gives conditional outputs $q_{\text{mix}}=0.07353192154110458$, an effective gap of 366.9359260048723 MeV, and $g_A=1.5686241046118607$. The effective gap is not the discovery of an observed new particle or nucleon excitation [5].

Chapter 79 Correlated physical load and energy pressure

Physical load is computed from the actual joint address-carrier state. Here p_n is the carrier state conditional on address n , A_s is a positive load operator, and $g_s(n)=1+\lambda_s \log(n)/\log(N)$, with $\lambda=(0,0.25,0.1)$. Channels are incorporated through the carrier marginal. A product of address moments and carrier expectations is valid only when addresses and the carrier are independent [6].

$$L_s = \sum_{n=2}^N w_n e_s(n) g_s(n) \text{Tr}(\rho_n A_s), \quad A_P = I, \quad A_D = K_c/2, \quad A_R = K_b/2.$$

$$E_s(a) = \eta_s V_0 a^{n_s} L_s, \quad V = V_0 a^3, \quad P_s = - \left. \frac{\partial E_s}{\partial V} \right|_{L_s} = - \frac{n_s E_s}{3V}, \quad (n_P, n_D, n_R) = (0, 0, 3).$$

The SI responses η are $(7.561582499072265, 7.285761328795971, 7.646102818811323) \times 10^{-10} \text{ J/m}^3$, with $V_0=1 \text{ m}^3$. K_c is the disclosed 128-site ring operator; the local perturbation matrix is $D=|0\rangle\langle 0|$. $K_b=(K_c+\epsilon D)/(1+\epsilon/(2N_{\text{carrier}}))$, where $N_{\text{carrier}}=128$. SOURCE N and carrier lattice size are distinct variables. Positivity of the operators and normalization of states ensure nonnegative loads. SI responses and volume exponents remain constitutive and calibration inputs.

Twelve cases test correlated diagnostic states grouped by $n \bmod 3$, two carrier states, $\epsilon=0,8$, and $a=0.5,1,2$. Passive joint permutations preserve outputs. Commutator evolution generated by the same energy Hamiltonian makes the derivative of total energy expectation zero. A positive calibration target with zero support is rejected; zero target and zero support yield $\eta=0$. A small positive occupancy of 10^{-12} is retained instead of being discarded by an arbitrary threshold [6].

$$E_{\text{before}} + E_{\text{supplied}} = E_{\text{after}} + W_{\text{record}} + E_{\text{supplier remaining}}.$$

The equation above is checked for 36 boundary ledgers combining three record branches and twelve states. Supplier and record terms are pressureless under this contract. Including both within the boundary propagates the same total energy and pressure; supplying energy from outside the boundary requires an explicit influx. Record energy is not declared nongravitating, and alternative internal models are not simultaneously added [6,7].

Chapter 80 Gravity rotation lensing and expansion from one state

Total energy density ρ and pressure P feed expansion, while resident-nonphenotype density ρ_D sets the shared transition acceleration a_T . In the equations below, ρ is energy density in J/m^3 , not mass density. The reference critical energy density is $u_{\text{crit}}=7.668947767821907 \times 10^{-10} \text{ J}/\text{m}^3$. Rotation and conditional lensing use the same a_T with no per-output recalibration [7].

$$\begin{aligned} \frac{H}{H_0} &= \sqrt{\frac{\rho}{u_{\text{crit}}}}, & q &= \frac{1}{2} \left(1 + \frac{3P}{\rho} \right), & a_T &= cH_0 \sqrt{\frac{\rho_D}{8u_{\text{crit}}}}. \\ g_b(r) &= \frac{GM}{(r^2 + b_0^2)^{3/2}}, & g(r) &= g_b(r) \nu(g_b/a_T), & \nu(y) &= \frac{1}{1 - e^{-\sqrt{y}}}, & v^2 &= rg. \\ \alpha_{\text{lens}} &= \frac{4}{c^2} \int_0^{z_{\text{max}}} g(\sqrt{b^2 + z^2}) \frac{b dz}{\sqrt{b^2 + z^2}}, & z_{\text{max}} &= \sqrt{R_{\text{patch}}^2 - b^2}. \end{aligned}$$

The local baryonic source is fixed as a Plummer mass $M=6 \times 10^{10}$ solar masses with scale $b_0=3$ kpc. Rotation is evaluated at 8.2 kpc, lens impact parameter $b=10$ kpc, and integration patch 200 kpc. The lens formula uses the disclosed weak-field condition $\Phi=\Psi$. This local renderer and homogeneous expansion connection do not constitute a derivation of complete four-dimensional gravitational dynamics [7].

Reference output	Value	Meaning
P density	$3.780791249536136 \times 10^{-11} \text{ J}/\text{m}^3$	Shared ledger
D density	$2.032185430065150 \times 10^{-10} \text{ J}/\text{m}^3$	Shared ledger
R density	$5.258550172595951 \times 10^{-10} \text{ J}/\text{m}^3$	Shared ledger
Pressure	$-5.258550172595951 \times 10^{-10} \text{ Pa}$	Volume derivative of the same E
H/H0	0.9999913260079337	Homogeneous reference branch
q	-0.5285585894376318	Deceleration parameter
aT	$1.191787531397112 \times 10^{-10} \text{ m}/\text{s}^2$	Shared transition scale
Rotation	207.5102418659153 km/s	Fixed local source
Lensing	0.5355822400088303 arcsec	Fixed weak-field patch

A separate 512-point Gaussian quadrature implementation agrees with the reference lens value to $1.3433698597964394 \times 10^{-14}$ arcsec for the same kernel. This is numerical integration consistency, not independent observational validation. When $\rho_D=0$, $a_T=0$ and the model reduces to baryonic rotation of 161.45051053125525 km/s and lensing of 0.2170452532181437 arcsec, consistent with the analytic reference. Zero total cosmic density gives $H=0$ and undefined q . A separately specified local baryonic source does not automatically disappear at this boundary [7].

Chapter 81 Validation results and disclosed construction

Stage	Connection improved	Reported checks
1	Finite-capacity input transport	Pinned report and hashes
2	Address assembly and channel load	52
3	Reduction and minimality scope	38
4	Selection phenotype internal states records	51 connection and 38 inherited replay
5	Correlated load and record boundary	23
6	Shared gravity and zero limits	20
7	Cross-stage evidence audit	6

These checks include unit, structural, inherited replay, and boundary tests with dependencies among them. Their counts are not summed into a number of independent scientific evidence items. The twelve macroscopic cases and thirty-six record boundaries define the executed input and branch contracts. The actual computation linking nucleon and macroscopic ledgers is assessed as an executed result, separately from physical uniqueness of the selected rules.

Four modes and eight frames, channel and three-family assignments, internal Δ and current calibration, load operators and volume laws, SI responses, local sources, and conditional gravity formulas remain disclosed specifications. The executed scope does not include CKM/PMNS mixing, unique SOURCE decoding of every species mass, natural supplier generation, or full four-dimensional gravitational dynamics. This scope statement neither cancels the completed six-stage improvement nor creates automatic additional stages [8].

Chapter 82 Falsification conditions and conclusion

Failure to reproduce reference outputs under identical frozen inputs, or failure of positivity, normalization, charge, or energy conservation, withdraws the corresponding connection verdict. The integrated verdict also fails if channel or address counts duplicate energy, the insufficient 20% supplier for the joint branch is concealed, pressure comes from a different energy function, or undisclosed per-output calibration is introduced. Extending the minimality proposition from the complete autonomous LTI continuation to uniqueness of every implementation invalidates that extension.

Internal gaps and unmeasured outputs from the current frozen SOURCE can be compared with experiment when the conditions identifying them with the same effective observable are met. Disagreement then requires revising the construction or response specification, or excluding the branch. Observations used for calibration are kept separate from comparison-only observations. This review part does not claim a new data fit or observational uncertainty analysis.

The originally defined six-stage weakness improvement is complete. Under disclosed inputs and branch supplier conditions, the conditional executable chain closes from finite SOURCE through phenotype, internal states, a single record, correlated physical load, shared energy and pressure, and macroscopic outputs. The method of minimal computation, common carrier, and phenotype operates within the stated computational scope. This conclusion concerns completion of an executable constructed model and does not extend to every possible physical phenomenon or necessity of every constitutive rule.

Chapter 83 Reproducibility and references

The original stage-seven distribution included editable sources, evidence snapshots and hashes. This is a provenance description, not an inventory of the present book package. Executable stages 1–6 are pinned to the commits below. Running `overall_audit.py` for stage seven reproduces the six evidence consistency checks, separately from a full rerun of preceding stages. The snapshots were generated in a Python 3.12, NumPy 2.3.5, SciPy 1.17.0 environment. Document preparation does not turn these numerical results into new observations.

- [1] Wonsik Choi and Jeongin Choi. Minimal Computing Cosmology 3.0. Cumulative integrated book source v0.4, 7 October 2026. Prior release DOI: <https://doi.org/10.5281/zenodo.23202732>. This identifies the prior public release, not a newly assigned DOI for this paper or v0.5.
- [2] WRRR/MCC 3.0 theory improvement stage 1. Finite SOURCE review. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/408602d166b3ec6635aff24803ef578580386112>
- [3] WRRR/MCC 3.0 theory improvement stage 2. Address structure and channel connection. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/d12ce729822cf78e4531c7303eef86507aa2238e>
- [4] WRRR/MCC 3.0 theory improvement stage 3. Reviewed scope of minimal computation. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/fccce122dc1b33fe06b5ced42ae2350f5d5f5621>
- [5] WRRR/MCC 3.0 theory improvement stage 4. Phenotype internal states and records. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/52e6c33c592350f89821ad3e7e5150de4a7a7b2c>
- [6] WRRR/MCC 3.0 theory improvement stage 5. Correlated load and boundary ledger. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/f5b6703a99248cbdcfb24cb0f95195fd660e276d>
- [7] WRRR/MCC 3.0 theory improvement stage 6. Shared gravity and zero limits. <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/dfa336c5526fe3e3923cee16e2a42dc2b3006412>
- [8] WRRR 3.0 theory final review stage seven v1.0. `FINAL_DECISION.json`, `EVIDENCE_MANIFEST.json`, `overall_audit.py`, and bilingual final reviews. 8 October 2026. This identifies `evidence_step7` in the earlier stage distribution, not a folder contained in the current book package. Consult the source archive ledger below.

Part XIV Current parent construction - paper v1.8

Wonsik Choi · Jeongin Choi

Independent Research, Seoul, Republic of Korea

Correspondence: janefather@gmail.com · ORCID 0009-0001-4263-9772

Paper v1.8 · 8 October 2026

Abstract

We develop a finite Wonsik Reality-Renderer Architecture (WRRRA) toy model to explore possible explanatory connections among quantum states and gravity, vacuum energy and expansion, and dark-matter and dark-energy effects. The purpose is model development and assessment of specified computational connections, rather than a claim to have solved these three problems or established WRRRA as the physical theory of the universe. The guiding hypothesis is that one nonprivileged universe can be described through minimal computation, a common carrier and phenotype. The implemented model maps a finite SOURCE and conditional carrier states through sector admission to information loads, calibrated energy, pressure and downstream gravity and expansion responses. Verified constants and observed sector shares are disclosed construction inputs. We identify what follows from those inputs and the WRRRA transformations, what is reproduced, and where the proposed connections remain incomplete. In a model with 1,014,999 active addresses and a 128-dimensional carrier, exact correlation-preserving reduction reproduces loads to 8.9×10^{-16} . Identical separate marginals need not preserve joint loads or calculated deceleration. A class-resolved lift repairs a concrete controlled-update failure. Applying the frozen WRRRA acceleration scale with the constitutive coefficient $\kappa=1/8$ through the credited empirical RAR kernel to 2,693 published SPARC points gives 0.13288 dex RMS without a new fit. These results substantiate the toy model's interface, identify its dynamical and observational limits, and document descriptive consistency. They do not derive microscopic quantum gravity or the small cosmological constant. The contribution is an executable, auditable model of a shared explanatory chain, including a precise account of which microscopic distinctions its present macroscopic outputs cannot resolve.

Keywords: WRRRA; finite toy model; model development; common carrier; quantum gravity; vacuum energy; dark sectors

84 Model development and the three explanatory questions

This study develops a finite toy model for investigating possible connections among three unresolved questions: quantum states and gravity, vacuum energy and expansion, and dark-sector effects. A toy model is a deliberately restricted construction in which assumptions, transformations and outputs can be inspected and tested. The research task is to implement and assess a proposed explanatory mechanism within that construction. It does not require that the complete physical questions have already been resolved.

The WRRRA hypothesis organizes one nonprivileged universe through minimal computation, a common car-

rier and phenotype. The model implements SOURCE → conditional carrier states → sector admission and phenotype → information loads → calibrated energy and pressure → gravity and expansion responses. The shared organization is the proposed explanatory structure. The calculations below determine its consistency, dependencies and present explanatory reach; they do not establish its unique physical realization.

The specific contribution is a worked WRRR interface result: replacing the correlated address-carrier state by its separate marginals can change the modeled pressure and deceleration, whereas retaining three weighted operators preserves every stipulated sector carrier readout. That snapshot representation nevertheless fails for the actual mod-3 controlled update; an explicit rank-six class lift restores closure. The model therefore identifies which information may be discarded, which correlations affect its gravity readouts, and which additional information its specified updates require. Calibration and inherited response laws are recorded separately from these consequences. The three physical questions motivate the construction; the results below assess its bounded explanatory mechanism.

Here “SOURCE” means the finite address measure and admission rule in Chapter 85; “phenotype” means the P-sector admission, not a derived catalogue of particle species. “Minimal computation” means preserving the declared readout/update contract while comparing information and implementation costs; it does not assert a global physical minimum. “Nonprivileged” is the motivating universality hypothesis, not an empirically established property of this particular arithmetic construction.

Observable-algebra reduction [2] organizes a state description around a restricted observable set. Exact dynamical reduction [3] additionally addresses preservation under a specified evolution. This work applies those distinctions to the disclosed WRRR coefficients: the concrete contribution is the actual-interface counterexample, its smallest linear class lift for the complete class-resolved arbitrary-carrier query family, and the resulting admissible macro domain. The general reduction identities are established mathematics. Their WRRR application makes a model-development decision testable: a snapshot cache cannot be used as the full state of this controlled evolution.

Evaluation of the three proposed connections

Quantum states and gravity. Verified inputs: normalized carrier states, disclosed operators and calibrated physical coefficients. WRRR transformation: the joint address-carrier state is mapped to sector-weighted carrier operators, loads, energy and the downstream gravity response. Outputs: exact preservation of specified correlated readouts and calculated expansion, rotation and conditional lensing; identical marginals can give different loads and deceleration. Falsification conditions: loss of state positivity or normalization, disagreement between direct and reduced readouts, failure of shared-energy bookkeeping, or disagreement with specified physical observations. This is an implemented connection between quantum state information and gravity readouts. It is not yet an internally calculated microscopic quantum-gravity theory; heterotic-string support is not counted as internal integration without the corresponding WRRR calculation.

Vacuum energy and cosmic expansion. Verified inputs: reference energy density, sector shares and disclosed dimensional calibrations. WRRR transformation: finite SOURCE admission and resident/return loads feed a finite energy function whose volume dependence determines pressure and expansion. Outputs: finite sector energy and pressure and a computed deceleration. Falsification conditions: inconsistent units, nonconserved exchange, disagreement between pressure and the stated energy derivative, or failure of the fixed model to reproduce the specified expansion observations. The finite construction avoids an unbounded ledger, but the observed small energy scale currently enters through calibration; finiteness by itself is not a derivation of that scale or a completed solution of the cosmological-constant problem.

Dark matter and dark energy. Verified inputs: reference shares 5%, 26.8% and 68.2%, source parameters and published acceleration measurements. WRRR transformation: arithmetic admission separates phenotype, resident nonphenotype and return sectors; their loads enter the shared energy and twist/local-gravity response. Outputs: reproduction of the calibrated shares and calculated expansion, rotation and conditional lensing, with a descriptive published-data acceleration comparison. Falsification conditions: violation of the common ledger, separately refitted outputs that no longer follow the same fixed state, or failure against specified rotation, lensing or expansion data. The RAR function is inherited and credited; the comparison is not a complete lensing or cosmological-data validation.

Taken together, these are a toy-model proposal whose specified connections are assessed by the computations below. They are not presented as three completed resolutions. Each output is assessed through its

disclosed transformation and failure conditions.

85 Verified inputs and the finite construction

Let the active address set be $n=2, \dots, N$, with $N=1,015,000$. Address 1 is inactive. Each address carries a normalized d -dimensional density operator ρ_n . Its classical weight is $w_n > 0$ with sum one. The complete classical-quantum state is

$$\Omega = \sum_{n=2}^N w_n |n\rangle \langle n| \otimes \rho_n, \quad \rho_n \succeq 0, \quad \text{Tr } \rho_n = 1.$$

This is a classical address register coupled to quantum carrier blocks; coherence between addresses is not part of this contract. Carrier coherence is allowed. In the numerical construction, $d=128$. Sector labels P, D, R denote phenotype, resident nonphenotype and return. Nonnegative admission effects $e_s(n)$ sum to one at every address. Define $f_s(n) = e_s(n)g_s(n)$, where g_s is a disclosed positive response. The general results below need no particular arithmetic classifier. The numerical instance uses the following rules.

$$w_n = \frac{n^{-\alpha}}{\sum_{m=2}^N m^{-\alpha}}, \quad g_s(n) = 1 + \lambda_s \frac{\log n}{\log N}, \quad (\lambda_P, \lambda_D, \lambda_R) = (0, 0.25, 0.1).$$

Primes have $(e_P, e_D, e_R) = (0, 0, 1)$; even composites have $(0, 1, 0)$. Odd composites use a calibrated four-mode, eight-frame admission:

$$d_k(n) = \sum_{j=1}^4 \frac{1}{2} \cos[\gamma_j(\log n + \xi k)], \quad t_k(n) = \frac{1}{1 + e^{-(d_k(n)-h)}},$$

$$e_P(n) = 1 - \prod_{k=0}^7 (1 - t_k(n)), \quad e_D(n) = 0, \quad e_R(n) = 1 - e_P(n).$$

The gamma values are 14.134725141734695, 21.022039638771556, 25.010857580145690 and 30.424876125859512. These are adopted spectral inputs; four-mode truncation is specified rather than physically derived. The common carrier is a periodic 128-site lattice. With S the cyclic shift and $D_0 = |0\rangle\langle 0|$, define

$$K_c = 2I - S - S^\dagger, \quad K_b = \frac{K_c + \epsilon D_0}{1 + \epsilon/(2d)}, \quad (A_P, A_D, A_R) = (I, K_c/2, K_b/2).$$

The denominator normalizes the trace of K_b to $2d$. These operators are nonnegative; they need not be bounded by I . The diagnostic $\epsilon=0$ and 8 changes the supplied carrier operator, without refitting dimensional coefficients.

Input	Value or rule	Status and role
Sector targets	0.05, 0.268, 0.682	Reference inputs; only two independent after normalization
N	$\text{lcm}(20, 250, 500) \times 29 \times 70$	Finite candidate based on rounded targets; no physical quantization law
α	1.8996877935161325	Joint calibration to address shares
h	1.4476744689338703	Joint calibration to address shares
ξ ; frames	0.1; 8	Specified admission rule
d ; ϵ	128; 0 or 8	Carrier size and diagnostic perturbation

Input	Value or rule	Status and role
η_P	$7.5615824991 \times 10^{-10} \text{ J/m}^3$	Frozen reference-state SI calibration
η_D	$7.2857613288 \times 10^{-10} \text{ J/m}^3$	Frozen reference-state SI calibration
η_R	$7.6461028188 \times 10^{-10} \text{ J/m}^3$	Frozen reference-state SI calibration
$V_0; n_P, n_D, n_R$	$1 \text{ m}^3; 0, 0, 3$	Reference volume and constitutive volume exponents
$H_0; u_{\text{crit}}$	$67.4 \text{ km/s/Mpc}; 7.6689477678 \times 10^{-10} \text{ J/m}^3$	Inherited reference constants
Local diagnostic	Plummer $M=6 \times 10^{10} M_{\text{sun}}, b_0=3 \text{ kpc}$	Specified source; not inferred galaxy profile
RAR comparison	Official RAR.mrt, 2,693 rows	Published observations; no fit in this revision

The rounded-target construction is not robust evidence for a unique capacity. For example, transferring 0.0001 from D to R changes the rational denominator and the candidate N to 20,300,000. Earlier frozen-response probes of large cutoffs are included in the reproduction package [7]. The model therefore retains a finite candidate and sensitivity analysis, not an observed measurement of the universe’s storage size. Pure address and SI-energy fractions also remain distinct ledgers: equal address weights do not imply equal physical energies.

Reference state and coefficient provenance

The address-share targets (0.05, 0.268, 0.682) and the inherited SI-energy targets (0.0493, 0.265, 0.6857) are distinct inputs. In `snapshots/source_step2/dependencies/original/m10_results.json`, the frozen 0.9 address measure with uniform carrier l/d has reference moments (0.05000000000000009, 0.2789373775449565, 0.687748727555431). Since $\text{Tr}[(l/d)A_s]=1$, the original calibration is $\eta_s = u_{\text{crit}} F_s / \mu_s(\text{ref})$, at $a=1$ and $V_0=1 \text{ m}^3$. This reproduces the η values above. Subsequent arithmetic and correlation probes retain these coefficients; their changed moments need not reproduce the original energy fractions exactly. Thus 5/26.8/68.2 is an address-admission calibration, not a second assertion about the same energy ledger.

$H_0=67.4 \text{ km/s/Mpc}$ is the adopted Planck 2018 base-Lambda-CDM reference [8]; $u_{\text{crit}}=3 H_0^2 c^2/(8 \pi G)$ is calculated from that anchor. The historical energy fractions are the project’s frozen rounded reference convention, not claimed here as the exact entries of a newly fitted Planck likelihood. The address targets are the project’s separate rounded construction targets. The frame offset is $x_i k$ with $x_i=0.1$ and $k=0, \dots, 7$; no independent $x_i k$ parameter is introduced.

86 Exact correlation-preserving interface

Define the three unnormalized sector-weighted carrier operators and scalar loads:

$$\Sigma_s = \sum_n w_n f_s(n) \rho_n, \quad L_s = \text{Tr}(\Sigma_s A_s).$$

Proposition 1 (readout sufficiency and positive realization). For fixed f_s , each Σ_s is nonnegative and exactly preserves every sector carrier query $\text{Tr}(\Sigma_s B)$, for arbitrary Hermitian B . The collection admits a completely positive trace-preserving realization when a failure branch is retained.

Proof. Positivity follows from nonnegative coefficients. Linearity of trace gives the query identity. Choose $M \geq \max_n \sum_s f_s(n)$. On the address-classical input, Kraus operators $J_{\{s,n\}} = \sqrt{f_s(n)/M} |s\rangle \langle n| \otimes I$ produce the flagged blocks Σ_s/M . Add $J_{\{F,n\}} = \sqrt{1 - \sum_s f_s(n)/M} |F\rangle \langle n| \otimes I$. Their adjoint products sum to the identity. The resulting state has unit trace and is a valid quantum channel output. Multiplying

the unnormalized flagged sector block by M recovers its weighted operator. If the carrier state is normalized after postselection, its sector probability must also be retained. Σ_s itself is an unnormalized bookkeeping operator, not a density matrix or an extra energy sector. ■

For G diagnostic classes with common conditional states τ_k and address class $k(n)$, one can stream

$$B_{sk} = \sum_{n:k(n)=k} w_n f_s(n), \quad \Sigma_s = \sum_{k=1}^G B_{sk} \tau_k.$$

This keeps correlations between sector response and carrier state. It does not replace the joint state by separate marginals. The grouping is a test input, not a derivation that all physical addresses fall into these classes.

Proposition 2 (restricted linear dimension). Let $F_{\{sn\}} = f_s(n)$ have rank r and fix strictly positive w_n . On the interior of the space of trace-one conditional carrier blocks, all-sector arbitrary-carrier readout variations have real dimension $r(d^2-1)$. An exact linear encoding for that entire query family requires at least this many variable real coordinates; r independent weighted operators attain the bound.

Proof. Expand each perturbation δp_n in a traceless Hermitian basis with d^2-1 elements. In each basis direction, $\delta \Sigma = F \text{diag}(w) \delta x$. Since $\text{diag}(w)$ is invertible, its image has dimension r . Directions are independent; sufficiently small perturbations around I/d preserve positivity. Thus the total image dimension is $r(d^2-1)$. A lower-dimensional linear encoding has a nonzero kernel that changes a target expectation; an appropriate Hermitian query separates that direction. Choosing a basis of the row space of F attains the dimension, with known fixed trace components. ■

This is a bound for a fixed query family and linear encodings. For three fixed A_s at one snapshot, just three scalar L_s suffice. Neither bound constrains nonlinear physical implementations or proves the physical necessity of the chosen arithmetic law. The dimension argument is standard linear algebra; its role is to define precisely what the WRRRA compression preserves.

87 Product-marginal error and dynamical closure

Let $\mu_s = \sum_n w_n f_s(n)$, the carrier marginal be $\rho_{\text{bar}} = \sum_n w_n \rho_n$, and $b_s(n) = \text{Tr}(\rho_n A_s)$. The product approximation is $L_s^{\text{prod}} = \mu_s \text{Tr}(\rho_{\text{bar}} A_s)$. Its error is exactly

$$L_s - L_s^{\text{prod}} = \text{Cov}_w(f_s, b_s), \quad |L_s - L_s^{\text{prod}}| \leq \frac{a_{s,\text{max}} - a_{s,\text{min}}}{2} \sqrt{\text{Var}_w(f_s)}.$$

The identity follows by subtracting the product of the two means. Cauchy–Schwarz bounds the covariance, and a random variable in the spectral interval of A_s has variance at most one quarter of that interval's squared width. This bound can be loose; the benchmark records both the observed error and bound. Zero spectral width makes the error zero, explaining preservation of the phenotype identity budget even when D or R loads change.

Snapshot sufficiency need not imply closed dynamics. A common carrier channel Φ independent of n obeys $\Sigma'_s = \Phi(\Sigma_s)$, so weighted operators remain sufficient. Address-controlled channels instead act before the weighted sum and generally require additional moments. In an equal-weight two-address qubit example, states $(|0\rangle\langle 0|, |1\rangle\langle 1|)$ and the swapped pair have the same average $I/2$ and the same initial Z expectation zero. Applying I at the first address and X at the second yields average $Z = +1$ and -1 , respectively. A marginal-only reduced state cannot replay this update.

Thus the correct contract is: fixed query operators at a snapshot, or a disclosed common carrier channel. An arbitrary new address-dependent update requires a fresh sufficiency or invariant-space check [3]. This prevents a compressed state from being described as a universally closed microscopic simulator. In the two-address example take all f_s constant: all three weighted operators are then equal between the two initial states. A stronger counterexample below uses the actual WRRRA coefficients.

Proposition 3 (class-controlled lift). Partition addresses into fixed disjoint classes C_k , keep w and f fixed, and let class k evolve by its disclosed common carrier channel Φ_k . Define

$$\Sigma_{sk} = \sum_{n \in C_k} w_n f_s(n) \rho_n, \quad \Sigma'_s = \sum_k \Phi_k(\Sigma_{sk}).$$

These class-resolved operators replay such updates exactly, and each block itself evolves as $\Phi_k(\Sigma_{sk})$. For the complete class-resolved carrier-query family, Proposition 2 gives minimum variable linear dimension $r_{\text{lift}}(d^2-1)$, where r_{lift} is the rank of the rows $f_s(n)1_{\{C_k\}}(n)$. This is sufficiency and a linear bound for that complete family; it is not a claim that every specified channel requires every class-resolved query. Proof: apply each linear channel inside its class sum and apply Proposition 2 to the lifted coefficient matrix. ■

Closure criterion and reusable audit. For arbitrary tuples of common CPTP maps on the specified classes, the original weighted interface is closed if the row space of F is invariant under multiplication by each class indicator. Equivalently, stacking the class-masked rows does not increase its rank. Then each Σ_{sk} is a linear combination of the original operators, so Proposition 3 supplies the update. Conversely, if a masked row leaves this row space, project it orthogonally onto the original null space. This yields a nonzero perturbation invisible to the original interface and visible to that class. Opposite small traceless carrier perturbations preserve positivity; an identity/bit-flip controlled update can distinguish the two states. Thus universal closure for this specified channel family cannot be asserted. This is an application of established invariant-space reduction [2,3], not a new general sufficiency theorem.

The audit has five reusable steps: (i) state the query family and form its address coefficient matrix F ; (ii) compute its rank and the retained operator family; (iii) mask its rows by the proposed control classes and compare the lifted rank; (iv) when rank grows, construct a null-space counterexample and retain class-resolved blocks; (v) choose a representation by measuring construction, storage and the required query workload. It applies to finite classical-quantum mixtures beyond arithmetic addresses. A snapshot cache is useful only when its contract matches the computation to be performed.

For the actual WRR f_s , use $C_1=\{n:n \bmod 3=0\}$, C_2 its complement. The initial rank is three, while the lifted rank is six. For $d=128$ the complete class-resolved family has 98,298 variable coordinates, versus 49,149 for the original three-operator family. An exact rank witness uses addresses 3,6,9 in the divisible-by-three class and 2,4,25 in its complement. In each class the prime, even-composite and odd-composite columns give a triangular 3×3 minor with strictly positive R , D and P pivots. The disjoint class supports combine these minors into rank six; no floating-point rank decision is needed. This answers a concrete implementation question: three scalar loads suffice for frozen cosmological outputs; three operators support changing snapshot carrier queries; class-resolved operators support the specified address-controlled updates. These are different information contracts, not interchangeable definitions of a universal minimum.

88 Shared energy, records and observable equivalence

The supplied energy and volume laws are

$$E_s(a) = \eta_s V_0 a^{n_s} L_s, \quad V = V_0 a^3, \quad \rho_s = E_s/V, \quad P_s = -n_s \rho_s/3.$$

They give pressureless P, D and constant-density R for frozen L . This matter-plus-constant-density behavior is a constitutive input, not a new derivation of accelerated expansion. Pressure is the derivative of the same E used for the density readout, at fixed L . Internal state evolution requires its own disclosed generator; this derivative is not the total derivative along evolving L .

Recording is checked in a boundary that includes its finite supplier. The same work cannot be counted both as new cosmic energy and as an already allocated phenotype resource:

$$E_{\text{sys,before}} + E_{\text{supplier,before}} = E_{\text{sys,after}} + E_{\text{record,after}} + E_{\text{supplier,after}}.$$

Here the record starts blank, its final energy is displayed separately, and supplier work equals the system-energy change plus record energy. The inherited effective-record, magnetic internal and joint internal models are alternatives, not simultaneously added particle species. The frozen supplier branches and their numerical budgets are documented in Appendix B. Let $p=\text{sum_sp_s}$ and $P=\text{sum_sP_s}$. The homogeneous and local diagnostic readouts use

$$H/H_0 = \sqrt{\rho/u_{\text{crit}}}, \quad q = \frac{1}{2}(1 + 3P/\rho), \quad a_T = cH_0\sqrt{\rho_D/(8u_{\text{crit}})}.$$

The local rule is $g(r)=g_b(r)v(g_b/a_T)$, $v^2=rg$, with $v(y)=1/[1-\exp(-\sqrt{y})]$. This is the existing empirical RAR function of McGaugh, Lelli and Schombert [4], evaluated here at the state-derived a_T . We do not claim the function itself is a WRRRA discovery. Conditional lensing integrates this same g under the stipulated weak-field condition $\Phi=\Psi$. Neither this local prescription nor the homogeneous relation supplies a complete four-dimensional gravitational theory.

Proposition 4 (three-load quotient and inverse within the renderer). With coefficients, volume, local source and response kernel fixed, any two microscopic states with equal L_P, L_D, L_R have identical H, q, a_T, v and conditional lensing. At positive density, the three densities are uniquely determined by H, q, a_T :

$$\rho = u_{\text{crit}}(H/H_0)^2, \quad \rho_R = \frac{1-2q}{3}\rho, \quad \rho_D = 8u_{\text{crit}} \left(\frac{a_T}{cH_0} \right)^2, \quad \rho_P = \rho - \rho_R - \rho_D.$$

Proof. The forward map factors through the three loads, establishing equivalence. Since $P=-\rho_R$, the q relation yields the second inverse; the a_T relation yields the third. Remaining density follows by subtraction. At fixed r and positive g_b , $g_b/[1-\exp(-\sqrt{g_b/a_T})]$ increases strictly with a_T , so a rotation value in its allowed range also determines a_T . Nonnegative recovered densities define the unrestricted scalar renderer domain. They are necessary but not sufficient to belong to the image of the fixed microscopic construction. ■

For fixed w, f and scale factor, $A_P=I$ fixes $L_P=\mu_P=\sum w_{nf_P}(n)=0.05$ independently of the conditional carrier states. Consequently the actual fixed-SOURCE macroscopic image has at most two variable loads. In this arithmetic instance, $f_D f_R=0$ at every address: D and R occupy disjoint supports. With otherwise arbitrary normalized conditional states, their exact attainable domain is

$$\begin{aligned} L_P &= \mu_P. \\ \mu_D \lambda_{\min}(A_D) &\leq L_D \leq \mu_D \lambda_{\max}(A_D). \\ \mu_R \lambda_{\min}(A_R) &\leq L_R \leq \mu_R \lambda_{\max}(A_R). \end{aligned}$$

Necessity follows from the spectral expectation bounds. Sufficiency follows by assigning independent extremal-state convex mixtures on the disjoint D and R supports. This rectangle is for the full conditional-state family, not for every constrained 32-group diagnostic family. At fixed a , convert the recovered densities to loads using the disclosed η and volume law and test this rectangle, including the fixed P equality. With $\epsilon=0$, both D and R operators range from zero to two; $\epsilon=8$ changes the R interval. The archive records these bounds and checks the reported states against them.

The unrestricted three-density renderer has information dimension at most three; its fixed-SOURCE microscopic image has at most two variable coordinates. A simpler three-load model is an exact baseline for these outputs. SOURCE structure is resolved by conditional address/carrier queries, not by declaring these already-equivalent macro numbers to be independent microscopic evidence. This equivalence is an explanatory constraint on what the present model can claim.

What the coupling explains within this model

Verified inputs are the same address marginal, the same carrier marginal, positive conditional states, and frozen η and volume exponents. The WRRRA transformation retains the covariance between sector response $f_s(n)$ and carrier expectation, passes its load changes into the same energy, and uses $P=-\rho_R$ in the common homogeneous response. The output is $q=-0.52856$ for the product replacement and $q=-0.54501$ with correlation retained (Chapter 89). This demonstrates a specific mechanism inside the toy model: an address-carrier correlation invisible to both separate marginals changes the return-sector density and hence the pressure-to-density ratio. It explains why marginal-only state descriptions are insufficient for the proposed gravity interface. It does not establish that the tested correlation is the measured cosmic state. The implementation fails if direct and retained-joint calculations disagree beyond tolerance or the covariance identity fails; observational adequacy is a separate empirical question.

A consequential constraint also follows from sharing the same ledger. For fixed SOURCE and scale factor,

define $p=\rho_P/u_{\text{crit}}$, $h=H/H_0$ and $t=a_T/(c H_0)$. Then

$$(1 + q)h^2 = \frac{3}{2}(p + 8t^2).$$

It follows by eliminating ρ_R from q and ρ_D from a_T . The 12 archived shared states satisfy this identity to 4.45×10^{-16} . Once p, h and t are fixed, q cannot be fitted independently; at fixed h , increasing the clustering contribution forces q upward under this law. This is a conditional consistency relation of the stipulated shared-energy response, not a newly derived gravitational field equation. Failure against a jointly specified physical state would reject that fixed response combination; introducing unrelated output-specific coefficients would change the model being tested.

Alternative constructions with the same calibrated shares

We bring the existing controlled architecture comparison into the main statement. All alternatives retain α and the SI coefficients and legitimately calibrate their admission threshold (or constant β) to the same address shares. Define $v_3(n)$ as the exponent of 3 in n and Q as the P -conditioned weight of addresses with odd $v_3(n)$. This is a conditional component-address query, not an observed particle abundance.

Admission	Q	q at $a=1$
One frame	0.4195836874	-0.5285597845
Eight frames	0.4020461347	-0.5285585894
Constant admission	0.3998832014	-0.5285589334

The same aggregate calibration does not make these conditional explanations identical. Eight frames and constant admission differ in Q by 0.0021629333, while their q values differ by only 3.44×10^{-7} and their current rotation outputs coincide. The exact eight-frame recurrence preserves Q and every address admission while reducing the direct trigonometric evaluations. This comparison supports retaining the phase architecture when the declared task includes that address/frame explanation; it does not empirically select eight frames. If only the macro outputs are requested, the load quotient remains the adequate smaller description. `snapshots/source_step3/structure_comparison.py` provides the construction and results.

Acceleration coefficient and sensitivity

Write $a_T = c H_0 \sqrt{\kappa \rho_D/u_{\text{crit}}}$. The inherited $\kappa=1/8$ is present in `snapshots/source_step2/connection_sources/carrier_compute.py` and in the frozen dependencies/ `original/run_stage5.py`. These files establish implementation provenance, but the supplied records do not establish a first-principles derivation or an independently documented original selection rationale for $1/8$. We therefore retain it explicitly as a constitutive choice. It is not derived from the eight admission frames, and its successful scale is not evidence that eight frames are physically necessary.

Holding ρ_D/u_{crit} fixed gives $d \ln a_T = d \ln H_0 + (d \ln \kappa + d \ln(\rho_D/u_{\text{crit}}))/2$. An H_0 change of ± 0.5 around 67.4 changes the scale by $\pm 0.742\%$; a $\pm 1\%$ density-ratio change changes it by about $\pm 0.5\%$. These are conditional sensitivities, not a propagated cosmological posterior. Alternative coefficient scenarios with all other inputs fixed give:

κ	a_T in m/s^2	RAR RMS in dex
1/16	8.42721×10^{-11}	0.142060
1/8	1.19179×10^{-10}	0.132875
1/4	1.68544×10^{-10}	0.146099

This three-point sensitivity sweep is not an optimization or proof that $1/8$ is unique. It quantifies how much the descriptive comparison relies on that construction choice. The supplied empirical kernel is responsible for the response shape.

The rank-six minimality statement covers the complete class-resolved query family and arbitrary allowed class channels. It is not a proof that six blocks are minimal for replaying only the single fixed I/X update. The controlled X updates in Chapter 87 remain computational dynamics. A separate inherited fixed-volume check uses the same total positive energy operator as a dimensionless-clock unitary generator and verifies that sector exchange sums to zero (`state_load.py`). Neither check supplies a measured SI-time evolution law or derives cosmic expansion from microscopic dynamics.

89 Numerical controls and published-data comparison

All tests preserve the supplied SI coefficients. A dense 256-address, eight-dimensional random pure-state reference is evaluated directly and via three Σ_s . Ninety additional positive-operator queries give maximum error 2.9×10^{-15} . The full test has $N-1$ active addresses, $d=128$ and 32 fixed classes $n \bmod 32$ with seeded complex pure carrier states. Loads agree to 8.9×10^{-16} . Storing all dense conditional matrices would require 266,075,897,856 bytes in complex128; storing three weighted matrices needs 786,432 bytes. This is a theoretical storage comparison, not measured allocation. The grouped input already has a much smaller representation, and source construction still costs $O(N)$. No total-runtime speedup over all alternative implementations is claimed.

The same-marginals diagnostic uses two extremal eigenstates of $K_c/2$ and address groups $n \bmod 3=0$ versus the complement. Its paired correlated blocks have exactly the same carrier marginal as the product baseline and the same address distribution. Positivity is checked. The retained covariance changes D by -0.000406237 and R by $+0.035378346$; the error identity is satisfied to 3.1×10^{-16} . This is a constructed state test, not an observed cosmic correlation.

Diagnostic	Product marginals	Correlation retained
P load	0.050000	0.050000
D load	0.278926	0.278519
R load	0.687743	0.723121
q	-0.52856	-0.54501
Rotation, km/s	207.510	207.487
Conditional lens, arcsec	0.535582	0.535435

Twelve inherited shared-state configurations are reproduced by the scalar quotient with maximum printed-output discrepancy 2.9×10^{-14} . Equal-load microscopic alternatives therefore remain indistinguishable under this renderer, while the product-marginal ablation does not generally preserve those loads. These are different tests and support different conclusions.

The exact same-marginals construction is as follows. Let P_0, P_1 be the minimum/maximum eigenprojectors of $K_c/2$, $C=(P_0+P_1)/2$ and $D=(P_0-P_1)/2$. With $p=\sum_{n \bmod 3=0} w_n=0.2894395421456772$ and $t=\min(1, (1-p)/p)=1$, set $p_1=C+tD$ and $p_2=C-ptD/(1-p)$. Their weighted average is C and both are positive and trace one. The product baseline assigns C to every address. The original twelve inherited cases instead use l/d at every address, or a packet on the mod-3 class and l/d elsewhere; the packet is $(|8\rangle+|16\rangle+|24\rangle)/\sqrt{3}$. They cover $\epsilon \in \{0,8\}$ and $a \in \{0.5,1,2\}$.

Rotation is evaluated at $r=8.2$ kpc for $g_b(r)=GM r/(r^2+b_0^2)^{3/2}$, $M=6 \times 10^{10}$ solar masses and $b_0=3$ kpc. Lensing uses impact $b=10$ kpc, patch radius $R=200$ kpc and

$$\alpha(b) = \frac{4}{c^2} \int_0^{\sqrt{R^2-b^2}} g\left(\sqrt{b^2+z^2}\right) \frac{b}{\sqrt{b^2+z^2}} dz.$$

Radians are converted to arcseconds. This is a finite-patch weak-field diagnostic, not a measured lens system. The inherited twelve cases also compare independent 512-node Gauss–Legendre and adaptive quadratures, agreeing below 10^{-9} arcsec.

A fair representation comparison. For the 32-class complex pure-state diagnostic, retaining 32 vectors and the 3×32 weight matrix costs 66,304 bytes; three dense weighted operators cost 786,432 bytes; three scalar

loads cost 24 bytes. The weighted representation is therefore larger than this grouped input. Ninety seeded, changing Hermitian carrier queries, evaluated for every sector, agree between grouped and weighted representations to 6.94×10^{-18} . The strengthened optimized workload sweep below evaluates practical utility including construction. Scalar loads cannot answer new carrier queries, while grouped states retain specified class-controlled information that a snapshot cache loses.

Workload sweep with construction included. The archived workload experiment tested $d \in \{32, 64, 128\}$, $G \in \{8, 32, 128\}$ and $m \in \{10, 90, 300\}$: 27 cases, five repetitions each, seeded complex pure states and positive $3 \times G$ weights. NumPy 2.3.5 uses OpenBLAS 0.3.30 on an AMD EPYC 9V74; OPENBLAS_NUM_THREADS, OMP_NUM_THREADS and MKL_NUM_THREADS are fixed to one before import. The maximum query discrepancy is 1.39×10^{-17} . Grouped queries use a direct BLAS matrix product followed by row contractions, with the conjugate buffer prepared once; contraction-path planning is not repeated. Reported retained bytes exclude working buffers for both methods. Query generation and address-weight compilation are common excluded inputs; cache construction is included. The table reports 90-query medians, in milliseconds. Full per-run variation is retained in workload_sweep_results.json.

d / G	Group / cache bytes	Group / cache total ms	Estimated break-even m
32 / 8	4,288 / 49,152	0.739 / 0.581	28
32 / 32	17,152 / 49,152	1.433 / 0.680	19
32 / 128	68,608 / 49,152	3.914 / 1.952	38
64 / 8	8,384 / 196,608	1.519 / 1.693	None
64 / 32	33,536 / 196,608	3.823 / 1.766	8
64 / 128	134,144 / 196,608	12.209 / 6.993	46
128 / 8	16,576 / 786,432	6.219 / 12.924	None
128 / 32	66,304 / 786,432	13.739 / 8.277	11
128 / 128	265,216 / 786,432	38.370 / 7.824	4

The cache including construction is faster in 16 of 27 local cases; it loses in 11 cases. Per-query saving and median build cost give estimated break-even counts from 4 to 49. This linear estimate is workload-dependent, not a promised crossover on other hardware. The five-run minimum and maximum values are archived rather than treated as inferential uncertainty. For this complex128-vector/float64-weight layout, $G < 3d^2/(d+1.5)$, the cache uses more storage; for larger G it can use less. No single representation dominates. These measured costs operationalize the audit rather than identifying a physical minimality law.

Actual-interface failure and repair. Let $A_{\{sn\}} = w_{nf_s}(n)$ on the first 256 active addresses, $z_n = +1$ in the mod-3 class and -1 elsewhere, and $u_n = A_{\{Rn\}} z_n$. Define $h = u - A^T(AA^T)^{-1}Au$ and $\delta = h / \max |h_n|$. Set $p_n^{\pm} = (I \pm \delta_n Z) / 2$ on a carrier qubit for those addresses and $I/2$ elsewhere. All initial Σ_s coincide because $A\delta = 0$; all traces are one and positivity is explicit from $|\delta_n| \leq 1$. Apply I in the mod-3 class and X elsewhere. The difference of weighted Z queries becomes $2A \text{diag}(z)\delta$. The initial maximum difference is 1.12×10^{-16} ; after the update the R-sector difference is 0.6668017457457079. This embeds in the 128-dimensional carrier. The six lifted operators from Proposition 3 retain the missing class information and replay the update. Thus the failure concerns all three WRRRA weighted operators, not merely an ordinary unweighted marginal. It gives a reproducible reason for retaining a larger state when moving from frozen outputs to address-controlled evolution.

90 Reproducibility and falsification

Current reproduction supplement (commit-pinned public archive): https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/blob/923469bc980341ec91a98aed1b2b7ff7749d43a7/publication/WRRRA_3_0_v1_8/WRRRA_3_0_Reproduction_v1.8.tar.xz

SHA256: b31a391c61e6843ef959a3e4ea82a8efa57cee6d9d2bdeb97478f2ffe7ff7dc1. Download the archive and extract it with `tar -xJf WRRRA_3_0_Reproduction_v1.8.tar.xz`, then run the listed Python commands from the extracted directory. This v1.8 revision changes presentation and access documentation only; the model, numerical sources and reported results are unchanged from v1.7.

Original location of the inherited v1.4 archive: https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/blob/2f2b1c5e9e4c0625870fcddb5e2216737665e629/publication/WRRRA_3_0_v1_4/WRRRA_3_0_Reproduction_v1.4.zip. Archive SHA256: ea30c242fd374a04f39b3562f76f7de26fd4659e75f18cf33d5a3cd39b79775b. The adjacent manifest maps every included source and input to its hash.

The archive contains the full inherited sources for stages 1–7, their nested frozen dependencies and input manifests, the new benchmark, the official data bytes and a one-command reproduction entry point. The reference environment is Python 3.12, NumPy 2.3.5 and SciPy 1.17.0; Matplotlib is needed only for Figure 1. From the archive root:

```
python reproduce_all.py
python benchmark.py
python rar_comparison.py
python workload_sweep.py
```

The first command runs 27 stage scripts in an isolated copy and records exit codes, stdout, stderr and expected PASS conditions. The inherited execution record reports that all completed successfully; the current release retains that record. Stages use their pinned dependency snapshots: this is a full rerun of the published stage checks, not a newly refitted model. Different floating-point libraries can change final digits and timing measurements. The second and third commands recreate the new numerical controls and comparison from fixed inputs. `benchmark_results.json` and `rar_results.json` provide the full machine-readable values, including spectral error bounds. SHA256 checks cover the input data and distributed sources.

Verification has three distinct levels. The proved identities require their stated mathematical assumptions. Numerical disagreement principally tests software conformance or reveals a violated assumption. Observational residuals test the physical renderer against data; physical selection additionally requires a specified comparison and uncertainty model. The following failure conditions describe implementation conformance, not observational refutation of a proved identity. The exact interface fails if an independently implemented direct joint readout differs from its weighted-operator counterpart beyond the declared numerical tolerance, or positivity is lost. The marginal-error result fails if its covariance identity or spectral bound fails. The three-load equivalence fails if the same fixed coefficients and equal loads produce different downstream outputs. Conversely, if address-controlled updates are introduced, snapshot sufficiency alone is no guarantee; the counterexample is a test that rejects that extension. Supplier execution is rejected below its maximum branch requirement, and energy duplication or separate output refits invalidate the shared-ledger contract.

For observational claims, the frozen response can be compared to specified data and residual summaries; its association with measured acceleration is explicit here. A stronger physical selection test would require galaxy-level nuisance parameters and covariance, a fixed test set and a comparison of inequivalent microscopic constructions. No universal numerical rejection threshold is invented retrospectively from the present residuals. Current observational consistency and microscopic identifiability are separate claims.

Current release and claim mapping

The source paper v1.8 and its then-current short book statement v0.10 use `WRRRA_3_0_Reproduction_v1.8.tar.xz`. It preserves the v1.4 source snapshots and adds the version map and `revision_audit.py`; the earlier URL identifies inherited provenance only. `RELEASE_MAP.json` maps this release, scripts and outputs, and `REPRO_MANIFEST_SHA256.json` lists file hashes. This paragraph records that source release. The current full book is v0.12; its historical chapters follow the adjacent corrections and execution grades.

Current claim	Script from archive root	Output
Exact interface, covariance and rank-six repair	<code>benchmark.py</code>	<code>benchmark_results.json</code>
Conditional architecture comparison	<code>snapshots/source_step3/structure_comparison.py</code>	<code>adjacent structure_comparison_results.json</code>
Shared energy and macro states	<code>reproduce_all.py</code>	<code>rerun/FULL_RERUN.json</code> and stage outputs

Current claim	Script from archive root	Output
RAR descriptive comparison	<code>rar_comparison.py</code>	<code>rar_results.json</code>
Optimized workload sweep	<code>workload_sweep.py</code>	<code>workload_sweep_results.json</code>
Calibration, kappa sensitivity, macro constraint	<code>revision_audit.py</code>	<code>revision_audit_results.json</code>

Run `python revision_audit.py` in addition to the commands above. The inherited 27-script stage rerun is retained with its original execution record. The source paper v1.8 reran the benchmark, RAR comparison and new audit; the present book cites that record, and historical timings are labeled as the previous measured workload rather than new measurements. No stage result is represented as an independently observed physical datum.

91 Conclusion

We have developed and audited a finite WRRa toy model for exploring a shared explanatory chain across quantum-to-gravity, vacuum/expansion and dark-sector questions. The implemented contribution is the connection from finite SOURCE and common carrier through admission and phenotype to correlated loads and the stated energy, pressure and gravity readouts. The model uses disclosed calibrated physical inputs; its reproduction of known values is a consistency result. Exact readout reductions, the same-marginals counterexample and the class-controlled repair establish what information this implementation needs and what its present dynamics can support. The published-data comparison documents descriptive consistency of the frozen acceleration scale in a credited empirical kernel.

This model-development result is distinct from resolving the three physical problems. The microscopic quantum-gravity dynamics and vacuum-energy scale are not derived here, and the stated macroscopic readouts do not uniquely identify the arithmetic SOURCE. Those results define the model's current explanatory reach and guide future extensions. The disclosed input-transformation-output chain and its failure conditions make the proposed model open to reproduction and further assessment.

Appendix A. Published-data consistency check

For an observational consistency comparison, we use all 2,693 rows of the official RAR.mrt release [5,6]. Its columns give log10 baryonic and observed acceleration and their marginal errors. The existing WRRa scale $a_T = 1.191787531397112 \times 10^{-10} \text{ m/s}^2$ is fixed before this calculation. We change no stellar mass-to-light ratio, distance, inclination, cosmic fraction or response coefficient. The comparison is descriptive reuse of a previously published dataset and an inherited empirical function, not a blind validation or an independent discovery.

$$r_i = \log_{10} g_{\text{obs},i} - \log_{10} \left[\frac{g_{\text{bar},i}}{1 - e^{-\sqrt{g_{\text{bar},i}/a_T}}} \right], \quad \text{RMS} = \sqrt{\frac{1}{m} \sum_i r_i^2}.$$

Acceleration prescription	Scale, m/s ²	RMS residual, dex	Median residual, dex
Fixed WRRa state scale	1.19179×10^{-10}	0.132875	-0.001555
Published empirical	1.20000×10^{-10}	0.132913	-0.002462
RAR scale			
Baryons only	None	0.513981	0.430000

Figure 1. Left: published points and fixed response curve. Right: observed-minus-calculated log residuals. The published empirical-scale curve differs from the WRRa-scale curve by at most 0.00143 dex and would




Figure 1: Published SPARC RAR points and fixed WRRR-scale residuals

almost overlap it. The RMS difference is not evidence of statistical superiority. Rows share galaxy-level nuisance parameters; this four-column release does not provide galaxy identifiers or a joint covariance. We therefore do not treat 2,693 rows as independent likelihood factors, quote a significance or estimate physical parameter uncertainty from this table. This result establishes descriptive consistency of the supplied scale and kernel with these data. It does not select the microscopic address law or validate conditional lensing.

Appendix B. Frozen supplier diagnostics

The three supplier branches are finite frozen inputs, not new particle constructions in this article. Their minimum budgets are maximum conditional branch-work requirements. At reference phenotype energy $E_P = 3.780791249536135 \times 10^{-11}$ J the retained values are:

Supplier branch	Mean work, J	Required budget, J
Effective record	$7.029276088428041 \times 10^{-12}$	$7.029707767554742 \times 10^{-12}$
Principal magnetic	$6.015266344305666 \times 10^{-12}$	$6.015700324113602 \times 10^{-12}$
Joint calibrated	$1.2033019830228727 \times 10^{-11}$	$1.2033440156388269 \times 10^{-11}$

The joint budget divided by E_P gives 31.8278354%. The first two suppliers are allocated budgets of 20% of E_P ($7.5615824991 \times 10^{-12}$ J); these allocations exceed the smaller required-budget thresholds listed in the table. The joint branch is allocated its disclosed required budget. Mean work alone is not sufficient for every conditional branch. `snapshots/source_step4/channel_record.py` and `mass_mixing_join.py` construct the three underlying branches; `snapshots/source_step5/record_load_join.py` tests their nested conservation. This paper takes these derived, pinned records as inputs and does not infer their microscopic origin from the conservation identity.

References

- [1] A. Jenčová and D. Petz. Sufficiency in quantum statistical inference. arXiv:math-ph/0412093. <https://arxiv.org/abs/math-ph/0412093>.
- [2] O. Kabernik, J. Pollack and A. Singh. Quantum State Reduction: Generalized Bipartitions from Algebras of Observables. Physical Review A 101, 032303 (2020). <https://doi.org/10.1103/PhysRevA.101.032303>.

- [3] T. Grigoletto, Y. Tao, F. Ticozzi and L. Viola. Exact Model Reduction for Continuous-Time Open Quantum Dynamics. *Quantum* 9, 1814 (2025). <https://doi.org/10.22331/q-2025-07-29-1814>.
- [4] S. S. McGaugh, F. Lelli and J. M. Schombert. The Radial Acceleration Relation in Rotationally Supported Galaxies. *Physical Review Letters* 117, 201101 (2016). <https://doi.org/10.1103/PhysRevLett.117.201101>; arXiv:1609.05917, Eq. 4.
- [5] F. Lelli, S. S. McGaugh and J. M. Schombert. SPARC: Mass Models for 175 Disk Galaxies with Spitzer Photometry and Accurate Rotation Curves. *Astronomical Journal* 152, 157 (2016). <https://doi.org/10.3847/0004-6256/152/6/157>.
- [6] F. Lelli, S. S. McGaugh, J. M. Schombert and M. S. Pawlowski. One Law to Rule Them All: The Radial Acceleration Relation of Galaxies. *Astrophysical Journal* 836, 152 (2017). <https://doi.org/10.3847/1538-4357/836/2/152>. Official published data: <https://astroweb.case.edu/SPARC/RAR.mrt>, retrieved 8 October 2026, SHA256 24aa7059dab7fa44787f7c11191052489899819370f6508621674769f3b72833.
- [7] W. Choi and J. Choi. Minimal Computing Cosmology 3.0, cumulative book and theory improvement stages. Previous public book release: <https://doi.org/10.5281/zenodo.23202732>. Frozen stage-six repository commit: <https://github.com/Wonsik-Choi-janefather/wrra-m-0.1/commit/dfa336c5526fe3e3923cee16e2a42dc2b3006412>. This revision includes snapshots for all preceding stages; no new DOI has been assigned to paper v1.8 or book v0.10.
- [8] Planck Collaboration. Planck 2018 results VI Cosmological parameters. *Astronomy & Astrophysics* 641, A6 (2020). <https://doi.org/10.1051/0004-6361/201833910>. Adopted reference H_0 under base-Lambda-CDM; project rounding conventions are stated separately.

Part XV Matter, gravity and records connected through seven papers

92 Starting from fractions without stopping at fractions

Verified inputs are the r11 cutoff of 1,000,000 addresses, the address classification, inverse-power weighting and the benchmark fractions 5%, 26.8% and 68.2%. The WRR transformation applies admission and return filters, reads binary ports of retained relations, and transfers sector probabilities to an energy account. Outputs are the inverse-square odd-composite share 4.876893353%, reproduction of the benchmark after joint calibration, and deceleration -0.523 under the declared volume law. Failure conditions are loss of positivity or completeness, a claimed target outside the admissible domain, disagreement between direct sums and the aggregate channel, or inconsistent pressure differentiation. The conclusion is conditional reproduction through an explicit explanatory connection [P1].

The opening question was why ordinary matter occupies about five percent. Once that number is reproduced, however, one must ask whether different generating processes can produce the same total. In r11, different finite drives reproduce the same totals while retaining different address distributions. How much remains and how the retained state is read are therefore different questions. Two ledgers with equal totals can answer a later query differently.

Normalization leaves two independent composition constraints. Joint calibration of two parameters is legitimate model construction and reproduction of known values is a consistency achievement. The three fractions are not three independent tests. The contribution lies in connecting address generation, selection, readout and energy through a disclosed transformation, rather than in claiming previously unknown numbers.

The homogeneous r11 branch assigns zero pressure to ordinary matter and the resident sector, and constant energy density to the return sector. Thus

$$q = \frac{1}{2}(0.05 + 0.268) - 0.682 = -0.523.$$

This is not a correction of the earlier book's approximately -0.52855 . That branch converts address-dependent loads using fixed SI coefficients. A comparison must align probability fractions, energy fractions, volume laws and normalization. This edition retains the two constructions separately. A microscopic particle refinement must respect the same distinction.

93 From number structure to particle selection

Verified inputs are four address preparations from P1, known electron and muon properties, and two stipulated pair-output channels. The WRR transformation reads prime multiplicities as a dimensionless statistic and maps it to conditional particle selection. Outputs are preparation-dependent probabilities, a record-preserving state extension, and criteria for energy and pressure aggregation. Failure conditions concern normalization, the energy account, volume differentiation and the specified state extension. The result is one executable possibility for the toy model, not physical confirmation that primes generate particles [P2].

For $n = \prod_p p^{\nu_p}$ and $\Omega(n) = \sum_p \nu_p > 0$, define

$$\chi(n) = \sum_p \left(\frac{\nu_p}{\Omega(n)} \right)^2.$$

This is the probability of matching labels when two prime-factor occurrences are drawn independently with replacement. For $12 = 2^2 \times 3$, $\chi = 5/9$; for $30 = 2 \times 3 \times 5$, $\chi = 1/3$. It measures concentration of multiplicities rather than numerical size and is invariant under relabelling primes. That symmetry alone does not select a unique statistic or species assignment.

The toy model assigns electron-pair probability χ and muon-pair probability $1 - \chi$. With normalized admitted-address weights π_n , the total electron-pair probability is $X = \sum_n \pi_n \chi(n)$. Reversed and constant-routing controls distinguish address effects from species assignment. Mass, charge and spin are supplied properties, not derived from this probability.

Energy and its derivative require separate checks. Recovery of mean energy from sector totals for every admissible distribution requires energy to be constant within each sector. Recovery of pressure requires its volume derivative to be constant within each sector. Equal energies at one reference volume need not have equal derivatives. Giving a light electron pair the total energy of a resting heavy muon pair places the difference in kinetic energy. The electron pair then carries positive kinetic pressure. Matching energy does not establish dust compatibility.

94 Can the same ledger survive particle decay?

Verified inputs are P2 routing probabilities, measured masses and muon lifetime, and a specified leading-order three-body decay kernel. The WRRRA transformation updates populations by a positive finite map and retains daughter momentum cohorts by birth time. Outputs are eventwise conservation, energy-work closure, evolving counts and pressure, and a criterion for dynamical aggregation. Failure conditions include negative populations, missing daughter energy, incorrect stress and failed convergence under time refinement [P3].

At fixed volume, muon rest energy can become daughter kinetic energy while total energy remains constant. Pressure need not remain constant. In the baseline preparation, PV/E_* rises from 0.246350 to 0.332735 over five muon lifetimes. These are outputs of the specified decay episode, not a calculation of cosmic composition through all epochs.

During expansion, birth times matter because older cohorts have experienced more redshift. Two positive cohort mixtures can have the same current number, energy and pressure but different future moments under identical expansion. Current totals are not automatically a sufficient state for future calculation.

Let T be a microscopic transition and C an aggregation. A closed aggregate transition $\bar{T}$ requires $CT = \bar{T}C$. The observables to be read must also be recoverable from the aggregate variables. Only on a preparation family satisfying both conditions can the smaller ledger replace the larger one. P3 removes frozen occupation for the declared decay episode; it does not turn hot daughters into pressureless dust.

95 How a common carrier implements the selection rule

Verified inputs are the two known pair channels, reference energy and two conditionally independent preparation registers. The WRRRA transformation connects an equality projector on factor-occurrence labels to one effective carrier interaction. Outputs are P2 species probabilities without separately fitting a coupling at each address, conversion changes under detuning, and a relation between control work and pressure. Failure conditions concern label equality, energy exchange, switching work and the compositional handoff to decay [P4].

With label probabilities $r_p = \nu_p/\Omega$, independent registers agree with probability $\sum_p r_p^2 = \chi$. A common pulse converts this statistic into two output paths. Common means that every address uses the same interaction rule, not that all addresses have identical preparations.

Independence is a substantive condition. Correlations can change the match probability even when both register marginals remain unchanged. Moreover, this effective implementation reproduces species probabilities, not the entire coherent output channel in P2. An identical probability table and an identical quantum channel are distinct achievements.

Away from reference-volume resonance, the energy mismatch changes conversion. Work performed when switching the pulse must remain in the ledger. Omitting it would misstate energy conservation. The result is a finite effective interaction candidate, not a derivation of the complete fifteen-channel carrier or the origin of the weak interaction.

96 Matching energy and pressure together

Verified inputs are the particle momenta and conversion from P2–P4, a fixed volume law, and the energy and records of an additional reservoir. The WRR transformation uses cold allocation or a correlated compensating sector to match one energy function and its derivative. Outputs are compatibility conditions, finite compression and reserve requirements, record capacity and post-decay update conditions. Failure conditions are energy or pressure residuals, negative reserves, departure from the admissible domain and mismatched daughter-cohort records [P5].

For frozen nonnegative relativistic populations, kinetic pressure is a sum of nonnegative occupied-momentum contributions. Exact dust pressure requires every occupied momentum to vanish. Cold allocation is possible, but unequal masses require an explicit change in reference energy, expected particle number or energy outside the particle sector.

For a retained hot allocation, a compensating sector may be specified. If record a carries particle energy $E_a(V)$, constant total energy C has the form

$$B_a(V) = C - E_a(V), \quad P_a(V) + P_{B,a}(V) = 0.$$

Writing this equation does not itself realize a physical reservoir. P5 constructs a finite reservoir correlated with records, and states the reserve and volume domain needed to keep energies nonnegative. It also excludes exact interval compensation of a hot kinetic sector by an independently convex additive reservoir. Cancellation at one point cannot be extended to an entire volume interval without checking the law; a constant-density negative-pressure term does not generally provide the required interval response.

As decay proceeds, records matched to the original pairs cannot remain unchanged. Reservoir records must follow the daughter momentum cohorts. This is an executed connection to P3. Background closure on the matching subspace nevertheless does not establish stability of all inhomogeneous perturbations or an autonomous physical controller.

97 Defining the weight of information through response

Verified inputs are a finite address-carrier preparation, load observables, equilibrium constraints, baseline stiffness and dimensional calibration. The WRR transformation builds a free energy from a finite load-coupled partition function and calculates response from its Hessian. Outputs are unequal susceptibilities for equal snapshots, a stability domain, mode-dependent responses and sufficient-reduction criteria. Failure conditions concern covariance identities, source-aware elimination, pressure differentiation or use outside the positive-Hessian domain as though stability persisted [P6].

Address count, entropy, mean energy and response coefficient are not interchangeable meanings of information. Two preparations share mean load 0.823167093659 but have joint-relaxation susceptibilities 1.016726892208 and 1.016710206812. These are calculated dimensionless examples, not measured gravitational differences.

Joint relaxation allows both address weights and internal modes to redistribute. Clamped-address relaxation fixes address weights while allowing internal modes to relax. The first uses global load covariance; the second uses the weighted sum of conditional address covariances. The response problem is incomplete until the allowed relaxation is declared.

Under joint relaxation, the local effective stiffness is

$$K_{\text{eff}} = K_0 - \beta \epsilon^2 \text{Cov}(b).$$

Where K_{eff} is positive definite, its inverse determines the local response to the signed source perturbation. Replacing all modes by one scalar gravitational enhancement requires a special proportionality between the covariance correction and the baseline stiffness. General responses are mode-dependent. Eliminating hidden field coordinates must also retain source corrections and the energy constant, not merely the reduced stiffness.

The motivating question is now precise: once the load, its energy coupling and the allowed preparation are fixed, a response can be calculated. Address counting alone has not derived Newton's constant, universal equivalence or the Einstein equation. Empirical dimensional calibration is a legitimate input; relative responses subsequently obtained through the fixed transformation are model outputs.

98 A sufficient present and accumulating residue

Verified inputs are sixteen four-bit histories, a parity preparation rule, a finite auxiliary and a three-cell octal residue tape. The WRR transformation retains response-relevant parity and transfers the remaining distinctions to the tape. Outputs are the distinction between sixteen states for exact reconstruction and two states sufficient for the specified response, an eight-state blank auxiliary requirement and a conditional overflow time. Failure conditions include a noninjective claimed lossless compression, omitted pointer or auxiliary resources, a failed finite thermal ledger and an allegedly successful transfer beyond the stated capacity [P7].

Only the present exists does not mean that distinctions from earlier events cannot remain in the present. Reconstructing all four-bit histories requires sixteen distinguishable states, while the stipulated P7 response requires only even and odd parity. Discarding the other three bits demands an account of their destination. In a reversible transfer, another finite resource owns those distinctions.

For a tape of m cells with b states per cell and q_0 initially occupied cells, allow every input string and exclude release or lossy compression of distinctions. The first impossible transfer in the specified protocol occurs at

$$n_* = m - q_0 + 1.$$

An initially blank three-cell tape reaches this event on the fourth transfer; a tape with one occupied cell reaches it on the third new transfer. This is a limit of the assigned tape and fixed controller output, not a calculation of the universe's total freely usable register capacity.

Post-saturation policies differ. STALL preserves distinctions but does not complete the next transfer. OVERWRITE permits a further action while losing old distinctions. Overwriting one octal cell is an eight-to-one map and is not reversible by itself. Capacity constrains subsequent actions without selecting a unique cosmic transition law.

Accumulated loss here means that distinctions inaccessible in the local record remain elsewhere and consume resources. Assuming fundamental destruction alone does not fill a reservoir. Physical damage or defect accumulation requires its own dynamics. The result therefore specifies a finite accumulation mechanism without deriving a Big Bang, cosmic restart or universal lifetime.

P7 uses `exact_response_states` with `query_tolerance=0` and removes the unused `epsilon` input. Its displayed response gap is $1.66853951 \times 10^{-5}$ and the minimax error for a single common response is $8.34269754 \times 10^{-6}$. These are finite-preparation outputs, not measured gravity. Loss refers first to access from the current response record, not global destruction of information in the reversible example. For the overflow statement the domains are integers $b \geq 2, m \geq 0, 0 \leq q_0 \leq m$.

99 One execution structure after seven papers

Verified inputs are the preparations, constants, volume laws and transitions declared by the seven studies. The WRRRA transformation links filters, address readout, effective conversion, decay cohorts and the records needed for energy, pressure and response. Outputs are conditions determining which distinctions can be discarded and which must remain. Failure occurs if incompatible units, states or preparations are presented as one execution, or an unexecuted connection is labelled complete.

Connection	Assessment in this edition	Remaining condition
Filters to totals	Conditional r11 reproduction	Physical realization of generation drives
Addresses to particle selection	Calculated stipulated readout	Realized species assignment and preparation
Selection to decay	Executed with supplied lifetime and kernel	Full cosmic creation and annihilation history
Common carrier to conversion	Effective-pulse probability reproduction	Full fifteen-channel and coherent-channel realization
Hot allocation to dust account	Conditional correlated-reservoir completion	Autonomous reservoir and controller dynamics
Information load to response	Finite susceptibility under specified equilibrium	Universal gravity coupling and relativistic dynamics
Records to overflow	Conditional finite-tape threshold	Cosmic saturation and a subsequent transition law

This table does not add seven papers into a completed universe simulator. Executed compositions include P4 conversion with P3 decay and P5 cohort-aware reservoir updates. P6 equilibrium response and P7 record control use separately declared preparations. Autonomous integration on one cosmic trajectory would additionally require a bath, controllers, clocks and spatial relations. Until those connections execute internally, they are not completed integration.

Minimal computation has acquired a sharper question. The largest possible ledger capable of reconstructing every past event is not required for every task. Distinctions needed for the declared next query and action must nevertheless remain. Sufficiency is query-relative: moving from energy to pressure, or from present readout to future evolution, can require a larger state. This is the calculable minimality established here.

The universe need not be privileged or perfectly restored; it may be an adequate state able to continue within its available resources. This is the interpretation proposed by Minimal Computing Cosmology. Treating completed physical infinity as Null motivates the finite account, rather than proving mathematical infinity inconsistent. Each finite theorem and calculation is assessed under its own assumptions. Reproducing known values counts as consistency and explanatory achievement. Unmeasured outputs computed after fixing verified inputs are conditional WRRRA predictions. A novel number is not a prerequisite for a structural contribution.

Source ledger for this edition

This revision integrates the archived manuscripts below. Reported computational check counts and numerical outputs belong to those studies; editing this book does not constitute rerunning all their experiments. Internal AI reviews of individual manuscripts are neither journal peer review nor a new review of this book. Submission, receipt and acceptance status were not newly verified or changed by this editing task.

- [B] Developing a finite WRRRA toy model to explore quantum gravity vacuum energy and dark sector connections, v1.8. The current parent construction, incorporated in Part XIV.
- [P1] Conditional Reproduction of Ordinary Matter Fractions, r11 English submission manuscript.
- [P2] From Number Structure to Particle Selection, archived v0.2-r1 package.

- [P3] Dynamic Decays and Nonequilibrium State Transitions in a Finite WRRRA Model, archived v0.2-r1 package.
- [P4] Effective Carrier Routing, Control Work, and Stress in a Finite WRRRA Model, v0.2 editorial revision submission package.
- [P5] Energy and Pressure Compatibility in Finite WRRRA Models: Cold Allocations and Compensating Sectors, v0.2.
- [P6] Information Weight and Effective Gravitational Coupling in a Finite WRRRA Model, v0.3.
- [P7] Record Capacity and Irreversibility in a Finite-State Model, v0.2, the 10 October 2026 Entropy preparation manuscript.

The P1–P7 sequence follows the manuscripts’ citation chain and P5’s explicit description as the fifth manuscript. Counting B in place of P1 would omit the composition study used as the starting point of the later works. Conflicting numbers are resolved by distinguishing preparation, volume law and readout before comparison. Author attribution and research genealogy are preserved.

Shared Historical Appendix G: Earlier Expanded English Twist Exposition

This identical reference appendix preserves the earlier expanded English exposition in both editions. It does not replace the common Chapters 19–27. Historical PASS/closure refers only to the specified calculations and adds no separate language-dependent current claim.

1 Overview of Minimal Computing Cosmology

1 1 Meaning of minimal computation

Minimal Computing Cosmology does not assume that the universe is optimized toward a goal. Our universe is not special; it is treated as a structure that remained among possible structures by enduring changing conditions. Minimal does not mean the absolutely smallest amount of computation. It means sufficient to avoid failure in every declared environment without requiring a more complex structure as a necessity. Survival takes priority over peak performance, and degrees of freedom that appear redundant under ordinary conditions may preserve stability under extreme change.

$$C_{\min} = \min\{C(M) : M \text{ survives all declared tests}\}$$

Equation (1)

The cost in Equation 1 includes the number of fields, independent functions, input constants, initial conditions, update frequency, and the cost of handling failure. A model that minimizes cost but collapses in one environment is not a minimal computation model. This principle explains why the present paper does not adopt a formula that fits galaxy rotation alone, but instead connects the same structure to lensing and the early universe.

1 2 Relationship between WRR Core and cosmology

WRR Core 1.0 is not a specific physical theory but a general framework for describing minimum execution. Law specifies permitted changes; Source is the updatable origin; Execution is the actual operation; Renderer converts an internal state into an observable expression; Reality is the currently realized state; and Record is what remains for the next comparison. Minimal Computing Cosmology applies this framework to the universe. This paper does not alter WRR Core.

Table 1 Levels of WRR Core and the twist stress model

Ledger — Level · Meaning in this paper · Status

- Level: WRR Core 1.0
 - Meaning in this paper: Domain independent rules for minimum execution
 - Status: Preserved

- Level: Minimal Computing Cosmology
 - Meaning in this paper: Execution model for cosmic states and observable expressions
 - Status: Overview only
- Level: Twist stress model
 - Meaning in this paper: Connects the dark sector Actual to the mass Phenotype
 - Status: Subject of this independent paper

1.3 Actual and Phenotype

A central distinction in WRRR separates the structure that owns a state from the expression that appears to an observer. In this paper, the mass of ordinary matter remains a property of conventional particles and fields. In the dark sector, spatial twist stress is the Actual. The Phenotype is the mass value inferred by an observer from the additional gravity that this stress leaves in the metric and in orbits.

$$\text{Actual } \mathcal{R} \rightarrow \text{Phenotype}$$

Equation (2)

The dark component is therefore not an entity whose every phenotype is zero. Its electrical and optical expressions are assumed to be negligible, but its gravitational expression is nonzero. It requires no volume assigned to an individual particle, yet its stress energy density has a spatial distribution. This distinction makes invisibility, absence of particle volume, and presence of mass mutually consistent.

2 Research Question and Model Assumptions

2.1 Research question

This paper asks one question. If ordinary matter is interpreted conventionally while only dark matter phenomena are rewritten as the mass phenotype of twist stress preserved in space, can galaxy rotation, gravitational lensing, and early structure formation survive simultaneously within one model? The question does not presuppose that particle dark matter is absent. If twist stress is insufficient, an additional particle owner must be returned to the cost ledger.

2.2 Assumptions and scope of claims

Table 2 Model assumptions and claim grades

Ledger — Symbol · Assumption · Claim grade

- Symbol: A1
 - Assumption: The conventional account of ordinary matter and the four forces is retained
 - Claim grade: Model boundary
- Symbol: A2
 - Assumption: The dark sector Actual is distributed spatial stress
 - Claim grade: Model choice
- Symbol: A3
 - Assumption: Matter and light couple universally to one metric
 - Claim grade: Condition for covariant completion
- Symbol: A4

- Assumption: The universe may have finite boundaryless topology but no particular topology is imposed
- Claim grade: Topological hypothesis
- Symbol: A5
 - Assumption: Time is a relation among change and records rather than an independent substance
 - Claim grade: Ontological interpretation
- Symbol: A6
 - Assumption: The cosmological background and galactic response share a matched normalization
 - Claim grade: Conditional closure

A4 and A5 are not theorems that follow automatically from the physical calculation. A6 is also not uniquely derived from the principle of minimal computation. This paper states these assumptions explicitly and identifies both the calculations they enable and the observations under which they fail.

3 The Present and the First Twist

3.1 The interpretation that only the present exists

The model does not require the past and future to exist as physical locations alongside the present. What exists is the present state. The past is a record remaining in the present, and the future is a possible update not yet realized. Time can be defined relationally by comparing the change of a repeatable clock variable with a reference rate of change.

$$\Delta t_{\text{clock}} \equiv \frac{\Delta C}{R_C}$$

Equation (3)

This interpretation does not deny change. States are updated and records differ. Consciousness reads differences among records as a continuous experience, but consciousness is not required for the computation of the universe. The covariant field tau used below is read as a label that orders states rather than as a substantive flowing time. Because its normalization enters the action and produces a physical degree of freedom, however, the tension with a fully timeless ontology remains in the cost ledger.

3.2 First quantization and the first twist

A completely undifferentiated state contains no direction, position, particle, or record. The first quantization distinguishes possible states, and the first twist creates differences among paths and directions. On this reading, the Big Bang is not primarily an explosion into an external empty space. It is the first distinction and asymmetry that opened the relational network of the present universe.

undifferentiated → distinguishable → relational universe

Equation (4)

Equation 4 is an interpretation of origin, not a microscopic process established by observation. The topological group and quantum degrees of freedom in which the first twist arose remain open questions. The paper assumes that origin and then asks whether the response of the present universe is calculable.

4 Finite Boundaryless Topology and Bidirectional Twist

4.1 Local geometry and global topology

A space that is almost flat locally may still be closed globally. Finite and boundaryless means that the space has finite volume while no point is an edge or a privileged center. It also removes any requirement that space expand into something external. To separate local fields from global information, the one form A is decomposed into Hodge components.

$$A = d\phi + \delta B + h, \quad dh = 0, \quad \delta h = 0$$

Equation (5)

$$\oint_{\gamma} h = \Theta_{\gamma} \neq 0$$

Equation (6)

The term $d\phi$ is the local gradient response, δB is the rotational response, and h is the harmonic sector that cannot be removed by local differentiation. If Θ_{γ} , the integral of h along a closed path γ , is nonzero, global gluing information remains even when it is invisible locally. Saying that every place is a gluing surface does not mean that a physical seam lies at every point. It means that no local patch is designated as the beginning or end of the universe.

A Mobius strip is a useful analogy for orientation reversal and gluing, but the strip itself has one boundary and therefore is not an exact boundaryless model of the universe. A physical candidate must be sought among closed three dimensional manifolds. This paper does not select a particular manifold and retains only the global information that can in principle be measured through holonomy.

4.2 Bidirectional twist

Left handed and right handed twists may coexist. Energy and total stress depend on the sum of the squared components rather than on the sign of direction, while orientation is recorded separately by their difference. Total stress may therefore remain nonzero even when the net directional bias is small.

$$\kappa^2 = \kappa_+^2 + \kappa_-^2, \quad \chi_T = \frac{\kappa_+^2 - \kappa_-^2}{\kappa_+^2 + \kappa_-^2}$$

Equation (7)

When χ_T is zero, the two directional stresses have equal magnitude. When its absolute value approaches one, one direction dominates. This parameter may be related to circularly polarized gravitational waves, parity odd correlations, or global directionality data, but this paper does not derive a specific signal amplitude.

4.3 Relationship between expansion and twist

What is observed is the redshift between galaxies, angular diameter distance, luminosity distance, and structure growth. The picture of space stretching outward is one interpretation of those relations. If the same distance relations can be reproduced by a changing global holonomy or gluing rule, expansion and twist may be observationally equivalent descriptions. Renaming the description alone would not constitute new physics.

Table 3 Relationship between expansion and twist interpretations

Ledger — Feature · Expansion description · Twist description · Discriminating criterion

- Feature: Background distance
 - Expansion description: Change of scale factor
 - Twist description: Change of relational gluing and holonomy
 - Discriminating criterion: Interpretively equivalent if identical
- Feature: Lensing and slip
 - Expansion description: Calculated from metric potentials
 - Twist description: Stress response contributes to the metric
 - Discriminating criterion: Distinct slip can discriminate
- Feature: Global topology
 - Expansion description: Curvature and topology supplied as inputs
 - Twist description: Residual twist along a closed path
 - Discriminating criterion: Repeated correlations and matched circles
- Feature: Local boundary effect
 - Expansion description: Usually absorbed into local fields
 - Twist description: Possible nonlocal response of global stress
 - Discriminating criterion: Environment dependent RAR residuals

Interpretive boundary

The model therefore does not reject the observed distance relations. It interprets their Actual as changing gluing and stress rather than as an expanding container. Distinguishing the two descriptions physically requires lensing slip, global topological correlations, or a nonlocal response to boundary conditions.

4.4 Identifiability of the size of the universe

$$L_\gamma = |\Theta_\gamma|_{H_0}^c \sqrt{\frac{\zeta}{6f_T}}$$

Equation (8)

Equation 8 shows that a representative closed path length L_γ can be calculated if the holonomy Θ_γ , stiffness ζ , and stress fraction f_T are known. Current observations, however, constrain only the product $\zeta \kappa^2$ rather than ζ and the twist rate κ separately. Calculating the size of the universe from twist alone therefore requires one additional independent measurement of stiffness or holonomy. Single values such as 73 Gly or 147 Gly are conditional results that require extra assumptions; they are not conclusions of this paper.

5 Spatial Stress and the Mass Phenotype

5.1 Minimal stress energy

A space with twist rate κ can store deformation energy. In the simplest quadratic response, the stress energy density is defined as follows. The dimensionless stiffness ζ is restricted to positive values for sign stability.

$$u_T = \frac{\zeta c^4 \kappa^2}{16\pi G}$$

Equation (9)

$$\alpha \equiv \frac{c\kappa}{H_0}, \quad f_T \equiv \frac{u_T}{\rho_c c^2} = \frac{\zeta \alpha^2}{6}$$

Equation (10)

The quantity ρ_c is the critical density. Once observation fixes the effective dark sector fraction f_T , the product $\zeta \alpha^2$ is fixed, but ζ and α remain inseparable. This degeneracy is not a calculation error. It identifies the combination that the data actually measure. Reporting stiffness and twist rate separately without another observation would overestimate what has been inferred.

5.2 Why stress appears as mass

Spatial stress need not carry electric charge or optical emission. If it modifies the metric and affects the paths of matter and light, however, observers translate that effect into a gravitational source. Moving the modified field response to the source side of the standard Poisson equation defines an effective dark density.

$$\rho_T^{\text{eff}} = -\frac{1}{4\pi G} \nabla \cdot [(1 - \mu)\mathbf{g}]$$

Equation (11)

The quantity ρ_T^{eff} is not a particle number density. It is a mass phenotype that expresses the same observational effect in the language of standard gravity. Because the mass of ordinary matter remains unchanged, the model does not claim that every mass in the universe is stress. Only the additional gravity classified as dark matter is reinterpreted as a stress Actual.

Table 4 Expression channels of matter and twist stress

Ledger — Observation channel · Ordinary matter · Twist stress

- Observation channel: Electrical interaction
 - Ordinary matter: Appears through charge and current
 - Twist stress: Assumed to have no direct expression
- Observation channel: Optical emission and absorption
 - Ordinary matter: Appears through electromagnetic states
 - Twist stress: Assumed to have no direct expression
- Observation channel: Individual particle volume
 - Ordinary matter: Particle and composite descriptions apply
 - Twist stress: Not required
- Observation channel: Spatial distribution
 - Ordinary matter: Distributed as matter density
 - Twist stress: Distributed as stress energy density
- Observation channel: Gravity and lensing
 - Ordinary matter: Source of the metric
 - Twist stress: Appears as an effective mass phenotype

Background stress may exist throughout the universe, including our local environment, but it need not be perfectly uniform. If baryonic distributions, boundary conditions, and global holonomy modify the local response, the center of effective density may differ among galaxies and galaxy clusters. Computing this

distribution is a central task for subsequent work.

6 Minimal Galactic Response

6.1 Static effective energy and field equation

The static weak field of a galaxy is described by a dimensionless potential ϕ and an acceleration scale a_T . Connecting the Newtonian and low acceleration regimes with one response function F can reduce the cost of assigning an independent halo profile to every galaxy.

$$E[\phi] = \int d^3x \left[\frac{a_T^2}{8\pi G} F(Y) + \rho_b c^2 \phi \right]$$

$$Y = \frac{c^4 |\mathbf{A}|^2}{a_T^2}, \quad \mathbf{g} = -c^2 \mathbf{A}$$

Equation (12)

$$\nabla \cdot \left[\mu \left(\frac{|\mathbf{g}|}{a_T} \right) \mathbf{g} \right] = -4\pi G \rho_b, \quad \mu = F_Y$$

Equation (13)

In the static limit, Equation 13 belongs to the same mathematical family as AQUAL or MOND type modified gravity [1,2]. This paper does not claim the equation itself as an original invention. Its distinct contribution is to interpret this response as an Actual of spatial stress and a mass phenotype, and to connect it with stress at cosmological scales.

6.2 Deep low acceleration regime

$$F(Y) = \frac{2}{3} Y^{3/2} \Rightarrow \mu(x) = x, \quad g = \sqrt{a_T g_b}, \quad v_f^4 = G M_b a_T$$

Equation (14)

Under spherical symmetry or a sufficiently isolated circular orbit approximation, Equation 14 produces the forms of both the observed radial acceleration relation and the baryonic Tully Fisher relation [3,4]. Because one a_T enters both relations, the computational cost is lower than assigning a separate set of dark halo parameters to every galaxy. Fits that include nonsphericity, gas distribution, external fields, and distance errors must nevertheless be tested through a hierarchical analysis of the full SPARC data set.

6.3 Static stability

$$\lambda_{\perp} = \mu > 0, \quad \lambda_{\parallel} = \mu + x\mu' > 0$$

Equation (15)

In the deep regime, $\mu = x$ gives the two positive eigenvalues x and $2x$. The ellipticity of the static field equation is therefore preserved. This does not prove that every mode of the full covariant theory is stable. Cosmological scalar modes, the ultra long wavelength Jeans sector, and nonlinear defects require separate tests.

7 Connecting Cosmological and Galactic Scales

7.1 Matched response condition

The model imposes a condition that the cosmological background stress and the galactic low acceleration response share the same normalization under a three dimensional isotropic projection. This is called matched response with $C = 1$. The cosmological Hubble combination and stress fraction then determine the galactic transition acceleration.

$$a_T = cH_0 \sqrt{\frac{f_T}{8}}, \quad f_T = 8 \left(\frac{a_T}{cH_0} \right)^2$$

Equation (16)

7.2 Numerical test

Table 5 Numerical substitution in the cross scale relation

Ledger — Input or output · Value · Meaning

- Input or output: H_0
 - Value: 67.4 km s⁻¹ Mpc⁻¹
 - Meaning: Cosmological scale of relational change
- Input or output: f_T
 - Value: 0.265
 - Meaning: Cosmological fraction of effective dark stress
- Input or output: cH_0
 - Value: 6.55×10^{-10} m s⁻²
 - Meaning: Cosmological acceleration combination
- Input or output: a_T
 - Value: 1.192×10^{-10} m s⁻²
 - Meaning: Galactic low acceleration transition scale
- Input or output: Inferred f_T
 - Value: approximately 0.269
 - Meaning: Value obtained from $a_T = 1.20 \times 10^{-10}$ m s⁻²

The calculated a_T is close to the empirical acceleration scale of approximately 1.2×10^{-10} m s⁻² used in galactic dynamics [3]. Conversely, substituting that empirical value gives f_T of approximately 0.269. The meeting of quantities inferred at different scales within one relation is relevant to interpretive closure.

The value $f_T = 0.265$, however, is imported from an analysis performed within standard cosmology [5]. The agreement in Equation 16 is therefore not presented as a fully model independent prediction. It is a conditional derivation under the matched response assumption. Discriminating power requires a simultaneous fit to independent likelihoods from galactic and cosmological data.

8 Gravitational Lensing

8.1 One metric

For the mass phenotype to replace a conventional dark matter interpretation, it must affect the deflection of light as well as stellar motion. The model assumes that matter and light couple universally to one metric and uses two scalar potentials in the static weak field limit.

$$ds^2 = -\left(1 + \frac{2\Phi}{c^2}\right)c^2 dt^2 + \left(1 - \frac{2\Psi}{c^2}\right)d\mathbf{x}^2$$

Equation (17)

$$\hat{\alpha} = \frac{1}{c^2} \int \nabla_{\perp} (\Phi + \Psi) d\ell$$

Equation (18)

8.2 Lensing mass and dynamical mass

$$\Phi = \Psi + \mathcal{O}(c^{-2}) \Rightarrow \eta_{\text{slip}} = \frac{\Psi}{\Phi} = 1 + \mathcal{O}(c^{-2}), \quad M_{\text{lens}} = M_{\text{dyn}}$$

Equation (19)

If Equation 19 holds, twist stress is not a nonmetric correction that changes rotation curves while missing light. The same effective mass phenotype appears in material orbits and light deflection. The condition $\Phi = \Psi$ must nevertheless be established by the covariant completion. If anisotropic stress or additional degrees of freedom generate a large slip, agreement between lensing mass and dynamical mass may fail.

Colliding galaxy clusters provide a stronger test [6]. When the baryonic gas and lensing centers separate, the model must solve how the spatial stress field moves and persists dynamically. A static spherically symmetric equation cannot by itself explain that separation. Nonspherical lensing shear and postcollision centroid motion are priority numerical experiments for the model.

9 Early Universe and Structure Formation

9.1 Covariant computational structure

Carrying the static galactic response into the early universe requires a covariant action. This paper uses mathematics close to the relativistic Khronon theory of Blanchet and Skordis as computational material [7]. This section imports a covariant completion and is not an original derivation of that completion.

$$S = \frac{c^3}{16\pi G} \int d^4x \sqrt{-g} [R - 2J(Y) + 2K(Q)] + S_m$$

Equation (20)

$$n_{\mu} = -\frac{c}{Q} \nabla_{\mu} \tau, \quad Q = c \sqrt{-g^{\mu\nu} \nabla_{\mu} \tau \nabla_{\nu} \tau}$$

$$A_{\mu} = c^2 n^{\nu} \nabla_{\nu} n_{\mu}, \quad Y = \frac{A_{\mu} A^{\mu}}{c^4}$$

Equation (21)

The function J governs the spatial stress response in galaxies, while K governs the cosmological response that allows stress to cluster like an almost pressureless component. τ may be read as a relational ordering label for states, but K gives physical meaning to its normalization Q . This covariant completion is therefore not identical to a purely reparameterization invariant and timeless ontology. The difference remains explicitly in the model cost.

9.2 DBI type response and minimum numerical test

$$K(Q) = \frac{2\mu^2}{\lambda_D} \left[1 - \sqrt{1 - \lambda_D(Q - 1)^2} \right]$$

Equation (22)

$$q(a) = Q - 1 = [\lambda_D + (a^3/r)^2]^{-1/2}, \quad r = \frac{l_0}{2\mu^2}$$

Equation (23)

$$w = \frac{\lambda_D q^2 + (1+q)\sqrt{1 - \lambda_D q^2} - 1}{2 + (1 + \lambda_D)q}, \quad c_{\text{ad}}^2 = \frac{q(1 - \lambda_D q^2)}{1 + q}$$

Equation (24)

$$c_s^2(a, k) = c_{\text{ad}}^2 \left[1 + \frac{c_{\text{ad}}^2 k^2}{4\pi G a^2 \rho_T (1 + w)} \right]^{-1}, \quad \Sigma_T = 0$$

Equation (25)

Using $H_0 = 67.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, a present stress fraction of 0.265, $\mu^{-1} = 22.3 \text{ Mpc}$, and $\lambda_D = 1$, and normalizing the integration constant l_0 with the present density, gives $r = 9.9913 \times 10^{-6}$.

Table 6 Minimum numerical test of the DBI type response

Ledger — Regime · q · w · c ad squared

- Regime: Very early $a \ll 10^{-3}$
 - q: approaches 1
 - w: approaches 0
 - c ad squared: approaches 0
- Regime: Recombination $a = 1$ divided by 1100
 - q: 0.999999997
 - w: 3.76×10^{-5}
 - c ad squared: 2.83×10^{-9}
- Regime: Present $a = 1$
 - q: 9.99×10^{-6}
 - w: 5.00×10^{-6}
 - c ad squared: 9.99×10^{-6}
- Regime: Relational variable $a = 10$
 - q: 9.99×10^{-9}
 - w: 5.00×10^{-9}
 - c ad squared: 9.99×10^{-9}

Near recombination, w and the adiabatic sound speed are very small, and the wavenumber dependent physical sound speed is smaller still. The anisotropic viscosity is zero. The component may therefore behave like cold dark matter in the background and in linear perturbations. This test checks only a minimum

survival condition for early structure formation. It does not replace a joint likelihood analysis of the full CMB angular spectrum, BAO, matter power spectrum, and growth rate data.

10 Measurability and Failure Conditions

10.1 What must be measured

Measuring only the mass phenotype makes a particle halo difficult to distinguish from spatial stress. Discrimination requires either a nonlocal response of the stress Actual or an environmental dependence that differs from a particle halo. If the local gravitational field responds when a global boundary condition B boundary changes, the following functional derivative may be nonzero.

$$\frac{\delta \mathbf{g}(\mathbf{x})}{\delta B_{\text{boundary}}(\mathbf{y})} \neq 0$$

Equation (26)

No present technology is shown to relax spatial stress artificially, and no expected signal amplitude has been derived. Local stress release is therefore an experimental proposal rather than an experimental result. A more practical first step is to use natural differences in boundary conditions supplied by astronomical environments.

Table 7 Measurement paths that can distinguish the twist stress Actual

Ledger — Test · Observable · Requirement of the twist model · Status

- Test: Galactic environment
 - Observable: RAR residuals and external field
 - Requirement of the twist model: Systematic change correlated with boundary conditions
 - Status: Testable
- Test: Nonspherical lensing
 - Observable: Shear and dynamical mass
 - Requirement of the twist model: One stress distribution explains both
 - Status: Testable
- Test: Colliding clusters
 - Observable: Gas stellar and lensing centroids
 - Requirement of the twist model: Dynamically reproduce motion and persistence of the stress field
 - Status: Open critical test
- Test: Global topology
 - Observable: Repeated CMB correlations and matched circles
 - Requirement of the twist model: Observable trace of nonzero holonomy
 - Status: Open test
- Test: Structure growth
 - Observable: CMB BAO $f\sigma_8$ and power spectrum
 - Requirement of the twist model: Maintain small sound speed and allowed growth rate together
 - Status: Calculation required
- Test: Local manipulation

- Observable: Gravitational field change without inertial mass change
- Requirement of the twist model: Direct response to a stress boundary condition
- Status: Experimental proposal

10.2 Failure conditions

A minimal computation model must avoid failure under every declared condition. Success in one domain does not compensate for failure in another. If any of the following failures is structural and can be patched only by extra parameters, the present model must be rejected or replaced by a higher cost version.

- One response function fails to preserve both the RAR and the baryonic Tully Fisher relation across diverse galaxies.
- The covariant completion produces gravitational slip beyond observational limits and cannot explain lensing and dynamical masses together.
- In colliding galaxy clusters, the model fails if it cannot place the stress centre at the observed lensing centre while separating it from the baryonic gas centre where required by observations.
- The effective sound speed or an unstable early universe mode damages both the CMB and structure growth.
- Reproducing observed distance relations in a finite boundaryless interpretation requires a separate gluing rule for each target.
- The covariant completion necessarily introduces an independent temporal degree of freedom that conflicts with the relational interpretation of time.

$$\text{Accept}_{2.2} \iff \text{survive}\{\text{galaxy, lensing, CMB, structure}\}$$

Equation (27)

11 Claim Grades and Position of the Paper

Table 8 Claim grade ledger for twist stress cosmology

Ledger — Claim · Grade · Exact meaning

- Claim: WRRR Core 1.0
 - Grade: Preserved
 - Exact meaning: Not modified in this paper
- Claim: Only the present exists
 - Grade: Ontological interpretation
 - Exact meaning: Time is interpreted as a relation among change and records
- Claim: First twist
 - Grade: Origin interpretation
 - Exact meaning: The Big Bang is read as the opening of a relational network
- Claim: Finite boundaryless universe
 - Grade: Topological hypothesis
 - Exact meaning: No specific three dimensional manifold has been selected
- Claim: The dark Actual is spatial stress
 - Grade: Model choice

- Exact meaning: Particle dark matter is not required as an owner
- Claim: Dark mass is a phenotype
 - Grade: Theoretical redescription
 - Exact meaning: Modified geometry is observed as an effective density
- Claim: Modified galactic response
 - Grade: Existing mathematics
 - Exact meaning: AQUAL and MOND type static limit
- Claim: Cross scale relation
 - Grade: Conditional closure
 - Exact meaning: Matched response condition $C = 1$
- Claim: Agreement of lensing mass
 - Grade: Leading order result
 - Exact meaning: Holds when $\Phi = \Psi$
- Claim: DBI cosmology
 - Grade: Imported covariant completion
 - Exact meaning: Khronon mathematics is used as computational material
- Claim: Full CMB fit
 - Grade: Open test
 - Exact meaning: Requires a Boltzmann code and likelihood analysis
- Claim: Local stress release
 - Grade: Experimental proposal
 - Exact meaning: Implementation and signal amplitude remain unknown

The interpretive result of this paper is to join known phenomena within one Actual and Phenotype structure and to identify a minimum computation path without a fatal conflict across scales. A new discriminating prediction can strengthen the theory, but it does not by itself determine whether the interpretation is legitimate. Conversely, the absence of an independent prediction does not permit assumptions to be stated as facts. The claim grades guard against both errors.

12 Conclusion

Twist stress cosmology preserves the conventional account of ordinary matter and interprets only the additional gravity attributed to dark matter as the mass phenotype of stress stored in space. Electrical and optical expressions are assumed to be negligible, and no individual particle volume is required, but a gravitational expression exists because the stress changes the metric. Saying that the dark component is distributed throughout the universe, including our surroundings, means that a stress field has both background and local structure; it does not mean that particles fill space uniformly.

A Hodge decomposition of local fields and global topology permits a finite boundaryless universe to coexist with ordinary local gravitational fields. Bidirectional twist can preserve total stress while keeping net directional bias small. The Mobius strip is only an analogy for orientation reversal, not an exact boundaryless universe. A representative size of the universe cannot be reduced to one number until stiffness and holonomy are known independently.

The static low acceleration limit uses existing MOND type mathematics to recover the RAR and baryonic Tully Fisher relation. Under the matched response condition, cosmological H_0 and fT give $aT = 1.192 \times 10^{-10} \text{ m s}^{-2}$. With one metric and small slip, lensing mass and dynamical mass agree at leading order. A covariant DBI type response produces small pressure and sound speed near recombination, but full CMB

and colliding cluster tests remain open.

This paper therefore does not declare a final theory superior to standard cosmology. Our universe is not special, and survival precedes any ranking of best or worst. The present calculation shows that one owner, twist stress, may avoid failure in galaxies, lensing, and the early universe. The next steps are a joint SPARC fit, a Boltzmann implementation of covariant perturbations, stress dynamics in colliding clusters, and a search for global holonomy. If the model fails those tests, it must be discarded or revised with the added cost made explicit. This is the closure required by Minimal Computing Cosmology.

Source archives and reproduction paths

The package's `evidence/` folder contains the unchanged originals below. Author lists, exact titles and existing DOIs remain those on each original title page and README. No new DOI is assigned to this book revision. Editorial checks are distinct from the scientific execution records in the sources.

ID	File in evidence/
B	WRRR_3_0_Paper_v1.8_Book_v0.10_Final.zip
P1	WRRR_M_r11_FoP_Submission_EN.docx
P2	WRRR_Address_Particle_Interface_v0_2_r1_Package.zip
P3	WRRR_Dynamic_Decays_v0_2_r1_Reviewed_Archive.zip
P4	WRRR_Carrier_Coupling_v0_2_Submission_Package.zip
P5	WRRR_Energy_Pressure_v0_2_Submission_Package.zip
P6	WRRR_Information_Weight_v0_3_Complete_Package.zip
P7	WRRR_Record_Capacity_v0.2_Source.zip

P7: `manuscript_en.md`, `manuscript_ko.md`, `reproduce.py`, `results.json`, `REVISION_NOTES.md`. B: nested `WRRR_3_0_Reproduction_v1.8.tar.xz` and its `RELEASE_MAP.json`.