WRRA Macroscopic Prediction Cases - Documentation Supplement v0.3 Wonsik Choi and Jeongin Choi | Independent Researchers | 15 October 2026

Status: This supplement adds a case registry and computation specifications to the frozen Shutter-SOURCE Three Targets v0.2. All original files are preserved byte-for-byte. No new vertical-kinematics, galaxy-lensing or growth-rate numbers are claimed. Existing outputs and proposed calculations are distinguished.

EXISTING MACROSCOPIC OUTPUTS Validation inputs: inherited 128-state SOURCE, static preparation, declared relative-path coupling g=1 and 1064 nm wavelength. WRRA transformation: exact finite-state average of relative optical phases. Outputs: visibility 0.3586975046 at the first minimum, 21.780958380 m; 0.6081539566 at partial recovery, 37.278764320 m; distance ratio 1.7115300286. Refutation: disagreement with the fixed preparation, exposure and visibility curve refutes that branch. Exposure distance is not automatically device size. Conclusion: an existing conditional macroscopic optical calculation.

Validation inputs for existing galactic rotation: simplified McMillan baryons, SPARC-calibrated response and fixed a_t=1.1918e-10 m/s^2. WRRA transformation: baryonic field to twist response to circular speed. Outputs from the 27 September 2026 v1.0 manuscript, not rerun here: 226.3 km/s at 8.2 kpc, 201.6 km/s at 25 kpc, RMS 2.42 km/s against the linearized Eilers benchmark over 8-25 kpc. This is not an individual-data likelihood result. Refutation: failure of the universal response or of the fixed cross-observable closure requires revision. Conclusion: inherited astronomical-scale outputs.

M1 - VERTICAL DISK KINEMATICS Validation inputs: fixed rotation-calibrated state, baryonic distribution, tracer density, selection function and explicitly declared boundary conditions. WRRA transformation: direction [n], rho_t and anisotropy q_t to Phi(R,z), then K_z=-dPhi/dz and vertical Jeans evolution including the velocity-tilt term. Outputs sought: K_z(R,z), sigma_z(R,z), thickness and flaring. Pending. The v1.0 manuscript explicitly leaves the q_t closure unspecified. A rotation curve alone does not uniquely determine the off-plane gravitational field. Refutation: failure of the declared fixed 3D state to explain rotation and vertical data, or the need for independent height-by-height tuning. Conclusion: registered candidate; establish the spatial-stress closure before computing. Calibration is allowed and its use of validation data is recorded.

M2 - LENSING FROM THE SAME GALACTIC STATE Validation inputs: the fixed 3D stress state, source/lens distances and geometry. WRRA transformation: derive or declare the metric potentials Phi and Psi from the same state; alpha_hat=(1/c^2) integral grad_perp(Phi+Psi) dl in the weak-field thin-lens approximation, followed by lens geometry. If Phi=Psi is adopted, state that GR closure explicitly; anisotropic stress need not imply equality. Outputs sought: deflection, convergence, shear and a conditional Einstein angle. Pending. A spherical equivalent mass inferred from rotation is not by itself a spatial stress distribution or an optical metric. No lensing number from a different prior model is relabeled as this galaxy's prediction. Refutation: failure of the fixed stress/metric state across kinematics and lensing, or the need for independent lensing amplitudes. Conclusion: registered candidate. Integration is credited after actual state-to-deflection computation, rather than citation of an external formula.

M3 - COSMIC STRUCTURE GROWTH Validation inputs: fixed background, composition, address weights, shutter, SOURCE preparation, initial perturbations and declared tracer treatment. WRRA transformation: SOURCE perturbations to conserved sector stress/exchange, metric perturbations and matter transfer; D(k,a) gives f=d ln D/d ln a; P_m(k,z) gives sigma_8(z) with an 8 h^-1 Mpc window and f sigma_8(z). Outputs sought: P_m(k,z), scale-dependent growth and f sigma_8(z). Pending. Seven cached CMB wavenumber transfers and their computed higher-order moments are not a substitute for the matter spectrum on a sufficient wavenumber grid. Refutation: conservation/constraint failure or disagreement of the fixed growth branch with specified observations; redshift-by-redshift retuning is recorded as failure of that fixed version. Conclusion: registered candidate; run the required matter-transfer calculation.

COMMON ASSESSMENT Order: validation inputs -> WRRA transformation -> outputs -> refutation -> conclusion. Verified constants may calibrate the model. Reproduction of known values counts as empirical consistency and explanatory performance. Outputs computed after freezing a calibrated model are attributed to that WRRA model; matching another theory's number does not remove attribution. The inherited 176 controls do not validate the three newly registered physical predictions.

SOURCES Shutter-SOURCE Three Targets v0.2, package-listed DOI 10.5281/zenodo.23380920. Galactic disk v1.0 manuscript, Wonsik Choi, 27 September 2026, sections 7.1, 7.4, 9 and 10. No claim is made that later public 3D closures were audited here. External cross-observable testing method: https://arxiv.org/abs/1812.08169. Live publication status must be verified at the record page rather than inferred from a reserved DOI embedded in archived files.

REVIEW REVISION v0.3-r1 Validation inputs: 11 frozen input files and the inherited actual CAMB transfers. WRRA transformation: rerun supply, photons, cmb_higher and review. Outputs: 39+16+65+56=176/176 controls passed. Stable numerical observables were compared with the frozen numerical reference. M1-M3 outputs remain pending. Refutation: numerical/reference, hash or ledger failure blocks this release. Conclusion: inherited calculation reproduced; documentation corrected.

M1: Validate the stationary/axisymmetric approximation, particularly for a warped or disequilibrium disk. If it fails, include time evolution. Declare covariance and acceptance criteria before the observational validation. M2: alpha_hat is the physical lens-plane deflection. Distinguish it from observed angles and specify angular-diameter-distance conventions and D_ls/D_s. A no-slip Phi=Psi closure is an explicit assumption, not an automatic derived result. M3: Scale-dependent growth need not admit a universal scalar f sigma_8. Use the survey wavenumber window and tracer model to define the effective estimator, or compare the calculated P_m(k,z) directly. Publication audit: the 23380920 record was a reserved draft when inspected before this release. Its public status is established by the live record and a separate publication receipt, not by the DOI printed in inherited files.