MAT185 Linear Algebra Assignment 1

Instructions:

Please read the MAT185 Assignment Policies & FAQ document for details on submission policies, collaboration rules and academic integrity, and general instructions.

- 1. Submissions are only accepted by Gradescope. Do not send anything by email. Late submissions are not accepted under any circumstance. Remember you can resubmit anytime before the deadline.
- 2. Submit solutions using only this template pdf. Your submission should be a single pdf with your full written solutions for each question. If your solution is not written using this template pdf (scanned print or digital) then your submission will not be assessed. Organize your work neatly in the space provided. Do not submit rough work.
- 3. Show your work and justify your steps on every question but do not include extraneous information. Put your final answer in the box provided, if necessary. We recommend you write draft solutions on separate pages and afterwards write your polished solutions here on this template.
- 4. You must fill out and sign the academic integrity statement below; otherwise, you will receive zero for this assignment.

Academic Integrity Statement:

| Full Name: JEON, Woohyun (Gordon) |
|-----------------------------------|
| Student number: 1008990451 |
| Full Name: ZHENG, Haoyan (Harvey) |

Student number: 1008715516

I confirm that:

- I have read and followed the policies described in the document MAT185 Assignment Policies & FAQ.
- In particular, I have read and understand the rules for collaboration, and permitted resources on assignments as described in subsection II of the aforementioned document. I have not violated these rules while completing and writing this assignment.
- I understand the consequences of violating the University's academic integrity policies as outlined in the Code of Behaviour on Academic Matters. I have not violated them while completing and writing this assignment.

By signing this document, I agree that the statements above are true.

Signatures: 1) Gordon Jeon

2) Harvey Zheng

1. Let $V = \{(x_1, x_2) \mid x_1, x_2 > 0, \text{ and } x_1 + x_2 = 1\}$. Is V a real vector space with respect to the usual entry-wise vector addition and scalar multiplication? Why or why not?

No. Axioms, AIII and AIV (Medici, p104) fail to hold in V.

AIII states, Zero: There exists a zero or null vector $0 \in V$ s.t. $\mathbf{u} + \mathbf{0} = \mathbf{u}$ (Medici pp104)

Therefore, by definition, there must be a zero or null vector, $\theta \in V$ s.t. $(x_1, x_2) + 0 = (x_1, x_2)$.

Accordingly, for the previous statement to be true, the zero vector must have values (0,0):

$$(x_1, x_2) + 0 = (x_1, x_2)$$

$$(x_1, x_2) + (0, 0) = (x_1, x_2)$$

$$(x_1 + 0, x_2 + 0) = (x_1, x_2)$$

$$(x_1, x_2) = (x_1, x_2)$$

However, $x_1, x_2 > 0$ in V, therefore, the real vector space V does not contain a zero vector, proving AIII does not hold in V.

AIV states, Negative: There exists a negative $-u \in Vs.t \ u + (-u) = 0$ (Medeci pp104)

Therefore, by definition, for an inverse vector to exist, the sum of the vector and itself needs to be the zero vector. However, as seen above, V does not contain a zero vector. Therefore, V fails in AIV.

2 Let $V = \{(x_1, x_2) \mid x_1, x_2 > 0, \text{ and } x_1 + x_2 = 1\}$. Define vector addition in V by

$$(x_1, x_2) + (y_1, y_2) = \frac{(x_1y_1, x_2y_2)}{x_1y_1 + x_2y_2}$$

and scalar multiplication in V by

$$c(x_1, x_2) = \frac{(x_1^c, x_2^c)}{x_1^c + x_2^c}$$

Then V is a real vector space.

(a) Verify that axiom AIII. (Medici, pp104) holds in V.

To verify AIII, you need the zero vector. Therefore, consider the following:

$$c(x_1, x_2) = \frac{(x_1^c, x_2^c)}{x_1^c + x_2^c}$$

In order for the expression to equal to 0, set c = 0, as values $x_1, x_2 > 0$ by definition:

$$c(x_1, x_2) = \frac{(x_1^c, x_2^c)}{x_1^c + x_2^c}, \ c = 0$$

$$= \frac{(x_1^0, x_2^0)}{x_1^0 + x_2^0}$$

$$= \frac{(1, 1)}{1 + 1}$$

$$= (\frac{1}{2}, \frac{1}{2})$$

$$\therefore (x_1, x_2) = (\frac{1}{2}, \frac{1}{2}) \ s.t \ c = 0$$

As shown above $(x_1, x_2) = (\frac{1}{2}, \frac{1}{2})$ and as per definition $\frac{1}{2} > 0$, therefore AIII holds in V.

(b) Verify that axiom AIV. (Medici, pp104) holds in V.

To verify AIV, there must exist a negative $-u \in V$ s.t. u + (-u) = 0. To find the zero vector, consider the following:

$$c(x_1, x_2) = \frac{(x_1^c, x_2^c)}{x_1^c + x_2^c}$$

In order for the expression to become the inverse of (x_1, x_2) , it must have values $(-x_1, -x_2)$ or $(-1)(x_1, x_2)$ (MIII, Medici, pp104) equivalently. Therefore, verify the scalar multiplication:

$$\begin{split} c(x_1,x_2) &= -1(x_1,x_2) \\ &= \frac{(x_1^{-1},x_2^{-1})}{x_1^{-1}+x_2^{-1}}, \ by \ definition \\ &= 1 \cdot \frac{(x_1^{-1},x_2^{-1})}{x_1^{-1}+x_2^{-1}}, \ by \ MIV \\ &= \frac{x_1,x_2}{x_1x_2} \cdot \frac{(\frac{1}{x_1},\frac{1}{x_2})}{\frac{1}{x_1}+\frac{1}{x_2}}, \ by \ Q.E.D \\ &= \frac{(x_2,x_1)}{x_2+x_1}, \ by \ MIII \\ &= \frac{(x_2,x_1)}{1}, \ by \ definition \\ &= (x_2,x_1) \end{split}$$

2. Let $V = \{(x_1, x_2) \mid x_1, x_2 > 0, \text{ and } x_1 + x_2 = 1\}$. Define vector addition in V by

$$(x_1, x_2) + (y_1, y_2) = \frac{(x_1y_1, x_2y_2)}{x_1y_1 + x_2y_2}$$

and scalar multiplication in V by

$$c(x_1, x_2) = \frac{(x_1^c, x_2^c)}{x_1^c + x_2^c}$$

Then V is a real vector space.

(c) Verify that axiom MIII. (Medici, pp104) holds in V. MIII states, Distributivity: (a) $(\alpha + \beta)\mathbf{u} = \alpha \mathbf{u} + \beta \mathbf{u}$ and (b) $\alpha(\mathbf{u} + \mathbf{v}) = \alpha \mathbf{u} + \alpha \mathbf{v}$ (Medici pp104)

To verify MIII, both (a) and (b) need to pass concurrently, consider the following:

$$(x_1,x_2) + (y_1,y_2) = \frac{(x_1y_1,x_2y_2)}{x_1y_1 + x_2y_2}, \ by \ definition$$

$$= (\frac{x_1y_1}{x_1y_1 + x_2y_2}, \frac{x_2y_2}{x_1y_1 + x_2y_2}), \ by \ definition$$

$$a[(x_1,x_2),(y_1,y_2)] = a(\frac{x_1y_1}{x_1y_1 + x_2y_2}, \frac{x_2y_2}{x_1y_1 + x_2y_2}), \ s.t. \ a \ is \ a \ scalar$$

$$a(x_1,x_2) + a(y_1,y_2) = (\frac{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}), \ by \ definition$$

$$(\frac{x_1^a, x_2^a}{x_1^a + x_2^a}) + (\frac{y_1^a, y_2^a}{y_1^a + y_2^a}) = (\frac{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}), \ by \ definition$$

$$(\frac{x_1^a}{x_1^a + x_2^a}, \frac{x_2^a}{x_1^a + x_2^a}) + (\frac{y_1^a}{y_1^a + y_2^a}, \frac{y_2^a}{y_1^a + y_2^a}) = (\frac{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{x_2y_2}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}), \ by \ definition$$

$$(\frac{(\frac{x_1y_1)^a}{(x_1^a + x_2^a)(y_1^a + y_2^a)}, \frac{(\frac{x_2y_2}{x_1}}{(x_1^a + x_2^a)(y_1^a + y_2^a)}}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}), \ by \ definition$$

$$(\frac{(\frac{x_1y_1)^a}{(x_1^a + x_2^a)(y_1^a + y_2^a)}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}), \ by \ definition$$

$$(\frac{(\frac{x_1y_1)^a}{(\frac{x_1y_1)^a}{(\frac{x_1y_1)^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}}), \ by \ definition$$

$$(\frac{(\frac{x_1y_1)^a}{(\frac{x_1y_1)^a}{(\frac{x_1y_1)^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})}, \frac{(\frac{x_2y_2}{x_1y_1 + x_2y_2})^a}{(\frac{x_1y_1}{x_1y_1 + x_2y_2})^a}}), \ by \ definition$$

Here, analyze the left-hand side (LHS) and right-hand side (RHS):

$$\begin{pmatrix} \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)}, \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \\ \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} = \begin{pmatrix} \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \\ \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix}$$

$$= \begin{pmatrix} \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \\ \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} \begin{pmatrix} \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \\ \frac{(x_1y_1)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} + \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} \begin{pmatrix} \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} \begin{pmatrix} \frac{(x_2y_2)^a}{(x_1^a+x_2^a)(y_1^a+y_2^a)} \end{pmatrix} \begin{pmatrix} \frac{(x_1y_1)^a}{(x_1y_1)^a+(x_2y_2)^a} \end{pmatrix} \begin{pmatrix} \frac{(x_2y_2)^a}{(x_1y_1)^a+(x_2y_2)^a} \end{pmatrix} \begin{pmatrix} \frac{(x_1y_1)^a}{(x_1y_1+x_2y_2)^a} \end{pmatrix} \begin{pmatrix}$$

As seen above the LHS and RHS are equivalent, proving MIII holds in V.

- **3.** Recall that $P_3(\mathbb{R}) = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, a_1, a_2, a_3 \in \mathbb{R}\}$, the set of polynomials of degree at most 3 with real coefficients, is a real vector space with respect to the usual polynomial addition and scalar multiplication.
- (a) Give an example of a subset S of $P_3(\mathbb{R})$ that is closed under vector addition but not under scalar multiplication. You should both state clearly your subset S and demonstrate that S satisfies the requirement of the question.

Consider,
$$S = \{a_1 x, a_2 x^3 | a_1, a_2 \in R\}$$

For closure in scalar multiplication, $a_1 \cdot a_2$ is always in the space of the subset S.

For closure in vector addition, the result is not always in the space of subset S.

Proof by counter-example:

 $a_0x \cdot a_1x^3 = (a_0 \cdot a_1)x^4$, which is outside of the highest power of 3 for the variable x defined in subset S.

- **3.** Recall that $P_3(\mathbb{R}) = \{a_0 + a_1x + a_2x^2 + a_3x^3 \mid a_0, a_1, a_2, a_3 \in \mathbb{R}\}$, the set of polynomials of degree at most 3 with real coefficients, is a real vector space with respect to the usual polynomial addition and scalar multiplication.
- (b) Give an example of a subset S of $P_3(\mathbb{R})$ that is closed under scalar multiplication but not under vector addition. You should both state clearly your subset S and demonstrate that S satisfies the requirement of the question.

Consider,
$$S = \{a_1, a_2 | \frac{\sqrt{2}}{2} < a_1 < 1, \frac{\sqrt{2}}{2} < a_2 < 1\}$$

For closure in scalar multiplication, $a_1 \cdot a_2$ is always less than 1, which satisfies the definition of subset S.

For closure in vector addition, the result is always greater than one, outside of the definition of subset S.