

# Experience-Dependent Neuroplasticity, Psychotherapy, and Long-Term Behavioral Change: A Synthesis

## 1. Introduction

Experience-dependent neuroplasticity refers to the brain's ability to reorganize its structure and function in response to life experiences, learning, and environmental influences. Psychotherapy leverages this plasticity to facilitate enduring changes in cognition, emotion, and behavior, offering hope for lasting recovery from mental health disorders (Beauregard, 2014; Kleim & Jones, 2008; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024). Recent research demonstrates that various forms of psychotherapy—including cognitive-behavioral therapy (CBT), mindfulness-based interventions, and psychodynamic therapy—can induce measurable neuroplastic changes in brain regions implicated in emotion regulation, self-referential processing, and social cognition (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Gkintoni et al., 2025; Piotrkowicz et al., 2021; Barsaglini et al., 2014). These neural adaptations are associated with symptom reduction and improved well-being, supporting the view that psychotherapy's effectiveness is rooted in its capacity to harness and direct neuroplastic processes for long-term behavioral change (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024). This review synthesizes the mechanisms, evidence, and implications of experience-dependent neuroplasticity in psychotherapy.

## 2. Methods

A comprehensive search was conducted across over 170 million research papers in Consensus, including Semantic Scholar, PubMed, and other databases. The search strategy targeted foundational models, neural mechanisms, clinical outcomes, and methodological advances linking experience-dependent neuroplasticity, psychotherapy, and long-term behavioral change. In total, 1027 papers were identified, 596 were screened, 360 were deemed eligible, and the top 50 most relevant papers were included in this review.

### Search Strategy




FIGURE 1 Flow of papers through the search and selection process.

Eight unique search groups were used, spanning neurobiological mechanisms, clinical interventions, long-term outcomes, and interdisciplinary perspectives.

### 3. Results

#### 3.1 Mechanisms of Experience-Dependent Neuroplasticity

Neuroplasticity encompasses synaptic remodeling, dendritic growth, neurogenesis, and changes in functional connectivity, all of which are modulated by experience and learning (Kleim & Jones, 2008; Sweatt, 2016; Voss et al., 2017; Socodato et al., 2023; Turner et al., 2017). Psychotherapy can induce these changes by providing new experiences, challenging maladaptive patterns, and fostering adaptive behaviors (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024). For example, CBT for social anxiety disorder has been shown to reduce amygdala volume and responsivity, correlating with decreased anxiety (Månsson et al., 2016). Similarly, long-term psychodynamic therapy normalizes activity in prefrontal-limbic circuits implicated in emotional regulation (Buchheim et al., 2012).

#### 3.2 Psychotherapy as a Driver of Neuroplastic Change

Functional and structural neuroimaging studies reveal that psychotherapy leads to normalization or reorganization of brain activity in regions such as the prefrontal cortex, amygdala, hippocampus, and cingulate cortex (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Weingarten & Strauman, 2015). These changes are associated with improvements in symptoms, cognitive flexibility, and emotional regulation (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024). Interpersonal and social processes, such as inter-brain synchrony between therapist and patient, have also been proposed as mechanisms of change, with evidence suggesting that increased synchrony predicts symptom reduction and well-being (Sened et al., 2025; Sened et al., 2022).

#### 3.3 Long-Term Behavioral Outcomes

Meta-analyses and longitudinal studies indicate that psychotherapy can produce enduring behavioral change, with effects persisting for months or years after treatment (Andersson et al., 2018; Cuijpers et al., 2021; Kline et al., 2018; Perry & Bond, 2012; Wilkinson et al., 2019; Gkintoni et al., 2025). The durability of these effects is linked to the degree and nature of neuroplastic changes induced during therapy (Månsson et al., 2016; Buchheim et al., 2012; Mason et al., 2017; Piotrkowicz et al., 2021). For example, changes in brain connectivity following CBT for psychosis predict recovery trajectories up to eight years post-treatment (Mason et al., 2017).

#### 3.4 Biological Markers and Modulators

Molecular and cellular markers such as brain-derived neurotrophic factor (BDNF) and epigenetic modifications have been implicated in psychotherapy-induced neuroplasticity (Piotrkowicz et al., 2021; Price & Duman, 2019; Olson, 2022). Pharmacological and somatic interventions (e.g., ketamine, psychedelics, brain stimulation) can enhance neuroplasticity, potentially creating windows of opportunity for psychotherapy to consolidate adaptive changes (Wilkinson et al., 2019; Calder & Hasler, 2022; Olson, 2022; Kupferberg & Hasler, 2024). Social and environmental factors, including support networks and enriched environments, further modulate neuroplastic potential and therapeutic outcomes (Song, 2024; McEwen, 2016; Simpkins & Simpkins, 2013).

**Key Papers**

Paper	Methodology	Population/Context	Key Results
(Månsson et al., 2016)	Multimodal neuroimaging, RCT	Social anxiety disorder	CBT reduces amygdala volume/responsivity, correlating with anxiety reduction
(Buchheim et al., 2012)	fMRI, longitudinal	Major depression	Long-term psychodynamic therapy normalizes prefrontal-limbic function
(Piotrkowicz et al., 2021)	Systematic review	Various disorders	Psychotherapy can increase BDNF, decrease gene methylation, supporting neuroplasticity
(Beauregard, 2014)	Review	Depression, anxiety	Psychotherapy normalizes brain activity, supporting neural mechanisms of change
(Sened et al., 2025)	fNIRS, proof-of-concept	Test anxiety	Inter-brain synchrony increases over therapy, associated with symptom reduction

FIGURE 2 Comparison of key studies on neuroplasticity, psychotherapy, and behavioral change.

**Top Contributors**

Type	Name	Papers
Author	G. Andersson	(Andersson et al., 2018; Cuijpers et al., 2021)
Author	A. Buchheim	(Buchheim et al., 2012)
Author	M. Beauregard	(Beauregard, 2014)
Journal	<i>Translational Psychiatry</i>	(Månsson et al., 2016; Mason et al., 2017; Alves et al., 2017; Ho & King, 2021)
Journal	<i>Journal of Clinical Medicine</i>	(Gkintoni et al., 2025; Piotrkowicz et al., 2021)
Journal	<i>World Psychiatry</i>	(Hayes & Hofmann, 2021; Cuijpers et al., 2021)







FIGURE 3 Authors & journals that appeared most frequently in the included papers.

**4. Discussion**

The evidence strongly supports a bidirectional relationship between experience-dependent neuroplasticity and psychotherapy: psychotherapy leverages neuroplastic mechanisms to drive long-term behavioral change, while neuroplasticity underpins the brain's capacity to adapt to therapeutic experiences (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024). The normalization of dysfunctional brain circuits, increases in neurotrophic factors, and changes in gene expression all contribute to sustained improvements in mental health (Månsson et al., 2016; Buchheim et al., 2012; Piotrkowicz et al., 2021; Price & Duman, 2019; Olson, 2022). However, the magnitude and durability of these changes depend on factors such as therapy type, individual neurobiological differences, social context, and the presence of supportive environments (Song, 2024; Mcewen, 2016; Simpkins & Simpkins, 2013; Voss et al., 2017).

Despite advances, challenges remain in identifying specific mediators and moderators of change, standardizing measurement of neuroplastic outcomes, and translating findings across diverse populations and disorders (Lemmens et al., 2016; Bruijnicks et al., 2021; Weingarten & Strauman, 2015). The integration of neuroimaging, molecular, and behavioral data is essential for refining models of therapeutic change and developing personalized interventions (Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Wilkinson et al., 2019; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024).

### Claims and Evidence Table

Claim	Evidence Strength	Reasoning	Papers
Psychotherapy induces experience-dependent neuroplasticity in brain regions linked to emotion and cognition	 Strong	Robust neuroimaging and molecular evidence across disorders and therapies	(Beauregard, 2014; Månsson et al., 2016; Buchheim et al., 2012; Barsaglini et al., 2014; Piotrkowicz et al., 2021; Price & Duman, 2019; Gkintoni et al., 2025; Song, 2024)
Neuroplastic changes are associated with long-term behavioral improvements	 Strong	Longitudinal and meta-analytic studies show durable effects linked to neural adaptation	(Andersson et al., 2018; Cuijpers et al., 2021; Mason et al., 2017; Kline et al., 2018; Perry & Bond, 2012; Månsson et al., 2016; Buchheim et al., 2012; Piotrkowicz et al., 2021)
Interpersonal synchrony and social context modulate neuroplasticity and outcomes	 Moderate	Emerging evidence from inter-brain studies and social neuroscience	(Sened et al., 2025; Sened et al., 2022; Song, 2024; McEwen, 2016; Simpkins & Simpkins, 2013)
Molecular markers (e.g., BDNF, gene methylation) reflect psychotherapy-induced plasticity	 Moderate	Systematic reviews and biomarker studies support this link, but findings are mixed	(Piotrkowicz et al., 2021; Price & Duman, 2019; Olson, 2022)
Pharmacological and somatic interventions can enhance neuroplasticity, augmenting psychotherapy	 Moderate	Evidence from combined treatment studies and animal models	(Wilkinson et al., 2019; Calder & Hasler, 2022; Olson, 2022; Kupferberg & Hasler, 2024)
Mechanisms of change in psychotherapy are complex and multifactorial	 Moderate	Mechanism research is heterogeneous; causal pathways remain difficult to isolate	(Lemmens et al., 2016; Bruijnicks et al., 2021; Weingarten & Strauman, 2015; Hayes & Andrews, 2020; Penedo et al., 2020)

**FIGURE** Key claims and support evidence identified in these papers.

## 5. Conclusion

Experience-dependent neuroplasticity is a foundational mechanism by which psychotherapy produces long-term behavioral change. Psychotherapy induces structural and functional brain changes, modulates molecular pathways, and leverages social and environmental factors to promote adaptive outcomes. While the evidence is robust, further research is needed to clarify mediators, optimize interventions, and personalize treatment.

## 5.1 Research Gaps

Key gaps include the need for standardized neuroplasticity measures, understanding individual variability in plasticity and response, and elucidating the interplay between biological, psychological, and social mechanisms.

### Research Gaps Matrix

Topic/Attribute	Neuroimaging	Molecular Markers	Social/Interpersonal	Longitudinal Outcomes	Combined Interventions
Depression/Anxiety	8	5	3	7	4
Psychosis/Severe Mental Illness	3	1	1	2	1
Social/Interpersonal Mechanisms	2	GAP	4	1	GAP
Combined Pharmacological/Psychotherapy	2	2	1	1	3

FIGURE Distribution of research across topics and study attributes, highlighting underexplored areas.

## 5.2 Open Research Questions

Future research should focus on integrating neurobiological, psychological, and social data to refine models of change and develop targeted, personalized interventions.

Question	Why
What are the specific neurobiological mediators of long-term behavioral change in psychotherapy?	Identifying these mediators will enable more precise and effective interventions for diverse populations.
How do individual differences in neuroplasticity influence psychotherapy outcomes?	Understanding variability can inform personalized treatment and improve efficacy.
Can combining neuroplasticity-enhancing interventions with psychotherapy optimize long-term outcomes?	Synergistic approaches may yield greater and more durable improvements in mental health.

FIGURE Key open questions for advancing research on neuroplasticity and psychotherapy.

In summary, psychotherapy leverages experience-dependent neuroplasticity to produce lasting behavioral change, but further research is needed to fully harness and individualize its transformative potential.

These papers were sourced and synthesized using Consensus, an AI-powered search engine for research. Try it at <https://consensus.app>

## References

Sened, H., Kaduri, K., Gamliel, H., Rafaeli, E., Zilcha-Mano, S., & Shamay-Tsoory, S. (2025). Inter-brain plasticity as a mechanism of change in psychotherapy: A proof of concept focusing on test anxiety.. *Psychotherapy research : journal of the Society for Psychotherapy Research*, 1-15. <https://doi.org/10.1080/10503307.2025.2451798>

- Sened, H., Zilcha-Mano, S., & Shamay-Tsoory, S. (2022). Inter-brain plasticity as a biological mechanism of change in psychotherapy: A review and integrative model. *Frontiers in Human Neuroscience*, 16. <https://doi.org/10.3389/fnhum.2022.955238>
- Calder, A., & Hasler, G. (2022). Towards an understanding of psychedelic-induced neuroplasticity. *Neuropsychopharmacology*, 48, 104 - 112. <https://doi.org/10.1038/s41386-022-01389-z>
- Song, Z. (2024). A contemporary analysis of the effects of neuroplasticity on well-being. *Armenian Journal of Health & Medical Sciences*. <https://doi.org/10.54235/27382737-2024.v4.1-46>
- Beauregard, M. (2014). Functional neuroimaging studies of the effects of psychotherapy. *Dialogues in Clinical Neuroscience*, 16, 75 - 81. <https://doi.org/10.31887/DCNS.2014.16.1/mbeauregard>
- Socodato, R., Almeida, T., Portugal, C., Santos, E., Tedim-Moreira, J., Galvão-Ferreira, J., Canedo, T., Baptista, F., Magalhães, A., Ambrósio, A., Brakebusch, C., Rubinstein, B., Moreira, I., Summavielle, T., Pinto, I., & Relvas, J. (2023). Microglial Rac1 is essential for experience-dependent brain plasticity and cognitive performance.. *Cell reports*, 42 12, 113447. <https://doi.org/10.1016/j.celrep.2023.113447>
- McEwen, B. (2016). In pursuit of resilience: stress, epigenetics, and brain plasticity. *Annals of the New York Academy of Sciences*, 1373. <https://doi.org/10.1111/nyas.13020>
- Wilkinson, S., Holtzheimer, P., Gao, S., Kirwin, D., & Price, R. (2019). Leveraging Neuroplasticity to Enhance Adaptive Learning: The Potential for Synergistic Somatic-Behavioral Treatment Combinations to Improve Clinical Outcomes in Depression. *Biological Psychiatry*, 85, 454-465. <https://doi.org/10.1016/j.biopsych.2018.09.004>
- Olson, D. (2022). Biochemical Mechanisms Underlying Psychedelic-Induced Neuroplasticity.. *Biochemistry*. <https://doi.org/10.1021/acs.biochem.1c00812>
- Kleim, J., & Jones, T. (2008). Principles of experience-dependent neural plasticity: implications for rehabilitation after brain damage.. *Journal of speech, language, and hearing research : JSLHR*, 51 1, S225-39. [https://doi.org/10.1044/1092-4388\(2008/018\)](https://doi.org/10.1044/1092-4388(2008/018))
- Gkintoni, E., Vassilopoulos, S., & Nikolaou, G. (2025). Mindfulness-Based Cognitive Therapy in Clinical Practice: A Systematic Review of Neurocognitive Outcomes and Applications for Mental Health and Well-Being. *Journal of Clinical Medicine*, 14. <https://doi.org/10.3390/jcm14051703>
- Price, R., & Duman, R. (2019). Neuroplasticity in cognitive and psychological mechanisms of depression: An integrative model. *Molecular psychiatry*, 25, 530 - 543. <https://doi.org/10.1038/s41380-019-0615-x>
- Månsson, K., Salami, A., Frick, A., Carlbring, P., Andersson, G., Furmark, T., & Boraxbekk, C. (2016). Neuroplasticity in response to cognitive behavior therapy for social anxiety disorder. *Translational Psychiatry*, 6. <https://doi.org/10.1038/tp.2015.218>
- Hayes, S., & Hofmann, S. (2021). “Third-wave” cognitive and behavioral therapies and the emergence of a process-based approach to intervention in psychiatry. *World Psychiatry*, 20. <https://doi.org/10.1002/wps.20884>
- Andersson, G., Rozental, A., Shafan, R., & Carlbring, P. (2018). Long-term effects of internet-supported cognitive behaviour therapy. *Expert Review of Neurotherapeutics*, 18, 21 - 28. <https://doi.org/10.1080/14737175.2018.1400381>
- Cuijpers, P., Quero, S., Noma, H., Ciharova, M., Miguel, C., Karyotaki, E., Cipriani, A., Cristea, I., & Furukawa, T. (2021). Psychotherapies for depression: a network meta-analysis covering efficacy, acceptability and long-term outcomes of all main treatment types. *World Psychiatry*, 20. <https://doi.org/10.1002/wps.20860>
- Mason, L., Peters, E., Peters, E., Williams, S., Kumari, V., & Kumari, V. (2017). Brain connectivity changes occurring following cognitive behavioural therapy for psychosis predict long-term recovery. *Translational Psychiatry*, 7. <https://doi.org/10.1038/tp.2016.263>
- Kline, A., Cooper, A., Rytwinski, N., & Feeny, N. (2018). Long-term efficacy of psychotherapy for posttraumatic stress disorder: A meta-analysis of randomized controlled trials.. *Clinical psychology review*, 59, 30-40. <https://doi.org/10.1016/j.cpr.2017.10.009>
- Perry, J., & Bond, M. (2012). Change in defense mechanisms during long-term dynamic psychotherapy and five-year outcome.. *The American journal of psychiatry*, 169 9, 916-25. <https://doi.org/10.1176/appi.ajp.2012.11091403>

- Buchheim, A., Viviani, R., Kessler, H., Kächele, H., Cierpka, M., Roth, G., George, C., Kernberg, O., Bruns, G., & Taubner, S. (2012). Changes in Prefrontal-Limbic Function in Major Depression after 15 Months of Long-Term Psychotherapy. *PLoS ONE*, 7. <https://doi.org/10.1371/journal.pone.0033745>
- Simpkins, C., & Simpkins, A. (2013). Neuroplasticity and Neurogenesis: Changing Moment-by-Moment. \*\*, 165-174. [https://doi.org/10.1007/978-1-4614-4842-6\\_13](https://doi.org/10.1007/978-1-4614-4842-6_13)
- Ho, T., & King, L. (2021). Mechanisms of neuroplasticity linking early adversity to depression: developmental considerations. *Translational Psychiatry*, 11. <https://doi.org/10.1038/s41398-021-01639-6>
- Hayes, A., & Andrews, L. (2020). A complex systems approach to the study of change in psychotherapy. *BMC Medicine*, 18. <https://doi.org/10.1186/s12916-020-01662-2>
- Penedo, J., Coyne, A., Constantino, M., Krieger, T., Hayes, A., & Holtforth, G. (2020). Theory-specific patient change processes and mechanisms in different cognitive therapies for depression.. *Journal of consulting and clinical psychology*. <https://doi.org/10.1037/ccp0000502>
- Turner, B., Kashima, D., Manz, K., Grueter, C., & Grueter, B. (2017). Synaptic Plasticity in the Nucleus Accumbens: Lessons Learned from Experience.. *ACS chemical neuroscience*, 9 9, 2114-2126. <https://doi.org/10.1021/acscchemneuro.7b00420>
- Alves, N., Correia, J., Patrício, P., Mateus-Pinheiro, A., Machado-Santos, A., Loureiro-Campos, E., Morais, M., Bessa, J., Sousa, N., & Pinto, L. (2017). Adult hippocampal neuroplasticity triggers susceptibility to recurrent depression. *Translational Psychiatry*, 7. <https://doi.org/10.1038/tp.2017.29>
- Lemmens, L., Müller, V., Arntz, A., & Huibers, M. (2016). Mechanisms of change in psychotherapy for depression: An empirical update and evaluation of research aimed at identifying psychological mediators.. *Clinical psychology review*, 50, 95-107. <https://doi.org/10.1016/j.cpr.2016.09.004>
- Sweatt, J. (2016). Neural plasticity and behavior – sixty years of conceptual advances. *Journal of Neurochemistry*, 139. <https://doi.org/10.1111/jnc.13580>
- Piotrkowicz, M., Janoska-Jażdżik, M., Kowesko, T., & Szulc, A. (2021). The Influence of Psychotherapy on Peripheral Brain-Derived Neurotrophic Factor Concentration Levels and Gene Methylation Status: A Systematic Review. *Journal of Clinical Medicine*, 10. <https://doi.org/10.3390/jcm10194424>
- Kupferberg, A., & Hasler, G. (2024). From antidepressants and psychotherapy to oxytocin, vagus nerve stimulation, ketamine and psychedelics: how established and novel treatments can improve social functioning in major depression. *Frontiers in Psychiatry*, 15. <https://doi.org/10.3389/fpsy.2024.1372650>
- Bruijniks, S., Meeter, M., Lemmens, L., Peeters, F., Cuijpers, P., & Huibers, M. (2021). Temporal and specific pathways of change in cognitive behavioral therapy (CBT) and interpersonal psychotherapy (IPT) for depression.. *Behaviour research and therapy*, 151, 104010. <https://doi.org/10.1016/j.brat.2021.104010>
- Voss, P., Thomas, M., Cisneros-Franco, J., & De Villers-Sidani, É. (2017). Dynamic Brains and the Changing Rules of Neuroplasticity: Implications for Learning and Recovery. *Frontiers in Psychology*, 8. <https://doi.org/10.3389/fpsyg.2017.01657>
- Weingarten, C., & Strauman, T. (2015). Neuroimaging for psychotherapy research: Current trends. *Psychotherapy Research*, 25, 185 - 213. <https://doi.org/10.1080/10503307.2014.883088>
- Barsaglini, A., Sartori, G., Benetti, S., Pettersson-Yeo, W., & Mechelli, A. (2014). The effects of psychotherapy on brain function: A systematic and critical review. *Progress in Neurobiology*, 114, 1-14. <https://doi.org/10.1016/j.pneurobio.2013.10.006>