

The Neuroscientific Basis for Why Emotionally Charged Events Are Remembered More Vividly

1. Introduction

Emotionally charged events are often remembered with greater vividness and persistence than neutral experiences. Neuroscientific research has revealed that this phenomenon is rooted in the interplay between the amygdala, hippocampus, and broader neuromodulatory systems, which together enhance the encoding, consolidation, and retrieval of emotional memories (Krenz et al., 2025; McGaugh, 2018; McGaugh, 2015; Qasim et al., 2021; Roesler et al., 2021; LaBar & Cabeza, 2006; Paré & Headley, 2023; Yonelinas & Ritchey, 2015; Todd et al., 2012; Hamann, 2001; McGaugh, 2004; Zhang et al., 2024; Hu et al., 2007; Sharot et al., 2004; Dolcos et al., 2004; Murty et al., 2010). Emotional arousal triggers the release of stress hormones and neuromodulators (notably norepinephrine and glucocorticoids), which activate the amygdala. The amygdala, in turn, modulates memory-related processes in the hippocampus and neocortex, leading to more robust and detailed memory traces. This review synthesizes evidence from neuroimaging, electrophysiology, pharmacology, and behavioral studies to explain the neural mechanisms underlying the vividness of emotional memories.

2. Methods

A comprehensive search was conducted across over 170 million research papers in Consensus, including Semantic Scholar, PubMed, and other sources. The search strategy involved 20 targeted queries grouped into 8 thematic clusters, focusing on the neural basis of emotional memory vividness, amygdala-hippocampal interactions, neuromodulation, sleep, and clinical implications. In total, 1,041 papers were identified, 672 were screened, 588 were deemed eligible, and the top 50 most relevant papers were included in this review.

Search Strategy




FIGURE 1 Flow diagram of the literature search and selection process.

Eight unique search groups targeted foundational theories, mechanistic studies, alternate terminology, critique, interdisciplinary expansion, methodological diversity, and citation graph exploration.

3. Results

3.1. Amygdala-Hippocampal Interactions

The amygdala is central to the emotional enhancement of memory. During emotionally arousing events, the amygdala becomes highly active and interacts with the hippocampus to prioritize the encoding and consolidation of emotional information (Krenz et al., 2025; McGaugh, 2018; McGaugh, 2015; Qasim et al., 2021; Roesler et al., 2021; LaBar & Cabeza, 2006; Paré & Headley, 2023; Yonelinas & Ritchey, 2015; Hamann, 2001; McGaugh, 2004; Zhang et al., 2024; Hu et al., 2007; Sharot et al., 2004; Dolcos et al., 2004; Murty et al., 2010). Direct brain recordings and stimulation studies show that increased amygdala and hippocampal activity during emotional encoding predicts better subsequent memory (Qasim et al., 2021; Roesler et al., 2021; Dolcos et al., 2004). The amygdala's influence is especially pronounced during the initial encounter with an emotional event, setting the stage for stable neocortical memory representations (Krenz et al., 2025; Dolcos et al., 2004).

3.2. Neuromodulatory Systems and Stress Hormones

Emotional arousal triggers the release of stress hormones (adrenaline, glucocorticoids) and neuromodulators (norepinephrine, dopamine), which act on the amygdala and related circuits (McGaugh, 2018; McGaugh, 2015; Paré & Headley, 2023; Provensi et al., 2018; Yonelinas & Ritchey, 2015; Schwabe et al., 2022; McIntyre et al., 2003; McGaugh, 2004; Roozendaal & Hermans, 2017; Hu et al., 2007; Gao et al., 2016; McIntyre et al., 2012). Noradrenergic activation, often via the locus coeruleus, enhances synaptic plasticity in the amygdala and hippocampus, facilitating the consolidation of emotional memories (McGaugh, 2018; Provensi et al., 2018; Yonelinas & Ritchey, 2015; McIntyre et al., 2003; Roozendaal & Hermans, 2017; Hu et al., 2007; Gao et al., 2016). The basolateral amygdala (BLA) is particularly important for modulating these effects (Roesler et al., 2021; Paré & Headley, 2023; McIntyre et al., 2003; McGaugh, 2004; Bocchio et al., 2017).

3.3. Network Integration and Sensory Vividness

Emotionally arousing events promote greater integration across large-scale brain networks, including the prefrontal cortex, parietal cortex, and sensory cortices (Park et al., 2025; Todd et al., 2012; Hermans et al., 2014; Murty et al., 2010). Emotional salience increases perceptual vividness at the time of encoding, which is reflected in stronger and more detailed sensory representations (Todd et al., 2012; Sharot et al., 2004). This heightened sensory encoding is mediated by amygdala-driven modulation of perceptual and mnemonic circuits (Todd et al., 2012; Sharot et al., 2004; Murty et al., 2010).

3.4. Sleep and Memory Consolidation

Sleep, especially REM and slow-wave sleep, further consolidates emotional memories by enhancing hippocampal and amygdalo-cortical connectivity (Payne & Kensinger, 2018; Sterpenich et al., 2007; Cairney et al., 2015). Emotional memories are more resistant to forgetting over time, a process supported by sleep-dependent systems-level consolidation (Payne & Kensinger, 2018; Sterpenich et al., 2007; Cairney et al., 2015).

Key Papers

Paper	Methodology	Key Neural Mechanism	Key Results
(Krenz et al., 2025)	fMRI, mediation analysis	Amygdala-neocortical	Initial amygdala response boosts stable neocortical patterns for emotional memory
(McGaugh, 2018)	Review, animal/human	Amygdala, stress hormones	Amygdala modulates consolidation via adrenergic/glucocorticoid activation
(Qasim et al., 2021)	iEEG, stimulation	Amygdala-hippocampus	High-frequency activity in both regions enhances emotional memory encoding
(Todd et al., 2012)	ERP, fMRI	Amygdala, sensory cortex	Emotional salience increases perceptual vividness and predicts later memory vividness
(McGaugh, 2004)	Review, animal/human	Basolateral amygdala	BLA modulates memory consolidation via noradrenergic/cholinergic systems

FIGURE 2 Comparison of key studies on the neural basis of emotional memory vividness.

Top Contributors

Type	Name	Papers
Author	J. D. McGaugh	(McGaugh, 2018; McGaugh, 2015; McIntyre et al., 2003; McGaugh, 2004; McIntyre et al., 2012; LaLumiere et al., 2017)
Author	B. Roozendaal	(Krenz et al., 2025; Roesler et al., 2021; Schwabe et al., 2022; McIntyre et al., 2003; McGaugh, 2004; Roozendaal & Hermans, 2017)
Author	E. Kensinger	(Payne & Kensinger, 2018; Bowen et al., 2017; Kensinger & Ford, 2020; Kensinger, 2009; Kensinger & Corkin, 2004; Kensinger, 2007)
Journal	<i>Nature Reviews Neuroscience</i>	(LaBar & Cabeza, 2006)
Journal	<i>Trends in Cognitive Sciences</i>	(Rouhani et al., 2023; Yonelinas & Ritchey, 2015; Hamann, 2001; Kensinger & Corkin, 2004)
Journal	<i>The Journal of Neuroscience</i>	(Krenz et al., 2025; Todd et al., 2012)

FIGURE 3 Authors & journals that appeared most frequently in the included papers.

4. Discussion

The vividness of emotional memories is underpinned by a network of neural mechanisms centered on the amygdala and its interactions with the hippocampus and neocortex (Krenz et al., 2025; McGaugh, 2018; McGaugh, 2015; Qasim et al., 2021; Roesler et al., 2021; LaBar & Cabeza, 2006; Paré & Headley, 2023; Yonelinas & Ritchey, 2015; Todd et al., 2012; Hamann, 2001; McGaugh, 2004; Zhang et al., 2024; Hu et al., 2007; Sharot et al., 2004; Dolcos et al., 2004; Murty et al., 2010). Emotional arousal triggers neuromodulatory cascades that enhance synaptic plasticity and memory consolidation, particularly through noradrenergic and glucocorticoid signaling in the amygdala and hippocampus (McGaugh, 2018; McGaugh, 2015; Paré & Headley, 2023; Provensi et al., 2018; Yonelinas & Ritchey, 2015; McIntyre et al., 2003; McGaugh, 2004; Roozendaal & Hermans, 2017; Hu et al., 2007; Gao et al., 2016; McIntyre et al., 2012). The amygdala's modulation of sensory and mnemonic circuits leads to more detailed and vivid memory traces, while large-scale network integration and sleep-dependent processes further support the persistence and vividness of these memories (Payne & Kensinger, 2018; Park et al., 2025; Todd et al., 2012; Hermans et al., 2014; Sterpenich et al., 2007; Cairney et al., 2015; Murty et al., 2010).

Notably, the subjective sense of vividness may be driven by strong memory for a few central details, rather than a greater quantity of contextual information (Phelps & Sharot, 2008; Sharot et al., 2004). Negative emotional events, in particular, are associated with increased sensory encoding and recapitulation at retrieval (Bowen et al., 2017; Kensinger, 2009; Sharot et al., 2004). However, the same mechanisms that enhance memory vividness can also contribute to memory distortions or maladaptive persistence, as seen in affective disorders (Krenz et al., 2025; McGaugh, 2015; Yonelinas & Ritchey, 2015; Zhang et al., 2024; LaLumiere et al., 2017).

Claims and Evidence Table







Claim	Evidence Strength	Reasoning	Papers
Amygdala activation enhances encoding and consolidation of emotional memories	 Strong	Convergent evidence from fMRI, iEEG, animal, and pharmacological studies	(Krenz et al., 2025; McGaugh, 2018; McGaugh, 2015; Qasim et al., 2021; Roesler et al., 2021; LaBar & Cabeza, 2006; Paré & Headley, 2023; Yonelinas & Ritchey, 2015; Hamann, 2001; McGaugh, 2004; Zhang et al., 2024; Hu et al., 2007; Sharot et al., 2004; Dolcos et al., 2004; Murty et al., 2010)
Noradrenergic and glucocorticoid systems mediate emotional memory enhancement	 Strong	Stress hormones and neuromodulators act on amygdala/hippocampus to boost memory	(McGaugh, 2018; McGaugh, 2015; Paré & Headley, 2023; Provensi et al., 2018; Yonelinas & Ritchey, 2015; McIntyre et al., 2003; McGaugh, 2004; Roozendaal & Hermans, 2017; Hu et al., 2007; Gao et al., 2016; McIntyre et al., 2012)
Emotional arousal increases perceptual vividness and sensory encoding	 Strong	ERP, fMRI, and behavioral studies show enhanced sensory detail for emotional events	(Todd et al., 2012; Sharot et al., 2004; Bowen et al., 2017; Kensinger, 2009; Phelps & Sharot, 2008; Murty et al., 2010)
Large-scale brain network integration supports emotional memory vividness	 Moderate	fMRI and graph theory studies show network cohesion during emotional encoding	(Park et al., 2025; Hermans et al., 2014; Murty et al., 2010)
Sleep further consolidates emotional memories	 Moderate	Sleep studies show enhanced hippocampo-cortical connectivity for emotional memories	(Payne & Kensinger, 2018; Sterpenich et al., 2007; Cairney et al., 2015)
Vividness may reflect strong memory for central details, not all context	 Moderate	Subjective vividness often tied to a few salient features	(Phelps & Sharot, 2008; Sharot et al., 2004; Bowen et al., 2017; Kensinger, 2009)

FIGURE Key claims and support evidence identified in these papers.

5. Conclusion

Emotionally charged events are remembered more vividly due to the coordinated action of the amygdala, hippocampus, and neuromodulatory systems, which together enhance the encoding, consolidation, and retrieval of emotional information. These neural mechanisms prioritize emotionally salient details, leading to robust and vivid memory traces.

5.1. Research Gaps

Despite significant advances, gaps remain in understanding individual differences, the role of positive versus negative emotion, and how these mechanisms may be targeted in clinical interventions.

Research Gaps Matrix

Topic/Attribute	Amygdala-Hippocampus	Neuromodulators	Sensory Encoding	Sleep	Clinical Populations
Negative emotion	12	10	8	6	7
Positive emotion	7	5	4	3	2
Individual differences	5	3	2	2	4
Network integration	6	4	3	2	2
Memory distortions	4	2	2	1	5

FIGURE Matrix of research topics and study attributes, highlighting areas with limited research coverage.

5.2. Open Research Questions

Future research should address the following questions to further clarify the neural basis of emotional memory vividness.

Question	Why
How do individual differences in amygdala reactivity influence emotional memory vividness and susceptibility to disorders?	Understanding variability can inform personalized interventions for affective disorders.
What are the distinct neural mechanisms for positive versus negative emotional memory vividness?	Most research focuses on negative emotion; positive emotion may engage different circuits.
How can neuromodulatory and network mechanisms be targeted to reduce maladaptive emotional memory persistence?	Insights could inform treatments for PTSD and related conditions.

FIGURE Open research questions for future investigation on the neural basis of emotional memory vividness.

In summary, the vividness of emotional memories is rooted in specialized neural mechanisms involving the amygdala, hippocampus, neuromodulators, and large-scale brain networks, which together prioritize and strengthen the encoding and retention of emotionally salient experiences.

These papers were sourced and synthesized using Consensus, an AI-powered search engine for research. Try it at <https://consensus.app>

References

- Krenz, V., Alink, A., Roozendaal, B., Sommer, T., & Schwabe, L. (2025). Memory Boost for Recurring Emotional Events Is Driven by Initial Amygdala Response Promoting Stable Neocortical Patterns across Repetitions. *The Journal of Neuroscience*, 45. <https://doi.org/10.1523/JNEUROSCI.2406-23.2025>
- McGaugh, J. (2018). Emotional arousal regulation of memory consolidation. *Current Opinion in Behavioral Sciences*, 19, 55-60. <https://doi.org/10.1016/j.cobeha.2017.10.003>
- Rouhani, N., Niv, Y., Frank, M., & Schwabe, L. (2023). Multiple routes to enhanced memory for emotionally relevant events. *Trends in Cognitive Sciences*, 27, 867-882. <https://doi.org/10.1016/j.tics.2023.06.006>
- McGaugh, J. (2015). Consolidating memories.. *Annual review of psychology*, 66, 1-24. <https://doi.org/10.1146/annurev-psych-010814-014954>
- Qasim, S., Mohan, U., Stein, J., & Jacobs, J. (2021). Neuronal activity in the human amygdala and hippocampus enhances emotional memory encoding. *Nature human behaviour*, 7, 754 - 764. <https://doi.org/10.1038/s41562-022-01502-8>
- Payne, J., & Kensinger, E. (2018). Stress, sleep, and the selective consolidation of emotional memories. *Current Opinion in Behavioral Sciences*, 19, 36-43. <https://doi.org/10.1016/j.cobeha.2017.09.006>
- Roesler, R., Parent, M., LaLumiere, R., & McIntyre, C. (2021). Amygdala-hippocampal interactions in synaptic plasticity and memory formation. *Neurobiology of Learning and Memory*, 184. <https://doi.org/10.1016/j.nlm.2021.107490>
- LaBar, K., & Cabeza, R. (2006). Cognitive neuroscience of emotional memory. *Nature Reviews Neuroscience*, 7, 54-64. <https://doi.org/10.1038/nrn1825>

- Bowen, H., Kark, S., & Kensinger, E. (2017). NEVER forget: negative emotional valence enhances recapitulation. *Psychonomic Bulletin & Review*, 25, 870 - 891. <https://doi.org/10.3758/s13423-017-1313-9>
- Park, J., Gollapudi, K., Ke, J., Nau, M., Pappas, I., & Leong, Y. (2025). Emotional arousal enhances narrative memories through functional integration of large-scale brain networks. *bioRxiv*. <https://doi.org/10.1101/2025.03.13.643125>
- Kensinger, E., & Ford, J. (2020). Retrieval of Emotional Events from Memory.. *Annual review of psychology*. <https://doi.org/10.1146/annurev-psych-010419-051123>
- Paré, D., & Headley, D. (2023). The amygdala mediates the facilitating influence of emotions on memory through multiple interacting mechanisms. *Neurobiology of Stress*, 24. <https://doi.org/10.1016/j.ynstr.2023.100529>
- Kensinger, E. (2009). Remembering the Details: Effects of Emotion. *Emotion Review*, 1, 113 - 99. <https://doi.org/10.1177/1754073908100432>
- Provensi, G., Passani, M., Costa, A., Izquierdo, I., & Blandina, P. (2018). Neuronal histamine and the memory of emotionally salient events. *British Journal of Pharmacology*, 177, 557 - 569. <https://doi.org/10.1111/bph.14476>
- Yonelinas, A., & Ritchey, M. (2015). The slow forgetting of emotional episodic memories: an emotional binding account. *Trends in Cognitive Sciences*, 19, 259-267. <https://doi.org/10.1016/j.tics.2015.02.009>
- Schwabe, L., Hermans, E., Joëls, M., & Roozendaal, B. (2022). Mechanisms of memory under stress. *Neuron*, 110, 1450-1467. <https://doi.org/10.1016/j.neuron.2022.02.020>
- Todd, R., Talmi, D., Schmitz, T., Susskind, J., & Anderson, A. (2012). Psychophysical and Neural Evidence for Emotion-Enhanced Perceptual Vividness. *The Journal of Neuroscience*, 32, 11201 - 11212. <https://doi.org/10.1523/JNEUROSCI.0155-12.2012>
- McIntyre, C., Power, A., Roozendaal, B., & McGaugh, J. (2003). Role of the Basolateral Amygdala in Memory Consolidation. *Annals of the New York Academy of Sciences*, 985. <https://doi.org/10.1111/j.1749-6632.2003.tb07088.x>
- Hamann, S. (2001). Cognitive and neural mechanisms of emotional memory. *Trends in Cognitive Sciences*, 5, 394-400. [https://doi.org/10.1016/S1364-6613\(00\)01707-1](https://doi.org/10.1016/S1364-6613(00)01707-1)
- Phelps, E., & Sharot, T. (2008). How (and Why) Emotion Enhances the Subjective Sense of Recollection. *Current Directions in Psychological Science*, 17, 147 - 152. <https://doi.org/10.1111/j.1467-8721.2008.00565.x>
- Hermans, E., Battaglia, F., Atsak, P., Voogd, L., Fernández, G., & Roozendaal, B. (2014). How the amygdala affects emotional memory by altering brain network properties. *Neurobiology of Learning and Memory*, 112, 2-16. <https://doi.org/10.1016/j.nlm.2014.02.005>
- Sterpenich, V., Albouy, G., Boly, M., Vandewalle, G., Darsaud, A., Baeteau, E., Dang-Vu, T., Desseilles, M., D'Argembeau, A., Gais, S., Rauchs, G., Schabus, M., Degueldre, C., Luxen, A., Collette, F., & Maquet, P. (2007). Sleep-Related Hippocampo-Cortical Interplay during Emotional Memory Recollection. *PLoS Biology*, 5. <https://doi.org/10.1371/journal.pbio.0050282>
- McGaugh, J. (2004). The amygdala modulates the consolidation of memories of emotionally arousing experiences.. *Annual review of neuroscience*, 27, 1-28. <https://doi.org/10.1146/ANNUREV.NEURO.27.070203.144157>
- Zhang, H., Skelin, I., , S., Paff, M., Mnatsakanyan, L., Yassa, M., Knight, R., & Lin, J. (2024). Awake ripples enhance emotional memory encoding in the human brain. *Nature Communications*, 15. <https://doi.org/10.1038/s41467-023-44295-8>

Bocchio, M., Nabavi, S., & Capogna, M. (2017). Synaptic Plasticity, Engrams, and Network Oscillations in Amygdala Circuits for Storage and Retrieval of Emotional Memories. *Neuron*, 94, 731-743.

<https://doi.org/10.1016/j.neuron.2017.03.022>

Kensinger, E., & Corkin, S. (2004). Two routes to emotional memory: distinct neural processes for valence and arousal. *Proceedings of the National Academy of Sciences of the United States of America*, 101 9, 3310-5.

<https://doi.org/10.1073/PNAS.0306408101>

Roozendaal, B., & Hermans, E. (2017). Norepinephrine effects on the encoding and consolidation of emotional memory: improving synergy between animal and human studies. *Current Opinion in Behavioral Sciences*, 14, 115-122. <https://doi.org/10.1016/j.cobeha.2017.02.001>

Hu, H., Réal, E., Takamiya, K., Kang, M., LeDoux, J., Huganir, R., & Malinow, R. (2007). Emotion Enhances Learning via Norepinephrine Regulation of AMPA-Receptor Trafficking. *Cell*, 131, 160-173.

<https://doi.org/10.1016/j.cell.2007.09.017>

Kensinger, E. (2007). Negative Emotion Enhances Memory Accuracy. *Current Directions in Psychological Science*, 16, 213 - 218. <https://doi.org/10.1111/j.1467-8721.2007.00506.x>

Sharot, T., Delgado, M., & Phelps, E. (2004). How emotion enhances the feeling of remembering. *Nature Neuroscience*, 7, 1376-1380. <https://doi.org/10.1038/nn1353>

Cairney, S., Durrant, S., Power, R., & Lewis, P. (2015). Complementary roles of slow-wave sleep and rapid eye movement sleep in emotional memory consolidation. *Cerebral cortex*, 25 6, 1565-75.

<https://doi.org/10.1093/cercor/bht349>

Gao, V., Suzuki, A., Magistretti, P., Lengacher, S., Pollonini, G., Steinman, M., & Alberini, C. (2016). Astrocytic β 2-adrenergic receptors mediate hippocampal long-term memory consolidation. *Proceedings of the National Academy of Sciences*, 113, 8526 - 8531. <https://doi.org/10.1073/pnas.1605063113>

Dolcos, F., LaBar, K., & Cabeza, R. (2004). Interaction between the Amygdala and the Medial Temporal Lobe Memory System Predicts Better Memory for Emotional Events. *Neuron*, 42, 855-863.

[https://doi.org/10.1016/S0896-6273\(04\)00289-2](https://doi.org/10.1016/S0896-6273(04)00289-2)

McIntyre, C., McGaugh, J., & Williams, C. (2012). Interacting brain systems modulate memory consolidation. *Neuroscience & Biobehavioral Reviews*, 36, 1750-1762. <https://doi.org/10.1016/j.neubiorev.2011.11.001>

LaLumiere, R., McGaugh, J., & McIntyre, C. (2017). Emotional Modulation of Learning and Memory: Pharmacological Implications. *Pharmacological Reviews*, 69, 236 - 255. <https://doi.org/10.1124/pr.116.013474>

Murty, V., Ritchey, M., Adcock, R., & LaBar, K. (2010). fMRI studies of successful emotional memory encoding: A quantitative meta-analysis. *Neuropsychologia*, 48, 3459-3469.

<https://doi.org/10.1016/j.neuropsychologia.2010.07.030>