

Supplementary Material

Comparison of PCS score between self-reported aphantasics (Study 2) and “blinded aphantasics” (Study 1) using extreme aphantasia cut-off (rescaled VVIQ = 0)

For the comparison of PCS between extreme aphantasics of Study 1 and 2 (characterized by mean rescaled VVIQ of 0), H1 for Bayes Factor was modelled using a uniform distribution from 0 to the mean PCS (1.19) of Study 1 “blinded” aphantasics. Mean PCS was higher for blinded extreme aphantasics ($M = 1.19$, $SD = 0.55$) than for self-reported extreme aphantasics ($M = 0.43$, $SD = 0.62$), $t(18.3) = 3.54$, $p = .002$, $SE = 0.21$, Hedges’ $g = 1.22$, 95% CI [0.31, 1.20], $B_{H(0,1.19)} = 169.58$, $RR_{BF<3} [0.14, 1.19]$, see Figure S1.




Figure S1. Mean PCS of extreme aphantasics: blinded aphantasics from Study 1 and self-reported aphantasics from Study 2. Dots represent individual participants.

Comparison of PCS score between Study 1 and Study 2 samples with a German internet volunteer sample (data: <https://osf.io/b7yfx/>)

To test whether demographic and recruitment differences may be driving the effect we observed in Study 2, we compared the PCS scores of the Study 1 student sample and the Study 2 internet sample from the general public with the PCS score of a sample recruited online from the general public and tested online on a German translation of the PCS, $n = 240$, age range 18-72, $M_{\text{age}} = 33.42$, $SD_{\text{age}} = 12.31$, females, 142, males, 97, other, 1. As Study 2 sample, the German sample was older than Study 1 sample, $t(243.39) = -17.46$, $p < .001$.

To calculate the BF for the comparison of PCS scores, we used a half-normal distribution with SD equal to half the mean difference between the PCS score of Study 1 and Study 2 (0.69). There was evidence for no difference between the German internet sample PCS score ($M = 1.88$, $SD = 0.71$), and the student sample of Study 1 ($M = 1.84$, $SD = 0.67$), $t(448.57) = -0.62$, $p = 0.53$, $SE = 0.05$, Hedges' $g = -0.05$, 95% CI [-0.14, 0.07], $B_{H(0,0.69)} = 0.10$, $RR_{BF<3} [0.19, 1.37]$, see Figure S2.

In contrast, there was evidence that the German sample PCS score was higher than Study 2 sample's PCS score (M difference = 1.41), $t(42.07) = -11.68$, $p < .001$, $SE = 0.12$, Hedges' $g = -2.01$, 95% CI [-1.65, -1.16], $B_{H(0,0.69)} = 7.55 \times 10^{11}$, $RR_{BF<3} [0.06, 1.37]$, see Figure S2.




Figure S2. Mean PCS of Study 1 sample, Study 2 sample, and German sample. Dots represent individual participants.