


Interaction landscape — evan_device.json
 $t_g=138.9$ ns, peak $|\eta|=1.00$



MODE INVENTORY

a	3.500 GHz	qubit a (target)
b	5.700 GHz	qubit b (target)
s	4.700 GHz	SNAIL coupler
sp1	4.600 GHz	spectator 1
pump	2.200 GHz	2w_p = 4.400 GHz
1/t_g	7.2 MHz	pulse bandwidth

INTERACTIONS

#	process	g (MHz)	det (MHz)	DRAG
T	target iSWAP (A-B exchange)	3.60	-0.0	--
1	SNAIL subharmonic	180.00	-300.0	4.4dB
2	spectator 1 subharmonic	18.00	-200.0	20.9dB
3	A $ 2\rangle$ leakage (via B)	5.09	-120.0	27.4dB
4	B $ 2\rangle$ leakage (via A)	5.09	-120.0	27.4dB
5	qubit A $ 1\rangle \rightarrow 2\rangle$ subharmonic	25.46	+780.0	29.7dB
6	qubit A subharmonic (emission)	18.00	-900.0	34.0dB
7	qubit A subharmonic	18.00	+900.0	34.0dB

DRAG column = suppression of that channel,
 $\sim g/|\text{det}|$ (blank = cannot help).

Which parasites DRAG can suppress

