


Interaction landscape — 2Gate4.9SNAIL.json
 $t_g=92.6$ ns, peak $|\eta|=1.50$



MODE INVENTORY

a	3.800 GHz	qubit a (target)
b	5.500 GHz	qubit b (target)
s	4.900 GHz	SNAIL coupler
sp1	4.650 GHz	spectator 1
pump	1.700 GHz	2w_p = 3.400 GHz
1/t_g	10.8 MHz	pulse bandwidth

INTERACTIONS

#	process	g (MHz)	det (MHz)	DRAG
T	target iSWAP (A-B exchange)	5.40	+0.0	--
1	qubit A $ 1\rangle \rightarrow 2\rangle$ subharmonic	57.28	-520.0	19.2dB
2	qubit A subharmonic (emission)	40.50	+400.0	19.9dB
3	qubit A subharmonic	40.50	-400.0	19.9dB
4	A-SNAIL spectator	54.00	+600.0	20.9dB
5	B $ 2\rangle$ leakage (via A)	7.64	-120.0	23.9dB
6	A $ 2\rangle$ leakage (via B)	7.64	-120.0	23.9dB
7	A-SP1 spectator	5.40	+850.0	43.9dB
8	B-SP1 spectator	5.40	-850.0	43.9dB

DRAG column = suppression of that channel,
 $\sim g/|\text{det}|$ (blank = cannot help).

Which parasites DRAG can suppress

