

Optimal synaptic strategies for different timescales of memory


Subhaneil Lahiri and Surya Ganguli

Stanford University, Applied Physics

February 26, 2016

What is a synapse?

Experimenters




[Klann (2002)]

What is a synapse?

Experimenters

Theorists



W_{ij}


[Klann (2002)]

Timescales of memory

Memories stored in different places
for different timescales

[Squire and Alvarez (1995)]

[McClelland et al. (1995)]




[Born and Wilhelm (2012)]

Also: Cerebellar cortex → nuclei.

[Attwell et al. (2002)]

[Cooke et al. (2004)]

Different synapses have different
molecular structures.



[Emes RD, Grant SGN, 2012.
Annu. Rev. Neurosci. 35:111–31]

[Emes and Grant (2012)]

Storage capacity of synaptic memory


A classical perceptron has a capacity $\propto N$, ($\#$ synapses).

Requires synapses' dynamic range also $\propto N$.

With discrete, finite synapses:

\implies new memories overwrite old,

\implies stability-plasticity dilemma.




[Petersen et al. (1998), O'Connor et al. (2005)]

When we store new memories rapidly, memory capacity $\sim \mathcal{O}(\log N)$.

[Amit and Fusi (1992), Amit and Fusi (1994)]


Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Recognition memory

Synapses given a sequence of patterns (pot & dep) to store



Recognition memory

Synapses given a sequence of patterns (pot & dep) to store




Later: presented with a pattern. Has it been seen before?

Compare $\vec{s} \cdot \vec{w}(t)$ to threshold.

[Sommer and Dayan (1998)]

Recognition memory

Synapses given a sequence of patterns (pot & dep) to store



Later: presented with a pattern. Has it been seen before?

Compare $\vec{s} \cdot \vec{w}(t)$ to threshold.

[Sommer and Dayan (1998)]


$$\text{SNR}(t) = \frac{\langle \vec{s} \cdot \vec{w}(t) \rangle - \langle \vec{s} \cdot \vec{w}(\infty) \rangle}{\sqrt{\text{Var}(\vec{s} \cdot \vec{w}(\infty))}}, \quad \overline{\text{SNR}}(\tau) = \int d\tau \frac{e^{-t/\tau}}{\tau} \text{SNR}(t).$$

Models of complex synaptic dynamics



Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak
● strong




States: #AMPA, #NMDAR, NMDAR subunit composition,
CaMK II autophosphorylation, activating PKC, p38 MAPK,...

[Fusi et al. (2005), Fusi and Abbott (2007), Barrett and van Rossum (2008)]


Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong




Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong




Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics

- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong


Potential event



Depression event

Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong



Models of complex synaptic dynamics


- Internal functional state of synapse \rightarrow synaptic weight.
 - Candidate plasticity events \rightarrow transitions between states
- weak ● strong

Potentiation




Depression

Proven envelope: memory frontier




Proven envelope: memory frontier

Upper bound on memory curve at *any* timescale.




Proven envelope: memory frontier



Upper bound on memory curve at *any* timescale.





Serial topology:





Models that maximise memory for one timescale



Models that maximise memory for one timescale



Models that maximise memory for one timescale




Synaptic structures for different timescales of memory


Short timescales



Intermediate timescales





Long timescales



Synaptic structures for different timescales of memory

Short timescales \longrightarrow Intermediate timescales \longrightarrow Long timescales





short topology



long topology

Synaptic structures for different timescales of memory

Short timescales \longrightarrow Intermediate timescales \longrightarrow Long timescales



short topology



long topology


strong transitions



weak transitions


Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.



Proposed Experimental design

Subject a synapse to a sequence of candidate plasticity events.
Observe the changes in synaptic efficacy.




EM algorithms:

Sequence of hidden states \rightarrow estimate transition probabilities


Transition probabilities \rightarrow estimate sequence of hidden states

[Baum et al. (1970), Rabiner and Juang (1993), Dempster et al. (2007)]


Simulated experiment




Simulated experiment




Simulated experiment



Simulated experiment



Simulated experiment



- We have formulated a general theory of learning and memory with complex synapses.
- We find a memory envelope: a single curve that cannot be exceeded by the memory curve of *any* synaptic model.
- We understood which types of synaptic structure are useful for storing memories for different timescales.

Acknowledgements

Thanks to:

- Surya Ganguli
- Stefano Fusi
- Marcus Benna
- David Sussillo
- Jascha Sohl-Dickstein

Funding:

- Swartz foundation
- Stanford Bio-X
- Genentech

References I



Eric Klann.

“Metaplastic Protein Phosphatases”.

Learning and Memory, 9(4):153–155, (2002) ,

<http://learnmem.cshlp.org/content/9/4/153.full.pdf+html>.

2 3



Larry R Squire and Pablo Alvarez.

“Retrograde amnesia and memory consolidation: a neurobiological perspective”.

Current Opinion in Neurobiology, 5(2):169–177, (April, 1995) .

4



James L McClelland, Bruce L McNaughton, and Randall C O'Reilly.

“Why there are complementary learning systems in the hippocampus and neocortex: Insights from the successes and failures of connectionist models of learning and memory.”, 1995.

4

References II



Jan Born and Ines Wilhelm.

“System consolidation of memory during sleep.”.

Psychological research, 76(2):192–203, (mar, 2012) .

4



Phillip J.E. Attwell, Samuel F. Cooke, and Christopher H. Yeo.

“Cerebellar Function in Consolidation of a Motor Memory”.

Neuron, 34(6):1011–1020, (jun, 2002) .

4



Samuel F Cooke, Phillip J E Attwell, and Christopher H Yeo.

“Temporal properties of cerebellar-dependent memory consolidation.”.

The Journal of neuroscience : the official journal of the Society for Neuroscience, 24(12):2934–41, (mar, 2004) .

4

References III



Richard D. Emes and Seth G.N. Grant.

“Evolution of Synapse Complexity and Diversity”.

Annual Review of Neuroscience, 35(1):111–131, (2012) .

4



Carl C. H. Petersen, Robert C. Malenka, Roger A. Nicoll, and John J. Hopfield.

“All-or-none potentiation at CA3-CA1 synapses”.

Proc. Natl. Acad. Sci. U.S.A., 95(8):4732–4737, (1998) .

5



Daniel H. O'Connor, Gayle M. Wittenberg, and Samuel S.-H. Wang.

“Graded bidirectional synaptic plasticity is composed of switch-like unitary events”.

Proc. Natl. Acad. Sci. U.S.A., 102(27):9679–9684, (2005) .

5

References IV



D. J. Amit and S. Fusi.

“Constraints on learning in dynamic synapses”.

Network: Computation in Neural Systems, 3(4):443–464, (1992) .

5



D. J. Amit and S. Fusi.

“Learning in neural networks with material synapses”.

Neural Computation, 6(5):957–982, (1994) .

5



Friedrich T Sommer and Peter Dayan.

“Bayesian retrieval in associative memories with storage errors.”.

IEEE transactions on neural networks / a publication of the IEEE Neural Networks Council, 9(4):705–13, (jan, 1998) .

6

7

8

9

10

11

12

13

14

References V



S. Fusi, P. J. Drew, and L. F. Abbott.

“Cascade models of synaptically stored memories”.

Neuron, 45(4):599–611, (Feb, 2005) .

16



S. Fusi and L. F. Abbott.

“Limits on the memory storage capacity of bounded synapses”.

Nat. Neurosci., 10(4):485–493, (Apr, 2007) .

16



A. B. Barrett and M. C. van Rossum.

“Optimal learning rules for discrete synapses”.

PLoS Comput. Biol., 4(11):e1000230, (Nov, 2008) .

16

References VI



LE Baum, T Petrie, George Soules, and Norman Weiss.

“A maximization technique occurring in the statistical analysis of probabilistic functions of Markov chains”.

The annals of mathematical statistics, 41(1):164–171, (1970) .

33 34 35 36 37 38 39



Lawrence R Rabiner and Biing-Hwang Juang.

Fundamentals of speech recognition, volume 14 of *Signal Processing*.

Prentice Hall, Inc., Upper Saddle River, NJ, USA, 1993.

ISBN 0-13-015157-2.

33 34 35 36 37 38 39

References VII

 A. P. Dempster, N. M. Laird, and D. B. Rubin.

“Maximum Likelihood from Incomplete Data via the EM Algorithm”.

Journal of the Royal Statistical Society. Series B (Methodological), (October, 2007) .

33

34

35

36

37

38

39

 J.G. Kemeny and J.L. Snell.

Finite markov chains.

Springer, 1960.

54

Technical detail: ordering states

Let \mathbf{T}_{ij} = mean first passage time from state i to state j . Then:

$$\eta = \sum_j \mathbf{T}_{ij} \mathbf{p}_j^\infty,$$

is independent of the initial state i (Kemeney's constant).

[Kemeny and Snell (1960)]


We define:

$$\eta_i^+ = \sum_{j \in \text{strong}} \mathbf{T}_{ij} \mathbf{p}_j^\infty, \quad \eta_i^- = \sum_{j \in \text{weak}} \mathbf{T}_{ij} \mathbf{p}_j^\infty.$$

They can be used to arrange the states in an order (increasing η^- or decreasing η^+). [back](#)

Technical detail: upper/lower triangular

With states in order:



Endpoint: potentiation goes right, depression goes left.

[back](#)