

[illegible]

Contents

1	Introduction	1
1.1	Background	1
1.2	Problem Analysis	1
1.3	Our Work	2
2	Assumptions and Notations	2
2.1	Assumptions and Explanations	2
2.2	Notations	4
3	Models	4
3.1	Model Development Explanation and Sequence	4
3.1.1	Notes for Models	4
3.1.2	Sequence	5
3.2	Model I: LVLG(Lokta-Volterra and Leslie-Gower) Model	5
3.3	Model II:	7
3.4	Model III:	7
3.5	Model IV:	7
4	Application of the Models	7
5	Sensitivity Analysis	7
6	Evaluation of the Model	7
6.1	Strengths	7
6.2	Weaknesses	7
7	Conclusion	7
	References	7

1 Introduction

1.1 Background

In the past few decades, with the rapid population growth, food supply has become one of the most pressing global issues. Under the conditions where scientific and technological advancements have not been fully adopted, the existing arable land area is insufficient to meet the food demand. As a result, many regions have resorted to deforestation for land conversion, as shown in Figure 1. Figure 2 illustrates that when the forest ecosystem is artificially disrupted, its complex spatial structure is destroyed.



Figure 1: Deforestation for Farming



Figure 2: Deforested Forest

Therefore, the converted forest area must undergo a long process of ecological reconstruction. However, due to the influence of human activities during this process, the post-clearing forest ecosystem can no longer return to its original state, but instead continuously undergoes succession into a stable agricultural ecosystem. To achieve both economic and ecological benefits and fulfill sustainable development goals, it is necessary to examine the ecological succession from converted forest to agricultural ecosystems and explore how green agriculture can bring dual benefits in terms of economic and ecological indicators.

1.2 Problem Analysis

The basic requirements of the project are to establish a model that reflects the ecological succession process from a converted forest area to a mature agricultural ecosystem over time. The model should incorporate both natural processes and human agricultural activities. Specifically, the requirements are as follows:

- **Develop an ecological model** for the converted forest area, particularly:
 - **Food web construction:** The model should include at least producers (e.g., plants) and consumers (e.g., herbivores and predators) and their interactions.
 - **Consideration of agricultural cycles and seasonality:** The impact of seasonal and cyclical agricultural activities should be taken into account.
 - **Impact of chemicals:** The model should account for the effects of chemicals such as herbicides and pesticides on plants, insects, bats, birds, and the stability of the ecosystem.

- **Reemergence of species during ecosystem maturation:** As the ecosystem matures, the model should consider the reemergence of two species and their impact on the ecosystem.
- **Impact of removing chemicals:** After the ecosystem matures, humans will attempt to remove chemicals. The model should assess the stability of the ecosystem after herbicides are removed, with the effects reflected through producers and consumers.
- **Introduction of bats into the food web:** The model should examine how bats, as insectivores and pollinators, interact with plants, insects, and predators, and how their inclusion influences the stability of the ecosystem. Additionally, the model should identify another species that could benefit the ecosystem and compare the effects of different species.
- **Analysis of the impact of organic farming:** The model should assess the impact of adopting organic farming practices, considering various scenarios and components. The evaluation should include the effects on the overall ecosystem and individual elements, such as pest control, crop health, plant reproduction, biodiversity, long-term sustainability, and cost-effectiveness. The model should analyze the impact of organic practices on pest management, soil health, and biodiversity, while weighing the economic costs and benefits. A comprehensive ecological and economic trade-off analysis should be provided to assess the feasibility of organic farming.

1.3 Our Work

- 1
- 2
- 3

2 Assumptions and Notations

2.1 Assumptions and Explanations

- **Accurate Data Assumption:** The model assumes that the data used are accurate.
Explanation: The data used in the model are sourced from official databases, and we believe the data to be accurate and reliable.
- **Geographic Applicability Assumption:** The model assumes that the applicable region is Southeast Asia, where two crops of rice are planted each year in the farmland.
Explanation: The climate of Southeast Asia is rather simple, with only two seasons-rainy and dry. Additionally, as is shown in Figure 3, the temperature variation within a year is minimal, which has a trivial effect on the ecosystem. Consequently, temperature can be considered as a constant. Due to such weather pattern, it aligns with the planting patterns commonly observed in Southeast Asia

to plant three crops of rice each year (showed in Figure 4), and the simplicity of crop types makes the model easier to establish.




Figure 3: Temperature Data in Southeast Asia [1]




Figure 4: Agricultural Cycle in Southeast Asia

- Stable Trait Assumption:** The model assumes that the traits of all organisms remain stable.
Explanation: Since the time span considered in the model is much shorter than the time required for evolutionary changes or mutations to occur, the traits of organisms are assumed to remain stable. This assumption also helps simplify the model.
- Stable Lighting Conditions Assumption:** The model assumes that the region under study experiences stable lighting conditions throughout the four seasons.
Explanation: Since the model focuses on tropical regions, the variation in day-light duration across different months within a year is minimal, thus the lighting conditions are treated as constant in the model.
- Stable Growth Environment Assumption:** The model assumes that no natural disasters, which could significantly impact the agricultural ecosystem, will occur during the time frame considered.
Explanation: Natural disasters are considered low-probability events in agricultural activities. To ensure the generalizability of the model, natural disasters should not be considered.

2.2 Notations

Table 1: Notations

Symbols	Description
\mathbf{X}	Vector $[N_{wd}, N_{crp}, N_{pst}, N_{ins}, \dots, C_{hc}, C_{pc}]^T$ to describe the system, etc.
wd	Subscription for weeds
crp	Subscription for crops
pst	Subscription for pest(who consumes crops)
ins	Subscription for other insects(who consume weeds)
bd	Subscription for small birds(herbivorous)
Bd	Subscription for huge birds(carnivorous)
bt	Subscription for bats
snk	Subscription for snake
frg	Subscription for frog
HC	Subscription for herbicide
PC	Subscription for pesticide
C_i	Concentration of certain chemical
N_i	Numbers of certain species
W_i	Biomass of certain species
w_i	Average mass of individuals of certain individuals
r_i	Natural growth gate of certain population
\mathcal{K}_i	Carrying capacity of certain population
α	The effect of chemical concentration on growth rate
$\beta_{i \rightarrow j}$	Interspecific competition factor
γ	Activity of decomposer
A_i, B_i	Effect of the predator-prey relationship on population i (specified coefficient).
N, P, K	Chemical elements

3 Models

3.1 Model Development Explanation and Sequence

3.1.1 Notes for Models

Before building all the models, here are some notes for all the models.

First, biomass is chosen over population density. This is because rice, as the primary producer in the agricultural ecosystem, has its population density artificially determined (i.e., it does not reproduce). Only the changes in its total biomass can reflect the developmental trend of the rice population.

Second, regarding interspecific competition, since the agricultural ecosystem is relatively simple compared to others, for the sake of model simplification, only interspecific competition between rice and weeds is considered.

Next, for the purpose of further simplifying the model, some parameters are combined. For example, the predation rate is incorporated into the predation coefficient of the prey

species, denoted as A .

Finally, since consumers are in a natural reproductive state with a relatively fixed age structure, the average individual biomass can be assumed to be constant. Therefore, biomass is directly proportional to the number of individuals in the population.

It should be noted that at the start time, all the variations and coefficients mentioned above are positive real numbers.

3.1.2 Sequence

Based on the conditions outlined above, we will gradually develop the model. Given the numerous factors involved in the final model, we adopt a step-by-step approach, starting from the simplest and progressing to more complex structures.

Initially, we will construct Model I: LVLG Model, based on the Lotka-Volterra model and the Leslie-Gower model, under the simplest food web conditions. This model will consider only producers, primary consumers, and secondary consumers.

Subsequently, as the ecosystem gradually develops, we will build Model II. In this model, we will incorporate the agricultural cycle (three crops per year), seasonal rotation (rainy and dry seasons), and the use of chemicals (herbicides and pesticides).

Next, we will consider the re-colonization of species and develop a more complex food web to form Model III. In this model, we will retain the features of Model II while considering the intricate interactions between multiple species and between species and the environment.

Finally, after the agricultural ecosystem matures, we will introduce human agricultural decisions, such as the removal of pesticides and the intentional introduction of new populations, to construct Model IV.

3.2 Model I: LVLG(Lotka-Volterra and Leslie-Gower) Model

We will first discuss the initial conditions of the model from the perspective of biological populations. The vertical structure of tropical forests is typically divided into several layers, including the canopy layer, understory, shrub layer, and ground layer. Each layer not only supports different plant species but also provides habitat and food sources for various animals. Deforestation will severely disrupt the vertical structure of the tropical rainforest food web. Therefore, our model assumes that at the initial time point following deforestation, the vertical structure of the ecosystem above the ground retains only a portion of the ground layer. All populations that previously depended on the canopy, understory, and shrub layers for habitat and food sources will have migrated out of the ecosystem.

Based on this assumption, the agricultural ecosystem in its initial food web retains only the following populations: plants, insects, and birds. First, to preserve interspecific competition and align with reality as closely as possible, we divide plants into two populations: rice and weeds. Second, although insects may exhibit food preferences, for simplicity, we assume that there is only one insect species that feeds on both crops and weeds. Finally, based on the feeding habits of birds in real life, we assume that birds feed

on both insects and the two plant species. Figure 5 presents a schematic diagram of the ecosystem, and Figure 6 shows the initial food web of the ecosystem.




Figure 5: Schematic map for LGVG Model




Figure 6: Food web in LGVG Model

Set February-when rice has just been planted-as the start time. Around the start time, in the simplest case, when climate, soil, and other conditions are favorable, only biological factors should be considered. If the population sizes of producers and primary consumers are used to describe the entire system, the Lotka-Volterra Model [2] and Leslie-Gower Model [3] can be applied as follows:

$$\begin{aligned}
 \frac{dW_{crp}}{dt} &= r_{crp}W_{crp} \left(1 - \frac{W_{crp} + \beta_{w \rightarrow c}W_{wd}}{\mathcal{K}_{crp}} \right) - \frac{A_{crp}W_{crp}W_{ins}}{1 + B_{crp}W_{crp}} \\
 \frac{dW_{wd}}{dt} &= r_{wd}W_{wd} \left(1 - \frac{W_{wd} + \beta_{c \rightarrow w}W_{crp}}{\mathcal{K}_{wd}} \right) - \frac{A_{wd}W_{wd}W_{ins}}{1 + B_{wd}W_{wd}} \\
 \frac{dW_{ins}}{dt} &= r_{ins}W_{ins} \left[1 - \frac{D_{ins}W_{ins}}{1 + E_{ins}(0.6W_{crp} + 0.4W_{wd})} \right] - \frac{A_{ins}W_{ins}W_{bd}}{1 + B_{ins}W_{ins}} \\
 \frac{dW_{bd}}{dt} &= r_{bd}W_{bd} \left[1 - \frac{D_{bd}W_{bd}}{1 + E_{bd}(0.2W_{ins} + 0.2W_{wd} + 0.6W_{crp})} \right]
 \end{aligned} \tag{1}$$

In general, these four equations introduce the natural growth term

$$r_i W_i \tag{2}$$

for population growth. The first two equations incorporate the interspecific competition term

$$\beta_{i \rightarrow j} W_i, \tag{3}$$

the larger this term, the more intense the interspecific competition, and the slower the biomass growth rate of population i . The first three equations consider the effect of predation through the term(e.g. the first one)

$$\frac{A_{crp}W_{crp}W_{ins}}{1 + B_{crp}W_{crp}}. \tag{4}$$

The last two equations(e.g. the third one) include the term

$$\frac{D_{ins}W_{ins}}{1 + E_{ins}(0.6W_{crp} + 0.4W_{wd})}, \tag{5}$$

which represents biomass reduction due to food scarcity. In the denominator, one factor is weighted according to the predation ratio: the less food available, the more predators there are, resulting in a slower growth rate of the predator population.

3.3 Model II:




Figure 7: Food web for NGLG Model

3.4 Model III:

3.5 Model IV:

4 Application of the Models

5 Sensitivity Analysis

6 Evaluation of the Model

6.1 Strengths

6.2 Weaknesses

7 Conclusion

References

- [1] World Bank Group. Indonesia-climatology-climate change knowledge portal, 2023. Accessed:2025-1-25.
- [2] Peter J Wangersky. Lotka-volterra population models. *Annual Review of Ecology and Systematics*, 9:189–218, 1978.

-
- [3] Lei Guo, Zi-Gen Song, and Jian Xu. Complex dynamics in the leslie–gower type of the food chain system with multiple delays. *Communications in Nonlinear Science and Numerical Simulation*, 19(8):2850–2865, 2014.