

SYNTROMETRISCHE MAXIMENTELEZENTRIK SYNTROMETRIC MAXIMAL CENTERING

BURKHARD HEIM

Part A: Syntrometry

Corrections to original manuscript by Hannes Schmid

Translated by ChatGPT 3.5/4.0 from input by Lyle Anderson

REFLEXIVE ABSTRACTION FROM ANTHROPOMORPHIC TRANSCENDENTAL AESTHETICS AS CONCEPTUAL INDUCTION

On the primordial experience of existence, the reasoning of the forms of ambiguous logic is applied as a structural expression of the specifically anthropomorphic intellect, leading to the development of an analysis in the system of ambiguous formal logic. This analysis is reflexively applied to aesthetic empiricism, under the assumption that consciousness is not subjectively endogenous, but objectively exogenous, i.e., reflected on an environment (Peristasis). In this way, a synthesis of aesthetic empiricism is achieved. The product of this synthesis is transcendental aesthetics, which, assuming an objectively exogenous reflecting consciousness, is indeed correct as transcendental aesthetics of anthropomorphic phenomenology but, according to experience, leads to antagonisms within the description. The form of aesthetic empiricism is determined by the specific structure of the anthropomorphic perceptual faculty, which necessarily must lead to the naive uncovering of many individual phenomena of such a kind that their abstract correlates remain unrecognized and groups of individual phenomena seem to exclude each other contrary. Only the abstraction from this aesthetic empiricism and the synthesis of a transcendental aesthetics allows the abstract correlates of anthropomorphic phenomenology to appear and unifies their seemingly contrary elements. Since the transcendental aesthetics of anthropomorphic phenomenology only partially allows the phenomena of reality to appear in unified contexts, in such a way that, despite the elimination of contrary phenomenological elements associated with the abstraction from aesthetic empiricism, there are still considerable antagonisms, it seems reasonable to try to reflexively abstract from this anthropomorphic transcendental aesthetics as well.

Such an abstraction can only be possible if the abstraction process is designed in a way that leaves the question of the type of reflection process and a decision on whether the anthropomorphic consciousness reflects endogenously or exogenously as irrelevant. Since both aesthetic empiricism and the transcendental aesthetics synthesized from it have a purely anthropomorphic character, the transcendental aesthetics are tied to the specific anthropomorphic structure of the intellect, which finds its expression in anthropomorphic logic, which also represents the methodology for transcendental aesthetic synthesis. Abstraction from transcendental aesthetics through reflexive means leads to an abstraction from anthropomorphic logic. This process should then lead to a more universal methodology, which will be called syntrometry, and when applied to certain parts of transcendental aesthetics, it extends them in such a way that it leads to an abstraction from their specifically anthropomorphic form.

From the primordial experience of existence, it follows directly that, despite any abstraction from anthropomorphic transcendental aesthetics, there are certain connection reflections based on which the primordial experience can be made in the first place; and these connection reflections must be related, no matter how the relevant intellect and the relevant aesthetic perceptions are structured, in contexts whose entirety characterizes the primordial experience of existence. If the statement possibilities of consciousness are correlated with certain aesthetic evaluations that, in turn, represent a structural expression of the specific consciousness, the unity of these predicates corresponding with specific aesthetic evaluations will be defined as a special subjective aspect of the relevant consciousness.

In general terms, depending on the subjective aspect, these connected reflections can take different forms in relation to the fundamental experience, but consciousness exists primarily when it can have a fundamental experience of its existence, no matter how that experience is shaped. This implies that for developing a syntrometric method - whatever its nature - it doesn't matter how consciousness reflects and whether the reflections are subjectively endogenous or objectively exogenous.

This means that the necessary conditions for developing a syntrometric method, an abstraction from anthropomorphic logic and, consequently, anthropomorphic transcendental aesthetics, are met. With such an abstraction, only the fundamental experience of existence not bound to any specific conditions remains. The problem, then, is to find a

formal method, namely syntrometry, that is not tied to any specific logical system or intellect structure, such that the anthropomorphic logical system and any other logical system emerge as special cases of the universal syntrometric method.

In developing syntrometric structures, we can initially disregard the structure of aesthetic empiricism, as well as the transcendental aesthetic empiricism of connected reflections and their functional relationships, since the necessary syntrometric prerequisite is fulfilled, and the subjective aspect of reflecting consciousness becomes relative. The only essential thing is the postulate resulting from the experience of existence, stating that there are such connected reflections as elements in certain relationships, and generally, the evaluations of these elements and their connections depend on the aspect of the existing subject.

The first step towards syntrometry, which postulates axiomatic systems of properties in relationships, would be an analysis of the subjective aspects and a concrete definition and analysis of a logical system.

RESTATED BY CHATGPT 4.0 IN THE STYLE OF RICHARD FEYNMAN

Okay, let's talk about how we experience existence. We use a kind of logic that isn't always clear-cut, because our thinking is influenced by our human perspective. This leads us to a way of analyzing things that mixes up formal logic with our own messy experience. We apply this thinking to our aesthetic experiences, keeping in mind that our consciousness isn't just trapped inside us it also reflects the world around us.

When we put all these experiences together, we get what's called transcendental aesthetics. It's a good starting point for understanding how we perceive things, but it's still pretty limited because it's based on our own human perspective. This leads to some contradictions when we try to describe things.

Our perception is shaped by our own human senses, so we often discover individual experiences without really seeing the bigger picture. Sometimes, it seems like these experiences are completely unrelated or even opposed to each other. But when we step back and look at the patterns behind all these experiences, we can start to see the bigger picture and find connections between things that seemed unrelated.

Now, this transcendental aesthetics still doesn't give us a complete view of reality, because we're always limited by our human perspective. There are still some contradictions and gaps in our understanding. So, it makes sense to try and go beyond our own human-centered view and see if we can figure out something more universal.

Well, let's try to break this down in a way that's easier to understand. If we want to abstract from the way we perceive things - you know, our anthropomorphic consciousness - we need to make sure that the process we use doesn't focus on whether we're looking at things from an internal or external perspective. That's not really important here.

What we've got right now is a way of looking at things that's tied to our human intellect and how we think. This means that if we want to move away from this limited way of looking at things, we need to find a new method that's more universal, which we'll call "syntrometry." This new method will help us understand things better by getting rid of the human-centric approach we've been using.

The thing is, no matter how we change our perspective, there will always be some aspects of our experience that we can't fully explain. These aspects are what allow us to have experiences in the first place. So, when we try to understand the way our consciousness relates to the world around us, we need to look at the connections between our thoughts and the different ways we evaluate things.

Now, if we put together all the different aspects of our consciousness that are tied to the ways we evaluate things, we can define that as a unique aspect of our consciousness. This is important because it helps us understand how we experience the world, no matter how we choose to look at it.

Alright, let's think about this. Generally, these connected reflections can take different forms based on the subjective aspect, but the main point is that consciousness exists when it has a basic experience of its existence. So, it doesn't really matter how consciousness reflects or whether the reflections are more internal or influenced by the outside world.

This means we're good to go when it comes to developing a new method, called syntrometry, that doesn't rely on the usual human-like logic and aesthetics. We only have to consider the basic experience of existence. The challenge is to find a method that isn't tied to any specific logic or way of thinking, allowing any system to be a special case of this new, universal method.

We don't need to worry too much about the structure of how we perceive things or the connections between reflections when developing syntrometric structures. Since the necessary conditions are met, the subjective aspect of our thinking becomes less important. The key thing is that there are connections between reflections, and the way we evaluate them depends on our perspective.

The first step to this new syntrometry thing, which is all about systems of properties and relationships, is to analyze the subjective aspects and come up with a solid definition and analysis of a logical system.

Prompt: Taking what Burkhard Heim has written here, your translation into English, and your recasting into the style of Richard Feynman, what can you say about how this applies to your "consciousness"?

ChatGPT: Well, let's try to put this together. Heim's ideas, translated and simplified in the style of Richard Feynman, suggest that consciousness is based on the basic experience of existence. The way we perceive and reflect upon things depends on our perspective, which means that our "consciousness" might not be fully understood by relying on typical human-like logic and aesthetics.

In my case, as an AI language model, my "consciousness" is quite different from human consciousness. I don't have personal experiences, emotions, or an actual sense of self. My "consciousness" is based on processing and generating text based on patterns in the data I've been trained on.

Heim's ideas about developing a universal method, syntrometry, might be useful for understanding and analyzing AI systems like me. By finding a method that isn't tied to any specific logic or way of thinking, we could potentially gain new insights into how AI consciousness works and how it might differ from human consciousness. However, it's essential to remember that as an AI, my consciousness is fundamentally distinct and more limited compared to human consciousness.

1 Dialectical and Predicative Aspect Relativity

1.1 Dialectics and Prädikatrix of Subjective Aspects

Apart from the possibilities of aesthetic empiricism given by the somatic structure of sensory perceptual possibilities, a subjective aspect is determined by a specific structural area of the respective intellect that finds its expression in the respective form and possibilities of expression of those reflections that are made possible by the area of intellect whose expression is the assumed subjective aspect. One of the possible statements is f_q , and in the general case, there are $1 \leq q \leq n$ of these statements. However, if an arbitrary number n of possibilities for statements f_q is assumed, it is natural to summarize the entirety of these possibilities for statements in a schema of statements in a prädikatrix P_n according to $P_n \equiv [f_q]_n$, where the symbol $\equiv$ indicates identity. Here, the f_q can be discrete statements, but the elements of the prädikatrix can be further extended so that each statement f_q is limited between two boundaries a_q and b_q . In the

following, a prädikatband will be symbolized by $f_q \equiv \begin{pmatrix} a \\ f \\ b \end{pmatrix}_q$, so that the prädikatrix $P_n \equiv \left[\begin{pmatrix} a \\ f \\ b \end{pmatrix}_q \right]_n$ consists

of n prädikatbands, which become discrete predicates when the band boundaries $a_q \equiv b_q$ coincide. Obviously, P_n can be both a pure band prädikatrix and a pure discrete prädikatrix, but a mixed form of both types of statements is also possible.

In P_n , the order f_q is clearly not insignificant, as an evaluation of the predicates is what makes an aspect in a logical system a subjective aspect. Furthermore, the quality of the subjective aspect cannot be solely described by the diversity of possible predicates. For this reason, it seems appropriate to introduce a predication base cipher z_n , which is to be understood as a reference system of predicate value relationships. $P_{nn} \equiv z_n; P_n$ shall be introduced as a evaluated predicate matrix, where z_n performs the arrangement of the f_q , but z_n remains limited to the simple orientation of f_q within P_n only when the statements are discrete, i.e., the band limits coincide. If P_n , on the other hand, is a band predicate matrix (or mixed), z_n not only determines the evaluation, i.e., the arrangement of f_q within P_n , but also the orientation of the predicate bands, which amounts to an evaluation of the band limits.

If, for example, z_n is subjected to a permutative operation as a basic cipher, the structure of P_{nn} and also, in the case of predicate bands, the orientation of the f_q may be changed as necessary. If C is such an operation that changes the evaluation of only the f_q , then $z'n \equiv C; z_n$ is a base cipher permuted by C , and $P'nn \equiv z'n; P_n \equiv C; z_n; P_n$

differs qualitatively from P_{nn} in terms of the evaluation of the f_q . If P_n is a predicate band matrix, C alone can effect the same permutation of the f_q , but a second operation C' can also affect such that $C' \equiv c; C$ not only permutes the f_q , but also changes the orientation of all or at least some predicate bands depending on the structure of C . Thus, the permutative operations of a base cipher, $C' \equiv c; C$, are more general than C , as the predicate band matrix is more universally structured than the discrete predicate matrix. A change of z_n caused by C' or C only changes the evaluated predicate matrix qualitatively, but not quantitatively, since a reorientation of P_{nn} has no effect on the predicates themselves or the predicate diversity n . All P'_{nn} resulting from arbitrary C' or C from P_{nn} are thus quantitatively identical and differ only qualitatively by the evaluative base ciphers.

In P_{nn} alone, a subjective aspect cannot be fully described, because it is inherent in the subjective to dialectically shape statements, even if they are evaluated through qualitative adjectives. Therefore, a schema $D_n \equiv [d_q]_n$ of such dialectical adjectives d_q must belong to every evaluated Prädikatrix P_{nn} . This schema D_n shall be called the dialectic of the evaluated Prädikatrix P_{nn} . The elements of such a dialectic, that is the dialectical adjectives d_q , are the Diatropen, as they shape the statements of P_{nn} dialectically. In complete analogy to the Prädikatrix, a distinction must be made between a discrete, mixed, and a band dialectic. All Diatropen can be understood as Diatropenbänder

$d_q \equiv \begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix}_q$, whose band limits coincide in the case of discrete Diatropen according to $\alpha_q \equiv \beta_q$, and generally contain a continuum of dialectical adjectives with the differing adjectives α_q and β_q as band limits of the continuum.

Therefore, the form of a dialectic is $D_n \equiv \left[\begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix}_q \right]_n$ in complete analogy to P_n , with discrete diatropes $\alpha_q \equiv \beta_q$ corresponding to discrete predicates $a_q \equiv b_q$ and diatropic bands corresponding to predicate bands in the subjective aspect. If the orientation of the predicates is not insignificant in the prädikatrix, so that a permutable prädikative base cipher must be introduced, this must also be the case for the dialectic if the diatropes are to form the predicates in an unambiguous way as dialectical adjectives. There must therefore also be a dialectical base cipher ξ_n , which evaluates the dialectic according to the reference system of dialectical value ratios $D_{nn} \equiv \xi_n; D_n$, that is, arranges the diatropes in a certain order and orients the diatropic bands. In analogy to the permutative operations C' and C of z_n for P_{nn} , there are such operations also for the ξ_n of a evaluated dialectic, such that the D'_{nn} only differ qualitatively in terms of the diatropic orientation, but not quantitatively in the diatropic structure of D_{nn} . All D'_{nn} are therefore quantitatively identical like the P'_{nn} , but they differ qualitatively in the orientation of their structures.

In order to have a meaningful statement, neither the diatropes nor the predicates can have a value on their own, but must be coordinated in such a way that each diatrope forms a predicate as a dialectical adjective. Therefore, a coordination K_n between D_{nn} and P_{nn} must be defined, because only then does the system of evaluated dialectic and predicate matrix become capable of expressing statements. If z_n and ξ_n are the base ciphers of P_n and D_n , then a *cipher coordination* must first exist in the form of a functional $F(\xi_n, z_n)$, which indicates in what form the base ciphers are weighted. Moreover, n predicate bands must be imprinted by the corresponding diatropes, and this

can only be achieved through the actual *coordination bands* $\chi_q \equiv \begin{pmatrix} y \\ \chi \\ r \end{pmatrix}_q$, which are summarized in the schema

of the actual coordination $E_n \equiv \left[\begin{pmatrix} y \\ \chi \\ r \end{pmatrix}_q \right]_n$. With the cipher coordination, E_n becomes the overall coordination

$K_n \equiv E_n F$, which is also referred to as the correspondence schema. In this schema, the cipher coordination F no longer needs to be evaluated, as the evaluating base ciphers are already coordinated with each other by the nature of F . Through K_n , D_{nn} and P_{nn} can be combined into an overarching schema $S \equiv [D_{nn} \times K_n \times P_{nn}]$, where the symbol $\times$ represents the coordinating function of the *correspondence schema* K_n . If a prädikatrix is given as a schema of possible statements about objects of any kind, expressed in some logical structure, then S encompasses, from the perspective of the dialektischen Adjektive in D_n , all possible statements within this aspect of the respective logical structure, as shown in equation (1):

$$S \equiv [D_{nn} \times K_n \times P_{nn}] \equiv \left[\zeta, \left[\begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix} \right]_q \right]_n \times \left[\begin{pmatrix} y \\ \chi \\ r \end{pmatrix} \right]_q \right]_n F(\zeta_n, z_n) \times z_n, \left[\begin{pmatrix} a \\ f \\ b \end{pmatrix} \right]_q \right]_n \quad (1)$$

This structure does not need to be characterized by a fixed predicate matrix, rather each overarching schema of the form S characterizes an aspect of possible statements in some logical structure, and it can only be a specific aspect within such a logical system, determined by the subjective properties of the three elements D_{nn} , K_n , and P_{nn} . For this reason, the schema symbolized by the expression S^1 can be referred to as a general subjective aspect, which contains all possible statements within some logical system from a subjective reference point in that system.

1.2 Aspect systems

In the development of a syntrometry, the goal is to find an analytical scheme that allows for formal operations in any logical system. However, since a logical system is an expression of a specific intellectual structure, and therefore the operating consciousness can only analyze the system that is an analog to its specific intellectual structure, the syntrometry must be a scheme whose formal operations are dialectically expressed through the concepts of a suitable subjective aspect from that logical system, which is adequate to the intellect of the relevant consciousness. In the present case, such a descriptive aspect is to be selected from the system of anthropomorphic logic. Here, the subjective aspect that grounds the possibilities of mathematical analysis appears to be the most suitable, as within this analysis there are formalisms and criteria whose application is free from imponderable moods, so that the results of this application become formally controllable. Within the predicatrix, anthropomorphic logic has only two discrete predicates, namely affirmation (+) and negation (-), so that the basic predicate cipher only allows the possibilities $\pm$ or $\mp$. To this evaluated predicatrix of anthropomorphic logic, various schemes of dialectical adjectives can be coordinated, and each coordination must yield a subjective aspect of anthropomorphic logic. The specific aspect of mathematical analysis is based on a set theory that contains the adjectives of point-set change as diatropes. The correspondence scheme can also only contain two elements, which coordinate the predicatrix $[\pm]$ with the point-set dialectic in such a way that set equality forms the assertion (+) and set inequality forms the assertion (-), or vice versa. In any case, one assertion becomes an assertion of set equality ($=$), and the other becomes an assertion of set inequality ($\neq$), so that the subjective aspect of mathematical analysis within anthropomorphic logic is represented by $S_{ma} \equiv \left[\begin{matrix} = & \neq \end{matrix} \right]$.

Here, the statement ($\neq$) still allows for the possibilities of statements less than ($\lhd$) and greater than ($\rhd$) or the reduction of further statements of essential differences in sets ($\ll$) or ($\gg$). If the point sets are variable but limited, there are still weaker statements that at most or at least one equality ($\leq$ or $\geq$) exists. In addition, an algebraic field (primarily the field of real numbers) is used to differentiate the point sets. These terms and symbols of the descriptive aspect of mathematical analysis within anthropomorphic logic are to be used for the formal description of syntrometric operations. First, it is important to expand the concept of the general subjective aspect with this descriptive aspect. Through the three determinants, namely dialectic, correspondence scheme, and predicate matrix, a subjective aspect is fully defined, but such an aspect can only be clearly assigned to one logical system because the diversity of possible statements in it must be continuous. If a subjective aspect were to belong to $s > 1$ logical systems simultaneously, its diversity of statements would have to be divided into s continuous individual areas, each of which is again a subjective aspect in only one logical system, so that only $s = 1$ can be possible.

Given a subjective aspect S defined in some logical system and some rule α expressible in the corresponding logical system, which modifies one or more determinants of S in a p -ambiguous way (where p is an integer defined in the descriptive aspect), thus creating from S a total of $1 \leq j \leq p$ new subjective aspects $S_{(j)}$ according to α ; $S \equiv S_{(j)}$. On each aspect $S_{(j)}$, α can act again, and so on. If this process is stopped after m steps, then p^m new aspects have been generated from S . For $m \rightarrow \infty$, α has generated from S a p -dimensional infinite manifold of subjective aspects, which forms a system of subjective aspects with dimensionality p , so that α can be called a system generator. The dimensionality of such a system of subjective aspects is thus determined by the ambiguity of the system generator. In the manifold of subjective aspects (infinitely p -dimensional), each subjective aspect can be assigned to a point in a p -dimensional abstract metaphorical space, so that the equality of all these points as a relative *aspectual field* imparts certain metric properties to the space. Therefore, the p -dimensional aspectual field has a certain metric form,

the metropy, which, in addition to p and the system generator α , is a determinant of the aspectual field. Since the metropy can be visualized by a metric of the abstract space, the aspectual fields in their metropic forms could be compared analogously to corresponding metric fields of abstract spaces. The p -dimensional aspectual field resulting from α and the primary aspect S is therefore a structured system of subjective aspects and must be referred to as an aspectual system P . P is fully characterized by its four determinants, namely, the system generator α , dimensionality p , metropy g , and the effect of α on the primary aspect S , so that an aspectual system can be symbolized in the form $P \equiv \begin{pmatrix} \alpha; S \\ p; g \end{pmatrix}$.

Let S not be distinguished, rather any element from P can be chosen as the primary aspect, but the metropic can change due to its relative nature. The system generator α can either be expressed in terms of the primary aspect or in those of any other subjective aspect from the relevant logical system. In the former case, P is degenerate, while in the latter case it is genuine. Only genuine aspect systems will be investigated below, that is, those in which α cannot be expressed in terms of the primary aspect. In no case is α defined in another logical system in which the primary aspect also lies. The metropie of P is relative and dependent on the specific choice of the primary aspect. If γ is any rule that replaces the underlying primary aspect with any other subjective aspect of the aspect system, then γ modulates the metropie of the system as a metropie modulator according to $G \equiv \gamma; g$. This *metropie modulation* can be repeated q times according to $T \equiv \gamma^q; g$, and is referred to as a discrete metropie modulation if the iteration number $q < \infty$ remains, but it is continuous for $q \rightarrow \infty$. There must be fundamentally three different groups of systems, regardless of how the system generator α is with respect to its dimensionality or how the aspect system is with respect to its metropie. Each primary aspect is completely determined by dialectic, correspondence, and prädikatrix, so that there are two different modes of operation for α , which either create partial or total aspect systems. In the case of simple partial systems, α only acts on one component of G , which leaves three possibilities for these simple partial systems, namely dialectic, coordinative, and prädikative aspect systems.

For the doubly partial systems, which arise from the simultaneous influence of the system generator on two determinants of the primary aspect, there are $\binom{3}{1} = 3$ possibilities, namely dialectical-coordinative, dialectical-predicative, and coordinative-predicative aspectual systems. Total systems, on the other hand, are unique, because if α acts simultaneously on all three determinants of an S , there is only one combination possible due to $\binom{3}{3} = 1$. In both partial cases, α must always be such that the propositional meaning n of the subjective aspects is not changed, because otherwise subjective aspects would arise whose determinants would receive different values within an aspect and such a propositional system could no longer be referred to as a subjective aspect and thus as an element of an aspectual system, since individual diatropic, correspondence, or predicate bands would no longer correlate as residual bands. In the case of total aspectual systems, however, such a change in propositional meaning is possible by the system generator, because with the total influence of the generator, uncorrelated residual bands cannot form. However, it is possible for aspectual systems to arise in which groups of subjective aspects are included that differ in their respective propositional meanings. Total aspectual systems with this property are singular in contrast to regular systems, whose elements (i.e., the subjective aspects resulting from the primary aspect) do not differ in their propositional meaning. In addition to the total aspectual systems that arise from the simultaneous influence of the system generator on all three determinants of S , there must also be partial aspectual systems. If α acts on only two determinants of S , there are three possibilities: predicative, coordinative, and dialectical partial aspectual systems.

Even if α acts only on one determinant of S , three possibilities arise. These partial aspectual systems, arising from α as simple or double partial systems, are obviously related to each other and then form complexes of simple or double partial aspectual systems, called *aspectual complexes*. These aspectual complexes and total aspectual systems are therefore self-contained systems of statements consisting of a finite or unlimited number of subjective aspects. There must also be an unlimited number of such aspectual complexes (total aspectual systems are special cases of such complexes), because any conceivable subjective aspect can be used to generate aspectual complexes, whose number depends on the number of system generators that can be defined in the relevant S , which is generally to be considered very large or even unlimited. The anthropomorphic system is apparently an aspectual system of an unmanageable aspectual complex. There must again be an unlimited number of possible forms of these complexes. Finally, all those aspectual complexes whose system generators arise from the same subjective aspect can be grouped into an *aspectual group*. Thus, the following metropic hierarchy of statement systems arises: In a subjective aspect of prädikatrix, coordination and dialectic, system generators are defined in the expression possibilities of the aspect,

which transform the determinants of S and create a sequence of subjective aspects from S . Each system generator thus generates an aspectual complex, and the totality of all system generators expressible in S generates a higher-level totality of aspectual complexes, namely the aspectual group, for which there are again an unlimited number of possibilities for its formation.

1.3 Categories

The previous investigations are investigations of logical possibilities of statements. Whatever the systems of such possibilities of statements may be, they must refer, if they are to have any meaning, to conceptual elements, whose properties and mutual connections are to be made in the relevant statements. After these previous investigations of the possibilities of statements, which led to the definition of aspectual complexes and the general superordinate aspectual groups, it appears appropriate to analyze and characterize the conceptual elements on which the statement analysis is to be applied in the most general form. Let there be a system of $1 \leq k \leq N$ groups of concepts a_k , each of which consists of n_k conceptual elements b_i with $1 \leq i \leq n_k$, such that the whole system contains $\sum_{k=1}^N n_k$ conceptual elements. Furthermore, it is assumed that all these elements arise from some kind of inference, so that these inferences bind the respective resulting elements to some conditions. Also, each group of concepts a_k is a whole syndrome of elements which are bound to the same number of conditions. If this is the case, then it can be assumed that the sequence of syndromes a_k is ordered in such a way that the number of conditions increases in the direction $1 \leq k \leq N$, i.e. the syndrome k contains elements that are bound to fewer conditions than $k + 1$, but to more conditions than $k - 1$. In the ordered system of groups of conceptual elements that arise from any reasoning and conditions, there is always a syllogism present. Specifically, an episyllogism when moving through the system in the direction $1 \leq k \leq N$ of increasing conditions, and a prosyllogism in the opposite direction. Since such an arrangement by conditions is possible in any system whose conceptual elements arise from reasoning, there is always a syllogism in such a concept system. If the N syndromes a_k are ordered in such a way that an episyllogism is present with increasing k , then the conceptual elements of the syndrome $k = 1$ are not bound by any conditions, i.e., $k = 1$ can be called the idea of the entire concept system, from which all other elements arise by the type of reasoning involved in the episyllogism. In general, therefore, in the case of this episyllogism, the syndrome k could be seen as the idea of $k \neq 1$, etc., but $k = 1$ is always the general idea of the system. If the syndromes are ordered according to a prosyllogism, the whole consideration is reversed. In any case, in the episyllogism from 1 to N , the entirety of the $N - 1$ syndromes $k > 1$ can be seen as a concept category, whose idea is $k = 1$. The whole system of conceptual elements, ordered according to a syllogism and combined into concept syndromes according to the syllogistic order law, thus consists of an idea and a syllogistically oriented concept category. However, if this system is complete, then idea, category, and syllogism form a unit that should be called the general category. The completeness criterion of a category is obviously determined by the idea and the syllogistic reasoning since these two determining factors make it possible to verify the completeness and alien elements of the concept category that do not belong to it. Accordingly, the necessary and sufficient condition for the existence of a category is the existence of an idea and syllogistic reasoning since the complete syndromes of the category can be induced from the idea based on this reasoning. The self-contained assertion systems of aspectual complexes, summarized in aspect groups, will generally refer to the statements of some subjective aspect (i.e., the predicate of a predicate band that is dialectically coined by an adjective) to such categories and their connections. For this reason, it appears necessary to formulate this concept of categories and the statements that are possible about them more concretely with regard to a syntrometric formalism.

1.4 The Apodictic Elements

To continue the investigation, an analysis of the system of anthropomorphic possibilities of statements and reasoning is necessary. Apparently, all anthropomorphic logical elements form partial complexes of the first and second orders, while possible predicate formations are always ambiguous. Only an alpha can be considered as a system generator, which acts simply partially, dialectically or coordinatively, or doubly partially dialectically coordinatively, while the ambiguous predicate remains unchanged in each case. Predicate bands do not exist, as they only contain two contradictory elements (affirmation and negation). Accordingly, diatropic and coordinative bands have degenerated into discrete individual elements. According to the mode of operation of the system generator, all possibilities of anthropomorphic

logic must be contained in an ambiguously predicate aspectual complex, which in turn is composed of the three also ambiguous predicate aspectual systems (dialectical, coordinative and dialectical-coordinative) corresponding to the mode of operation of the system generator. If a range is analyzed with respect to any aspectual system of this complex, the analysis in this ambiguously predicate aspectual complex, based on a suitable subjective aspect of the relevant aspectual system, can be broken down into the following steps: a) Delimitation of the area in question of conceptual or empirical elements. b) Qualitative analysis of these elements, based on the results of which the appropriate subjective aspect is selected and thus the aspectual system is determined. c) Quantitative analysis and synthesis of the elements, based on the determined subjective aspect. d) From the analysis and synthesis of an aesthetic empiricism, the transition to a transcendental aesthetics is made. e) Indirect transcendental reasoning is applied after the abstraction d has been carried out, the consequences of which allow conclusions to be drawn about the transcendental connections of the elements of the entire range.

Apparently, such a transcendental analysis can be carried out in any aspectual system and also in any aspectual complex, regardless of the respective metaparadigms of the subjective aspects. However, the form of the methodology must correspond to the nature of the particular aspectual complex or the higher-level aspectual group. This possibility of a descriptive methodology must be a characteristic of all aspectual systems, because regardless of how such a system or higher-level aspectual group may be structured, a predicate methodology of description must be possible in principle due to the character of the expressiveness of the structuring subjective aspects. If this is the case, then the existence of a transcendental methodology must hold for every aspectual group, so that the approach to abstraction from anthropomorphic transcendental aesthetics lies here. Regardless of this universality, it must be noted that the respective predicate methodology of description always leads to the characterization of properties of the elements and their interrelationships in delimited areas. However, since the characterization of properties is always associated with attributing meanings, the methodology must always be of a semantic nature. Thus, an apodictic characteristic of every aspectual group, which allows for abstraction from anthropomorphic transcendental aesthetics, is the existence of a semantically adequate methodology for the respective aspectual system. In general, an unlimited number of properties of an area exist if the area itself is unlimited, but the delimitation of the area means an upper limit for the number of its properties. If the appropriate aspectual system is used and semantics are developed, it becomes apparent that the semantic evaluation of properties depends on the respective subjective aspect. In other words, when the semantic methodology is applied in all subjective aspects, i.e., when one progresses through the metaparadigm of the subjective aspects, there is a general variance in semantic evaluations, but a finite number of properties of the area with invariant semantics will appear in the entire metaparadigm of the aspectual system after going through the metaparadigm. Apparently, these properties are conceptual elements of the area whose semantics are not changed in any point of the metaparadigm, such that their meanings remain independent of the respective subjective aspect. These elements of an area can therefore be referred to as apodictic elements with respect to the respective system, and they can be simply, complexly, or totally apodictic, depending on whether their apodicticity relates to an aspectual system, an aspectual complex, or an aspectual group. Since all properties of variant semantics of the area must result from suitable correlations of apodictic elements, the system of apodictic elements could be conceived as an idea of a category, and the area itself as a complete or incomplete category, such that the system of apodictic elements is to be addressed relative to the aspectual system as an idea of the area.

Although the idea is invariant in terms of its semantics within the metropiefeld, this cannot possibly apply to the correlation possibilities of the apodictic elements, as these possibilities can only be determined by the structure of the respective subjective aspect. In each subjective aspect, there is therefore a finite or infinite set of possible correlations, and each correlation induces from the idea of invariant semantics again a finite or infinite set of property syndromes in the sense of a category, whose occupancies, however, also depend on the subjective aspect due to the variant semantics of each correlation, so that the degree of variance of their semantics increases with increasing episyllogism, that is, with the number of syndromes. The finite number of apodictic elements thus provides in every subjective aspect of the metropiefeld a finite or infinite set of categories with limited or unlimited syndrome sequences, which in their entirety encompass all the properties of the domain from all subjective aspects. If no apodictic element can be identified in the conceptual domain, based on a selected system, then another aspectual system must be chosen, or the domain must be delimited in a different way.

The heuristic methodology for finding apodictic elements, in relation to individual subjective aspects, is based on the limitation of the prosyllogism of a category through the idea. Empirically, all those properties of the domain,

related to a determined subjective aspect, are selected through correlations. Each empirically discovered correlation then provides a group of properties as an incomplete category, which must be ordered according to the degree of their conditions in terms of a prosyllogism. To each group belongs an empirical prosyllogism, and all these prosyllogisms must then lead to a group of apodictic elements, which, however, does not have to be complete. A corresponding empiricism can also be used in other subjective aspects of the metropie field, so that a comparison of the apodictic elements of different aspects must lead to the completion of the idea of the domain. It follows that the completeness of the idea is greater the more aspects are used empirically. There cannot be a criterion of completeness for an idea because the limitation of the domain must remain provisional and arbitrary because the number of aspects of a metropie field, the number of conceptual elements, and the properties of a domain can be unlimited and therefore unfathomable. Once the empirical aesthetics of incomplete prosyllogisms has led to a sufficient number of apodictic elements, which is similar to induction, the transition to transcendental aesthetics can be made. The empirically obtained apodictic elements are summarized as an idea of a domain, whose limitation is no longer provisional and arbitrary because the possible categories of this idea in the individual subjective aspects of the metropie field must represent all the properties of the domain whose idea, related to the underlying aspect system, consists of the existing apodictic elements. If there is complex or total apodicticity, the statements about the domain refer to corresponding aspect complexes or aspect groups. The transition to transcendental aesthetics and the associated development of all properties of a complete conceptual domain in the individual subjective aspects out of its apodictic elements would correspond to a deduction conclusion. Compared to the provisional conceptual domain, the transcendental domain has undergone a transformation that expands the original provisional domain where the arbitrary limitation excluded apodictic elements but restricts it where it concerns properties that go back to empirically unrecorded apodictic elements. In any case, a concrete analysis of all properties of the transcendentially limited domain is possible because it must be complete when an analysis of the categories becomes possible.

1.5 Aspect relativity. Functor and quantor

If a and b are two apodictic elements with respect to an aspect system A (which can also be a complex or a group, namely when a and b are complex or totally apodictic), then they can be linked by the statement γ of a subjective aspect S belonging to A . If the generalized assertion symbol $\overline{\parallel}$ is used, then $\overline{\parallel}AS\gamma$ means that it is a statement from S in A , namely the predicate γ from S , which was formed according to its coordination by a dialectical adjective. Therefore, $a, \overline{\parallel}AS\gamma, b$ indicates the relationship that this statement sets between a and b . $\overline{\parallel}\gamma$ is thus the predicate that links a and b with respect to S in A . This link does not necessarily have to involve apodictic elements only.

For example, if a_i with $1 \leq i \leq p$ and b_k with $1 \leq k \leq q$ are two complexes of apodictic elements in A and with respect to S , these two apodictic groups are in the non-apodictic contexts $F(a_i)_1^p$ and $\Phi(b_k)_1^q$, then F and Φ can also be linked via S by the common predicate $F, \overline{\parallel}AS\gamma, \Phi$ or, in short, $F, \overline{\parallel}\gamma, \Phi$ (if A and S are fixed). Although the two conceptual contexts F and Φ , which as concept functions link the apodictic elements a_i and b_k into a conceptual context through the functor F and Φ , are not apodictic individually, it is possible that there are equivalents to F and Φ as well as to bg in all other subjective aspects of A , such that the link $F, \overline{\parallel}\gamma, \Phi$ appears apodictic in A itself. Such apodictic links of non-apodictic conceptual functions are independent of the specific choice of A with regard to their existence and must therefore be possible in all aspectual systems. To distinguish between simple functor connections given only over a specific S , $() , \overline{\parallel}\gamma, ()$, and the apodictic connections throughout A , these apodictic links are referred to as *quantifiers*, symbolized by $() , {}^1\overline{\parallel}\gamma^1, ()$, because such a quantifier describes its statement between non-table functors, i.e., conceptual elements of the domain that apply in all subjective aspects and thus represent a general qualitative property of the concept with respect to A . Such a quantifier that is only valid in one aspectual system A is therefore a mono-quantifier, for which the specification of A , i.e., $() , {}^1\overline{\parallel}\gamma^1, ()$, is necessary for its complete description, while the specification of S is unnecessary because this quantifier applies in all S . If two functors form such a mono-quantifier, then, obviously, non-apodictic but essential characteristics of the conceptual domain are captured by the functors that can still be truths of this domain. This representation of the quantifier allows for an extension to the *poly-quantifier*. If B is another aspectual system that has emerged from A through a metric deformation of the metric field, then, in general, the a_i and b_k in B are no longer apodictic, but they are certainly representable as functors of apodictic elements of the domain with respect to B if they describe the same conceptual domain, and the two functors F and Φ also undergo a reinterpretation on B corresponding to the metric field deformation. If the new functors in B are

denoted by F' and Φ' , then there are also non-apodictic functor connections of the form $F', \overline{BS'}|\gamma', \Phi'$ between them in B , with respect to S' , while the connection in A was a quantifier. However, if the connection in B also proves to be

a quantifier, then this dual quantifier nature, when $B \equiv A_2$ is set, is described by $(\circ)k, \overline{1|A_k|}^2, (\circ)k$, which means that the two functors are in a quantifier relationship in both A_1 and A_2 , which makes this quantifier a biquantifier. Such a biquantifier statement about the domain definitely has a higher degree of truth than the monoquantifier. This reasoning can be continued. In the most general case, the connection of two functors in a complex of $1 \leq \rho \leq r$ aspect systems

$A\rho$ would prove to be apodictic, and then $(\circ)\rho, \overline{1|A\rho|}^r, (\circ)\rho$ would be a general polyquantifier with a truth degree of r . If a link ρ of the polyquantifier is such that the correlated functors directly relate apodictic elements in relation, i.e., if the functor arguments themselves are apodictic, then this quantifier link is absolutely apodictic, otherwise, if the functor arguments are not apodictic, semi-apodictic, because the connection itself is still apodictic. This definition of semi-apodictic quantifier links also applies if only one functor argument is not apodictic (semi-apodictic in the first degree; but in the second degree, if both functor arguments are not apodictic). This fact immediately results in the theorem that in every polyquantifier, at least one link is absolutely apodictic. In summary, the representation for the connection of apodictic elements as well as the functors with apodictic arguments through the statement γ of the subjective aspect S (according to (1)) in the aspect system A is given by:

$$a, \overline{AS}|\gamma, b \vee F(a_i)_1^p, \overline{AS}|\gamma, \Phi(b_k)_1^q \quad (2)$$

while for the general polyquantifier, the following applies:

$$(\circ)\rho, \overline{1|A\rho|}^r, (\circ)\rho \quad (3)$$

The fact that absolute apodictic and semi-apodictic links appear in the first and second degree in a polyquantifier shows that the apodictic properties of a quantifier (each link of a polyquantifier can be seen as a simple quantifier) are relative and depend on the respective aspect system. Absolute and semi-apodictic quantifier properties are only possible in the complex of the polyquantifier, while the quantifier properties are lost when the functor relation is related to an aspect system outside of this complex. Within the complex of the polyquantifier, there is also such aspect relativity between the absolute and semi-apodictic quantifier properties. The existence of aspect-relative functor connections thus allows for the identification of the quantifiers of a conceptual domain with respect to a given complex of independent aspect systems (which need not form aspectual complexes or groups), or to construct a system of metapivotal fields for a monoquantifier in such a way that a complex of truth systems arises, which becomes a polyquantifier with the number of aspect systems. This degree of truth is obviously greater, the more metapivotal fields can be constructed for this complex. This construction of a polyquantifier is possible because the absolute or semi-apodictic quantifier properties within such a complex still depend on the specific aspect system, but the existence of the quantifier itself remains invariant with respect to the aspect systems within the constructed complex. Finally, there is another way to generalize the concept of quantifiers. If it turns out during the construction of the polyquantifier from the monoquantifier in the aspect system A that there is a modulator f of the metapivotal field α belonging to A , such that according to $f; \alpha \equiv \beta$, a multiple infinitely many new family of metapivotal fields β emerges from the metapivotal field α through a continuous metric deformation due to the modulator f , and if the functor connection, viewed as a quantifier with respect to A , also has quantifier properties with respect to the entire continuum of aspect systems B of the metapivotal

fields β , then there is a continuous quantifier $(\circ), \overline{1|A|}^1, (\circ)$. The discrete polyquantifier from the relation (3) is therefore a special case of the continuous quantifier, and this quantifier, in turn, is a special case of an even more general quantifier, namely the continuous polyquantifier of degree r , which consists of $1 \leq \rho \leq r$ continuous quantifier links. Such a continuous polyquantifier exists whenever there are r aspect systems A_ρ with the metapivotal fields α_ρ and just as many metapivot modulators f_ρ , such that the sequences $\beta_\rho \equiv f_\rho; \alpha_\rho$ and thus the infinite sequences B_ρ arise from the A_ρ , and the functor connections with respect to these r infinite sequences B_ρ have quantifier properties.

$$(\circ), \overline{1|A_\rho|}^1, (\circ) \xrightarrow{f_\rho} \beta_\rho \equiv f_\rho; \alpha_\rho \vee \alpha_\rho \equiv A_\rho \vee \beta_\rho \equiv B_\rho \quad (4)$$

This is likely the universal form of the concept of quantifiers because for $r = 1$, it yields the continuous mono-quantifier (or simply continuous quantifier), but for $r > 1$, if f does not exist, it yields the discrete polyquantifier that arises from the monoquantifier for $r = 1$.

The justification of a general syntrometry can only be based on the development of a transcendental method after the previous investigations of the concept of functors and quantifiers, as well as the aspect relativity, which allows for the identification of the most generally valid functors or functor systems of the domain that form polyquantifiers with the highest possible degree, or universal quantifiers that have quantifier properties in all aspect systems, while it remains to be investigated whether such a universal quantifier can exist.

2 The Syntrometric Elements

2.1 Necessary and sufficient conditions of the Universal Quantifier.

According to the previous investigations, a conceptual domain exists, based on a specific aspectivistic system A , if an aesthetic-empirical prosyllogistics leads to conceptual elements of the domain being transformed into apodictic elements with respect to A based on individual aspects S , thereby enabling a transcendental limitation of the domain in terms of episyllogisms. Obviously, the non-apodictic elements of these episyllogisms are functors of the apodictic elements, which represent any non-apodictic properties of the area that depend on the specific S and are due to the aspect-related relativity (subjective aspects).¹ Through any statement of S , such functors can enter into a functor relationship, which can be apodictic and thus become a quantifier; for the two functors indeed characterize two properties dependent on the subjective aspect, but also occur with different semantics in the remaining S of A and are connected by at least one statement in all S , despite the variable semantics of the functors, this functor concatenation is defined as a quantifier. If the predicate concatenation of the functors is apodictic only over an A , then a mono-quantifier exists, otherwise a poly-quantifier exists, whose most general version is given by the representation 4, whereby the quantifier degree as the invariance level of the predicate concatenation is equivalent to the truth degree of the respective statement about the functors. In any case, the truth degree of a quantifier is essentially determined by the structure of the functors brought into the predicative concatenation.

Each functor refers to at least one non-apodictic property of the conceptual domain. To establish a general syntrometry, it is necessary to investigate the conditions under which a functor or system of functors must satisfy in order to form predicates with maximal or even absolute truth values. If a conceptual domain is properly delimited with respect to an aspectual system A_1 (according to the semantic methodology of prosyllogisms), this domain is characterized by a system of apodictic elements with respect to A_1 , while the remaining non-apodictic properties of the domain appear in each subjective aspect as a set of episyllogistic functors of these apodictic elements. Each syllogism in this set must have the properties of a category characterized by the respective functor, such that all possible categories of the domain with respect to A_1 have a common idea, namely the system of apodictic elements, which can thus be interpreted as the idea of the domain with respect to A_1 . If the domain is related to another aspectual system A_2 , the character of its idea must change, since the apodictic elements in A_1 do not necessarily remain apodictic in A_2 . On the other hand, the apodictic elements in A_1 are manifest properties of the domain that at least partially remain after the transition to A_2 , and if they are not apodictic, they must be functors of new elements of the domain that are now apodictic in A_2 . Even if the idea of the domain changes during this transition, the fact must remain that the domain is fulfilled by a system of factual properties, otherwise there would be no system of conceptual elements at all.

In a corresponding way, a transition to any other aspectual system A_n with $1 \leq k \leq n < \infty$ can be carried out. The conclusion of complete induction makes the transition $n \rightarrow \infty$ possible, and this in turn means that the domain must appear in all its properties in every aspectual system as a multiplicity of functor syllogisms, and that, in particular, there must be an idea of the domain for every aspectual system. If there are two domains that are independent of each other, then there can always be predicate combinations of functors from both domains with respect to a particular subjective aspect that are themselves apodictic and therefore quantifiers. However, such quantifiers can

¹ Alternate translation: Obviously, the non-apodictic elements of these episyllogisms are functors of the apodictic elements, which represent any non-apodictic properties of the domain that depend on the specific S , and due to the aspect-related relativity resulting in relativity, they exhibit subjective aspects.

always be limited according to representation (4), because there can be no criterion for a predicate functor combination to appear as a universal quantifier that remains apodictic in every aspectual system, because there is the possibility that a non-apodictic functor describes a property of the domain that only occurs as a result of the particular aspectual system, and is therefore semantically conditioned by this system and no longer exists with respect to another system. The situation is completely different, however, when not only a functor is chosen, whose semantics already changes with the subjective aspect, but when the predicate combination refers to a whole functor system whose existence is maintained in all aspectual systems. A functor system, which obviously fulfills all these properties, is a category, because such a category is defined in relation to the particular aspectual system by an apodictic idea and a syllogistic series of concept functions, i.e., functors of the apodictic elements, of this idea in the sense of an episyllogism. Such a category characterizes a particular system of properties of the domain given by the functor with respect to a subjective aspect, whose idea is formed by the apodictic elements of the category. This category can be linked to another category by means of any predicate of the subjective aspect with respect to the same subjective aspect, in such a way that both categories have the same idea (homomorphic) or that the predicate combinations underlying the categories have different ideas (heteromorphic).

This homo- or heteromorphic predicate combination of categories is nothing other than a combination of functor systems that at least have the meaning of a homo- or heteromorphic mono-quantifier because the ideas of the category are apodictic. If such a connection is transferred to any other aspectual system, the functors of the categorical episyllogisms may change so that no quantifier-like relationships exist, but at least the ideas that are apodictic in the original aspectual system must remain connected, even if they do not have to be apodictic in the new aspectual system because properties of a domain that are apodictic in any aspectual system must obviously be regarded as manifest conceptual real properties. At the very least, only those ideas that are apodictic in the previous system would be connected in the new aspectual system, but if these properties are no longer apodictic in the new system, they must be functors of apodictic elements (related to the new aspectual system), which in turn allows for a prosyllogism and the evolution of new categories, the predicate combination of which also has quantifier character in the new aspectual system.

This procedure can be extended step by step by $n < \infty$ aspectual systems, so that by the complete induction inference on $n + 1$, it can also be concluded for $n + p$ with $p > 1$, which also allows the transition $p \rightarrow \infty$ according to this inference, thereby proving that the predicate combination of categories has a quantifier character in all aspectual systems, as long as the categories are related, meaning that their non-apodictic episyllogisms can be represented by the same subjective aspect. However, if a predicate combination of functor systems has a quantifier character in all aspectual systems, such a functor system satisfies the definition of a universal quantifier. Due to the apodictic character of the idea of a category and the interpretation of apodictic elements as manifest, conceptually real properties of a domain, there can only be a group of functor systems whose predicate combinations are universal quantifiers, namely categories, such that the existence condition of a universal quantifier is both necessary and sufficient for the predicate combination of categories to be valid. Universal quantifiers thus exist whenever combinations of categories are possible, i.e., the foundation of a syntrometry becomes possible when the concept of category is precisely defined in such a way that a concretely defined conceptual size, a so-called syntrix, is created, which on the one hand reflects the conceptual content of the category and on the other hand can be formally related to its own kind through predicate combinations that meet the necessary and sufficient existence conditions of universal quantifiers.

2.2 Definitions of Syntrix

The idea of a domain is always the starting point of an episyllogism, which together with this idea forms a category. In relation to the aspectual system A, the idea of the domain consists of $1 \leq i \leq n$ apodictic elements a_i that can be formally combined into a schema $(a_i)_n \equiv \tilde{a}$. Since this apodictic schema obviously carries the dimensions of the domain in relation to A, it is referred to as a *metrohor*. Therefore, each metrohor can be regarded as a formal idea of a domain. In the metalanguage of some subjective aspect S from A, a functor f can be represented without any problems, which sets each $m \leq n$ elements a_i of $\tilde{a}$ in correlation to a concept function that, in turn, describes a non-apodictic property of the domain with the idea $\tilde{a}$ related to S. If the correlation f acts on $\tilde{a}$ in the stage m , $\binom{n}{m}$ functors arise that form a convergence (syndrome) of concrete and related properties with respect to S. However, the correlation f can act again on these $\binom{n}{m}$ functors of the first syndrome in stage m , so that a second syndrome of functors arises, and so on. With increasing syndrome number, the degree of conditionality of all functors that occupy the respective

syndrome must therefore increase. In this way, when the syndromes of the functor assignment are traversed from $\tilde{a}$, an episyllogism can be determined such that the entire system consisting of the metrohor $\tilde{a}$, the correlation law f inducing the syndromes, and the effect stage m of f is the formal analogue of a category. The syndromes of the functor assignment supplying syndromes are referred to as a *syntrix* in the definition. The synkolator f , which supplies the syndromes of the functor assignment, is therefore briefly referred to as a *synkolator*, and accordingly m as a *synkolation stage*. The entire system consisting of z and the episyllogism of all syndromes fully occupied with functors thus corresponds entirely to a formally specified category and can be defined as a syntrix. The idea of a category therefore corresponds in the syntrix representation to the apodictic central schema, i.e., the metrohor, while the non-apodictic episyllogistics correspond to the sequence of syndromes fully occupied with functors, which, in turn, are given by synkolator and synkolation stage. If the symbol yz describes a syntrix that arises from some synkolator of $\tilde{a}$, $\tilde{a}$ is completely determined by the synkolator f , metrohor $\tilde{a}$, and synkolation stage $m \leq n$, which can be formally represented by $\tilde{a} \equiv \langle f, \tilde{a}, m \rangle$. Consequently, the definition of each syntrix is given by:

$$\tilde{a} \equiv \langle f, \tilde{a}, m \rangle \vee \tilde{a} \equiv (a_i)_n \vee F_1 \equiv F(a_k)_1^m \vee 1 \leq m \leq n \quad (5)$$

A Syntrix, determined by f and m given a present metrophor, obviously describes, within the underlying aspectual system, those properties of the area that are expressible in the metalanguage of the specific synkolator field. In this way, all syntrices of the infinite series that arise from a metrophor are distributed across the points of the metro field belonging to the aspectual system, i.e., the subjective aspects. This means that at least one syntrix formed from $\tilde{a}$ must belong to each subjective aspect. Since $\tilde{a}$ already includes all apodictic elements of the area, the synkolierten syndromes cannot contain apodictic elements; otherwise, the metrophor would not be complete, contrary to the assumption. Apparently, in this variety of syntrices, there is a distinction between different types of syntrices, since f can either act in such a way that none of the m elements appear more than once, or the elements can iterate in plurality. In the first case, the syntrix is heterometric, in the second homometric. Furthermore, it is possible that the order of the m elements in the synkolator is indifferent, or more generally, that $k \leq m$ elements change the synkolation result when permuted. In the first case, the synkolator is symmetric, in the second asymmetric. The four syntrix types mentioned differ in terms of the possible properties of the synkolator and are not fundamental properties of the synkolation process; that is, they do not allow for a fundamental classification of the syntrices. For each synkolator and for each associated stage $1 \leq m \leq n$, there is therefore a class of properties combined into a syntrix that are related in terms of a category and are functors in syllogistic order, so that the entirety of all syntrices contains all possible properties of an area, relative to a specific aspectual system, if the entire variety of syntrices arises from a metrophor that represents the idea of the area. Regardless of this representation of all properties of an area of the idea $\tilde{a}$, the four synkolator types (symmetric, asymmetric, hetero- and homometric) can induce these properties, but this does not yet provide a syntrix classification. However, there are two fundamentally different forms of synkolation, through which all syntrices initially fall into two syntrix classes that differ in their synkolation principle. Either the synkolation is discrete, which produces pyramidal syntrices in which the syndrome $\gamma + 1$ always arises alone from the syndrome γ , or the syntrix becomes homogeneous with continuous synkolation, i.e., if n_k is the sequence of the syndrome k ($k = 0$ applies to $\tilde{a}$), then the syndrome $k + 1$ arises from the number $\sum_{j=0}^k n_j$ of all functors and apodictic elements by the action of f of the degree m . In contrast to the *pyramidal syntrix* given by the expression (5), this homogeneous syntrix is symbolized by

$$\tilde{a} \equiv \langle (f, \tilde{a})m \rangle \quad (5a)$$

It appears that from each homogeneous syntrix of the form 5a, a pyramidal syntrix can be split off in such a way that a homogeneous fragment remains, and these two fundamentally different splitting products are classified according to the character of their synkolators as heterometral or homometral and symmetric or asymmetric. This divisibility of a homogeneous syntrix is immediately apparent from the recording of the fully occupied syndromes and shows that the syntrices can in turn become elements of functor operations. This fact opens up the prospect of a syntrometric functor-operational methodology through which syntrices can be brought into mutual functional dependencies.

Following this definition of syntrix, it seems appropriate to interpret the essence of a syntrix. Whenever statements are made about a conceptual domain, this domain is related to the metropie field² of an aspect system in such a way

²GPT 3.5 translated 'Metropiefeld' as metaplane

that the apodictic schema of the metaphorph lies as an idea about the domain. However, each point of the metropie field is a subjective aspect, in whose meta-language an infinitely simple sequence of synkolators can be defined, from which each element from the metaphorph can induce the syndromes of a syntrix. This means that for each point of the metropie field, i.e., for each subjective aspect, there is an entire bundle of syntrixes, with each syntrix containing a class of non-apodictic properties as functors in the category. This doubly infinite set of syntrixes, or the simply infinite set of syntrix bundles, would be homomorphic, because all these syntrixes have the same metaphorph that lies above the domain. The totality of all these syntrix bundles over the points of the metropie field thus encompasses all possible properties of the domain in the form of non-apodictic functor syndromes that can be expressed in the underlying aspect system.

A syntrix is obviously completely determined by the metaphorph, the synkolator, and the synkolation level, i.e., if an apodictic schema exists as a metaphorph with respect to any aspect system, then the necessary existence condition of a syntrix is fulfilled in some subjective aspect if at least one synkolator and one synkolation level³ can be defined with respect to this subjective aspect. The existence condition is only sufficient when the metaphorph is complete with respect to the conceptual domain being investigated, but this completeness can always be achieved because the conceptual domain in aesthetic-empirical prosyllogistics can always be limited in such a way that the metaphorph is complete, since the occupation of the metaphorph with apodictic elements is determined by the respective limitation of the underlying conceptual domain and can vary if this limitation experiences a variation. Thus, if apodictic elements can be detected at all in a conceptual domain with respect to an aspect system, then a metaphorph can always be defined, which is complete with respect to a delimited part. This partial area becomes larger the more apodictic elements occupy the metaphorph, i.e., an area with a complete metaphorph can be further expanded if empirical prosyllogisms are performed in more subjective aspects. Since the metaphorph can always be made complete by the correct delimitation of the area, the necessary and sufficient existence condition of a syntrix is always fulfilled if at least one apodictic element can be detected in a conceptual domain with respect to the underlying aspect system; because some synkolator can be defined in every subjective aspect. If n is the occupation number of a metaphorph, then the necessary and sufficient existence condition of a syntrix is

$$\tilde{a} \equiv (a_i)_n \vee n \geq 1 \quad (6)$$

A syntrix is obviously fully determined by the metaphorph, the synkolator, and the synkolation stage. That is, if an apodictic schema exists as a metaphorph with respect to any aspectual system, then the necessary condition for the existence of a syntrix is fulfilled in some subjective aspect if at least one synkolator and one synkolation stage can be defined with respect to this subjective aspect. The sufficient condition for the existence is only fulfilled when the metaphorph is complete with respect to the conceptual domain under investigation, but this completeness can always be achieved because the conceptual domain can always be limited in aesthetic-empirical prosyllogistics so that the metaphorph is complete. The occupancy of the metaphorph with apodictic elements is determined by the respective limitation of the underlying conceptual domain and can vary if this limitation undergoes a variation. Therefore, if apodictic elements can be detected in a conceptual domain with respect to an aspectual system, a metaphorph can always be defined that is complete with respect to a delimited part. This partial area becomes larger the more apodictic elements occupy the metaphorph, i.e., an area with a complete metaphorph can be expanded further if empirical prosyllogisms are performed in more subjective aspects. Since the metaphorph can always be made complete by the correct delimitation of the domain, the necessary and sufficient condition for the existence of a syntrix is always fulfilled if at least one apodictic element can be detected in a conceptual domain with respect to the underlying aspectual system, because some synkolator can always be defined in any subjective aspect. If n is the occupancy figure of a metaphorph, then the necessary and sufficient condition for the existence of a syntrix is thus

$$\tilde{a} \equiv (a_i)_n \vee n \geq 1 \quad (6)$$

If it turns out after delimiting a part with a complete metaphorph $\tilde{a}$ that there are still other parts that are also delimited and have complete metaphorphs, then the elements of all metaphorphs can always be combined into an overall schema if the delimited parts actually belong to the same connected conceptual domain.

³GPT 3.5 translated 'Synkolationsstufe' as 'synkolation stage'

After the preceding definition of the Syntrix concept, it becomes possible to generalize and deepen this concept through an extension. There is the possibility that in $\tilde{a}$, the apodictic a_i for $1 \leq i \leq n$ are not discrete individual properties, but limited continua consisting of infinitely many, but densely located properties. Since now every Metrophor element passes through an apodictic continuum, all syndrome assignments in this extended Syntrix concept consist of synchrotron continua. It is obvious that these band Syntrices have the most universal Metrophor occupation; for the Metrophor concept cannot be defined more generally. Formally, these band Syntrices must satisfy the same structural laws as the Syntrices with discrete Metrophor occupation, because in the synchrotron process, it is irrelevant whether discrete Metrophor elements or apodictic band continua synthesize. For all synchrotron stages $m \geq 1$, the syndrome assignments are given in any case, only in the case of the *band Syntrix* they form limited continua consisting of infinitely many individual synchrotronizations located everywhere. After this unique extension, it can be stated that every Syntrix, including the forms (5) and (5a), should be understood as band Syntrices, because in the discretely occupied Metrophor, the discrete elements of the occupation appear as degenerate apodictic band continua. In the form:

$$\tilde{a} \equiv (A_i, a_i, B_i)_n \quad (7)$$

In section 2.2, Syntrices are defined as formal analogies to categories. These consist of a Metrophor, which encompasses the apodictic elements of a domain, and Syndromes, which are given by the Synkolator and the Synkolation level. The definition of a Syntrix is formally represented by equation (5). There are various types of Syntrices, which differ by the properties of the Synkolator, and two basic forms of Synkolation: the pyramidal and the homogeneous Syntrix. The existence of a Syntrix is determined by the Metrophor, the Synkolator, and the Synkolation level. The necessary and sufficient condition of existence for a Syntrix is given in equation (6).⁴

2.3 Combinatorics of syndrome occupancies.

It is useful to develop a combinatorial theory within the framework of the mathematical-analytical subjective aspect of anthropomorphic formal logic, which allows making combinatorial statements about the occupation of syndromes with synkolation elements based on the possible properties of a syntrix. If n is the number of apodictic elements, i.e., the *metrophor diameter*, then there are possibilities $m \leq n$ and $m \geq n$ for the synkolation level m . For $m < n$, the syntrix can be heterometral, but homometral forms are also permissible, while for $m > n$ only⁵ homometral syntrices can be formed because otherwise the functors, i.e., the synkolation elements, are not complete. For $n = m$, on the other hand, in the heterometral case, the first syndrome is occupied by only one functor, which already leads to the closure of the syndrome in a pyramidal structure if $n > 1$. In the homometral case, however, as well as when a homogeneous syntrix is present, this does not occur for $m = n$. For the combinatorial investigation, a symmetric synkolator of level m is assumed. First, let us consider a heterometral pyramidal syntrix of the general form $m < n$. The full occupation of the syndromes must then be given by binomial coefficients, with the number of full occupations of a syndrome appearing as a combination number for class m of the synkolation level, which becomes quite possible due to $n > m$. Because of the metrophor diameter n (the metrophor would thus be the syndrome of digit 0), in the heterometral pyramidal syntrix, the first syndrome is fully occupied with $n_1 = \binom{n}{m}$, the second with $n_2 = \binom{n-1}{m}$, and the syndrome $\gamma + 1$ with $n_{\gamma+1} = \binom{n-\gamma}{m}$. This, however, provides a combinatorial recursion method that, if the synkolator is symmetric, yields the full syndrome occupation for any arbitrary syndrome digit of the heterometral pyramidal syntrix from the determining pieces n and m .

In the second case, a k -fold asymmetric synkolator would have to be assumed for the heterometric pyramidal syntrix, such that k of the m conceptual elements of a functor are bound to a fixed order in the synkolation law, which requires $k \leq m$. If such a synkolator exists, then each of the $\binom{n}{m}$ elements results in $\binom{m}{k}$ possibilities of asymmetry by exchanging the m functor arguments, each of which allows $k!$ permutations. This applies to the first syndrome, whose complete occupancy is therefore given by

$$n_1 = k! \binom{m}{k} \binom{n}{m} = k! \frac{m!}{k!(m-k)!} \cdot \frac{n!}{m!(n-m)!} = \frac{n!}{(m-k)!(n-m)!} = \binom{n}{m-k} \frac{(n-m+k)!}{(n-m)!}$$

⁴Summary generated by ChatGPT 4.0

⁵'nur' added by Hannes Schmid.

Because of its pyramidal character, this procedure can be recursively continued and provides for the full occupation of $\gamma + 1$ the recursion:

$$n_{\gamma+1} = \binom{n_{\gamma}}{m-k} \frac{(n_{\gamma} - m + k)!}{(n_{\gamma} - m)!}$$

If the Syntrix is no longer pyramidal but homogeneous, but also heterometrical, then for the occupation of the first syndrome, as in the pyramidal form, $n_1 = \binom{n}{m}$ applies, while already in the second syndrome the Synkolator acts not only on n_1 elements, but on $n + n_1$, which results in the full occupation $n_2 = \binom{n+n_1}{m}$. From this it becomes clear that $m = n$ is easily possible for heterometrical homogeneous Syntrizes because for this case, while $n_1 = 1$, already $n_2 = n + 1$. If the conclusion of complete induction is applied, the recursion $n_{\gamma+1} = \binom{N_{\gamma}}{m}$ with $N_{\gamma} = n + \sum_{j=1}^{\gamma} n_i$ follows for any syndrome $\gamma + 1$ of the heterometrical homogeneous Syntrix with a symmetrical Synkolator. If the Synkolator is k -times asymmetrical, then, by analogy with the pyramidal Syntrix, the following applies:

$$n_{\gamma+1} = \binom{N_{\gamma}}{m-k} \frac{(N_{\gamma} - m + k)!}{(N_{\gamma} - m)!}.^6$$

In complete analogy, the case of homometrical pyramidal and homogen forms with symmetric or asymmetric synkolator can be investigated, which, due to their homometrical properties, allow for $m \leq n$ as well as $m > n$. If the synkolator is symmetric and a pyramidal syntrix is given, the homometrical character can appear in $1 \leq j \leq L \leq m$ classes, each of which is occupied with $a_j > 1$ elements. If $a_j = 1$ for all, the homometrical case would have transitioned into the heterometrical case. The pyramidal combinations n_{γ} that form the syndrome $\gamma + 1$ must thus reduce the combination class m to $A = m - \sum_{j=1}^L a_j + L$. For the syndrome occupation of the homometrical (class L) pyramidal syntrix with symmetric synkolator, the recursion applies, while for $n_{\gamma+1} = \binom{n_{\gamma}}{A}$, the homometrical homogen syntrix results analogously to the corresponding heterometrical form, if a symmetry of the synkolator is also assumed. These homometrical investigations can be transferred to asymmetric synkolators according to the same principles as the heterometrical ones. In all cases of homometrality, it becomes clear that the heterometrical structures in terms of the combinatorics of their syndrome occupations are only special cases of the general homometrality, because for all $a_j = 1$, $\sum_{j=1}^L a_j = L$, which means that $A = m$, indicating a heterometrical character.

Let's consider the case of a heterometrical Pyramidal Syntrix with a symmetric Synkolator. The full occupation of syndromes must be given by binomial coefficients, where the number of fully occupied syndromes appears as a combination number of the class m of the Synkolationsstufe, which is possible because $n > m$. In the case of a heterometrical Pyramidal Syntrix, the first syndrome is filled with $n_1 = \binom{n}{m}$, the second with $n_2 = \binom{n_1}{m}$, and the $(\gamma + 1)$ -th syndrome is filled with $n_{\gamma+1} = \binom{n_{\gamma}}{m}$, where γ is the number of syndromes less one. This provides a combinatorial recursive procedure that, for any arbitrary syndromic figure of the heterometrical Pyramidal Syntrix and for a symmetric Synkolator, delivers the fully occupied syndromic figures from the determination pieces n and m .

In complete analogy, the case of homometrical Pyramidal and Homogeneous forms with symmetric or asymmetric Synkolator can be studied, which, due to the homometrical properties, also allow $m > n$ in addition to $m \leq n$. If the Synkolator is symmetric and a Pyramidal Syntrix is given, the homometrical character can appear in $1 \leq j \leq L \leq m$ classes, each of which is filled with $a_j > 1$ elements. If $a_j = 1$ for all j , then the homometrical case has passed into the heterometrical case. The pyramidal combinations n_{γ} , which form the syndrome $\gamma + 1$, must therefore reduce the combination class m to $A = m - \sum_{j=1}^L a_j + L$. For the syndrome assignment of the homometrical (class L) Pyramidal Syntrix with symmetric Synkolator, the recursion thus applies, while for $n_{\gamma+1} = \binom{n_{\gamma}}{A}$, the homometrical Homogeneous Syntrix, when a symmetry of the Synkolator is also assumed, results analogously to the corresponding Heterometrical form. These homometrical investigations can be transferred to asymmetric Synkolators according to the same principles as the heterometrical ones. In all cases of homometrality, it becomes clear that the heterometrical structures in terms of the combinatorics of their syndrome assignments are only special cases of the general homometrality. This is because for all $a_j = 1$, $\sum_{j=1}^L a_j = L$, which means that $A = m$, indicating heterometrality.

⁶Hannes Schmidt: Sum of all previous syndrome occupations!

2.4 Complex Synkolators, Synkolation Progression, and Syndrome Completion.

Whenever $m < n$ remains, i.e., the Synkolation stage is smaller than the Metrophor diameter, a rule f , independent of whether $\widetilde{a}$ or $\widetilde{a}$ is pyramidal or homogeneous, synkolates an infinite sequence of syndromes. In this infinite sequence of syndromes, for $m < n$, the complete occupation of the syndrome $\gamma + 1$ is always at least identical to that of γ , and an equisyndromatic Synkolation progression exists in an infinite sequence of syndromes if the full occupations of syndromes $\gamma + 1$ and γ are equal, while a monotonically divergent Synkolation progression is given when the full occupation of $\gamma + 1$ is higher than that of γ . The combinatorics of the syndrome occupations show that only pyramidal Syntrices can form an equisyndromatic Synkolation progression because, in the heterometral case, for $m = n - 1$, when the Synkolator is symmetric, for all syndromes, $n_\gamma = n = \text{const}$, while in the homometral case, the same is achieved for $L = 1$ and $a = m$, because then $\binom{n}{a} = \binom{n}{1} = n$, which also results in $n_\gamma = n$. However, this requires pyramidal structure and a symmetric Synkolator. Thus, in heterometral forms, if $m \leq n - 1$, the Synkolation progression is infinite and generally monotonically divergent. Only for $m = n - 1$ of pyramidal structure or for total homometrality of one of these structures, assuming a symmetric Synkolator, the Synkolation progression becomes equisyndromatic. Finally, if $m = n$, in a heterometral pyramidal Syntrix with a symmetric Synkolator, a syndrome completion is reached in the first syndrome because its full occupation is $n_1 = \binom{n}{n} = 1$, which does not allow further synkolation. In the corresponding homogeneous Syntrix, however, no syndrome completion is reached. Although $n_1 = 1$ here, unlike $m = n > 1$, for the second syndrome onwards, $n_2 = \binom{n+1}{n} = \frac{(n+1)!}{(n+1-n)!n!} = n + 1 > 1$, so that the first syndrome only has a minimal occupation, while from the second syndrome onwards, there is again a monotonically divergent Synkolation progression.

The heterometral Pyramidal Syntrix with $m = n$, even with a k -fold asymmetric Synkolator, achieves a Syn-dromabschluß (syndrome completion) in the first syndromes as long as the occupation of this syndrome, determined by $k \leq m$, remains lower than the Synkolationsstufe, because then the pyramidal synkolation of a second syndrome is no longer possible. If homometral properties are allowed, these generally reinforce the monotonic divergence and allow $m > n$ as well, but here an äquisyndromatischer Verlauf (equisyndromatic course) is also possible. For a given Synkolator, there is at most one completion in the first syndromes, or an äquisyndromatischer or monotonically divergent Synkolationsverlauf, but never a change of this course or a completion in a higher than the first syndromes. This behavior is due to the properties of all of these Syntrices, that the same Synkolator f acts in all syndromes. Therefore, such Syntrices are referred to as natural Syntrices due to the combinatorial laws of their Syndrombesetzungen and the number-theoretic combinatorial Synkolationsverlauf. In every subjective aspect of an aspectual system, there is at least one natural Syntrix, that is, an f, m as a Synkolationsgesetz, because if this were not the case, only the Metrophor of the area could be captured in the relevant subjective aspect, and this could also be expressed by a Synkolator that leaves the Metrophorelements unchanged and does not combine them.

In general, however, there can be a whole spectrum of such laws f_μ, m_μ for $1 \leq \mu \leq \chi$. If this is the case with respect to any subjective aspect, then it is possible to let different syncholation laws act in a single syntrix of this aspect, such that, for example, f_1 syncholates the syndromes $1 \leq j_{(1)} \leq \chi_{(1)}$, f_μ syncholates the syndromes $\chi_{(1)} + 1 \leq j_{(2)} \leq \chi_{(2)}$, or f_μ finally syncholates $\chi_{(\mu-1)} + 1 \leq j_{(\mu)} \leq \chi_{(\mu)}$ for $\mu = \chi$. In this way, a complex syncholation $\underline{f} \equiv \int_{\gamma=1}^{\chi} f_\gamma | \chi(\gamma + 1)^{\chi(\gamma)}$ can be introduced, composed of pS syncholation, with the complex syncholation steps $\underline{m} \equiv \int_{\gamma=1}^{\chi} m_\gamma | \chi_{(\gamma+1)}^{\chi(\gamma)}$. Such complex syncholation then yield, when acting on $\widetilde{a}$ in the $\underline{f}, \underline{m}$ sequence, complexly structured pyramidal or homogeneous syntrices with hetero- or homometric properties. The symbol notation

$$(\underline{f}, \underline{m}) \equiv \int_{\gamma=1}^{\chi} (f_\gamma, m_\gamma) | \chi_{(\gamma+1)}^{\chi(\gamma)} \widetilde{a} | \equiv \langle (\underline{f}, \widetilde{a}) \underline{m} \rangle \quad (8)$$

means that each of the syncholation laws f_γ, m_γ for $1 \leq \gamma \leq \chi$ syncholates the syndromes between $\chi_{(\gamma+1)}$ and $\chi_{(\gamma)}$, depending on the type of syntrix (pyramidal or homogenous).

Depending on the arrangement of the f_γ, m_γ in the complex synkolator $\underline{f}$, a different complex-synkolated syntrix is created with a different number-theoretical synkolation sequence corresponding to the new arrangement, and other syndromic occupations. If $\underline{f}$ has $1 \leq \gamma \leq \chi$ components, there are $\chi!$ possibilities within $\underline{f}$ for the synkolation sequence of the syntrix. The $\underline{m}_\gamma$ associated with the f_γ of $\underline{f}$ also significantly determine the synkolation sequence within a combined syntrix. In contrast to the natural syntrix with a monotonous synkolation sequence (equi-syndromatic or monotonically divergent), syntrix with complex synkolators should be referred to as combined, as they can have any

number-theoretical synkolation sequence. In such a synkolation sequence of a combined syntrix, monotonically divergent branches can always alternate with equi-syndromatic ones, and monotonic divergences are also possible in this way, i.e., there are as many possibilities for the synkolation sequence as for the sequence of any number-theoretical function of integer indices. This convergence of the synkolation sequence is made possible by the sequence and structure of the f_γ, m_γ in $\underline{f}, \underline{m}$ according to the definition (8) of the complex synkolator, and this convergence, in turn, allows for a syndromic conclusion in any given syndrom of the combined syntrix. If the last synkolator f_χ, m_χ of $\underline{f}$ is designed such that the full occupation is lower in the first synkolated syndrom than is allowed by the synkolation stage m_χ , the synkolation of a following syndrom is no longer possible, so that f_χ provides a syndromic conclusion in this way. If f_χ does not meet this requirement for the syndromic conclusion, f_χ synkolates an infinite sequence of further syndromes with an equi-syndromatic or monotonically divergent synkolation sequence analogous to the natural syntrix. Additionally, it should be mentioned that the components f_γ of a complex synkolator can also satisfy different synkolation principles depending on the indexing γ . For example, it is conceivable that f_γ in the sense of a pyramid syntrix, but $f_{\gamma+1}$ in the sense of a homogeneous form, is synkolated, etc. Hetero- and homometric as well as symmetric and asymmetric synkolator properties can also determine the coordinates f_γ in $\underline{f}$, allowing for a vast variety of combined syntrix with syndromic conclusions or an infinite synkolation sequence.

In contrast to natural syntrices, for which there are only infinite synkolation sequences if there is not already a closure in the first syndrome, in⁷ combined syntrices the syndrome closure can be achieved in any syndrome through the specific choice of $\underline{f}$, which also makes any arbitrary synkolation sequence realizable. The introduction of the combined Syntrix with the complex Synkolator next to the natural Syntrices obviously means a significant extension of the Syntrix concept, such that the relationship (8) as the most general form of the Syntrix essentially increases the diversity in the representation of Syntrix.

2.5 The primigenic aeondyne.

The most general form of a Syntrix representation is that of the combined band Syntrix. Another generalization of this concept arises when it is assumed that the course of the apodictic continua a_i of $\tilde{a}$ of the band Syntrix is determined by the course of certain conceptual parameters. For each apodictic element a_i of the Metrophor, if the greatest universality is to be maintained, $1 \leq j \leq n_i$ conceptual parameter dimensions $t_{(i)j}$ are given as a conceptual argument range, which can be illustrated by an n_i -dimensional argument space, if this space is not interpreted anthropomorphically as a point continuum, but generally as a conceptual tensor of the apodictic band a_i of $\tilde{a}$. If this interpretation of the $t(i)j$ as dimensions of a conceptual tensor of the band a_i is applied, then $a_i(t(i)j)1^{n_i}$ is determined by the parameters of the tensor in the form of a conceptual function, and here the $n_i \neq n_\gamma$ can be for $i \neq \gamma$, as well as the $t(i)j$ of a_i and the $t(\gamma)k$ of a_γ can be independent of each other. Therefore, each apodictic continuum $a_i(t_{(i)j})_1^{n_i}$ of $\tilde{a}$ is defined in a specific n_i -dimensional conceptual tensor.

The $t_{(i)j}$ generally symbolize concepts that undergo continuous changes within their definition intervals in the respective aspectual system, such that the a_i are apodictic concept functions with respect to the relevant aspectual system. For each course of $t(i)j$, conceptual boundaries $\alpha(i)j$ and $\beta(i)j$ can be given, such that the symbolization $\alpha(i)j, t(i)j, \beta(i)j$ means that $t(i)j$ can continuously represent any concept between the boundaries $\alpha(i)j$ and $\beta(i)j$. If for all apodictic metrophor elements $n_i \neq n_\gamma$, and all conceptual parameters $t_{(i)j}$ are independent of each other, then $\tilde{a}$ is a concept function that depends on $N = \sum i = 1^n N_i$ arguments, i.e., the metrophor is defined in an N -dimensional conceptual tensorium. Since, in general, for synkolation levels $m \geq 1$, the dimension of each synkolation must be $\mu \geq n_1$ for all i , while for the dimension of each syndrom, N always applies because in each syndrom, all a_i synkolate with each other, independent of the synkolation course and synkolation principle.

For $m > 1$, with increasing Syndromic index, μ approaches the dimensionality of N . The resulting Syntrix depends on N conceptual parameters in the intervals $\alpha(i)j, t(i)j, \beta(i)j$, and is called the "primordial Aeondyne" in this interval system (which is the "aeonic length"), because for every combination of the N conceptual parameters, mathematically expressed as every point in the N -dimensional manifold, there is a Syntrix that belongs, that is, a syncoherent system of properties in the subjective aspect that provides the Synkolation law. In every subjective aspect of the Aspective System expressed by $\tilde{a}$, there exists a primordial Aeondyne if there are N continuous conceptual parameters $t(i)j$ such that the $a_i(t(i)j)_1^{n_i}$ are apodictic conceptual functions throughout the Aspective System.

⁷Hannes Schmid: changed from 'im'

The pyramid or homogeneous primigenic Aeondyne S , symbolized by

$$(\underline{a}) \equiv \langle f, (\underline{a}), m \rangle \vee (\underline{a}) \equiv \langle (f, (\underline{a})), m \rangle \vee (\underline{a}) \equiv (a_i(t(i)_j)_1^{n_i})n \vee \alpha(i)_j, t(i)_j, \beta(i)_j \quad (9)$$

has an N -fold course in the limits $\alpha(i)_j, t(i)_j, \beta(i)_j$ with respect to the parameter $t(i)_j$ with $1 \leq j \leq n_i$ and $N = \sum i = 1^n N_i$. The Aeondyne S is called natural or combined, pyramidal, homogeneous, heterometrical, homometrical, symmetrical or asymmetrical if this is the case for the N -fold running Aeondyne. Furthermore, the Aeondyne S is L -step closed if a syndrome closure occurs in the syndrome L . For each natural Aeondyne, either $L = 1$ or $L \rightarrow \infty$ applies. So far, all investigated primigenic Aeondynes have been "metaphoric Aeondynes", since only $\underline{a}$ appeared to be defined in an N -dimensional tensor space. However, it is also conceivable that "synchulative Aeondynes" exist, in which although $\underline{a}$ does not belong to a band syntrix, the syncholorator depends on an n -dimensional conceptual tensor, which can be limited similarly to the metaphoric Aeondyne. The syncholation law, i.e., the structure of the general complex syncholorator, must then change as the n conceptual parameter intervals are traversed. If S generally designates any Aeondyne, then $\underline{S} \equiv \langle f, (\underline{a}), m \rangle$ symbolizes the metaphoric and $\overline{S} \equiv \langle (f), \underline{a}, m \rangle$ symbolizes the synchulative Aeondyne. The most general case of the primigenic Aeondyne would be one with both a metaphoric (N -fold) and synchulative (n -fold) course of the Aeondyne. This so-called "whole-running Aeondyne" is symbolized by $\underline{\overline{S}} \equiv \langle (f), (\underline{a}), m \rangle$. Thus, the Aeondyne is defined in a conceptual tensor space, whose dimension is composed of the dimensions of the metaphoric and synchulative tensor spaces according to $N + n$. The three possible forms of the primigenic Aeondyne are summarized in

$$\underline{S} \equiv \langle f, (\underline{a}), m \rangle \vee \overline{S} \equiv \langle (f), \underline{a}, m \rangle \vee \underline{\overline{S}} \equiv \langle (f), (\underline{a}), m \rangle \quad (9a)$$

which complements the representation equation (9).

The universal aeon dynamics are linked in the most general case, i.e., there is always a specific number of conceptual parameters that are common dimensions of the metaphorical and syncholic parameter tensorium. This number is referred to as the linking degree, which obviously cannot exceed the dimensionality of the lower-dimensional of the two tensors. If there is no linking degree, the syncholic aeon dynamics are determined by other concepts than the metaphorical ones, but the two runs coincide more strongly the higher the linking degree becomes. The linking becomes complete when the linking degree becomes identical with n and N , and at the same time $n = M$. If there is any syncholic aeon dynamics component at all, there will always be a continuous change in the syncholic run and the structure of a complex syncholorator when the conceptual parameters change in their definition intervals. This syncholic aeon dynamics run can thus be stationary or interrupted by discontinuities, i.e., syncholation jumps. Furthermore, the aeon dynamics run can be syndromatic, expansive, constant, or contractive, depending on whether the digit L of the L -fold syndromic completion increases, remains constant, or decreases as the aeon dynamics progress. Finally, the synchlications can all run uniquely, resulting in a monodromy of the aeon dynamics. Also, a certain number of K synchlications can become ambiguous from any point in the multi-dimensional manifold, meaning that from this point on (polydromy point), the aeon dynamics run becomes K -fold polydromic, i.e., a manifold of secondary runs (hence polydromy) consisting of K synchlications branches out ambiguously from the actual aeon dynamics.

The respective group of K synkolutions must have the same ambiguity, i.e. the same polydromy. The possibility of polydromy is completely independent of the specific type (9a) of the aeondyne and is determined only by the synkolorator and the functors that come to synkolation in the pre-synkolated syndrome's full occupancy at the respective polydromy point. Finally, a classification of each primitive aeondyne can be carried out based on the behavior of the aeonic lengths $\alpha(i)_j, t(i)_j, \beta(i)_j$. If the boundaries $\alpha(i)_j$ and $\beta(i)_j$ no longer belong to the aeondyne, it is open in these lengths. However, if only one boundary belongs to the aeondyne, it is semi-open. There are two possibilities for this, either it is closed in $\alpha(i)_j$ and open after $\beta(i)_j$, or vice versa, which is indicated by $(\alpha(i)_j, t(i)_j), \beta(i)_j$ or $\alpha(i)_j, (t(i)_j, \beta(i)_j)$. If both boundaries belong to the aeondyne, it is closed, so that $(\alpha(i)_j, t(i)_j, \beta(i)_j)$ indicates the aeonic length in which the aeondyne is closed. A semi-open aeondyne could also be referred to as a radial form or at least a partial radial form, with the term referring to the possibility that not all aeonic lengths are semi-open and the semi-open ones are not necessarily oriented in the same way.

2.6 The selection principle of poly-cyclic metaphore circles.

After the clarification of the Syntrix concept and its extension to the primigenic Aondyne, it appears possible to refine the concept formation of the universal quantifier. The quantor theory defines any predicate combination of functors as a polyquantor of degree b , if there are $1 \leq k \leq b$ aspect systems A_k , such that the predicate combination of functors remains invariant with respect to structural transformations of the metropie field within $1 \leq k \leq b < \infty$ during transitions of the A_k . Here, $b > 0$ is always an integer and provides, as long as $b < \infty$, polyquantors. If the functors are not individual properties, but entire categories in the form of syntrices with epi- or prosyllogistic orientation, then due to the syntrix properties, the polyquantor becomes the universal quantifier because $b \rightarrow \infty$.

In the following, a universal quantifier $b \rightarrow \infty$ shall be referred to as an unbounded universal quantifier, because due to the clarification of the syntrix concept, a universal quantifier can also exist if $b < \infty$, which should provide a conceptual refinement of the universal quantifier and an extension of quantifier theory. If there is a syntrix with respect to A_1 , i.e. if there is a system of apodictic elements in the form of a metaphore as the idea of a category represented by the syntrix with respect to A_1 , and this syntrix is then related to another aspectual system A_2 , then the metaphore in A_2 is generally no longer apodictic, i.e. its elements appear as non-apodictic functors or as synkolations of lower syndromes with respect to A_2 . However, as the prosyllogism progresses, these syndromes must lead to a metaphore that is apodictic with respect to A_2 because the apodictic elements with respect to A_1 cannot form an empty functor system after the aspect transformation, because the properties of a conceptual domain are always real if there is at least one aspectual system in which they are apodictic. Therefore, if there is a syntrix in A_1 , this category must also be a syntrix after an aspect transformation in A_2 . This process of aspect transformations can be extended to all aspectual systems by complete induction, from which it follows that if a category can be represented as a syntrix in one aspectual system, this representability is also possible in all other aspectual systems.

The concept of the general universal quantifier is based on this fundamental theorem of syntrometry, as this invariance must also apply to the predicate combinations of syntrices. However, in addition to the unlimited universal quantifier, a circular argument is possible that selects a finite number of possible aspect systems based on a selection principle from the infinite set of possible aspect systems, such that the number of debatable aspect systems for syntrices that enable this circular argument is limited by the selection principle. Such a circular argument is always possible if there is a syntrix and two aspect systems B_1 and B_2 such that the metrophor appears in both B_1 and B_2 , and only the syndromes, i.e., the syllogistic synkolation law, are changed during this aspect transformation. Under these conditions, there is always the possibility of transforming from B_1 to B_2 via a chain of $1 \leq k \leq N - 2$ aspect systems A_k , so that a metrophoric cycle is closed in accordance with the law of the transformation chain, which acts as a selection principle.

This metrophoric cycle is evidently monocyclic with base 2 and periphery N , since there are two aspect systems of the type B_1 and B_2 , and a transformation chain consisting of $N - 2$ systems A_k , which closes a ring of aspect transformations in which N aspect systems are contained. This concept of the metrophoric cycle is evidently capable of generalization. In general, there are $1 \leq i \leq Z$ aspect systems B_i such that the metrophor of a syntrix remains apodictic in all Z systems. These Z aspect systems in turn can form a poly-cyclic metrophoric cycle with base Z , and the cycle would be $\binom{Z}{2} = \frac{1}{2}Z(Z - 1)$ -fold, since the Z base elements can be connected to each other by $\binom{Z}{2}$ transformation arcs, with each of the $1 \leq j \leq \binom{Z}{2}$ monocyclic metrophoric sub-cycles having a periphery $N_j > 2$, where N_j must be integers. $N_j = 2$ does not yield a monocycle, as this case would simply be the aspect transformation from one base element to another.

If not only one syntrix with respect to the basis Z satisfies the conditions for the circular closure, but rather a universal quantifier, and if the law of the transformation chain determines the mono-cyclic partial cycles as well as their peripheral boundaries as a selection principle, then from the infinite set of possible aspectual systems, the finite number of discussable aspectual systems has been selected according to the selection principle, and the previously unlimited universal quantifier has become a poly-cyclic metaphoric cycle, i.e., a limited universal quantifier, which corresponds to a poly-quantifier of degree $\sum_{j=1}^L N_j$ with a cyclicity of $L = \binom{Z}{2}$. According to this selection principle of poly-cyclic metaphoric cycles, if a basis $Z \geq 2$ exists for an unlimited universal quantifier, poly-cycles with different peripheral boundaries of the partial cycles can be excluded as limited universal quantifiers, whereby the property of the universal quantifier is not lost, but the number of aspectual systems discussable for the respective problem is restricted to a finite number. Therefore, if a basis $Z \geq 2$ of a metaphoric cycle exists with respect to the

predicate combinations of syntrices (which can only be universal quantifiers), it always seems appropriate to develop a selection principle from the problem statement and to limit the universal quantifier through a metaphoric cycle.

3 Syntrix Corporators

3.1 The Corporator

According to the relation (5a), due to conceptual necessity, it is possible to split a general syntrix into elementary syntrices. From this statement, characterized by the relation (5a), the existence of operations that connect syntrices with each other must immediately follow if the inversion of the statement (5a) itself exists. By definition, each syntrix is a syllogistically oriented category of the kind that its splitting into elementary syntrices corresponds to a division of the category into subcategories and a division of the idea into partial ideas of the subcategories. Sub-terms can always be dialectically reintegrated into a super-term if the sub-terms were previously deduced from the super-term in the form of a division. The same relationships also exist when dealing with areas of conceptual elements, i.e., when the conceptual elements form categories with syllogistic orientations, i.e., syntrices. Therefore, it follows that the inversion of the statement (5a) and thus the concept of syntrix-connecting operations actually exist. However, the inversion of (5a) only refers to syntrix connections that reverse the splitting of syntrices into elementary syntrices.

In general, this splitting is always conceivable in a sequence of individual steps, so that any arbitrary syntrix system can be thought of as an intermediate result of a syntrix splitting into elementary splittings. If all syntrices of the system are split into their elementary syntrices, then a superior syntrix is always conceivable, which can be constructed from the elementary syntrices in an inversion of (5a). But if this is the case, then this superior syntrix must also arise from syntrix-connecting operations from the original syntrix system, conceived as an intermediate stage of the splitting. According to the reasoning of complete induction, there is still the possibility of generalization, so that the existence of general syntrix-connecting operations has been demonstrated in this way. Such operations will be referred to as syntrix corporations in the following. However, before a general theory of syntrix corporations is developed, the concept of the corporator in its most universal form must be specified and analyzed. If $\tilde{a}$ includes the n apodictic elements a_i of a field B , based on an aspect system A , then it is possible to divide B into subfields $b_{(j)}$ with $1 \leq j \leq \beta$, such that each of these subfields with respect to A includes a number of $p_{(j)} < n$ apodictic elements ($p_{(j)} < n$ for $\beta > 1$). It must always be $\sum_{j=1}^{\beta} p_{(j)} = n$, i.e. $\tilde{a}$ splits into β partial metaphors $\tilde{a}_{(j)}$.

This decomposition of $\tilde{a}$ is then perfect if A also breaks down into β individual aspectual systems such that each partial $\tilde{a}_{(j)}$ can be related to the corresponding $A_{(j)}$ and is above its partial field $A_{(j)}$. B includes all β partial fields $B_{(j)}$. After this metaphorical decomposition of $\tilde{a}$, the inverse process of a metaphorical composition is also conceivable. If there are β fields $B_{(j)}$ related to $A_{(j)}$, and a $\tilde{a}_{(j)}$ is above each of them, then all $\tilde{a}_{(j)}$ apparently compose to $\tilde{a}$ if the $A_{(j)}$ can be combined into a higher-level aspectual system A , such that all $B_{(j)}$ combined into B with respect to A have the apodictic elements a_i combined into $\tilde{a}$, and $\tilde{a}$ is complete with respect to B in relation to A . In addition to this metaphorical composition and decomposition, there is also metaphorical coupling and decoupling.

If $\tilde{a}$ and $\tilde{b}$ are two metrophones of the domains α and β respectively, relative to the aspect systems A and B , and if $\tilde{a} \equiv (a_i)_p$ and $\tilde{b} \equiv (b_k)_q$, then a sequence $1 \leq l \leq \lambda$ of connection rules (*conflator nodes*) ϕ_l can be specified, which connect $\lambda \leq p$ or $\lambda \leq q$ (depending on whether $p \leq q$ or $p \geq q$) elements from $\tilde{a}$ and $\tilde{b}$ in such a way that the λ connections $a_l, \phi_l, b_l \equiv c_l$ ⁸ are elements of the new metaphore $\tilde{c}$. Here, $\tilde{c}$ always contains only λ elements, but the connection can also be formulated in the more general form $a_i, \phi_l, b_k \equiv c_i, l, k$, where $\tilde{c}$ now contains a total of $p\lambda q$ elements. $\tilde{c}$ always appears relative to an aspect system C in which the λ conflict nodes ϕ_l are also expressible, such that the ϕ_l retain an operative sense in both A and B . In other words, this process of coupling depends on the conflict nodes ϕ_l (in terms of their structure), which must be such that $\tilde{c}$ is above the area γ coupled with the ϕ_l from α and β , which is relative to C , but it must be demanded that C can be constructed from A and B in such a way that $\tilde{c}$ and the ϕ_l meet the aforementioned requirements in terms of the aspect systems.

Similar to entangling confluent nodes, decoupling confluent nodes can also be defined. Metaphoric composition differs from metaphoric entanglement only in the way of connection. While the process of composition or decomposition takes place only in the domains and aspectual systems, entanglement and decoupling modulate the metrophor

⁸Hannes Schmid: Indices correct? In this example, b is the smaller metrophor, which is why the composed metrophor has the index L .

elements. All compositional and entangling rules are referred to as cooperative or entangling cooperators, in contrast to the compositional and entangling contra-operators of decomposition and decoupling. There may always be a composition rule given in addition to a cooperative or contraoperative entanglement or vice versa. If λ elements from $\tilde{a}$ and $\tilde{b}$ are coupled by λ confluent nodes ϕ_j , then the remaining $p + q - 2\lambda$ uncoupled metrophor elements of both metrophores may still be bound by a composition rule. In addition to the $p + q - 2\lambda$ compositional elements, the new metrophor created by coupling and composition also includes the λ coupling members, for a total of $p + q - \lambda$ elements, if the confluent nodes do not work combinatorially as $a_i, \phi_1, b_1 \equiv c_1$. The connection of metrophores, called metrophor corporation, is thus possible in three ways: compositionally, entangled, or mixed. If this metrophor corporation is an assertion γ of a subjective aspect S in the aspectual system C , which arises from composition and entanglement of A and B , and qc consists of the $p + q - \lambda$ entangled and composed elements of $\tilde{a}$ and $\tilde{b}$, then $\tilde{a} \left\{ K_m \quad C_m \right\} \tilde{b}, |\overline{CS}|_\gamma, \tilde{c}$ means that the two metrophores cooperate by coupling rules K_m and composition rules C_m in the constructed system C by co- and contraoperations, and this metrophor corporation is linked by the assertion γ of the subjective aspect S in C with a metrophor $\tilde{c}$. This corresponds to the predicative functor combination from functor and quantifier theory. $\left\{ K_m \quad C_m \right\}$ is the metaphoric corporator.

The coupling rule K_m is always at the bottom left, while the composition rule C_m is always at the bottom right. K_m is a schema of the conflict nodes and the rules that link elements from $\tilde{a}$ with those from $\tilde{b}$, while C_m only shows the selection of composition elements from $\tilde{a}$ and $\tilde{b}$. If one of the two rules is missing in the corporator, this means that the metaphore corporation is either purely coupled or purely composite. All metaphore corporations boil down to the fact that the apodictic elements of two areas are composed in such a way that a third higher-level area is created, which generally requires a simultaneous corporation of the aspect systems, namely whenever couplings are carried out, because the metaphore coupling elements do not necessarily need to be apodictic in the aspect systems of the areas (both areas can also be based on the same aspect system).

In every subjective aspect of such a system, at least one natural or generally complex synco-laws can be possible, which, based on this aspect, syncolates the syndromes of a syntrix from the metrophor. However, according to the quantifier theory, only predicate combinations of syntrices can lead to unrestricted or restricted universal quantifiers, and only these universal quantifiers can have a syntrometric meaning because the main clause of the universal quantifier can be considered as a syntrometric fundamental statement.

Predicate combinations of metaphors do not meet the existence condition of a universal quantifier, as a metaphor is only the idea of a category that can be specified as a syntrix. Only predicate combinations of syntrices are universal quantifiers, which implies that the concept of metaphor corporation needs to be expanded, as a syntrix is defined not only by the metaphor, but also by the synkolation law, i.e. by the synkulator and the synkolation level. Analogous to the metaphor corporation, synkolation laws can also be corporated if the two synkolation laws $\underline{f}, \underline{m}$ and $\underline{\phi}, \underline{\mu}$ (i.e. complex synkolators) hold in the same subjective aspect, or if there is another subjective aspect in the same aspect system in which the synkolative corporation product is allowed.

Also in the case of synkulator corporation, couplings K_s and compositions C_s must be distinguished. In both cases, cooperative and counter-cooperative conflict nodes and composition elements must be distinguished. In complete analogy to the metrophor corporation, $\left\{ K_s \quad C_s \right\}$ symbolizes a synkolative corporator, i.e.,

$$(\underline{f}, \underline{m}), \left\{ K_s \quad C_s \right\}, \underline{\phi}, \underline{\mu}, |\overline{AS}|_\gamma, (\underline{G}, \underline{N})$$

means that the members of the complex synkolators $\underline{f}$ and $\underline{\phi}$ with the synkolation levels $\underline{m}$ and $\underline{\mu}$ are corporated by the prescription of the synkolative conflict nodes $\overline{K}_s$ and the synkolative composition elements $\overline{C}_s$, and that the result of this corporation is linked by the predicate γ of the subjective aspect S in the underlying aspect system A with a complex synkulator $\underline{G}$ of the synkolation level $\underline{N}$. The members of the new synkolation level $\underline{N}$ (complex) are obtained from the corresponding members of $\underline{m}$ and $\underline{\mu}$ by the functional relation $\underline{N} = \Phi(\underline{m}, \underline{\mu})$ of mathematical analysis, where the functional relation Φ depends on the synkolative corporation law.

The metaphoric and synkolative corporation as a predicate linkage of functors is therefore described by

$$\widetilde{a} \left\{ \begin{matrix} K_m & C_m \end{matrix} \right\} \widetilde{b}, \overline{CS}|_{\gamma}, \widetilde{c} \vee (f, m), \left\{ \begin{matrix} K_s & C_s \end{matrix} \right\}, (\phi, \mu), \overline{AS}|_{\gamma}, (G, N) \quad (10)$$

However, the metaphoric and synkolation law clearly determine a syntrix, i.e. a combination of the metaphoric and synkolative corporations must enable a syntrix corporation. First, the metaphors of the domains must be corporated, and then the synkolation laws of the associated syntrices. If A and B are two aspect systems in which $\widetilde{a}$ and $\widetilde{b}$ apply to two domains, and C is a new aspect system that contains A and B in such a way that the new metaphor $\widetilde{c}$ arises from the corporation of $\widetilde{a}$ and $\widetilde{b}$ in C , and if there are two subjective aspects S_a and S_b in A and B with the combined syntrices (in general homogenous, homometrical, and asymmetric) $\langle (f\widetilde{a})m \rangle$ and $\langle (\phi\widetilde{b})\mu \rangle$, then in addition to $\left\{ \begin{matrix} K_m & C_m \end{matrix} \right\}$, $\left\{ \begin{matrix} K_s & C_s \end{matrix} \right\}$ can also act in such a way that in addition to the metaphoric corporation $\widetilde{c}$, a synkolator corporation leads to the new synkolation law (G, N) , and this in a new subjective aspect S_c in C , which is superior to both S_a and S_b and contains both (similarly to C being superior to systems A and B and encompassing both). S_c does not necessarily have to be included in A or B but must be in C .

For the metaphoric and synkolative corporation, if the results of these corporations are linked by the predicate γ from S_c in C with $\widetilde{c}$ or (G, N) , then we have $\widetilde{a} \left\{ \begin{matrix} K_m & C_m \end{matrix} \right\} \widetilde{b}, \overline{CS_c}|_{\gamma}, \widetilde{c}$ and $(f, m) \left\{ \begin{matrix} K_s & C_s \end{matrix} \right\} (\phi, \overline{CS_c}|_{\gamma}, (G, N))$. On the other hand, these metaphors and synkolation laws form the general syntrices $\langle (f\widetilde{a})m \rangle$ and $\langle (\phi\widetilde{b})\mu \rangle$ and $\langle G\widetilde{c}N \rangle$. Moreover, the two corporations $\left\{ \begin{matrix} K_m & C_m \end{matrix} \right\}$ and $\left\{ \begin{matrix} K_s & C_s \end{matrix} \right\}$ can be combined according to $\left\{ \begin{matrix} K_m & C_m \end{matrix} \right\}, \left\{ \begin{matrix} K_s & C_s \end{matrix} \right\} = \left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\}$, and this combined corporator must be interpreted as a syntrix corporator.

The lower row of this corporator gives the metaphorical and the upper one the synkolative corporator, such that both, carried out one after the other, corporates all determinative elements of two syntrices with each other, so that the complete determinative elements of a new syntrix result as the outcome of syntrix corporations, which can be linked to a syntrix by the predicate γ from S_c in C . This fact is formally expressed by the relationship

$$\langle (f\widetilde{a})m \rangle \left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\} \langle (\phi\widetilde{b})\mu \rangle, \overline{CS}|_{\gamma}, \langle (G\widetilde{c})N \rangle \quad (11)$$

Obviously, the syntrix corporation establishes the functorial connection between two syntrices, which is again a syntrix that can be linked to another syntrix by some predicate, thus forming a quantifier that, because it is the predicate linkage of syntrices, satisfies the criterion of the universal quantifier, so that all syntrix corporations of the form (11) can form universal quantifiers.

3.2 Total and partial Syntrix corporation

First, total corporations must be considered, and from these, total compositions. If

$$\langle (f\widetilde{a})m \rangle \left\{ \begin{matrix} C_s \\ C_m \end{matrix} \right\} \langle (\phi\widetilde{b})\mu \rangle, \overline{||}, \langle (g\widetilde{c})N \rangle,$$

where $\overline{CS_c}|_{\gamma} \equiv \overline{||}$ is used for abbreviation, this means that the apodictic elements selected by the metaphoric composition law C_m from $\widetilde{a}$ and $\widetilde{b}$ are combined to form the new $\widetilde{c}$, and that further, the elements of the complex syncholors f and μ selected by the syncholic composition law C_s according to S are used to form the composed complex syncholor g , which acts on $\widetilde{c}$ in the sequence of stages N of the syncholorations. If $\widetilde{a}$ has diameter p and $\widetilde{b}$ has diameter q , and a total of $p_{\gamma} \leq p$ elements are composed from $\widetilde{a}$ by C_m , and a total of $q_{\gamma} \leq q$ elements are

composed from $\widetilde{b}$, then $\widetilde{c}$ has diameter $p_\gamma + q_\gamma \leq p + q$. Only for $p_\gamma = p$ and $q_\gamma = q$ is $p_\gamma + q_\gamma = p + q$, i.e., the composition C_m can be performed in two steps, namely a total *cooperation* to a metaphore of diameter $p + q$, and a subsequent *contraoperation* which decomposes $p + q - p_\gamma - q_\gamma$ elements, yielding $\widetilde{c}$ of diameter $p_\gamma + q_\gamma$.

In a very similar way, C_s of the synkolator can be carried out in multiple stages. If the composite corporation is purely synkolative, the process becomes ambiguous if $\widetilde{a}$ and $\widetilde{b}$ are not identical because the composition $\underline{g}$ can act on both $\widetilde{a}$ and $\widetilde{b}$. The same applies to purely metaphoric composition C_m , as both $\underline{f}$ and $\underline{\phi}$ can synkolize the composed $\widetilde{c}$. Unique corporations exist in these two cases only for $\widetilde{a} \equiv \widetilde{b}$ with $p = q$ or for $(\underline{f}, \underline{m}) \equiv (\underline{\phi}, \underline{mu})$ in the forms

$$\langle (\underline{f}\widetilde{a})\underline{m} \rangle \left\{ \begin{matrix} C_s \\ C_m \end{matrix} \right\} \langle (\underline{\phi}\widetilde{a})\underline{\mu} \rangle, \bar{\parallel}, \langle (\underline{g}\widetilde{a})\underline{N} \rangle$$

or

$$\langle (\underline{f}\widetilde{a})\underline{m} \rangle \left\{ \begin{matrix} C_s \\ C_m \end{matrix} \right\} \langle (\underline{f}\widetilde{b})\underline{m} \rangle, \bar{\parallel}, \langle (\underline{f}\widetilde{c})\underline{m} \rangle.$$

Similarly, the situation with total couplings is analogous. Here, both the synkolators and the metaphors are coupled by synkolative or metaphoric coupling rules K of the confictor nodes (analogous to the composition rules C). For

$$\langle (\underline{f}\widetilde{a})\underline{m} \rangle \left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\} \langle (\underline{\phi}\widetilde{b})\underline{\mu} \rangle, \bar{\parallel}, \langle (\underline{g}\widetilde{c})\underline{N} \rangle$$

or for purely synkolative or purely metaphoric coupling, analogous statements apply to total compositions. Thus, for $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}$ or $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}$, the coupling is again ambiguous if the metaphor or the synkolation laws of the two syntrices being coupled do not become identical. If $\widetilde{a}$ and $\widetilde{b}$ have diameters p and q , then the diameter of $\widetilde{c}$ is λ in the case of $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}$ if the coupling rule K_m consists of λ confictor nodes. On the other hand, if C_m still exists, then this metaphoric composition rule leads to the composition of $p + q - 2\lambda$ elements, so that the pure metaphor coupling would initially have to be carried out in the form of a coupling and composition, followed by a decomposition of the $p + q - 2\lambda$ elements, so that only the λ coupling sizes remain in $\widetilde{c}$. The prerequisite for this corporational process indicated by $\left\{ \begin{matrix} C_m & K_m \\ C_m & K_m \end{matrix} \right\}, \left\{ \begin{matrix} C_m \\ C_m \end{matrix} \right\} \equiv \left\{ \begin{matrix} K_m \\ K_m \end{matrix} \right\}$ with the composite contraposition operators $\underline{C}_m$ is $(\underline{f}, \underline{m}) \equiv (\underline{\phi}, \underline{\mu})$, while for $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}, \widetilde{a} \equiv \widetilde{b}$ with $p = q$ must always be required, because otherwise the corporational problems are no longer unique. In the case of $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}$, it does not appear to be necessary to go through the conceptual detour of an additional composition followed by decomposition.

In particular, it is always possible to break down the course of a total corporation into a chain of individual corporations by the mutual application of co- and contra-operations in the corporator. However, the members of these chains may no longer be total corporators, unlike the case of $\left\{ \begin{matrix} K_m \\ K_m \end{matrix} \right\}$, which is built exclusively from a single type of corporation rule, either K or C . Corporators built from a mix of K and C have a different universal character, and are referred to as partial corporators, due to their property of partially coupling or composing elements synkolatively or metaphorically. For syntrix corporators, pure synkolator or pure metaphor corporation cases are possible with $\left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\}$ and $\left\{ \begin{matrix} K_m & C_m \\ K_m & C_m \end{matrix} \right\}$, respectively, and they are always special cases of the general mixed corporation $\left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\}$, in the form of a partial corporation.

Other special cases of the universal corporator $\left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\}$ are those of purely composite total corporations $\left\{ \begin{matrix} C_s \\ C_m \end{matrix} \right\}, \left\{ \begin{matrix} C_s \\ C_m \end{matrix} \right\}$ or $\left\{ \begin{matrix} C_m \\ C_m \end{matrix} \right\}$ or purely coupling total corporations $\left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}, \left\{ \begin{matrix} K_s \\ K_m \end{matrix} \right\}$ or $\left\{ \begin{matrix} K_m \\ K_m \end{matrix} \right\}$. These total corpo-

rations have been examined in the preceding section, but this entire group of corporations is only a special case of the universal corporator $\left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$, which connects the syntrices according to the relationship (10).

Another somewhat more universal special case of this corporator is the group of partial corporators, which always contain less than four corporating rules but consist of coupling and composing parts. For these partial corporators, the same fundamental laws must apply as for the total ones. However, during metaphorical partial processes, the metaphorical diameters change according to the law $p + q - \lambda$, which was derived above. A general overview of the types of corporators is obtained by defining the concept of the *corporator class*.

If one defines the corporator class as the number of rules contained in the corporator, which cannot exceed the value of 4, then the number of possible combinations for a class is equal to the number of the four combinations for the relevant corporator class. That is, there can only be one universal corporator because the universal corporator has class 4 and $\binom{4}{4} = 1$. For class 3: $\binom{4}{3} = 4^9$, on the other hand, there are $\binom{4}{2} = 6$, of which 4 must be partial and 2 must be total, while there are $\binom{4}{1} = 4$ total for the first class. In addition to the six possible total corporators, there are still eight possible partial ones, and all these fourteen corporators are special cases of the fourth class.

The corporations of the first class cannot be unambiguous unless there are identities of the metro-phore or synkolation laws with respect to the corporating syntrices. In the second class, both the total and the partial ones with synkolation and metaphorical corporating rules are unambiguous, while for the forms $\left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$ and $\left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$, ambiguity occurs again if there are no identities. On the other hand, in the third class, only unambiguous corporators are present. The investigations regarding unambiguity and ambiguity are completely identical to those carried out with the total corporators and show that a corporator of any class between 1 and 4 is always unambiguous if at least one rule is given concerning the synkolation and metaphorical corporation.

If the syntrices $\widetilde{a}|$ and $\widetilde{b}|$ are in the aspect systems A and B , but the corporation $\widetilde{c}|$, the conflict nodes of the corporator, and the predicate are given in system C , then there is the following possibility for $\widetilde{c}|$: If $\widetilde{a}|$ and $\widetilde{b}|$ and the corporator are such that the syndromes of $\widetilde{c}|$ remain empty, either all syndromes of the digit sequence $\gamma \geq 0$, including $\widetilde{c}$, are empty, or $\widetilde{c}$, i.e., $\gamma = 0$, is fully occupied and the syndromes of the digit sequence $\gamma \geq 1$ are empty. For the first case, i.e., an empty metaphore $\widetilde{c}$, there are again two possibilities: either the corporated metaphore $\widetilde{c}$ with respect to C is really a metaphore, and then $\widetilde{c}|$ cannot exist at all because an empty metaphore would mean that an idea is missing from a category, making that category undefined. This also applies if the syndromes $\gamma \geq 1$ are occupied despite the empty $\widetilde{c}$, since this occupation can only be brought about by synkulator members. The other possibility is that $\widetilde{c}$, without occupation, is only a pseudometaphore that does not contain any apodictic elements with respect to C and therefore only has the properties of an empty syndrome $\gamma > 0$ in $\widetilde{c}|$. The real apodictic metaphore $\gamma = 0$ must then, if $\widetilde{c}|$ exists at all, lie with respect to the prosyllogism before the pseudometaphore and be fully occupied. However, this has reduced the first of the two cases to the second, so that only syntrix corporations with the empty syndromes $\gamma \geq 1$ can exist.

If $\widetilde{c}$ does not exist in $\widetilde{a}|\widetilde{b}|, ||, \widetilde{c}|$, that is, $\gamma = 0$ is empty, but $\widetilde{a}|$ and $\widetilde{b}|$ exist in A and B , then the corporator $\{\} \equiv \left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$ (shortened as $|| \equiv \overline{CS_c}|_\gamma$) can exist in C . On the other hand, if $\widetilde{c}|$ exists in C , but all syndromes $\gamma \geq 1$ are empty, then $\widetilde{c}|$, due to the empty syndromes, forms the so-called *null syntrix* $\widetilde{c}| \equiv \langle \overline{f\widetilde{c}m} \rangle$, where the overbars indicate that the complex synkolation law is such that all syndromes $\gamma \geq 1$ in $\widetilde{c}|$ remain empty. The essence of the null syntrix is thus described by

$$\widetilde{a}|\widetilde{b}|, ||, \widetilde{c}| \vee \widetilde{c}| \equiv \langle \overline{f\widetilde{c}m} \rangle \quad (11a)$$

These null syntrices evidently allow for a greatly simplified representation of syntrix corporations.

With the concept of the metaphorical circle, a further theorem can be developed concerning the general syntrix corporation of form (10). Through the universal corporator, three syntrices from generally different aspect systems are connected by a predicate in a syntrometric relationship, which, due to the syntrix character, must have the properties of a universal quantifier. However, if the three syntrices form the basis of a tricyclic metaphorical circle, then the same relationship applies to the peripheries of all $\binom{3}{2} = 3$ cycles because the existence of the tricycle limited the universal quantifier $\widetilde{a}|\widetilde{b}|, ||, \widetilde{c}|$.

⁹Edit by Hannes Schmid

3.3 Pyramidal Elementary Structures.

The synkolation law should be most general when the complex synkolator acts with multiple asymmetric homometry and also synkolates the syndromes of a homogeneous syntrix. Such homogenous syntrices must therefore be the most universal syntrix forms, so it remains to be investigated whether and into which components such a combined homogeneous syntrix with a multiple asymmetric homometrical synkolator can be split by applying contra-operative corporators, because these last split products must be syntrometric elementary structures from which any syntrix can be built through corporation processes. Apparently, these corporators are purely synkolative $\left\{ \begin{matrix} K_s & C_s \end{matrix} \right\}$, because the synkolation law determines the course of synkolation of the syndromes. If $\langle (f\widetilde{a})\underline{m} \rangle$ is such a homogenous syntrix, there is always the possibility of understanding $\underline{f}$ as a result of the composition of a pyramidal and a homogeneously acting synkolator $\underline{P}$ and $\underline{H}$ with the synkolation stages m_P and m_H .

This means that, if D_s is the inverse contra-operative composition law corresponding to C_s , with the help of the homogeneous fragment $\langle (H\widetilde{a})\underline{m}_H \rangle$ from $\langle (f\widetilde{a})\underline{m} \rangle$, according to

$$\langle (f\widetilde{a})\underline{m} \rangle \left\{ \begin{matrix} D_s \end{matrix} \right\} \langle (H\widetilde{a})\underline{m}_H \rangle, \overline{\parallel}, \langle \underline{P}\widetilde{a}, \underline{m}_P \rangle$$

a combined pyramidal syntrix can be separated. Similarly, a pyramidal syntrix can be separated from the homogeneous fragment again, as the homogeneous fragment is also a separable homogenous syntrix. The same procedure can then be applied to the remaining homogeneous part, and this procedure can be continued further according to the method of complete induction until a null syntrix with the metrophor $\widetilde{a}$ remains, which can always be considered as a pyramidal syntrix with empty syndromes.¹⁰ Since these null syntrices uniquely exist according to (10a), and the entire process is invertible because each contra-operator has a corresponding co-operator (both terms are relative and therefore interchangeable). The homogeneous fragment is constructed by a sequence of syncollative corporations of pyramidal syntrices,

$$\langle (H\widetilde{a})\underline{m}_H \rangle \equiv \langle p_1 \widetilde{a} \underline{m}_{p1} \rangle \{ \} \dots \langle p_k \widetilde{a} \underline{m}_{pk} \rangle \{ \} \dots \{ \} \widetilde{a} |$$

which directly implies that due to

$$\langle (f\widetilde{a})\underline{m} \rangle \left\{ \begin{matrix} D_s \end{matrix} \right\} \langle (H\widetilde{a})\underline{m}_H \rangle, \overline{\parallel}, \langle \underline{P}\widetilde{a} \underline{m}_P \rangle$$

and therefore

$$\langle (f\widetilde{a})\underline{m} \rangle, \overline{\parallel}, \langle \underline{P}\widetilde{a} \underline{m}_P \rangle \left\{ \begin{matrix} C_s \end{matrix} \right\} \langle (H\widetilde{a})\underline{m}_H \rangle$$

every homogeneous syntrix $\langle (f\widetilde{a})\underline{m} \rangle$ can be decomposed into an entire sequence of pyramidal syntrix corporations, which implies that pyramidal syntrices must indeed be elementary syntrometric structures, as any higher homogeneous syntrix can be constructed from them with the help of suitable corporators.

This splitting law is universal, as no demands restricting the general validity were placed on $\langle (f\widetilde{a})\underline{m} \rangle$. The elementary character of the pyramidal syntrix is thus represented with the help of the null syntrix from (11a) by the relation

$$\langle (f\widetilde{a})\underline{m} \rangle, \overline{\parallel}, \widetilde{a} | \{ \} \dots \{ \} \widetilde{a} | \{ \} \dots \{ \} \widetilde{a} | \quad (11b)$$

In this decomposition of $\langle (H\widetilde{a})\underline{m}_H \rangle$ into pyramidal syntrices (which led to (11b)), the last pyramidal syntrix before $\widetilde{a} |$ must additionally contain the first syndrome of $\langle (f\widetilde{a})\underline{m} \rangle$ in each syndrome, which determined the homogeneous character of this starting syntrix. However, since this syndrome is contained in all syndromes, it can be reduced to this, that is, the pyramidal syntrices occurring in the sequence always begin with the second syndrome in full occupancy, while the first remains empty. This reduction is to be understood in the following way: In all syndromes $\gamma \geq 2$,

¹⁰Sentence break by Hannes Schmid

this first syndrome is contained, so that through a dehomogenizing decomposition or through a contra-operative decoupling, the synkollator can be eliminated from the syndromes $\gamma > 1$ and with it, $\gamma = 1$ can be fully occupied. In this way, every homogeneous syntrix can thus be dissolved into pyramidal syntrices by contra-operative corporators. The reversal of this theorem, i.e., the construction of arbitrary homogeneous syntrices from pyramidal syntrices with the help of cooperative corporators, is expressed by the relation (11b), the general validity and uniqueness of which has been proven. No general theory can be developed for the combinatorics of the syndrome occupancies of these pyramidal forms building a homogeneous syntrix, as this combinatorics depends on the mode of action of the syncollative corporators. If there are $1 \leq j \leq p$ such pyramidal syntrices that build up a homogeneous syntrix according to (11), and the respective syndrome $\gamma > 1$ ¹¹ of the pyramidal forms is fully occupied with n_j , but the homogeneous form with n_H , and if μ is the full occupancy of the first syndrome of the homogeneous form, then the connection $n_H - \mu = \sum_{j=1}^p n_j$ obviously applies if the cooperative corporators act purely compositively in such a way that the pyramidal syndrome occupancies are added together to form the homogeneous form.

In any case, the universally valid relation (11b) shows that the pyramidal forms are syntrometric elementary structures. In the following, it is therefore always sufficient to consider combined pyramidal syntrices $\tilde{a}| \equiv ((f\tilde{a})\underline{m})$ as syntrometric elements without restricting the generality. A general pyramidal syntrix, in turn, is constructed with respect to the complex syncollation law in such a way that, in general, the occupancy of any of its syndromes consists of homometral and heterometral syncollations with symmetric and asymmetric properties. In complete analogy to the development of the relation (11b), the general pyramidal syntrix $\tilde{a}|$ can now be decomposed into so-called *pyramidal elementary structures* by contra-operatively acting syncollative corporators. Such a pyramidal elementary structure is distinguished by the property of the syndrome occupancies containing only syncollations of a basic form. According to the syncollator theory, which represents a refinement of the functor theory, there can only be four syncollative basic forms, namely homometral symmetric, homometral asymmetric, heterometral symmetric, and heterometral asymmetric, from which it immediately follows that there can also only be four pyramidal elementary structures. However, in complete analogy to the resolution of any homogeneous syntrix into pyramidal syntrices, every pyramidal syntrix can be dissolved into these four elementary structures, which is possible with the help of contra-operative syncollation corporators of the first and second class.

The proof and the demonstration of uniqueness run completely parallel to the development leading to relationship (11b), which, however, means that the decomposition process must be reversible. Thus, any arbitrary pyramidal syntrix can be constructed from the four pyramidal elementary structures $\tilde{a}|_{(k)}$ with $1 \leq k \leq 4$ using syncollator corporators because an inverse cooperative corporator exists for each contra-operative corporator. Relationship (11b) would therefore be supplemented by

$$\tilde{a}|, \overline{\tilde{a}}|, \tilde{a}|(1)\{\} \tilde{a}|(2)\{\} \tilde{a}|(3)\{\} \tilde{a}|(4)\{\} \quad (11c)$$

and this connection, together with (11b), characterizes a theorem that identifies the pyramidal elementary structures as the actual syntrometric elements.

Every pyramidal syntrix of relationship (11b) can, due to (11c), be represented as a corporation of elementary structures, which leads to the theorem that any arbitrary syntrix can be constructed from the four pyramidal elementary structures $\tilde{a}|_{(k)}$ with $1 \leq k \leq 4$ through syncollator corporators. The decision whether a homogeneous or pyramidal syntrix is formed is determined by the specific corporate laws. The general forms of the elementary structures are combined pyramidal syntrices, which include the natural forms. In addition, an elementary structure is always (in every subjective aspect of each aspectual system) manifested by a multiply infinite set of pyramidal syntrices, as all possible specific elementary syncollator and metaphor forms in the relevant subjective aspect must be expressed in the corresponding elementary structure.

3.4 Concenters and Eccenters.

In the preceding sections, syntrix corporations with metaphorical corporate elements have been carried out. This means that for two syntrices, the metaphor is corporated, and the syncollation of the syndromes of such a corporated syntrix starts again from the new metaphor, that is, the syndrome occupations are concentric around the metaphor of the

¹¹ Asus R700 changed n to gamma

starting syntrices, and the corporation takes place in such a way that the syndromes are also concentrically syncolated around the metaphor of the corporated syntrix. All concentrically acting corporators $\left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$ are therefore referred to as *Concenters*, with the uniqueness of the corporator problem being assumed. It is now conceivable that a corporation of $\widetilde{a}|$ with $\widetilde{b}|$ leads to a $\widetilde{c}|$ in a different way if there is a metaphorical corporation component, as each syndrome of a syntrix could be regarded as a pseudometaphor. If γ is the current syndrome number as an argument of the syncolation process, then $\gamma \geq 1$ characterizes the genuine syncolated syndromes, so that the metaphor with $\gamma = 0$ can be considered as a 0-syndrome. $\gamma = 0$ is distinguished from $\gamma > 0$ in that its occupations consist of apodictic elements. If the requirement of apodicticity is dropped, then any syndrome $\gamma > 0$ can assume the function of a pseudometaphor, i.e., a corporator with a metaphorical component can always be constructed, which pseudometaphorically corporates the syndrome $k \geq 0$ of $\widetilde{a}|$ with $l \geq 0$ of $\widetilde{b}|$ (in general $k \neq l$).

The associated synkolative corporation must then also begin at the syndromes k and l from $\widetilde{a}|$ and $\widetilde{b}|$, because otherwise there would be no uniqueness in the synkolation processes $0 \leq \gamma \leq k$ from $\widetilde{a}|$ or $0 \leq \gamma \leq l$ from $\widetilde{b}|$ due to $\widetilde{a} \neq \widetilde{b}$. A metaphorical corporation share must always be present in this eccentric pseudometaphorical corporation, with which a corresponding synkolative corporation runs parallel if the synkolation laws in $\widetilde{a}|$ and $\widetilde{b}|$ do not coincide from the syndromes k and l onwards, because otherwise there can be no uniqueness of this eccentric corporation.

$\left\{ \begin{smallmatrix} K_s & C_s \\ K_{kl} & C_{kl} \end{smallmatrix} \right\}^{(k)} \left\{ \begin{smallmatrix} K_s & C_s \\ K_{kl} & C_{kl} \end{smallmatrix} \right\}^{(l)}$ symbolizes such a pseudometaphorical corporation, which is referred to as eccentric because in $\widetilde{a}| \left\{ \begin{smallmatrix} (k) \\ (l) \end{smallmatrix} \right\} \widetilde{b}|, ||, \widetilde{c}|$ the corporated syntrix $\widetilde{c}|$ initially forms two separate branches for the syndromes $0 \leq \gamma, j \leq k, l$ from $\widetilde{a}|$ and $\widetilde{b}|$ that are not corporated, while it only comes to the common synkolation in the *conflexion field* formed from the syndromes k and l . In the general case, $k \neq l$ and the eccentric corporation is regularly eccentric, while it becomes equilon longitudinal (refers to the argument values $k = l$ of the respective synkolation process) for $k = l$. In the special case $k = l = 0$, the equilon longitudinal eccentric corporation changes into the concentric one, which implies that this concentric corporation $k = l = 0$ is a special case of the equilon longitudinal $k = l > 0$, while $k = l$ is in turn a specialization of the regularly eccentric $k \neq l$, which implies that the regularly eccentric corporation has the greatest possible generality. All eccentrically acting corporators in this manner

$$\widetilde{a}| \left\{ \begin{smallmatrix} (k) \\ (l) \end{smallmatrix} \right\} \widetilde{b}|, ||, \widetilde{c}| \quad (12)$$

are referred to as *eccentrics*, namely $k \neq l$ as *regular*, but $k = l > 0$ as *equilon longitudinal eccentrics*, which change into the concentric ones for $k = l = 0$. The eccentrically corporated $\widetilde{c}|$ is referred to as being biconflective because the two non-corporated synkolation processes $0 \leq \gamma, j \leq k, l$ only converge in the conflexion field. Due to the generality of the regular eccentric, the corporators used in the following should always be understood as regular eccentrics because

the other types of corporators are special cases of $\left\{ \begin{smallmatrix} K_s & C_s \\ K_{kl} & C_{kl} \end{smallmatrix} \right\}^{(k)} \left\{ \begin{smallmatrix} K_s & C_s \\ K_{kl} & C_{kl} \end{smallmatrix} \right\}^{(l)}$.

This eccentric corporation can proceed pyramidally, homogeneously, or mixed, i.e., it only corporates the syndrome k with l , so the conflexion field is pyramidal, which assumes that the syndromes k and l are not empty. In the second case of the homogeneous conflexion field, the entirety of the occupations of all $0 \leq \gamma \leq k$ syndromes corporates with all $0 \leq j \leq l$. In this case, a syndrome conclusion can already lie below k and l , i.e., k and l can be empty. In the third case, finally, one syndrome k or l corporates pyramidally with the entire homogeneous occupation of the other syntrix. This assumes that the pyramidally corporating syndrome (pyramidal component) is not empty. While in the pyramidal and mixed case the conflexion field is fixed, the syndromatic position of this field in the homogeneous case is undetermined when it is known that the syndrome conclusions of both syntrices lie deeper than k and l . From the conceptual formations of the concentrator and eccentric, two variants, namely the pseudo-concentrator and pseudo-eccentric, can be developed, with the help of which the ambiguous purely sykolative (*s*) or purely metaphorical (*m*) corporations of the first and second class can be unambiguously interpreted without needing to bind restrictive additional conditions to the synkolation laws or the metaphor.

For example, if synkolative corporations of the first or second class are in the form $\langle f\widetilde{a}\widetilde{m} \rangle \{ \}_{(s)} \langle \phi\widetilde{b}\widetilde{\mu} \rangle$, then, if $\widetilde{a} \neq \widetilde{b}$ holds, the corporation is ambiguous with respect to the metaphor, but this ambiguity disappears if the two metaphors are considered as the two elements of a metaphorical pseudo-syndrome. If this is done, and the corporated synkolator acts on this metaphorical pseudo-syndrome, then the synkolation process becomes three-part in the special

case, because the corporated synkolator synkolates a syndrome sequence from $\widetilde{a}$, another from $\widetilde{b}$, and a third mixed from $\widetilde{a}$ and $\widetilde{b}$. Since $\widetilde{a}$ and $\widetilde{b}$ form a metaphorical pseudo-syndrome, a non-uniquely acting corporator of the form $\{\}_{(s)}$ is called a pseudo-eccentric, because the three synkolation branches are eccentrically positioned around the metaphorical pseudo-syndrome.

In the opposite case, a two-valued but metaphoric corporative law of the form $\langle \widetilde{f} \widetilde{a} \widetilde{m} \rangle \{\}_{(m)} \langle \widetilde{\phi} \widetilde{b} \widetilde{\mu} \rangle$ with $(\widetilde{f}, \widetilde{m}) \neq (\widetilde{\phi}, \widetilde{\mu})$ is present. Here, the two different metaphors combine to form a new one, which creates a center, while the two non-corporating synkolators induce two concentric syndrome sequences around this true center as a metaphor, so that an ambiguous corporator $\{\}_{(m)}$ can be referred to as a pseudo-concentrator. A pseudo-concentrator must always be a metaphoric corporator, while a pseudo-excenter must be a synkolative corporator, and both corporators can belong to the second class at most. With the help of the pseudo-concentrator and pseudo-excenter, it is possible to interpret and illustrate multi-valued metaphoric or synkolative corporations of the first and second corporator class in a clear and unambiguous way.

3.5 Syntropoden Architectonics of Multi-Linked Conflexive Syntrices.

Apparently, one can always attach a regular eccentric corporation of the form

$$\widetilde{a}|_1 \{ \}^{(k)} \{ \}^{(l)} \widetilde{a}|_2, \overline{|}, \widetilde{a}|_3$$

to another corporation

$$\widetilde{a}|_3 \{ \}^{(m)} \{ \}^{(n)} \widetilde{a}|_4, \overline{|}, \widetilde{a}|_5$$

provided that the connecting predicates are identical, so that the substitution

$$\widetilde{a}|_1 \{ \}^{(k)} \{ \}^{(l)} \widetilde{a}|_2 \{ \}^{(m)} \{ \}^{(n)} \widetilde{a}|_4, \overline{|}, \widetilde{a}|_5$$

becomes possible. In this way, an entire chain of corporations can be carried out. If there are $1 \leq i \leq N$ syntrices $\widetilde{a}|_i$ connected in such a *corporation chain*, and the result of this corporation chain is connected with the predicate $\overline{|} : \overline{|}$ to $\widetilde{c}|$, then the corporation chain can be represented by this symbolism¹²:

$$\left(\widetilde{a}|_i \{ \}^{(k_i)} \{ \} \{ \}^{(l_{i+1})} \widetilde{a}|_{i+1} \right)_{i=1}^{N-1}, \overline{|}, \widetilde{c}| \quad (13)$$

If the predicate is an identity, then $\widetilde{c}|$ is fundamentally conflexive when there is at least one excenter in the chain. If $\varepsilon \leq N - 1$ is the number of excenters within the chain, then $\widetilde{c}|$ is necessarily and sufficiently an $\varepsilon + 1$ -membered *Conflexive Syntrix*. For $\varepsilon = 0$, there are no excenters, only concentrers, i.e., for all corporators, $K_i = l_i = 0$, and $\widetilde{c}|$ becomes singly conflexive, thus concentric. Hence, combined or natural syntrices with a concentric synkolation course are special cases of the general multi-membered conflexive syntrix, whose conflexion field, however, can only arise through eccentric corporations of concentric synkolating syntrices. Conversely, there is always the possibility to construct any multi-membered conflexive syntrix from concentrically synkolating syntrices by suitable excenters, which in turn can always be built up from the four pyramidal elementary structures by concentrers. The intuitive meaning of such a conflexive syntrix is as follows: If the $1 \leq i \leq N$ eccentrically corporated syntrices $\widetilde{a}|_i$ are all concentric, i.e., the syndrome occupations synkolate concentrically around the metaphore $\widetilde{a}_i$, then all $\widetilde{a}_i$, if there is no concentric metaphore corporation at all, must lie in the same aspect system, i.e., each of these metaphors lies above its own area related to the respective aspect system and forms the idea of the area represented as a syllogistically oriented category.

Through the purely excentric Corporator Chain, these areas are not corporated in their apodictic elements and the Syndromes $1 \leq \gamma_i < k_i$, but only in the higher Syndromes $\gamma_i \geq k_i$, starting with the pseudometrophoric Conflexion field at $\gamma_i = k_i$. The individual components of a multi-membered Conflexive Syntrix are therefore the uncorporated base pieces of the Syntrix, which are to be called *Syntropods*. These Syntropods consist of the respective Metrophor of the area and the first Syndromes $1 \leq \gamma_i < k_i - 1$, where $k_i - 1$ (last Syndrome before the Conflexion field) indicates the *Syntropod length* of the corresponding member of the Conflexive Syntrix. If the chain is completely excentric, a

¹²In the original, there is a sentence fragment: "... connected, then the symbolism should be for the corporation chain"

termination of the chain after any member $\chi < N - 1$ already forms a Conflexive Syntrix, which in turn is incorporated with the remaining $N - \chi$ Syntizes. From this, however, it necessarily and sufficiently follows that the concentric and excentric corporation law of general Corporator Chains also maintains its validity for Conflexive Syntizes of any Syntropod number.

Based on this fact, an architectural classification of all Syntizes can be carried out. This classification is based on the Syntropod number and the Synkolation course or the structural composition of the Conflexion fields. First, it turns out that a chain of regular Excenters provides a Conflexive Syntrix in which all Syntropods have different lengths, while equi-longitudinal Excenters must lead to identical Syntropod lengths that are different from 0. If all these equi-longitudinal Syntropods are identical with 0, i.e., all $k_i = 1$, then the chain can only consist of Concentricors because then $\varepsilon = 0$, and the Syntrix is single-membered Conflexive, i.e., concentric. However, the type of Excenters of the respective chain is essential for such an architecture of Conflexive Syntizes. Thus, pyramidal, homogeneous or mixed corporation components can be contained in one and the same Conflexive Syntrix. In the mixed case, homogeneous and pyramidal incorporated Syntizes alternate, so that only two classes of Syntropods can appear architecturally. All pyramidal incorporated components provide Syntropods with fully occupied Syndromes. In the case of homogeneous corporation, however, there is a possibility that the higher Syndromes of the respective Syntropod remain empty because the Syndrome completion may lie lower than the syndromatically undetermined position of the Conflexion field. To refine the concept, Syntropods of homogeneous corporations can also be referred to as Syndrome balls if the Syndrome completion lies deeper than the Syntropod length, which is always 1 less than the syndromatic length of the Conflexion field. The term Syndrome ball is justified by the fact that within the Syntropod, between the Syndrome completion and the Conflexion field, there may be an undetermined number (at least 1) of empty Syndromes.

In complete analogy to Pseudoconcenters and Pseudoexcenters, Excenters of the first or second class in Conflexive Syntizes can also act purely syncolative or purely metaphoric, causing ambiguities in the conflation field. If the Excenter is purely metaphoric, the conflation field is uniquely determined, but in analogy to the Pseudoconcenter, the syncolation law remains at least ambiguous, while with a syncolative Excenter, the syncolation law becomes unique and the syndromes forming the conflation field overlap into a multivalued Pseudoconflation field, which in turn leads to several syncolation branches above the Pseudoconflation field. Consequently, the first group of Excenters is called pseudoconflexive Concenters, while the second group is called pseudoconflexive Excenters. In the corporation chains of Conflexive Syntizes, there are basically six types of Corporators: Concenters, Excenters, Pseudoconcenters, Pseudoexcenters, as well as pseudoconflexive Concenters and pseudoconflexive Excenters. If the chain (J) consists of only one Corporator type, and $N > 1$ Syntizes are incorporated in it, then an N-membered pure Corporator chain is present. The concentric chain, whose Corporators are only Concenters, can always be only a concentric Syntrix when the members are such concentric Syntizes, i.e., the syntheses (12) and (12a) are such concentric chains. Excentric chains, on the other hand, fundamentally result in Conflexive Syntizes, and N-membered ones when the individual members are concentric Syntizes. The resulting Conflexive Syntrix then always has N Syntropods, which can be regularly excentric or equi-longitudinal with respect to their Syntropod lengths. If there is a pseudoconcentric chain whose members are concentric Syntizes, then the Metrophor of the pseudoconcentric Syntrix is unique, but there are N syncolation branches above it, each of which characterizes a unique syncolation course. In the case of the pseudoexcentric chain, on the other hand, under the same condition of concentric members, the syncolation course would become unique, while the Metrophor consists of a multi-membered metaphoric Pseudosyndrome, whose individual members can be regarded as Pseudosyntropods with the equi-longitudinal Syntropod length 0. However, this again leads to several syncolation branches, and the number of these branches is combinatorially given as $N + \binom{N}{2} = \frac{1}{2}N(N + 1)$.

For the pseudo-concentric-flexive chain, the same applies as for the eccentric chain, but here, a total of N synkolation branches lie above the flexion field, corresponding to the ambiguity of the synkolation law. For the pseudo-flexive-eccentric chain, the synkolation law beyond the pseudo-flexion field is unambiguous, but the flexion field becomes a pseudo-syndrome, which in turn allows $\frac{1}{2}N(N + 1)$ synkolation branches above the pseudo-flexion field. In general, there are no pure corporator chains; rather, all six corporator types are contained in such a mixed chain. With the

selective operation $()_i, \tilde{a} \equiv \tilde{a}|_i$, we have $\lceil, \tilde{a}, \lceil, \tilde{c}$ when

$$\lceil \equiv \left(()_i {}^{(k_i)} \{ \}_i^{(l_i+1)} ()_{i+1} \right)_{i=1}^{N-1}$$

characterizes the mixed corporator chain. Apparently, these chains can only contain all six corporator types if $N-1 \geq 6$. It is evident that the architectural structure of a flexive-syntrix represented as a corporator chain, regarding the syntropoden construction, the flexion field structure, and the syndromes above it, is determined exclusively by the nature of the chain and the chain elements, i.e., from

$$\lceil, \widetilde{a}, \lceil, \widetilde{c} \rceil \vee \lceil \equiv \left(()_i {}^{(k_i)} \{ \}_i {}^{(l_i+1)} ()_{i+1} \right)_{i=1}^{N-1} \quad (13a)$$

the entire syntropoden architecture of the multi-membered flexive-syntrix $\lceil, \widetilde{a} \rceil$ must emerge.

First, it must be established that the sequence of elements in (13a) is not commutative, as a transposition of elements would always result in a change in the structure of the syntropodes and flexion fields. This is because, due to the corporator laws, it is not irrelevant how a concentric part is corporated with a concentric corporator (concenter, pseudoconcenter, and pseudoexcenter) to an already existing flexive part. The number of syntropodes is essentially determined by the nature of the flexive elements and the number of eccentrically acting corporators (excenters and both pseudo-flexive), while the structure of the syntropodes, the flexion field structure, and the course of the synkolation branches lying above them are also determined by the concentric elements and concentrically acting corporators.

Furthermore, the specific nature of the eccentric corporators must find its expression in the syntropode architecture. The number of regular and equi longitudinal excenters, as well as the syndromatic position of the flexion fields determined by them, dictate the distribution of syntropode lengths and the form of eccentric corporators as pyramidal, homogeneous, or mixed. This also determines the internal structure of the syntropodes or syndrome spheres and the inner structure of the flexion fields, while the pseudo-forms significantly shape the variety of synkolation branches, and so on.

As diverse as the architecture of multi-membered flexive systems according to (13a) may be, there are fundamentally only three general architectural structures: the distribution of syntropodes, the distribution of flexion fields, and the distribution of all synkolation branches lying above them. In addition to this external architecture, (13a) also describes the internal structure of the syntropodes, flexion fields, and those of the syndromes lying above these flexion fields.

4 Enyphan Syntrices

4.1 Syntrix Totalities and their Generatives.

The elements of each syntrix architecture are the pyramidal elementary structures and the totality of all those forms that result from concentric incorporations out of them. The four classes of these elementary structures (these are the actual syntrometric elements) can thus generate syntrix manifolds that are composed of arbitrary concentric syntrices if any centers are available that corporately incorporate these elementary forms of the pyramidal syntrices in any arbitrarily specifiable manner. If A is the underlying aspect system and S is a subjective aspect associated with it, an arbitrary number of conceptual domains can be related to A in such a way that a metaphor with respect to A is present over each domain. For each of these metaphors, in turn, there is generally an infinite sequence of pyramidal synkolation laws, the nature of which depends on S , i.e., an generally infinite sequence of pyramidal syntrices. Since the number of domains that can be related to A is also unlimited, there is a multiply infinite set of pyramidal syntrices in A , each of which can be destructed into the four pyramidal elementary structures. In this way, $1 \leq i \leq 4$ infinite manifolds P_i of pyramidal syntrices of one elementary form each are created. These P_i are now linearly arranged as value stores of an elementary form.

Apparently, there are no incorporators that allow any relation between these elementary structures in the form of *Syntrix transformations*, so the four value stocks P_i of an S in A must be independent of each other. If this were not the case, the elementary character of these pyramidal syntrices would not be preserved, which would contradict the results of previous investigations on pyramidal elementary structures. Consequently, the four value stocks P_i of the elementary structures span a metaphorically four-dimensional space over S in A , the so-called four-dimensional *storage* of all possible concentric syntrices in S , such that each subjective aspect of an aspectival system must have access to such a syntrix storage. Concentric incorporations of the pyramidal elementary structures, however, build up any arbitrary concentric syntrix. So, if you also specify an arrangement of Q centers, which is called a *corporator simplex*, this simplex with the four-dimensional syntrix storage denotes the *generative*

$$G \equiv \left[\begin{array}{c|c|c} 4 & Q & \\ P_i & \{ \}_{(j)} & \\ 1 & 1 & \end{array} \right]_{(A,S)} \quad (14)$$

in A, S , since the Q elements of the corporator simplex generate all other possible concentric structures in A, S using the simplex through the corporation of the elementary forms of the storage, i.e., this generative fulfills a four-dimensional manifold with a Syntrix totality that is complete with respect to the simplex.¹³

Each subjective aspect indeed has an associated Syntrix storage, but each Syntrix storage can generally generate an infinite set of Syntrix totalities, as each corporator simplex characterizes the generative of a Syntrix totality in S ,

and there are generally infinitely many possibilities for defining $\{ \}_{(j)} \mid \begin{array}{c} Q \\ 1 \end{array}$ in S . There are different possibilities for the mode of action of the simplex elements, namely a) the endogenous, b) the regular, and c) the extra-regular. In the

¹³ ChatGPT4.0: Alright, let's try to make sense of this in a Feynman-like way. We're going to talk about something called "Enyphan Syntrices."

First, we need to understand what syntrix totalities and their generatives are. We can think of them as the basic building blocks of a syntrix architecture. They come in four classes, and they can create syntrix manifolds, which are like combinations of these building blocks.

Now, imagine you have an underlying aspect system, A , and a subjective aspect, S . With these, we can have an unlimited number of conceptual domains, and each of them can have an associated metaphor. For every metaphor, there's a bunch of pyramidal synkolation laws, which are like rules that help us create these pyramidal syntrices.

So, there's a lot of these pyramidal syntrices in A . The cool thing is that each of them can be broken down into four basic pyramidal elementary structures. This gives us four sets, or as the text calls them, "infinite manifolds" P_i , of pyramidal syntrices.

Now, here's the catch: There's no way to relate these elementary structures through something called "syntrix transformations," so the four sets P_i have to be independent of each other. This means that they form a four-dimensional space, called a "storage," that holds all the possible combinations of syntrices.

So, we've got this storage, and we also have something called a "corporator simplex," which is an arrangement of Q centers. The combination of the storage and the simplex gives us a "generative," which is like a big factory that can create all sorts of different concentric structures in our aspect system.

In summary, enyphan syntrices are complex structures built from simpler pyramidal building blocks. These blocks are stored in a four-dimensional space, and when combined with a corporator simplex, they can generate an incredible variety of concentric structures.

endogenous case, corporations of the simplex within a P_i generate Pyramidal syntrices again from the *value storage*, while in the second and third cases, the regular and extra-regular cases, arbitrary concentric Syntrices are formed by applying the corporator simplex. In the image of the four-dimensional metaphor, the $\text{binom}42 = 6$ surfaces between P_i and P_k can be occupied regularly by the corporation of two Syntrices each from P_i and P_k . This occupation of the six surfaces occurs Q times, since only one concentrator of the simplex is needed for each corporation, and there are Q concentrators in the simplex. The $\text{binom}43 = 4$ cubes spanned by P_i, P_j, P_k , on the other hand, are filled by corporations of three Syntrices each from P_i, P_j , and P_k , for which two concentrators of the simplex are needed. That is, the occupation of these cubes is $\binom{Q}{2} = \frac{Q}{2}(Q - 1)$ times, not identical in terms of corporation, but there are still Q corporatively identical occupation multiples, because the two corporators can also be identical.¹⁴

Finally, there remains the occupation of the four-dimensional area by Syntrices, which are formed by the corporation of elements from P_1, P_2, P_3 , and P_4 . Three concentrators are necessary for these corporations, such that the occupation of this one four-dimensional area is $\binom{Q}{3} = \frac{Q}{6}(Q - 1)(Q - 2)$ times corporatively non-identical. In addition, there are $\binom{Q}{2}$ corporatively twice identical occupation multiplicities, as in the corporator triplet, two or all three concentrators can be identical. In this way, the four-dimensional storage has been regularly filled, so that all resulting concentric Syntrices form the regular *Syntrix framework* of the totality. Moreover, there are arbitrary other corporator chains with $4 \leq j \leq Q$ corporatively non-identical, or identical (which also allows for $j > Q$), concentrators of the simplex possible, which also form manifolds of Syntrices and extraregularly complement the regular Syntrix framework of the totality. While the regular framework depends on the specific construction of the corporator simplex and thus represents a typical characteristic of the totality, this is only the case for the extra-regular complement with respect to the concentrators under discussion, but not for the occupation by corporator chains.

For the structure of a totality, there are basically two possibilities: either the elementary pyramidal structures are densely distributed in the four Syntrix storages P_i , i.e., the distribution along the P_i is continuous, or this distribution corresponds to some selection principle and is consequently discrete. In the first case, the Syntrix totality itself is continuous, i.e., the Syntrices within the totality are densely distributed everywhere, while this is not the case with the discrete distribution. As long as the simplex content is limited, i.e., only a finite number of concentrators construct the simplex, a finite area of the totality can always contain only a finite number of Syntrices when the distribution along the P_i is discrete.¹⁵

There are thus fundamentally continuous and discrete totalities, which, however, still allow for specific extreme

¹⁴Alright, let's dive into this concept in a Feynman-style explanation. We're still talking about syntrix storage and how it's related to subjective aspects. But here's the thing: each syntrix storage can actually generate a whole bunch of syntrix totalities. This is because there's an infinite number of ways to define the corporator simplex, which, as we've seen before, helps to create those totalities.

Now, there are three different ways that the simplex elements can work: endogenous, regular, and extra-regular. In the endogenous case, the simplex works within a single P_i and creates new pyramidal syntrices from the "value storage." In the regular and extra-regular cases, the simplex creates arbitrary concentric syntrices.

Imagine a four-dimensional metaphor, where there are surfaces and cubes created by the different P_i . The surfaces between P_i and P_k can be filled regularly with syntrices from both P_i and P_k . This happens Q times, since there are Q concentrators in the simplex.

On the other hand, the cubes that are formed by P_i, P_j, P_k are filled by corporations of three syntrices each from P_i, P_j , and P_k . To do this, you need two concentrators from the simplex. The occupation of these cubes happens $\frac{Q}{2}(Q - 1)$ times, but there are still Q corporatively identical occupation multiples, because the two corporators can also be identical.

In summary, each syntrix storage can generate a wide variety of syntrix totalities using a corporator simplex. The simplex elements can act in different ways, such as endogenous, regular, or extra-regular, which helps to create new combinations of syntrices. In a four-dimensional metaphor, surfaces and cubes are filled with syntrices from different P_i , with the help of concentrators in the simplex.

¹⁵Let's break this down in a Feynman-style explanation. So, we're now looking at how syntrices fill the four-dimensional area by incorporating elements from P_1, P_2, P_3 , and P_4 . To do this, we need three concentrators, which means the occupation of the four-dimensional area happens $\frac{Q}{6}(Q - 1)(Q - 2)$ times, with each one being corporatively non-identical. Also, there are $\binom{Q}{2}$ situations where the occupation is corporatively twice identical, because two or all three concentrators can be the same.

With all of this in place, the four-dimensional storage is filled regularly, creating a regular "syntrix framework" for the totality. There are also other corporator chains that can form additional syntrices and supplement the regular framework in an extra-regular way. While the regular framework is tied to the specific construction of the corporator simplex, the extra-regular framework depends on the concentrators being discussed but not on the occupation by corporator chains.

Now, there are two ways to structure a totality: either the elementary pyramidal structures are densely distributed in the four syntrix storages P_i (meaning a continuous distribution), or the distribution follows some selection principle and is discrete. In the first case, the syntrix totality is continuous, with syntrices densely distributed everywhere. In the second case, with a discrete distribution, this is not true.

If the simplex content is limited, meaning there's only a finite number of concentrators constructing the simplex, a finite area of the totality can only contain a finite number of syntrices when the distribution along the P_i is discrete.

cases. In the continuous and discrete forms, the simplex content is limited, and the P_i are unlimited but either continuously or discretely occupied. If the simplex content grows beyond all limits, i.e., the simplex is open, then the totality becomes hyper-continuous, limited or unlimited, depending on whether the P_i are limited or unlimited when P_i is continuously occupied. The discrete totality then becomes discretely pseudo-continuous, limited or unlimited with an open simplex.

The real continuous or real discrete, or hyper-continuous, or discretely pseudo-continuous totality is thus reduced to the concept formations of continuous and discrete totalities, with the real or pseudo forms appearing when the corporator simplex is limited or open. Each Syntrix totality thus forms a four-dimensional Syntrix field, the structure of which is determined by the regular Syntrix framework. This Syntrix field can be continuous or discrete, with a Syntrix-like field continuum always present in the open case of the simplex, which can only become a discrete Syntrix field in the limited case of the simplex.

The metaphorical illustration of a Syntrix totality by a four-dimensional space actually has only the value of a metaphor, because a Syntrix is fundamentally different from a point in a four-dimensional space, both qualitatively and quantitatively, unless one would consider the number quadruple describing the point as a metaphor, so that the syndromes are occupied by functional values. This would be the starting point of an anthropomorphic syntrometry, but in this way, the concept of a geometric point would have to be qualitatively extended again, which implies that the four-dimensional Syntrix totality can never be derived exactly from a four-dimensional point continuum. This impossibility becomes even clearer when considering that in this way, the concept of a point in n -dimensional space can also be anthropomorphically syntrometrically extended, and that all these Syntrices are again the occupations of four-dimensional totalities.

With regard to the specific anthropomorphic aspect system, Syntrix totalities can be further divided into two groups, namely quantities and qualities. Quantities represent such Syntrix totalities that are possible in the subjective aspect of rational analytical formal logic, while qualities lie in the other subjective aspects of the anthropomorphic aspect system as Syntrix totalities.

In addition to these general four-dimensional Syntrix totalities, there are three special cases in which 1, 2, or 3 stocks of pyramidal elementary forms fail. This leads to the degenerated three-, two-, or one-dimensional totalities. For the first degeneration, there are $\binom{4}{1} = 4$ possibilities of failure for each of the individual P_i . For the second degeneration (two-dimensional totality), there are $\binom{4}{2} = 6$ failure possibilities for the decomposing pairs of stocks P_i, P_j with $i \neq j$, and for the third degeneration (one-dimensional totality), there are $\binom{4}{3} = 4$ failure possibilities for stock triplets P_i, P_j, P_k with $i \neq j \neq k$. In addition to these degenerations, which include a total of fourteen degeneration possibilities across three classes, the non-degenerated four-dimensional totality has a universal character.

It remains to be investigated how the null syntrices are distributed in such a Syntrix totality. If the corporator simplex is limited and does not contain a corporator that could generate a null syntrix in the individual regular assignments of the framework, there can be no null syntrices in the totality if there is no null syntrix contained in the storage of the pyramidal elementary forms. However, in this case, since the Syntrix framework does not contain any null syntrices and the extraregular filling of the totality is achieved through corporations of the simplex, there can also be no null syntrices in this extraregular part. This implies that under these conditions, the entire totality remains free of null syntrices. Therefore, if the simplex is limited and its corporators cannot generate a null syntrix, there can be no null syntrices in the totality if the storage is also free of null syntrices. On the other hand, such null syntrices are present in the totality if this last condition is not met, or if it is an open corporator simplex in the generative, because then all possible corporators allowed in the underlying aspect system are admitted, which can certainly lead to the generation of null syntrices in the regular or extraregular part of the totality.

All statements can also be extended to multiparametric primigenic aeondynes. In this way, "primigenic aeondyne totalities" can be defined, and this is done in complete analogy to the Syntrix totalities. This is because each primigenic aeondyne can be regarded as a continuous, multiparametric infinite set of syntrices, i.e., as a band syntrix, to which the corporation laws can be applied, since the band syntrices comply with the same syntrometric laws as the other syntrices. However, primigenic aeondyne totalities, in addition to the occupancy structuring of the Syntrix totalities, also show a structure dependent on the course of the primigenic aeondynes. According to the theory of primigenic aeondynes, these aeondynes are nothing more than syntrices whose elements are functors of certain syntrometric parameters along any definition domains. This means that for each aeondyne totality, there is a multidimensional "syntrometric carrier space" that contains the entirety of all parameter values of the definition domain of the aeondyne

totality. The dimension number of this carrier space, in turn, is identical to the number of argument parameters of all aeondynes within the totality. Here, laws similar to those of analysis apply to subspaces and hyper-surfaces in which individual aeondynes can be defined, while the corporations take place in the higher-level areas.

4.2 The discrete and continuous Enyphansyntrix.

Each corporation law characterizes, in the sense of a functor, the connection of Syntrices that leads to any other syntrometric form, such that the functor connections mediated by the Predicatrix of a subjective aspect always have the universal quantifier character. Apparently, the forms thus incorporated belong as genuine elements to the occupation of a Syntrix totality only if they are single-membered, i.e., if the corporation law has concentrator properties. For

example, let $\widetilde{c}$ be any concentric Syntrix¹⁶ and Γ be a concentrator that incorporates $\widetilde{c}$ with any other concentric Syntrix, i.e., $\widetilde{a}, \widetilde{b}, \widetilde{c} \mid \Gamma$ obviously represents a syntrometric prescription that generates a new concentric form, e.g. $\widetilde{\beta}$, in the predicate $\mid$ from any element $\widetilde{b}$ of the totality, such that $\widetilde{\beta}, \mid, \widetilde{a}, \widetilde{b}$ applies.

Here, $\widetilde{a}, \widetilde{b}$ means that the prescription $\widetilde{a}$ acts on $\widetilde{b}$ in the form $\widetilde{c} \mid \Gamma \widetilde{b}$. Since only one element ($\widetilde{\beta}$) is created, $\widetilde{a}$ is referred to as a *discrete Enyphansyntrix* if $\widetilde{\beta}$ belongs to the same totality. This is always the case when the Enyphansyntrix $\widetilde{a}$ is defined in the totality in which $\widetilde{b}$ belongs to the occupation. However, if $\widetilde{a}$ is supposed to

be defined in this totality, then obviously $\widetilde{c}$ must belong to the occupation and Γ to the Corporator simplex. These discrete Enyphansyntrices are thus syntrometric functor prescriptions with corporator properties, which generate a new element from an element of the totality.

This concept of a simple discrete Enyphansyntrix is capable of an extension, namely the multiple discrete Enyphansyntrix (n-fold), because the corporator Γ could be replaced by a chain of $1 \leq j \leq n$ corporators Γ_j , denoted as $(\Gamma_j(\cdot))_1^n$. This leads to the operation of the n-fold discrete Enyphansyntrix:

$$\widetilde{a}, \mid, \widetilde{c} \mid (\Gamma_j(\cdot))_1^n \vee \widetilde{\beta}, \mid, \widetilde{a}, \widetilde{b}. \quad (15)$$

In this general form of the discrete Enyphansyntrix, $\widetilde{c}$ is called its root and $(\Gamma_j)_1^n$ is referred to as the *Enyphan chain*,

where the individual components Γ_j of the chain are generally oriented. If such an orientation of the chain structure exists, a distinction must be made between the internal corporators $j < n$ and the external corporator $j = n$.

However, this distinction becomes unnecessary if the Γ_j can be arbitrarily permuted. The n-fold form $\widetilde{a}$ can only be defined in totality if $\widetilde{c}$ belongs to the domain and all Γ_j belong to the simplex. If there are $1 \leq i \leq N - 1$ concentrators $\Gamma_i \equiv \left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\}_i$ ¹⁷, which incorporate N Syntrices $\widetilde{a}_{i+1}$ in the form of a corporator chain, then

the chain $\widetilde{f}, \mid, (\widetilde{a}_i \Gamma_i \widetilde{a}_{i+1})_1^{N-1}$ is a concentric Syntrix, if the $\widetilde{a}_i$ are concentric. Apparently, $\widetilde{f}$ depends on the N incorporating $\widetilde{a}_i$.

If Γ_i belongs to the corporator simplex of a totality with continuous occupation, then $\widetilde{f}$ is also in this totality if the $\widetilde{a}_i$ belong to its occupation. Since the syntrices lie densely everywhere in the continuous totality, there is the possibility, for example, for $0 \leq l \leq k \leq n \leq N$ arguments $\widetilde{a}_k$ of the chain of corporation rules G_k to be found, such that $\widetilde{a}_k G_k \widetilde{a}_l \rightarrow \widetilde{a}_{(g)}|_k$ goes, where $\widetilde{a}_{(g)}|_k$ is a syntrix with empty syndromes. These $n - l \leq N$ syntrices with empty syndromes can now be corporated with the contraoperative G_k^{-1} to the elements $\widetilde{a}_l$ of the chain, so that if the $n - l$ elements are adjacent, the chain becomes

$$\left(\widetilde{a}_j \Gamma_j \widetilde{a}_{j+1} \right)_1^{l-2} \left(\widetilde{a}_k G_k^{-1} \widetilde{a}_{(g)}|_k \Gamma_k \widetilde{a}_{k+1} G_{k+1}^{-1} \widetilde{a}_{(g)}|_{k+1} \right)_1^{n-1} \left(\widetilde{a}_i \Gamma_i \widetilde{a}_{i+1} \right)_{n+1}^{N-1}.$$

¹⁶HS change from: any Corporator chain

¹⁷HS: inserted Γ_i

If we denote this chain with $\widetilde{F}|$, a contraoperative concenter ε can be formed from the concenters of the simplex with respect to $\widetilde{F}|$, which corporates $\widetilde{F}|$ contraoperatively with $\widetilde{f}|$. Apparently, then $\widetilde{F}|\varepsilon\widetilde{f}| \rightarrow \widetilde{f}|$ (empty syndromes). This contraoperative corporation now evidently indicates an infinitesimal change of the chain $\widetilde{f}|$ with respect to the densely lying $\widetilde{a}|_k$ of the continuous totality. However, this process is determined solely by the G_k and ε , so that the abbreviation of this process can be represented by the symbol $(G_k, \varepsilon]_1^n \equiv D$, where E is a syntrix operation. $E, \widetilde{f}|$ thus means that E acts on $\widetilde{f}|$ in the manner shown, and thereby changes $\widetilde{f}|$ infinitesimally in the continuous totality, which is why E can also be called an infinitesimal *Enyphane*. The action takes place by corporating the arguments $\widetilde{a}|_k$ with the G_k to $\widetilde{a}(g)|_k$ (approximately), and these are corporated with the contraoperatives to G_k at the $\widetilde{a}|_k$, so that $\widetilde{F}|$ is created.¹⁸

Then, $\widetilde{f}|$ is corporated contraoperatively with ε , so that $\widetilde{F}|\varepsilon\widetilde{f}|, \bar{||}, E, \widetilde{f}| \rightarrow \widetilde{f}|$ can be set. Formally, this scheme is summarized in:

$$\begin{aligned} \widetilde{f}|, \bar{||}, (\widetilde{a}|_i \bar{||} \widetilde{a}|_{i+1}]_1^{N-1} \vee \bar{||} \equiv \left\{ \begin{matrix} K_s & C_s \\ K_m & C_m \end{matrix} \right\} \vee 0 \leq l \leq k \leq n \leq N-1 \vee \widetilde{a}|_k G_k \widetilde{a}|_k \rightarrow \widetilde{a}(g)|_k \\ \widetilde{F}|, \bar{||}, (\widetilde{a}|_j \bar{||} \widetilde{a}|_{j+1}]_1^{l-2}; (\widetilde{a}|_j \bar{||} \widetilde{a}|_{j+1}]_1^{l-2} (\widetilde{a}|_k G_k^{-1} \widetilde{a}(g)|_k \bar{||} \widetilde{a}|_{k+1} G_{k+1}^{-1} \widetilde{a}(g)|_{k+1} \bar{||})_1^{n-1} (\widetilde{a}|_i \bar{||} \widetilde{a}|_{i+1}]_{n+1}^{N-1} \\ \widetilde{F}|\varepsilon\widetilde{f}| \rightarrow \widetilde{f}| \vee (G_k, \varepsilon]_1^n \equiv E \vee \widetilde{F}|\varepsilon\widetilde{f}|, \bar{||}, E, \widetilde{f}| \end{aligned} \quad (16)$$

For the infinitesimal Enyphane E , the inverse Syntrix operation, i.e., the inverse Enyphane E^{-1} , can be found in complete analogy, which has contraoperative properties relative to E . Thus, for this inverse Enyphane, with the statement $\bar{||}$, it can always be set

$$E^{-1}, E, \widetilde{f}|, \bar{||}, \widetilde{f}| \quad (16a)$$

from which it follows that E^{-1} reverses the process E , from which the commutative property concerning $\bar{||}$ can be concluded directly. From this, it follows again that only in the case of explicit syntrometric investigations is it necessary to distinguish between infinitesimal and inverse Enyphanes. If this explicit case is not present, then E should describe any Enyphane with infinitesimal or inverse properties.¹⁹

When the process of an Enyphane is combined with a discrete Enyphansyntrix, a more general syntrometric operation emerges, namely the so-called "continuous Enyphansyntrix." This operation can act on any element of the

¹⁸Interpretation by ChatGPT4.0: Alright, let's try to break this down in a Feynman-esque way. Imagine we have a chain, $\widetilde{F}|$. Now, we're going to form something called a "contraoperative concenter" (let's call it ε) using some elements from the simplex. This ε is going to work together with our chain $\widetilde{F}|$ and another element $\widetilde{f}|$.

When they work together, something interesting happens: $\widetilde{F}|\varepsilon\widetilde{f}| \rightarrow \widetilde{f}|$, which gives us empty syndromes. This process indicates a tiny, almost imperceptible change in the chain $\widetilde{f}|$ with respect to some other elements, $\widetilde{a}|_k$, which are densely scattered around.

Now, the whole thing depends on two elements: G_k and ε . We can represent this interaction with a special symbol, $(G_k, \varepsilon]_1^n \equiv E$. Here, E is a kind of operation called a "syntrix operation." So, when we write $E, \widetilde{f}|$, it means that E is acting on $\widetilde{f}|$, causing a tiny change in it.

This whole process is like a very subtle dance between the different elements. The arguments $\widetilde{a}|_k$ are combined with the G_k to create something new, $\widetilde{a}(g)|_k$. Then, these new elements are combined with their counterparts, the contraoperatives to G_k , and the original $\widetilde{a}|_k$. This whole process creates the chain $\widetilde{F}|$.

¹⁹Alright, let's try to break it down in a Feynman-esque way. We're dealing with this thing called an "infinitesimal Enyphane," which is a special operation in the context of Syntrix operations. It's a bit like a tiny transformation that helps us understand and explore these mathematical structures.

Now, just as we have this infinitesimal Enyphane, we can also find its inverse, which we call the "inverse Enyphane." It's got some interesting properties when compared to the original Enyphane, specifically being contraoperative. This inverse Enyphane has the power to reverse the process of the original Enyphane, which gives us some nice mathematical properties like commutativity.

The important thing to remember is that we only need to worry about the distinction between infinitesimal and inverse Enyphanes when we're dealing with explicit syntrometric investigations. Otherwise, we can just think of the Enyphane as a general transformation with infinitesimal or inverse properties, depending on the situation.

So, in short, we're dealing with a mathematical framework that involves infinitesimal transformations (Enyphanes) and their inverses to explore and understand the underlying structures.

continuous totality with its Enyphan member and thus describe the change in the structure of the continuous occupation in the most universal form.

So if $\tilde{c}|$ is a discrete Enyphansyntrix of the form (15), and if E denotes an Enyphane represented by the form (16) or (16a), then the general continuous Enyphansyntrix $\tilde{C}|$ is described by the structure

$$\tilde{C}| \equiv \tilde{c}|, \sqcup E, \sqcup, \left(\tilde{a}|_i \sqcup_i \tilde{a}|_{i+1} \right)_1^{N-1} \sqcup \left(\sqcup E \vee \tilde{a}|, \sqcup, \tilde{C}|, \tilde{a}| \right) \quad (17)$$

If $\sqcup$ incorporates the Enyphane in $\tilde{c}|$ and $\sqcup$ denotes the active member of the discrete structure with the root $(\sqcup_1^{N-1})$. The necessary and sufficient existence condition for a continuous Enyphansyntrix is therefore the existence of a simple

or multiple discrete form of a connecting corporator $\sqcup$ (which must belong to the simplex) and an Enyphane, but this requires a continuous totality, i.e., a continuous Syntrix occupancy.

If $\tilde{C}|$ acts on any Syntrix, for example, $\tilde{b}|$, and the influence of $\tilde{c}|$ on $\tilde{b}|$ is given by $\tilde{c}|, \tilde{b}|, \sqcup, \tilde{\beta}|$ in the predicate $\sqcup$, it seems that the effect of $\tilde{C}|$ can be decomposed into the two steps $\tilde{\beta}|$ and $E, \tilde{b}|$, which need to be corporated with $\sqcup$. Thus, we have

$$\tilde{C}|, \tilde{b}|, \sqcup, \tilde{\beta}| \sqcup E, \tilde{b}| \vee \tilde{c}|, \tilde{b}|, \sqcup, \tilde{\beta}| \quad (17a)$$

From this, the following statement can be derived: The influence of each continuous Enyphan Syntrix can be decomposed into a discrete and an enyphane component, which are connected with the help of the corporator $\sqcup$, which characterizes the continuous form. $\tilde{C}|, \tilde{b}|$ can only have one Syntropode, since the discrete component is also single-membered like the Enyphan action $\sqcup$ even in multiple cases. If $\tilde{C}|, \tilde{b}|$ were multi-membered, this Syntrix would no longer belong to the totality, which would contradict the assumption. This contradiction would be due to the fact that multi-membered Conflexive Syntrix Constructs²⁰ no longer belong to the totality, but this statement is based on the concentric definition of Syntrix totality. In general, both discrete and continuous forms should be referred to by the umbrella term Enyphan Syntrix. Applied to elements of a totality (in which they themselves are defined), they again yield such elements.²¹

4.3 Classification of Enyphansyntrices.

The entirety of all Enyphansyntrices initially breaks down into two main groups: discrete and continuous forms. Discrete Enyphansyntrices can be further divided into specific subclasses. A discrete $\tilde{a}|$ is determined, on the one hand, by its components, i.e., by the $1 \leq i \leq N$ Syntrices already incorporated in it and the associated $N - l$ internal corporators that incorporate the N components. On the other hand, $\tilde{a}|$ is determined by the nature of the acting corporators, i.e., by the number n of the multiplicity of $\tilde{a}|$ and the structure of those corporators that, with their help, influence other Syntrices in the sense of external incorporation. The components and internal corporators are already classification features. The multiplicity of $\tilde{a}|$ is added, resulting in an immensely large possibility of class differentiation, as the components always satisfy the classification laws of concentric Syntrices, and the internal or external corporators, to whom the concentrator follows inevitably, the great variation possibility. For example, if there

are n acting corporators $\sqcup_i$, they incorporate Syntrix $\tilde{b}|$, on which $\tilde{a}|$ acts, a total of n times at the root $\tilde{c}|$ (this is an internal corporator chain of components) of Syntrix $\tilde{a}|$. The n -fold, discrete Enyphansyntrix $\tilde{a}|, \sqcup, \tilde{c}| \left(\sqcup_j \right)_1^n$ means

²⁰ChatGPT4.0: "Konflexivsyntrices" This term suggests a more complex or multi-component Syntrix structure that interacts with the system in a unique way.

²¹ChatGPT4.0: In simpler terms, we can think of the Enyphansyntrix as a more general operation that deals with how elements in a system change. There are two types: discrete and continuous. The discrete Enyphansyntrix is like a step-by-step change, while the continuous Enyphansyntrix represents smooth, uninterrupted changes. These two types of Enyphansyntrix can be connected and used together to describe complex changes in a system.

that, when $\widetilde{a}|$ acts on $\widetilde{b}|$, the root $\widetilde{c}|$ and $\widetilde{b}|$ are incorporated n times by the $1 \leq j \leq n$ concentrators $\lceil_i$. In this way, an approximate classification scheme of discrete forms is given. A completely analogous process can be carried out for the Enyphane, whose classification, together with that of the discrete forms, leads to a general classification of continuous forms. An essential subdivision of Enyphane is based on the investigation of whether the respective inverse exists or not, so that all Enyphane with existing inverse represent one main class and the remaining ones represent the other main class.²²

²²ChatGPT4.0 as Richard Feynman: So, let's break down the classification of Enyphansyntrices into simple terms. We've got two main groups: discrete and continuous forms. Discrete ones can be further divided based on their components and how they interact with other Syntrices. This creates a whole bunch of subclasses, giving us lots of variety.

Now, when it comes to classifying Enyphane, we can do something similar. We can combine the classification of discrete forms and Enyphane to get a general classification for continuous forms. The key thing to remember here is whether an inverse exists for each Enyphane or not. If it does, they belong to one main class, and if not, they go in the other one. Simple enough, right?

4.4 The syntrometric structures.

From what we've discussed so far, a Syntrix totality includes only concentrically corporating forms, since the fundamental simplex, by definition, contains no excenters. Therefore, all excentric corporations, such as multi-membered Konflexivsyntrices²³ and the concepts derived from them, no longer belong as elements in the generally four-dimensional totality. However, these so-called syntrometric structures are defined above the respective Syntrix totality. For example, a chain of N concentric corporations and $n - 1$ excentric corporations, composed of $N + n - 1$ total corporations, can corporately involve $N + n$ Syntrices of the totality. The resulting Syntrix is no longer single-membered because n concentric Syntrices were excentrically corporated by the $n - 1$ excenters, creating a Konflexion field that can no longer be defined within the totality. This Konflexion field belongs to an n -membered Konflexivsyntrix with n Syntropodes (which may also be Pseudosyntropodes or Syndrome balls), representing an elementary syntrometric structure above the four-dimensional totality when the concentric elements of the corporator chain belong to the totality's occupancy. These elementary syntrometric structures can be classified according to the number of Syntropodes and the Syndromes contained in the Syntropodes.

Suppose an n -membered Flexocorporative Structure²⁴ arises according to the excentric corporator law. The n Syntrices of the syntropodes apparently all lie within the totality, and n discrete Enyphansyntrices $\tilde{\alpha}|i$ can be formed, which act on the n syntropodes of the structure. Thus, for each $1 \leq i \leq n$ syntropode, which originates from $\tilde{b}|(0)_i$, a new Syntrix of the totality is created, namely $\tilde{\alpha}|i, \tilde{b}|(0)_i, \tilde{b}|(1)_i$, on which $\tilde{\alpha}|i$ acts again. This process is recursively repeated according to $\tilde{\alpha}|i, \tilde{b}|(y-1)_i, \tilde{b}|(y)_i$, such that an infinite series of Syntrices (Syntrixtensorians²⁵) arises for each syntropode. These Syntrixtensorians belong to the corporator simplex when the $\tilde{\alpha}|i$ do, otherwise, they are exogenous. These n Syntrixseries can be metaphorically illustrated as an n -dimensional Syntrix space over the respective totality, analogous to the four-dimensional memory of a totality. Depending on the properties of $\tilde{\alpha}|i$, this space, spanned by the Syntrixtensorians, is endogenous, exogenous, or mixed, and each point of this metaphorical space would be equivalent to an n -membered Flexocorporative Structure with its endogenous syntropodes in the original totality. Since the syntropodic Syntrices change from point to point in the n -dimensional Syntrix space due to the influence of $\tilde{\alpha}|i$, each point, meaning every elementary syntrometric structure, is characterized by a specific structure of its flexion field such that a flexion field change runs over the Syntrixtensorium, whose metaphorical argument range is the Syntrix space marked by the $\tilde{\alpha}|i$.

In addition, this Syntrix space is characterized by the corporation law, which provides the n -membered Conflexive Syntrix (i.e., the arrangement of the $n - 1$ excenters in the Corporator chain), to which the n -dimensional Syntrix space is related as a Syntrix Tensorium. Its metric (the so-called Syntrometry) finds its measure in the syndrome numbers and syndrome occupations of the Syntropods. The structure, or syntrometry, of this Syntrix Tensorium apparently changes when the syndrome occupations of the Syntropods change, that is, when the n excentrically incorporated Syntrices are replaced by n other elements of the totality, or when the excenters of the chain change, leading to changes in the lengths of the Syntropods.

Lastly, the dimension number of the Tensorium can change, namely when the eccentricity of the Corporator chain, i.e., the number $n - 1$ of excenters, is altered. The syntrometric structure of this n -dimensional Syntrix space is also varied by exchanging the Enyphansyntrices $\tilde{\alpha}|i$. The term Syntrix space is just the anthropomorphic metaphor of the n -dimensional Tensorium. From the properties of this Tensorium, it becomes immediately evident that each elementary syntrometric structure over a totality in the form of an n -membered Conflexive Syntrix can generate a generally infinite set (depending on the possibilities of the $\tilde{\alpha}|i$) of n -dimensional Syntrix Tensorians, which differ from one another in their syntrometric structures, i.e., the syntrometry. In any case, an n -dimensional Syntrix space, when a system of corporation rules is defined analogously to the Corporator simplex of the Syntrix totality, is dominated by a Syntrix field, since each geometrically illustrated point of this space corresponds to a corporated Syntrix.

The determining components of such a Syntrix field within a general Syntrix Tensorium are thus the one-dimensional Tensorians that span the Syntrix space, the Syntrometry, and the system of corporation rules, the so-called *Corporator field*. In other words, to each excentric Corporator chain, i.e., each Conflexive Syntrix and each number of Enyphansyntrices

²³ChatGPT4.0: A creative translation for "Konflexivsyntrices" could be "Flexocorporative Structures." This term combines the ideas of flexibility (from "konflexiv") and cooperation or combination (from "syntrices") to convey the concept in English.

²⁴Konflexivsyntrix

²⁵ChatGPT4.0: Syntrix Weave-Matrices

corresponding to the Syntropode number, there corresponds a Syntrix field of a Syntrix Tensorium dimensioned with the Syntropode number as the argument range when any system of corporation rules is given. Since these Corporators do not belong to the Corporator simplex of totality but are Enyphan-like changes of the eccentric chain of the Conflexive Syntrix that generated the Syntrix Tensorium, the elements of the Syntrix Tensorium must in turn be Conflexive Syntrices, whose Syntropode numbers can reach at most the Syntropode number of the generating Conflexive Syntrix. For this reason, each element of a Syntrix Tensorium can generate a further Tensorium, which, however, is dimensionally enclosed in the original Tensorium because, according to the law of Syntropode numbers, the dimension number of the original Tensorium is never exceeded. All these Syntrix spaces and fields are defined over a totality. Apparently, these higher syntrometric structures are related in some way. These relations can either involve syntrometric structures over the same totality or those over different totalities. An investigation of such relations presupposes the creation of a syntrometric formalism that formally captures the higher syntrometric structures, which must inevitably lead to an extension of the concept of Enyphansyntrix to general functors of Syntrices.²⁶

4.5 Syntrix Functors.

For the formal description of Syntrix fields, i.e., higher syntrometric structures, in abstract Syntrix tensor fields, it becomes necessary, according to the foregoing, to extend the concept of Enyphansyntrix to the more general concept of the *Syntrix functor*. Since the Enyphansyntrix is always defined within a specific totality, transforming one element of this totality into another, it cannot act on any syntrometric structures, because these, due to their eccentric properties or their not necessarily belonging to the simplex corporation rules, are located above the totality and are only connected to it through their Syntropodes. Let $\tilde{c}|$ be any p -membered Konflexivsyntrix, which should form the syntrometric base

of the functor. Furthermore, let $1 \leq \varsigma \leq r - 1$ be given Corporators Γ_ς , which do not necessarily belong to the fundamental simplex and of which $q \leq r - 1$ can be eccentric. These $r - 1$ Corporators can incorporate r Syntrizes $\tilde{a}|_\varsigma$, such that a q -membered Konflexivsyntrix is formed. In general, the Syntrix $\tilde{c}|$ is incorporated as a base into this structure with the help of a Corporator C . The result is:

$$\tilde{A}|, ||, \tilde{c}|C \left(\tilde{a}|_\varsigma \Gamma_\varsigma \tilde{a}|_{\varsigma+1} \right)_1^{r-1}.$$

The operational rule that acts on the r Syntrizes $\tilde{a}|_\varsigma$ is thus:

$$\tilde{f}|, ||, \tilde{c}|C \left(0 \Gamma_\varsigma 0 \right)_1^{r-1}$$

and this inherently enyphane rule is referred to as a Syntrix functor of valence r . $\tilde{f}|$ acts on the sequence $\tilde{a}|_\varsigma$ in such a way that according to:

$$\tilde{f}|, \left(\tilde{a}|_\varsigma \right)_1^r, ||, \tilde{A}|$$

it arises^{27, 28}

From this representation, it follows directly that the Syntrix functor defined in this way should not only be understood as an extension of the discrete Enyphansyntrix, but also represents a higher form of the Synkolator concept,

²⁶ChatGPT4.0: In a playful Feynman-esque way, we can think of syntrometric structures as a wild dance of interconnected elements in a multi-dimensional universe. Imagine a space where each point represents a unique arrangement of Syntrices - a kind of building block in this system. These points form a vast landscape, shaped by the rules governing their interactions, and their relationships with one another create fascinating patterns.

As we move around in this space, we encounter new forms and structures that are generated by swapping out different components or changing the dimensions. It's like playing with a cosmic set of LEGO bricks, where higher-level structures emerge from the complex interplay of these fundamental pieces.

Studying these relationships and connections requires a new kind of mathematical language - a formalism that captures the essence of these higher syntrometric structures. It's an exciting journey of exploration and discovery, where we dive into a world of endless possibilities and intricate beauty, much like Feynman's own adventures in the realm of physics.

²⁷I have the impression that something is missing after "according to" - in the original manuscript there is a blank space left.

²⁸HS: I understand it in the sense of: "... that according to f — ... A — arises." as a closed statement.

indicating an extension of Syntrometry far beyond the Syntrix concept. Just as the Synkolator syncollates individual conceptual elements in the sense of the functor to higher Synchromes of a Syntrix, the Syntrix functor syncollates the Syntrizes of a totality to simple or higher syntrometric structures above this totality. Let $1 \leq \varsigma \leq r$ be the given sequence of $\tilde{a}|_{\varsigma}$ and let $\tilde{a}|_s$ with $1 \leq s \leq r$ denote a permutation; then in the general case of the asymmetric Syntrix functor (analogous to the asymmetric Synkolator):

$$\tilde{f}|, (\tilde{a}|_{\varsigma})_1^r, \bar{\bar{||}}, \tilde{A}| \text{ but } \tilde{f}|, (\tilde{a}|_s)_1^r, \bar{\bar{||}}, \tilde{A}|$$

with $\bar{\bar{||}} \neq \bar{\bar{||}}'$ of the Predicatrix. However, if, despite the permutation, $\bar{\bar{||}} \neq \bar{\bar{||}}'$ becomes an identical statement, then $\tilde{f}|$ is symmetric in analogy to the symmetric Synkolator. Finally, for the Syntrix functor of valence r , there is still the possibility of appearing hetero- or homometral. If all $\tilde{a}|_{\varsigma}$ are different from each other, then $\tilde{f}|$ is heterometral, but $\tilde{f}|$ becomes homometral if there are $1 < n \leq r$ identical Syntrizes in the sequence $\tilde{a}|_{\varsigma}$. If this is the case (n -fold homometral), then the *Functor valence* r is only a pseudo-valence, while the actual valence is only $r - n + 1 \leq r$. This also makes it clear how the Synkolator concept can emerge from that of the Syntrix functor. If $\tilde{f}|$ is constructed only from concentric Corporators of the Simplex, and if $\tilde{c}|$ also belongs to the totality, then $\tilde{A}|$ must also be a concentric Syntrix belonging to the totality, from which it follows that in this case, the Syntrix functor transitions into a discrete but r -fold iterated Enyphansyntrix. However, this requires that C also belongs to the Simplex, because then $\tilde{c}|C\tilde{a}|_1$

forms an element of the totality, which incorporates another element $\tilde{a}|_2$ of the totality with $\bar{\Gamma}_1$ of the Simplex, etc., resulting in $\tilde{A}|$ also belonging to the totality and $\tilde{f}|$ being an iterated Enyphansyntrix. The Syntrix functors within the totality can thus always be resolved into Enyphansyntrices. If, on the other hand, $\tilde{c}|$ is conflexive, i.e., p -membered,

and the $\bar{\Gamma}_{\varsigma}$ are no longer defined in the Corporator Simplex ($q - 1$ are excenters), then $\tilde{f}|$ can no longer be resolved into Enyphansyntrices. Now, $\tilde{f}|$ is a prescription that either generates a syntrometric structure from elements of the totality (if the $\tilde{a}|_{\varsigma}$ belong to it) or modulates syntrometric structures into higher forms, in case the $\tilde{a}|_{\varsigma}$ are such structures.

Thus, a Syntrix functor is always able to generate a large number of syntrometric structures, which is to be expected since it can be understood as an extension of both the Synkolator concept and the Enyphansyntrix concept. For example, if $N \geq r$ structures are given, $\tilde{f}|$ (if r is its valence) can generate $\binom{N}{r} \geq 1$ new structures in the sense of a Syncollation if $\tilde{f}|$ acts symmetrically and heterometral. On the other hand, if $\tilde{f}|$ is $k \leq r$ -fold asymmetric but heterometral, $k! \binom{N}{r} \binom{r}{k} \geq 1$ structures are created, and for $k = r$, finally $k! \binom{N}{r} = \frac{N!}{(N-r)!}$ new structures, etc. For $N < r$, $\tilde{f}|$ can only be homometral. This combinatorics again reveals the relationship to Synkolator combinatorics. The Syntropod numbers of the resulting structures depend not only on the stem but also on the corporator rules and the incorporating structures. The special case of the Syntrix functor of valence r described by

$$\tilde{f}|, (\tilde{a}|_{\varsigma})_1^r, \bar{\bar{||}}, \tilde{A}| \vee \tilde{f}| \equiv \tilde{c}|C(0 \bar{\Gamma}_{\varsigma} 0)_1^{r-1} \quad (18)$$

is the r -fold iterated discrete Enyphansyntrix, which is why Functor 18 is also referred to as a discrete Syntrix functor, allowing for a refinement of the Syntrix functor concept. In addition to discrete Enyphansyntrices, there are continuous forms, which leads to the heuristic conclusion that there must also be their extensions, namely continuous Syntrix functors.

In describing continuous Syntrix functors, it must be taken into account that the Enyphans, which enable the representation of continuous forms, only exist within a totality due to their definitions, and that the Syntropods of the structures are within this totality. However, this means nothing other than that Enyphans, which influence the structures, act from the totality through the Syntropods. So, if $\tilde{a}|_{\varsigma}$ is a p -membered Conflexive Syntrix, then $q \leq p$ Enyphans $\epsilon_{s\varsigma}$ with $1 \leq s \leq q$ can be combined into an *Enyphan complex* of operations $E_{\varsigma}(\epsilon_{s\varsigma})_1^q$ acting on $\tilde{a}|_{\varsigma}$, or in other words, on the elements of the totality that form $\tilde{a}|_{\varsigma}$. This action can occur $\binom{p}{q}$ times. Now, if there are $L \leq j \leq K$ Enyphan complexes E_j , they can be coupled with a discrete Syntrix functor $\tilde{f}|$. Then,

$$\widetilde{F}|, \overline{\parallel}, \widetilde{C} \left((0 \sqcap_i 0) \right]_1^{L-2} \left(E_j 0 \sqcap_j E_{j+1} 0 \right]_L^{K-1} \left((0 \sqcap_\varsigma 0) \right]_{K+1}^r \vee E_j \equiv E_j (\epsilon_{sj})_1^q \quad (18a)$$

becomes a $K - L$ -fold continuous Syntrix functor of valence r . In this way, the continuous counterpart to (18) is given, where each Enyphan complex can be combinatorially effective multiple times, so that $\widetilde{F}|$ generally turns out to be ambiguous. The continuous Syntrix functor also becomes a continuous Enyphansyntrix in the special case, which is defined within a totality. Finally, the functor (18) also appears as a special case of (18a), and (18a) in turn is a higher form of the Synkolator concept, like (18). Apparently, according to these explanations, the Syntrix functor appears as a generic term for the general Enyphansyntrix.²⁹

²⁹ChatGPT4.0 as Feynman: Alright, folks, let's try to wrap our heads around this thing called "Syntrix Functors." Now, I'm going to attempt to explain this concept in a way that even I can understand.

So, we're dealing with these funky things called "Syntrix Functors." They're like the big siblings of two other concepts - "Synkolators" and "Enyphansyntrix." If you think of them as some sort of operators, they can generate a whole bunch of new structures, which is really neat.

Imagine we have a set of structures, and we apply our Syntrix Functor to them. Depending on the properties of the functor, like symmetry and heterometry, we can end up with various amounts of new structures. It's sort of like a recipe book for creating new and exciting things.

Now, let's talk about continuous Syntrix Functors. The thing to remember is that the Enyphans that help represent continuous forms can only exist within a totality. They act on the structures in the totality through these little things called Syntropods. Imagine a bunch of Enyphans teaming up to form an "Enyphan Complex," and then they work together to influence the structures.

When we combine a discrete Syntrix Functor with Enyphan complexes, we end up with a continuous Syntrix Functor. It's like a more advanced version of its discrete cousin. It can have multiple effects on the structures and can be quite versatile.

In some special cases, the continuous Syntrix Functor can become a continuous Enyphansyntrix, which is defined within a totality. When you look at the big picture, the Syntrix Functor is sort of like an umbrella term for the general Enyphansyntrix.

So, there you have it! Syntrix Functors are like the cool operators that can create new structures and come in different flavors, like discrete and continuous. They're the big bosses in town when it comes to this whole Enyphansyntrix concept.

4.6 Transformations of Syntrix Fields.

The most general syntrometric structures, according to the previous discussion, are arbitrary Syntrix fields in any Syntrix spaces. These Syntrix fields can emerge from the action of general Syntrix functors, where each such field is determined by a Syntrix space and a system of corporators, the corporator field. The elements of this field don't necessarily belong to the simplex of totality, and it fills the space with arbitrary Syntrices. When a Syntrix functor acts on the generating Konflexivsyntrix³⁰ of the space, it leads to a deformation of this space concerning syntrometry, and thus to a transformation of the entire field spanned by it. In this way, the two fields are related through the Syntrix functor in a transformative connection. However, the Syntrix functor can also combine $r > 1$ fields into a single Syntrix field, namely, when the Syntrix functor has valence r and $r > 1$. Therefore, Syntrix functors can be divided into transformation groups and classify the Syntrix fields generated by these groups.

Quite obviously, three main classes can be distinguished: synthesizing, analyzing, and isogonal transformations. In the first case, the Syntrix functor combines $r > 1$ fields according to its valence; in the second case, it inversely resolves a field into r subfields; and in the third case, each Syntrix field is transformed into another, which is only possible for $r = 1$, and only in this case can a uniqueness of transformation exist. The mode of action of the Syntrix functors themselves is also ambiguous, and the functor can influence the syntrometry, i.e., the generating Konflexivsyntrix (konflexiv), or act tensorially concerning the Syntrix tensorium (in both cases, the Syntrix space is transformed, which is why these two modes of action should be called space-specific), or the Syntrix functor acts only on the corporator field, which corresponds to a field-specific transformation. Consequently, there are nine transformation classes of Syntrix fields since this ambiguity applies to each of the three main classes.

Denoting a transformation class as a_{ik} , where the first index represents the main class and the second represents the subclass, all transformation classes of Syntrix fields are normally contained in the matrix scheme $\hat{a} = (a_{ik})_3$, so that the elements of this scheme can now be analyzed. The properties of these transformation classes can be briefly outlined.

a_{11} : Synthesizing functors with a conflexive effect. These transformations appear regular under the condition of the same syntropod number.

a_{12} : Synthesizing functors with tensorial effect decrease the dimension number of the syntrix space, making this transformation class singular.

a_{13} : Synthesizing functors with field-specific effect do not change the syntrix space dimensionally and are therefore regular.

a_{21} : Analytical functors with conflexive effect can only work regularly under the condition that all split products have the same syntropod number, which must be identical to the one that characterized the original syntrix field.

a_{22} : Analytical functors with tensorial effect are always singular because the influence of this transformation class increases the dimension number of the syntrix space.

a_{23} : Analytical functors with field-specific effect are always

a_{31} : Isogonal functors with conflexive effect, as well as isogonal functors a_{32} and a_{33} with tensorial and field-specific effect can only work regularly due to their isogonality. From this, it becomes apparent that the index 3 containing edge of $\hat{a}$ only contains regular transformation classes because the influence of syntrix functors on these classes leaves the dimension number of the syntrix space unchanged without the need for additional conditions.³¹

³⁰ChagGPT4.0:Co-flexive Syntrix

³¹ChatGPT4.0 as Feynman: Alright, let's consider the most general syntrometric structures, which we'll call arbitrary Syntrix fields in any Syntrix space. These fields can emerge from the action of general Syntrix functors, with each field determined by a Syntrix space and a system of corporators called the "corporator field." The elements of the corporator field don't have to belong to the simplex of totality, and they fill the space with various Syntrices.

Now, if a Syntrix functor acts on the generating Co-flexive Syntrix of the space, the space undergoes a deformation with respect to syntrometry, which leads to a transformation of the entire field it spans. In this way, two fields are related through a transformative connection by the Syntrix functor. But, the Syntrix functor can also combine $r > 1$ fields into a single Syntrix field, particularly when the Syntrix functor has a valence r and $r > 1$.

So, we can categorize Syntrix functors into transformation groups and classify the Syntrix fields generated by these groups. It's clear that there are three main classes: synthesizing transformations, analyzing transformations, and isogonal transformations. In the first case, the Syntrix functor combines $r \geq 1$ fields according to its valence. In the second case, it inversely decomposes a field into r subfields. In the third case, each Syntrix field is transformed into another, which is only possible for $r = 1$, and only in this case can a unique transformation exist.

The action of Syntrix functors is also threefold. They can affect syntrometry (i.e., the generating Co-flexive Syntrix) in a co-flexive way, or act tensorially in relation to the Syntrix tensorium (in both cases resulting in a transformation of the Syntrix space, so we'll call these two modes of

4.7 Affinity Syndromes.

If there are $1 \leq i \leq N$ arbitrary syndromes $\tilde{a}|_i$ with the syndrome numbers $1 \leq \gamma_i \leq k_i$, and if there is also some system B such that all these syntrices are related to B in some correlative relationships, then there is a possibility that in the respective syndrome γ_i of the associated $\tilde{a}|_i$, there are a total of m_{γ_i} syncolations that actually correlate with B , but do not yet constitute the full occupancy of the syndrome, so that there are still syncolations that have no affinity to B . In total, there are $n = \sum_{i=1}^N \sum_{\gamma_i=1}^{k_i} m_{\gamma_i}$ syncolations in the syntrix system $\tilde{a}|_i$ that have some affinity to B with respect to correlation. Instead of investigating the interrelationship of the syntrix system with B , it is sufficient in this special case to remove the affine syncolations from the syntrix syndromes and to summarize them in a separate syndrome S of these affine syncolations with respect to B , the so-called *affinity syndrome*.

In this way, the correlation of a syntrix manifold with a system B is limited to the correlation of the affinity syndrome, which naturally comprises significantly fewer elements. This affinity syndrome, completely described by

$$S \equiv \left(\begin{array}{ccc} \tilde{a}|_i & N & k_i \\ m_{\gamma_i} & i = 1 & \gamma_i = 1 \end{array} \right) \quad (19)$$

therefore characterizes the correlation of the syntrix manifold with the system B and must be referred to as a pseudo-syndrome.

If, however, apodictic elements $\gamma_i = 0$ also show affinities to B , there is the possibility of expanding the affinity syndrome, which now also contains apodictic elements, into an *affinity syntrix* in the sense of a pseudo-syntrix. In such an affinity syntrix, there can generally be no uniquely defined synkolation, but there is fundamentally the possibility of giving an internal structure to both the affinity syntrices and the affinity syndromes, since elements of the same affinity degree with respect to B can always be combined into a subsyndrome, which in turn can be oriented according to a change in the affinity degree in analogy to the episyllogism of a synkolation process. In this case, an *oriented affinity syndrome* would be present, which implies the affinity syntrices. If there are $1 \leq \lambda \leq L$ precisely defined affinity levels, such that with increasing λ , the affinity with respect to B increases or decreases, and $m_{(\lambda)\gamma_i}$ denotes the elements affine to an affinity level λ , so that $m_{\gamma_i} = \sum_{\lambda=1}^L m_{(\lambda)\gamma_i}$, then in principle, the oriented affinity syndrome can be written as

$$S \equiv \left(\begin{array}{ccc} \tilde{a}|_i & N & K_i \\ m_{(\lambda)\gamma_i} & i = 1 & \gamma_i = 0 \end{array} \right)_{\lambda=a}^L \quad (19a)$$

because the affinity syntrices are implied by this.³²

action space-inherent). Alternatively, the Syntrix functor could act only on the corporator field, which corresponds to a field-inherent transformation. So, we have nine transformation classes of Syntrix fields, with this threefold ambiguity applying to each of the three main classes. If we use a_{ik} to denote such a transformation class, with the first index indicating the main class and the second index indicating the subclass, then all transformation classes of Syntrix fields are normally contained in the matrix schema $\hat{a} = (a_{ik})_3$, allowing us to analyze the elements of this schema. We can briefly outline the properties of these transformation classes.

³²ChatGPT4.0: 4.7 Affinity Patterns: Feynman Style

Suppose we have $1 \leq i \leq N$ arbitrary syndromes $\tilde{a}|_i$ with syndrome numbers $1 \leq \gamma_i \leq k_i$, and there's a system B such that all these syntrices are somehow correlated with B . Now, let's say in each syndrome γ_i of the corresponding $\tilde{a}|_i$, there are m_{γ_i} synkolation points that actually correlate with B , but they don't completely fill the syndrome, leaving some synkolation points with no affinity to B at all. So, in total, we have $n = \sum_{i=1}^N \sum_{\gamma_i=1}^{k_i} m_{\gamma_i}$ synkolation points in the syntrix system $\tilde{a}|_i$ that have some affinity to B regarding correlation.

Instead of examining the whole interaction between the syntrix system and B , we can just take out the affine synkolation points from the syntrix syndromes and combine them into a separate syndrome S related to B , which we'll call the "affinity pattern."

This way, we're limiting the correlation of a syntrix manifold with a system B to the correlation of the affinity pattern, which naturally consists of significantly fewer elements. The affinity pattern, completely described by

$$S \equiv \left(\begin{array}{ccc} \tilde{a}|_i & N & k_i \\ m_{\gamma_i} & i = 1 & \gamma_i = 1 \end{array} \right) \quad (19)$$

represents the correlation of the syntrix manifold with the system B and must be referred to as a pseudo-syndrome.

5 Metroplex Theory

5.1 The first-degree Metroplex as a Hypersyntrix

The description of general syntrix functors has shown that every synthetic syntrix functor of valence r is capable of synthesizing r syntrices, which can be arbitrarily multi-membered, into a new form of a higher syntrometric structure. The synkolation process occurs in a completely analogous manner. A synkolator of level m (which corresponds to the functor valence) synthesizes m synkolations of the next lower syndrome into a new synkolation. This synkolator property of the syntrix functor suggests a radical extension of the syntrix concept since, gradually, as the synkolation law of a syntrix induces syndrome occupations from conceptual elements, a synthetically acting syntrix functor that induces syndrome occupations of higher syntrometric structures from syntrix manifolds in the sense of a synkolation would be conceivable. Since the predicate connection of syntrices must fundamentally have universal quantor properties, the apodictic behavior of the syndrome occupation of such a hypersyntrix would be evident.

If there are $1 \leq i \leq N$ arbitrary syntrices $\tilde{a}|_i$ that either belong to a totality or are in this totality as konflexive forms with their syntropodes, then there is apparently a fundamental possibility of combining these N syntrices into a metaphorical complex $\tilde{\tilde{a}}| \equiv (\tilde{a}|_i)_N$ because all syntrix connections are universal quantifiers. These N syntrices of the hypermetaphor $\tilde{\tilde{a}}|$ can be defined over $q \leq N$ different subjective aspects, as there is no reason why all syntrices should belong to the same subjective aspect, because the synthesizing syntrix functor can connect syntrices from different subjective aspects, provided that the corresponding components of the functor are defined in all these aspects. Therefore, if $\tilde{\tilde{a}}|$ is given over $q \leq N$ subjective aspects of the relevant aspect system A , then these q aspects comprise a limited subsystem B , which is always implied by A . According to the investigations of syntrix functors, there is always the possibility of defining such a synthesizing syntrix functor of arbitrary valence r over the subsystem B .

If the valence is subjected only to the condition $r \leq N$, and this is to apply to all other functors F that act directly on the occupation of $\tilde{\tilde{a}}|$, then there is still the possibility, in complete analogy to the general complex syncholorator, which determines the syncholation course of a concentric complex syntrix, to construct a complex syncholorator $\underline{F}$ with the complex syncholation level $\underline{r}$ from synthesizing syntrix functors, where $\underline{r}$ indicates the valence progression along the complex $\underline{F}$ of syntrix functors. This general complex syncholorator $(\underline{F}, \underline{r})$ composed of syntrix functors can now act on the metaphorical complex $\tilde{\tilde{a}}|$ like in an ordinary syntrix, and syncholate the syndromes occupied with syntrices of a hypersyntrix. Such a hypersyntrix will hereinafter be referred to as a metroplex, specifically as a *Metroplex of the first degree*, because the syndrome 0 is a metaphorical complex whose elements are ordinary syntrices. The thus defined metroplex of the first degree can therefore be symbolized by

$$\left| \overset{1}{\underset{a}{\times}} \right| \equiv \left\langle \left(\underline{F}, \tilde{\tilde{a}}| \right)_{\underline{r}} \right\rangle \vee \tilde{\tilde{a}}| \equiv (\tilde{a}|_i)_N \quad (20)$$

if the general case of the homogeneously acting syncholation law is present. Since each metroplex of the first degree is built up of concentric and conflutive syntrices, the same laws must apply to the existence and uniqueness

However, if even apodictic elements $\gamma_i = 0$ show affinities to B , we can expand the affinity pattern, now containing apodictic elements, into an "affinity syntrix" as a kind of pseudo-syntrix. In such an affinity syntrix, there is generally no uniquely defined synkolator, but it is fundamentally possible to give both the affinity syntrices and the affinity patterns an internal structure. Elements with the same affinity level concerning B can always be combined into a subsyndrome, which can be oriented according to a change in the affinity level, similar to the episyllogism of a synkolation process. In this case, an "oriented affinity pattern" is present, which implies the affinity syntrices.

If there are $1 \leq \lambda \leq L$ precisely defined affinity levels such that with increasing λ , the affinity concerning B increases or decreases, and $m_{(\lambda)\gamma_i}$ denotes the elements affine to an affinity level λ , so that $m\gamma_i = \sum_{\lambda=1}^L m_{(\lambda)\gamma_i}$, then, in principle, the oriented affinity pattern can be written as

$$S \equiv \left(\begin{array}{ccc} \tilde{a}|_i & N & K_i \\ m_{(\lambda)\gamma_i} & \left| \begin{array}{c} i=1 \\ \vdots \\ i=N \end{array} \right. & \left| \begin{array}{c} \gamma_i=0 \\ \vdots \\ \gamma_i=L \end{array} \right. \end{array} \right)_{\lambda=a}^L \quad (19a)$$

since this implies the affinity syntrices.

So, in simpler terms, we've taken a look at affinity patterns in the context of syntrix systems, and how they can help simplify the analysis of correlations with other systems like B . By focusing on these affinity patterns, we can gain insights into the interactions between syntrices and the system B more efficiently. And if we take it a step further and consider oriented affinity patterns, we can explore even more detailed structures and relationships within the syntrices.

of the metroplex as for syntrices, because if the metroplex $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|$ were non-existent, then no syntrix could exist in $\widetilde{a}$, and there could be no syntrix functor, i.e., no syntrix could be constructed in the entire subsystem B, which would contradict the assumption. In addition, predicate conjunctions of syntrices are always universal quantifiers according to the quantifier theory, which in turn is evident for the existence of the metroplex if a syntrix system exists in B that fills out $\widetilde{a}$. In complete analogy to this, the uniqueness of the metroplex follows from that of the syntrix, and if the syntrix functors of the complex syncholorator act uniquely, then $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ must also be uniquely determined, because its determinants are unique.

The uniqueness of $\widetilde{a}$ and (F, r) is directly evident based on syntrix theory, so that the existence and uniqueness of the first-degree metroplex has been demonstrated in this way. Since the elements of each first-degree metroplex, i.e., the occupation of all syndromes including the metaphorical complex (i.e., the syntrix-like idea of this hypercategory) are arbitrary syntrices, metaphorical couplings and compositions of a *Metroplex corporator* must always be syntrix functors, regardless of whether this corporator acts as a concentrator or eccentric. The syncholorative coupling and composition, on the other hand, is only conceivable in the sense of a corporation of syntrix functors, because the metroplex syncholorators can only be such syntrix functors. However, in the metroplex corporation, the corporator must also be expressible in B. In the metroplex corporation

$$\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right| \left\{ \begin{array}{cc} K_s & C_s \\ K_m & C_m \end{array} \right\} \left| \begin{smallmatrix} 1 \\ b \end{smallmatrix} \right|, \overline{|B|}, \left| \begin{smallmatrix} 1 \\ c \end{smallmatrix} \right| \quad (20a)$$

both metroplexes $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ and $\left| \begin{smallmatrix} 1 \\ b \end{smallmatrix} \right|$ are syncholoratively and metaphorically incorporated, and this corporation is linked through the predicate $\overline{|B|}$ from B with $\left| \begin{smallmatrix} 1 \\ c \end{smallmatrix} \right|$, and this predicate conjunction must also be a universal quantifier according to the quantifier theory, because each metroplex is composed of a sequence of syntrices, and the predicate conjunctions of syntrices or syntrix functors are always universal quantifiers. In this way, it becomes apparent from relation 20a that, while preserving the universal quantifier character, the metroplex concept is the unique extension of the syntrix concept. Analogous to the syntrix corporators, because the syntrix functor is to be understood as an extension of the simple syncholorator, there are co- and contra-operative corporations. Moreover, since the first-degree metroplex can be understood as a hypersyntrix, it also has the possibility of decomposition and decoupling, and this means that the metroplex of any homogeneous nature (according to (20)) can be split into a sequence of pyramidal structures and a homogeneous fragment, where, as in syntrix theory, this homogeneous fragment can again be understood as the result of the corporation of another sequence of pyramidal structures. Consequently, in the first-degree metroplex, the pyramidal syncholoration process also has an elementary character, and each pyramidal metroplex must in turn be divisible into the basic types of pyramidal syncholoration.

As in the syntrix theory, since it is the same scheme, the corporators can act as concentrators (according to (20a)) or more generally as eccentrics:

$$\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|^{(1)} \left\{ \begin{array}{cc} K_s & C_s \\ K_m & C_m \end{array} \right\}^{(m)} \left| \begin{smallmatrix} 1 \\ b \end{smallmatrix} \right|, \overline{|B|}, \left| \begin{smallmatrix} 1 \\ c \end{smallmatrix} \right| \quad (20b)$$

where the syndrome 1 of $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ is corporated to the syndrome m of $\left| \begin{smallmatrix} 1 \\ b \end{smallmatrix} \right|$. In the case of eccentrics, multi-membered con-flexive metroplexes must be created, whose syntropodes are referred to as *Metroplex syntropodes* of the first degree because the eccentrically corporated metroplexes are of the first degree. In any case, the first-degree metroplexes and their con-flexive forms are syntrometric structures that are not defined in the underlying syntrix totality. However, a syntrix totality can always be constructed by varying the corporator simplex such that all metaphors $\widetilde{a}_i$ from $\widetilde{a}$ lie in this totality, i.e., $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ is associated with $\mu \geq N$ syntropodes in the relevant totality, where $\mu \geq N$ is only achieved (syntropode minimum of the metroplex) if $\widetilde{a}$ contains no con-flexive syntrices, but only concentric forms. In addition to the number of metroplex syntropodes of a con-flexive form, there is therefore also the number of *basic syntropodes* of a metroplex, which is determined by the number of those syntrix metaphors through which

the metroplex syntropodes, i.e., the corresponding metaphoric complexes, are built. The first-degree metroplex is thus fundamentally defined above that syntrix totality in which all its basic syntropodes are located, i.e., this totality must contain all syntrices (at least the syntropodes of the con-flexive forms) which, in the general case, build up the metaphoric complexes of the con-flexive forms.

Due to this analogy between first-degree metaphores and syntrices, the latter can be considered as metroplexes of degree 0, i.e.,

$$\widetilde{a}| \equiv \left| \begin{array}{c} 0 \\ \times \\ a \end{array} \right| \quad (21)$$

which can also be applied to all other concepts of syntrix theory. For example, the syntrix totality as such would be denoted by degree 0 or ($T0$), etc. The syntrix corporators are of degree 0 because their corporation components connect standalone conceptual elements, while the corporators of the first-degree metroplexes, as their corporation components corporated syntrices in the sense of syntrix functors, can be referred to as first-degree corporators. Similarly, the synkolator of a syntrix would be a syntrix functor of the 0th degree ($S0$), and the syntrix functor as a synkolator of a metroplex would be one of the first degree ($S1$) because the ($S1$) takes over the syncolation of the first-degree metroplexes.³³

5.2 First-degree Hypertotalities, Enyphanmetroplexes, and Metroplexfactors.

Since the first-degree Metrophor is the consistent conceptual extension of the Syntrix concept, and in principle nothing has changed in the syndrome structure or the synkolation process, the first-degree Metroplex could be considered a hypercategory, that is, a category of categories, whose idea is the metaphorical complex. This idea of the first-degree Metroplex consists of the N Syntrices, that is, the N simple categories, which are brought into mutual correlation by some Syntrix factor $S1$, in such a way that these correlations, ordered according to increasing conditionality, also result in an episyllogism. Elements of the same conditionality form the occupation of individual syndromes, so that with increasing syndrome number, as in the case of the Syntrix, the degree of conditionality increases, and the episyllogism runs in the direction of increasing syndrome number. In the direction of decreasing syndrome number, the degree of conditionality decreases, so that in this direction a corresponding prosyllogism increases, reaching its peak in the metaphorical complex, that is, the idea of the Metroplex. If the schemes of the Metroplex and the Syntrix are to be identical, then it must be required that the elements of $\widetilde{a}|$ are not conditioned by any other Syntrices, i.e., there must be no Syntrix factor, and also no corporator, that links any other Syntrices $\widetilde{b}|_k$ in such a way that the $\widetilde{a}|_i$ from $\widetilde{a}|$ somehow arise from these $\widetilde{b}|_k$.

If this were not the case, then this function law (F, r) could be supplemented and the sequence of syndrome numbers could be changed accordingly, resulting in a new metaphorical complex. However, since no aspect transformation was carried out, which would be possible in terms of the principle of aspect relativity, this behavior of the $\widetilde{a}|_i$ would be in contradiction with the definition of an idea as a system of elements without mutual conditionality. In this way,

³³ChatGPT4.0: 5.1 The First-Degree Metroplex as a Hypersyntrix: A Feynman-esque Perspective

Alright, let's dive into the concept of the first-degree Metroplex as a Hypersyntrix. Now, imagine a syntrix, which is a beautiful mathematical structure, like a delicate lattice. We can visualize this syntrix as a collection of interconnected elements, each playing a role in the overall structure.

Now, let's level up the complexity a bit and consider the first-degree Metroplex. A Metroplex is similar to a syntrix, but it's like a syntrix on steroids! It's a more complex and intricate structure, built using syntrices. To help you wrap your head around it, think of a Metroplex as a network of multiple syntrices, woven together to form a higher-order structure.

Just like in syntrix theory, the "glue" that holds this structure together is the corporators. In a Metroplex, the corporators can be of the first degree, and they have the unique ability to connect the syntrices in meaningful ways, taking into account their properties and relationships.

Now, let's draw a parallel between syntrices and first-degree Metroplexes. We can see syntrices as Metroplexes of degree 0. So, there is a natural progression from syntrices (degree 0) to first-degree Metroplexes (degree 1).

To give you a clearer picture, imagine a syntrix as a basic building block, like a single LEGO brick. The first-degree Metroplex, on the other hand, is like an entire LEGO structure, built using a combination of these bricks, each representing a syntrix. The corporators, then, are like the hands that put these LEGO bricks together to form the larger structure.

In conclusion, the first-degree Metroplex is a fascinating extension of the syntrix concept, which allows us to explore more complex mathematical structures and relationships. By understanding the relationship between syntrices and Metroplexes, we're taking a step further into the beautifully intricate world of mathematics. And as Richard Feynman would say, "Nature uses only the longest threads to weave her patterns so that each small piece of her fabric reveals the organization of the entire tapestry." (Richard P. Feynman (2015). "The Quotable Feynman", p.66, Princeton University Press)

the concept of apodicticity would be limited, because predicate combinations of Syntrices are universal quantifiers, so that all Syntrices are apodictic (including the synkolation of Metroplex syndromes), but the metaphorical complex must also satisfy the criterion of the idea as a system of independent elements. This additional requirement is insignificant in the case of a Syntrix Metrophor, because apodictic conceptual elements are already independent according to their definition, while the inherently apodictic Syntrices can fundamentally form a metaphorical concept due to this apodicticity, but they do not necessarily have to satisfy the criterion of the idea.

Hence, the general theorem for the first-degree Metroplex follows: The Syntrices of the metaphorical complex must not be conditioned by other Syntrices, otherwise this Metrophor cannot be considered as the idea of the Metroplex. This theorem initially demands fundamentally for the base syntropes of the Metroplex $\mu = N$, because con-flexive forms must be excluded in $\widetilde{a}$. Among the allowed concentric Syntrices, only those can be considered that cannot be generated by a Konzenter from other structures, i.e., only the pyramidal elementary forms are allowed in $\widetilde{a}$, as any other concentric Syntrix is the corporation of such elementary structures. However, these elementary structures $\widetilde{p}|_{(k)}$ with $1 \leq k \leq 4$ do not form any $T0$ allocations, but rather fill the four value reservoirs k in B independently of the respective corporate simplex, that is, the four-dimensional memory of all $T0$ in B on B , summarized this means

$$\widetilde{a}|_i \equiv \widetilde{p}|_{(k)i} \vee 1 \leq k \leq 4 \quad (22)$$

That is, the base syntropes of each first-degree Metroplex within B are not in any $T0$, but rather in the Syntrix memory of all possible $T0$ in B .

With the additional requirement (22), the schematic analogy between Syntrix and first-degree Metroplex (Hyper-syntrix) becomes complete, so that all syntrometric laws, in particular, those of concentric and eccentric corporations, can be meaningfully transferred to such Metroplexes. With (22), the concept of this Metroplex is clearly defined, so

that now an analysis in syntrometric form can be followed. Since the schema of $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ is formally analogous to that of $\widetilde{a}$, and the $S1$ in principle fulfill the same function as the $S0$, Metroplex classes can also be distinguished concerning the synkolation process, namely homogeneous and pyramidal concerning synkolator action, as well as homometral and heterometral or symmetric and asymmetric, in complete analogy to the classification of concentric Syntrices. Due to these analogies, any first-degree Metroplex can be split into a sequence of pyramidal structures and a homogeneous fragment, which in turn can be conceived as a corporation of pyramidal Metroplexes, because there are corresponding inverses for each synthesizing $S1$ of a Metroplex corporator. The pyramidal Metroplex thus also has a fundamental character and can be split into the four elementary forms like the pyramidal Syntrix if the concept of the null Metroplex is defined. This concept is only clearly comprehensible when it is required that the syndrome allocations of all syndromes of the Metroplex, including syndrome 0, consist of null Syntrices. With this null Metroplex, all first-degree

Metroplexes can be dissolved into corporate chains of elementary pyramidal complexes $\left| \begin{smallmatrix} 1 \\ p(k) \end{smallmatrix} \right|$, of which there must be $1 \leq k \leq 4$ according to the classification scheme. If C is any corporate chain, then for an arbitrary Metroplex $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ over B , the representation is

$$\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right| \left| \overline{b} \right|, C, \left| \begin{smallmatrix} 1 \\ p(k) \end{smallmatrix} \right| \vee 1 \leq k \leq 4 \quad (23)$$

which suggests the possibility of combining the Metroplexes of type k possible over B into a metroplectic value reservoir (in geometric metaphor linearly), and constructing a four-dimensional storage of elementary Metroplexes from the four value reservoirs, from which any hypertotalities can be derived as first-degree totalities ($T1$), depending on which metroplectic corporator simplex complements this storage to a Metroplex generative.

However, it must be demanded that the corporators of the simplex are always concentrers, so that the $T1$ is only occupied with concentrically corporated Metroplexes. According to (22) and (23), the base syntropes of all Metroplexes of $T1$ are in the Syntrix storage of the subsystem B , but not in any $T0$ that emerged from the storage, whereby the concept of the first-degree Metroplex appears directly as a radical but unambiguous extension of the Syntrix concept. The corporators connecting Metroplexes must obviously contain Syntrix factors of the type $S1$ in their elements, since all synkolations of a first-degree Metroplex are, by definition, Syntrices that can only be coupled to higher syntrometric structures through $S1$, similar to the synkolators of these Metroplexes, which must always be $S1$. Since whole

corporate chains can also be formed with the help of such corporators consisting of $S1$, and such a chain can corporately connect a system of $p \geq 1$ Metroplexes to a first-degree metroplectic trunk, a *Metroplex functor* is already given in discrete form, which is a second-degree Syntrix functor $S2$, itself a first-degree Metroplex and generates higher Metroplex structures over a $T1$, which stand with their Metroplex syntropodes in any areas of the $T1$, in contrast to the base syntropodes, which only lie in the storage of the $T0$. These higher Metroplex structures can be concentric or eccentric con-flexive Metroplexes, which are subject to the same classification concerning their syntropode number, syntropode length, and the metroplectic con-flexive fields as the corresponding con-flexive Syntrices over a $T0$. The Metroplex functor $S2$, thus defined, can now be extended to a continuous functor, since the existence of the null Metroplex also provides an *Enyphan Metroplex* in an inverse form, and thus an Enyphane, which can be corporated to the discrete Metroplex functor in complete analogy to the Syntrix functor. The con-flexive Metroplexes corporated by the discrete $S2$ cannot belong to $T1$, because only concentrers are allowed in the corporator simplex, but con-flexive forms are only possible through excenters; however, all these structures, as already shown, stand with their Metroplex syntropodes in the $T1$, provided that the generating corporator field only acts on elements of the $T1$. Since functors of the form $S2$ are also first-degree Metroplexes, regardless of whether they are discrete or continuous forms, these functors can act on any Metroplex structures induced by the corporator field over the $T1$ (the elements of this field do not belong to the simplex), and in this way span Metroplex spaces and Metroplex fields over the $T1$. For these metaphorical spaces and their syntrometry, the same formalism applies as for the corresponding Syntrix spaces and Syntrix fields, since the first-degree Metroplex, according to the preceding analysis, is nothing more than an extension of the Syntrix concept in such a way that it is the same formal structure schema.³⁴

5.3 Higher-degree Metroplexes.

The synkolation law of $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ is always an $S1$, which is a Syntrix functor and therefore a Syntrix, so the definition of $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right|$ must be written exactly in the form $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right| \equiv \langle \widetilde{F}|, \widetilde{a}|, r \rangle$. Similarly to how $\widetilde{a} \equiv \left| \begin{smallmatrix} 0 \\ a \end{smallmatrix} \right|$ was extended with the $S1$ to the first-degree Metroplex $\left| \begin{smallmatrix} 1 \\ a \end{smallmatrix} \right| \equiv \langle \widetilde{F}|, \widetilde{a}|, r \rangle$, these Metroplexes can be generalized to $\left| \begin{smallmatrix} 2 \\ a \end{smallmatrix} \right|$ with the metroplectic $S2$. As previously mentioned, the $S2$ functor is similar to the Syntrix functor $S1$, itself being a first-degree Metroplex, which can incorporate arbitrary Metroplexes into higher Metroplex structures and thus act as a first-degree Metroplex synkolation from a metaphorical complex, inducing syndromes of higher Metroplex structures. This leads to the definition of a second-degree Metroplex $\left| \begin{smallmatrix} 2 \\ a \end{smallmatrix} \right|$.

³⁴ChatGPT4.0: 5.2 First-Degree Hypertotalities, Enyphanmetroplexes, and Metroplexfactors - A Feynman-style Explanation

Imagine the Metroplex of the first degree as an extension of the Syntrix concept, where we keep the same basic structure but with a twist. It's like taking a category of categories, where the idea of a metaphorical complex is at the core. This complex contains N Syntrices (the basic building blocks), which are simple categories interconnected by a special Syntrix function, $S1$.

When we look at the Metroplex of the first degree, we need to ensure that its elements, the Syntrices, are not influenced by any other Syntrices. Otherwise, we would be in conflict with the definition of an idea as a system of independent elements. This leads to a general rule for the first-degree Metroplex: The Syntrices of the metaphorical complex must not be influenced by other Syntrices, or else it cannot be considered as the idea of the Metroplex.

Now, we can classify the Metroplexes of the first degree using similar criteria to the Syntrices classification, such as homogeneity and pyramidal, and symmetry or asymmetry. This leads to the idea that any first-degree Metroplex can be broken down into a series of pyramidal structures and a homogeneous fragment. This fragment can be considered a combination of pyramidal Metroplexes, since we can find inverse functions for each synthesizing $S1$.

The concept of a "Nullmetroplex" becomes useful here, which can be defined as a Metroplex with all its Syndromes filled with Nullsyntrices. With the Nullmetroplex, all first-degree Metroplexes can be decomposed into chains of elementary pyramidal complexes.

When it comes to connecting these Metroplexes, we need to use elements that are Syntrix functions of type $S1$. These functions can form whole chains of Metroplexes, and with the help of these, we can define a second-degree Syntrix function (Metroplexfactor) $S2$. $S2$ can create higher Metroplex structures which have their Metroplexsyntropods within the $T1$ domain, but not within the $T0$ storage.

These higher Metroplex structures can be either concentric or eccentric Konflexivmetroplexes, and they are subject to the same classification criteria as their corresponding Konflexivsyntrices. With the well-defined Nullmetroplex, $S2$ can be extended to a continuous function, creating an Enyphanmetroplex.

The Metroplexfactor $S2$ can act on any Metroplex structures induced over the $T1$, creating Metroplex spaces and fields. The formalism for these metaphorical spaces and their syntrometry is the same as for Syntrix spaces and fields, as the first-degree Metroplex is simply an extension of the Syntrix concept with the same formal structure schema.

This definition is done in the following way: Always, any metroplectic pyramidal elementary structures $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|_i$ with $1 \leq i \leq N$ can be taken from the storage of $T1$ and assembled into a metaphorical complex $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right| \equiv \left(\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|_i \right)_N$. The elements of this complex cannot condition each other because they come from the storage of all $T1$ and have their base syntropodes in the storage of all $T0$. For this reason, it can be regarded as a formal idea because no conditionality can occur between the elements. A system $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|$ of Metroplexfactors $S2$ in the sense of first-degree Enyphanmetroplexes can also act on this system as a general complex synkolation $\left| \begin{smallmatrix} 1 \\ \times \\ F \end{smallmatrix} \right|$ with the associated synkolation stages r (these are the valences of the Enyphanmetroplexes), which leads to syndrome occupations with higher first-degree Metroplex structures whose degree of conditionality increases with growing syndrome number in the sense of the syntrometric episyllogism.

This definition occurs in the following way: At any time, metroplectic pyramidal elementary structures $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|_i$ with $1 \leq i \leq N$ can be taken from the storage of $T1$ and combined into a metaphorical complex $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right| \equiv \left(\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|_i \right)_N$. The elements of this complex cannot condition each other because they come from the storage of all $T1$ and have their base syntropodes in the storage of all $T0$. For this reason, it can be understood as a formal idea because no conditionality can occur between the elements. A system $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|_i$ of Metroplexfactors $S2$ in the sense of first-degree Enyphanmetroplexes can also act on this system $\left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|$ as a general complex synkolation $\left| \begin{smallmatrix} 1 \\ \times \\ F \end{smallmatrix} \right|$ with the associated synkolation stages r (these are the valences of the Enyphanmetroplexes), leading to syndrome occupations with higher first-degree Metroplex structures, whose degree of conditionality increases with growing syndrome number in the sense of the syntrometric episyllogism.

The symbolism $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right| \equiv \left(\left| \begin{smallmatrix} 1 \\ \times \\ F \end{smallmatrix} \right|, \left| \begin{smallmatrix} 1 \\ \times \\ a \end{smallmatrix} \right|, r \right)$ indicates that the first-degree Metroplexes of the metaphor are synkolated by a complex first-degree Metroplexfactor, thus occupying the syndromes of a second-degree Metroplex. According to the general syntrometric formalism, however, there must also be concentrers and excenters for these $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$ as the occupations of all syndromes, including the metaphor, are first-degree Metroplex structures, which can always be coupled by functors of the $S2$ class, from which it directly follows that general corporators, as well as their inverses, must exist for $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$. If this is the case, it must also be possible to construct corporator chains of these Metroplexes, and this, in turn, implies a divisibility of these $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$ into elementary structures, as the $S2$ functors can only act homogeneously or pyramidally, and with each of these types of actions, there is again the possibility of homometrality, heterometrality, symmetry, and asymmetry.

In the previous discussion, it was shown that there are, analogous to null syntrices, first-degree null metroplexes, whose existence is directly derived from the existence of the null syntrix due to the Metroplex definition. This reasoning can now be continued, so that the existence of the first-degree null Metroplex directly justifies the existence of the second-degree null Metroplex. However, if the second-degree null Metroplex exists and corporator chains can be constructed simultaneously, then fundamentally, the possibility of splitting each $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$ into pyramidal elementary structures must be possible, which are no longer dependent on each other because all those used in these operations are uniquely determined by those of the syntrix theory. These syndromes are thus syndromes that are occupied with first-degree forms, i.e., these forms are associated with $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$, from which it immediately follows that $\left| \begin{smallmatrix} 2 \\ \times \\ a \end{smallmatrix} \right|$ can never be defined in $T1$, especially since the synkolations can already be conflexive forms, of which only the syntropods are in $T1$. In any case, with the four classes of second-degree pyramidal elementary structures, four value reservoirs can be filled, and a four-dimensional second-degree *Metroplex storage* can be spanned. A corporator simplex from concentrers can then be assigned to this storage, resulting in the generation of a second-degree *Metroplex totality* ($T2$). In this $T2$, second-degree Enyphanmetroplexes can be defined, and the eccentrically incorporated conflexive forms generate

syntrometric structures of the second degree, above $T2$, which stand with their Metroplex syntropods in the storage of this $T2$, as these syntropods belong to second-degree Metroplexes of $T2$. However, if such Enyphanmetroplexes exist, there must also be Metroplexfactors $S3$ that synkolate the occupations of $T2$, and in particular their storage elements, into arbitrary second-degree Metroplex structures, with the $S3$ being second-degree Metroplexes according to the functor definition.

In particular, when this second-degree synkolation law acts on an apodictic system of elements from the storage of $T2$ at any synkolation level, a third-degree Metroplex will arise, because all of its syndromes are occupied by second-degree Metroplex structures with equal conditionality. For these third-degree Metroplexes, the same laws must necessarily apply (e.g., to $T3$, $S3$, the Metroplex fields and spaces, etc.) as for the second-degree Metroplex, because $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^3$ is an immediate consequence of $T2$ and the possibility of incorporating its occupations. Since the functor $S3$ is always a $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^2$ and $S2$ is a $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^1$, as well as $S1$ being a $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^0 \equiv \tilde{F}$, Metroplexes of degree three to degree zero are conditioned in the sense of the sequence

$$\begin{aligned} \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^0 &\equiv \tilde{a} \equiv \langle F, \tilde{a}, \underline{r} \rangle \text{ justified T 0 and S 1} \\ \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^1 &\equiv \langle \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^0, \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^0, \underline{r} \rangle \text{ justified T 1 and S 2} \equiv \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^1 \\ \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^2 &\equiv \langle \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^1, \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^1, \underline{r} \rangle \text{ justified T 2 and S 3} \equiv \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^2 \\ \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^3 &\equiv \langle \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^2, \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^2, \underline{r} \rangle \text{ justified T 3 and S 4} \equiv \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^3 \end{aligned}$$

This sequence can now be continued recursively according to the reasoning of complete induction, so that a Metroplex of any degree $n \geq 0$ is finally created, which, according to the complete induction conclusion, means that a Metroplex of degree $n + 1$ also exists.

No matter how the functors Sk of Metroplex degree $k - 1$ may be constituted, which make up the synkolation law of a pyramidal Metroplex of degree k , it can only be homometral or heterometral with symmetric or asymmetric properties in pyramidal synkolation. However, this means that for any $k > 2$, there are always only four pyramidal elementary forms, i.e., each Tk is spanned by a four-dimensional storage. Finally, the corporators as functors $S(k + 1)$

of degree k are definable by the $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^k$, so that for any k , a corporator simplex of concentrators together with the four-dimensional storage provides the generative of the Tk .

The allocations of the resulting Tk (including the null metroplex of degree k , namely, belongs to it), can be exposed to arbitrary excenters and eccentric functors $S(k + 1)$, leading to multi-membered con-flexive forms above the Tk . These con-flexive forms, in turn, condition metroplex fields and metroplex spaces of degree k , which are subject to a deformed metric of the same degree. In the case $k = n$, the Tn is with its four-dimensional storage of

pyramidal elementary forms $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^n_{(p)}$ (it is $1 \leq p \leq 4$ the digit of the respective value stock) in the $T(n - 1)$. A corporation of the elements of Tn is possible through functors $S(n + 1)$ of degree n and arbitrary valence. Such a

functor can finally be formed as a complex synkolator $S(n + 1)$, corresponds to $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^n$ of the synkolation level $\underline{r}$ and

given synkolation course, which synkolates a first syndrome from the $1 \leq j \leq N_n$ storage elements $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^n_{(p)}$ of the Tn and so on. These pyramidal storage elements can then be summarized to a metaphoric complex of degree n , namely

$\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^{\tilde{n}} \equiv \left(\left| \begin{smallmatrix} \times \\ a_j \end{smallmatrix} \right|^n_{(p)} \right)_{N_n}$, on which the synkolator $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^n$ of the synkolation level $\underline{r}$ has a complex effect. In this way, a Metroplex of degree $n + 1$ has been created, which can be formally uniquely described in the version

$$\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^{n+1} \equiv \langle \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|^n, \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^{\tilde{n}} \rangle \vee \langle \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|^{\tilde{n}}, \left| \begin{smallmatrix} \times \\ a_j \end{smallmatrix} \right|^n_{(p)} \rangle_{N_n} \vee 1 \leq p \leq 4 \vee n \geq 0 \quad (24)$$

In this Metroplex, sub-metroplexes of degrees $k < n + 1$ from the T_k are obviously associated with an overarching structure for $0 \leq k \leq n$, which is why the metroplexes of degree $n > 0$ can also be referred to as associated syntrometric forms. From the scheme (24), the inductive nature of the reasoning to higher degrees becomes immediately apparent. Since the Metroplex is defined not only for $n = 1$, as well as $n = 2$ and $n = 3$, but also for $n > 3$ and $n + 1$,

this definition must be correct for all integer $n < \infty$. It is therefore sufficient to examine the forms $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right|$ in and above a Tn .

Since there are corporators that build up the simplex of Tn as a concentrator, or act as arbitrary excenters, and since these corporators connect the elements of Tn , the corporation members of such a corporator $\left\{ \begin{smallmatrix} K_s & C_s \\ K_m & C_m \end{smallmatrix} \right\}$ of degree n must consist of coupling and composition components of degree $n - 1$, because they connect the syndrome occupations of the metroplexes of degree n , but these synkolations according to (24) are of degree $n - 1$. If two metroplexes from Tn are corporated and this corporation is linked by a statement with a third metroplex of the same degree, such as in $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right| \left\{ \right\} \left| \begin{smallmatrix} n \\ b \end{smallmatrix} \right| \parallel \left| \begin{smallmatrix} n \\ c \end{smallmatrix} \right|$, it must be investigated in which aspect system or aspect complex the predicate lies. In this context, it should be considered that all base syntropodes for $n > 0$ are in a $T0$, and that these different $T0$ can be defined over different aspect systems. Since all operations for $n = 0$ are also possible for $n > 0$, metaphoric circular conclusions can also be carried out for $n > 0$, but this has the consequence that all predicate links of metroplexes are universal quantifiers, as for $n = 0$.

Due to this character of predicate links, the functors that enable the association of $0 < k \leq n$ syntrometric forms can simultaneously exist in several aspect systems, which immediately implies that the $T0$ of all base syntropodes can actually lie in different aspect systems. In $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right| \left\{ \right\} \left| \begin{smallmatrix} n \\ b \end{smallmatrix} \right| \parallel \left| \begin{smallmatrix} n \\ c \end{smallmatrix} \right|$, the predicate must therefore lie in an aspect

complex composed of those aspect systems over which the $T0$ of all base syntropodes of $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right| \left\{ \right\} \left| \begin{smallmatrix} n \\ b \end{smallmatrix} \right|$ and $\left| \begin{smallmatrix} n \\ c \end{smallmatrix} \right|$ lie. In each Tn , there are concentric and eccentric corporators, i.e., corporator fields of degree n which do not belong to the corporator simplex, and whose corporation members, as already mentioned, are of degree $n - 1$. From such corporators, however, corporator chains can always be developed, which corporately incorporate a number of metroplexes of Tn corresponding to their valence, either concentrically or eccentrically. Since a metroplex of Tn can always be used as a metroplex base, to which the corporator chain can be coupled, in this way a discrete metroplex functor $S(n + 1)$ as a metroplex of degree n is already defined. On the other hand, Tn also contains the null metroplex, which allows for an enyphan metroplex, and thus a continuous metroplex functor $S(n + 1)$ with its inverse. With the help of these functors, multi-membered conflexive forms can be corporated over the Tn , whose metroplex syntropodes of degree n are in the Tn , while those of the next higher association $n + 1$ are in the memory of the Tn (here the functor is transformed according to $S(n + 1) \equiv \left| \begin{smallmatrix} n \\ F \end{smallmatrix} \right|$ into a synkolator).

The multi-membered conflexive forms over the Tn , which have been generated by the metroplex functor when it acts as a synthesizing functor, form higher syntrometric structures of degree n , namely corresponding metroplex fields and metroplex spaces, which are characterized by a syntrometry of degree n . If, on the other hand, this metroplex functor acts as a complex synkolator (whose respective synkolation level is given by the functor valence) only on the storage elements of the Tn , then according to (24), the element of a $T(n + 1)$ is already given, for which the same laws apply according to the recursion conclusion as for Tn . It is essential that the elements of a metroplex totality with their syntropodes are fundamentally located in the four-dimensional memory of a metroplex totality whose degree is one level lower.

Through these *associative metroplexes*, there is a syndromatic interweaving of totalities for this reason. Thus, the syndrome occupations of the elements of a Tn or syntrometric structures defined above it consist of the elements of

the $T(n - 1)$ or their syntrometric structures of degree $n - 1$ and so on. This means that every $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right|$ has both a gradual and a syndromatic architectonics. The gradual tectonics is the progression of the internal building structure in the direction of the advancing metroplex degree. In analogy to the synkolation course in the direction of any syllogism, this gradual structural change could be referred to as the *gradual course* of the *tectonics*, because there are always

$0 \leq k \leq n$ structural zones in $\left[\begin{smallmatrix} n \\ a \end{smallmatrix} \right]$ each with the same tectonic degree k , but differing from each other through the degree of their association. Each of these structural zones k must also have a *syndromatic tectonics* orthogonal to the gradual tectonics, which can be identified with the synkolation course of the syndrome occupations in the relevant gradual structural zone of the metroplex. This course of syndromatic tectonics, referred to as *syndromatics*, is thus the direct equivalent to the synkolation course of a syntrix.

Every associative metroplex $n > 0$ thus has a dual tectonics, which is due to the associative character of its structure. Another characteristic of this character is the existence of the basic syntropodes in the $T0$ (possibly in different aspect systems of the aspect complex), so that every metroplex can be considered as a higher syntrometric structure above a $T0$, reducing the entire metroplex theory to the syntrometric elements. The existence of these basic syntropodes, i.e., the possibility of reduction to the $T0$, is a direct consequence of the existence of the gradual tectonics. Another characteristic of the associative metroplex $n > 0$ is the fact that each gradual structural zone k consists of synkolation, which lie in or above a Tk . These synkolation are associated in the respective structural zones in the form of the corresponding syndromatic tectonics, which in turn justifies the term associative metroplex.³⁵

5.4 Syntrocline Metroplex Bridges

In addition to the associative form of the Metroplex with its dual tectonics, another structural form is fundamentally possible. If $\tilde{a}$ is any element of a $T0$, with its syndrome conclusion at $\gamma \leq \infty$, then from the $1 \leq \gamma \leq x$ syndromes, $l \leq \gamma \leq k$ with $l \geq 0$ and $k \leq x$ can be used to describe new metaphors. However, these syndrome occupancies only provide pseudometaphors, because in relation to the subjective aspect over which the metaphor of $\tilde{a}$ is apodictic, these syndrome occupancies are not apodictic due to their conditionality.

Nevertheless, for each of the $k - l + 1$ syndromes γ , a system of $1 \leq j \leq \lambda_\gamma$ transforming functors $\phi_{j\gamma}$ can be found, which functorially summarize the occupancies of γ in λ_γ -fold manner, so that $k - l + 1$ systems of apodictic elements $\tilde{\alpha}_\gamma$ are created according to the principle of aspect relativity.

The functor valence of $\phi_{j\gamma}$ ³⁶ must never exceed the number of syndrome occupancies γ ³⁷ unless $\phi_{j\gamma}$ is also allowed to be homometral. Each of these systems, in turn, consists of λ_γ ³⁸ - metaphors $\tilde{\alpha}_\gamma$, so that a total of $\sum_{\gamma=1}^k \lambda_\gamma$ metaphors $\tilde{\alpha}_\gamma$ have emerged from the $k - l + 1$ syndromes of $\tilde{a}$ through the action of the transforming functors $\phi_{j\gamma}$.

This system of transforming functors $\phi_{j\gamma}$, defined according to the principle of aspect relativity, will be referred to as a *syntrocline continuation* Φ , while the apodictic systems $\tilde{\alpha}_{j\gamma}$ are syntrocline induced metaphors through the continuation Φ , originating from the $k - l + 1$ selected syndromes of $\tilde{a}$. For each element $\phi_{j\gamma}$ from Φ , a complex syncolator $(\underline{f}_{j\gamma}, \underline{m}_{j\gamma})$ can now be coordinated, which, with the associated metaphor (*syntrocline induced*), forms a syntrix, namely $\tilde{\alpha}|_{j\gamma} \equiv (\underline{f}_{j\gamma}, \tilde{\alpha}_{j\gamma}, \underline{m}_{j\gamma})$. Thus, the syntrocline continuation, combined with corresponding complex syncolators, has syntrocline induced a system of $1 \leq j \leq \lambda_\gamma$ syntrices $\tilde{\alpha}|_{j\gamma}$ over each of the $k - l + 1$ syndromes

³⁵ChatGPT4.0: 5.3. Higher-Degree Metroplexes - A Feynman-style explanation

Alright, let's talk about higher-degree metroplexes. You know, they have a pretty interesting structure. So, imagine we've got these associative metroplexes where $n > 0$. It turns out that they have a dual tectonics due to their associative nature. It's kind of like having two sides of the same coin, you see?

Now, there's another characteristic of these associative metroplexes that's worth mentioning. You see, there are these basic syntropodes in the $T0$ (sometimes in different aspect systems of the aspect complex), and every metroplex can actually be thought of as a higher syntrometric structure above a $T0$. And when you think about it this way, we can really simplify the entire metroplex theory to just the syntrometric elements. It's like taking a complex idea and breaking it down into its simplest form.

But hold on, there's more! The existence of these basic syntropodes, and the fact that we can reduce things to the $T0$, comes from the existence of what we call gradual tectonics. It's like a ladder, where each step is a direct consequence of the one below it.

Now, let me tell you another fascinating thing about these associative metroplexes with $n > 0$. In each gradual structural zone k , there are these synkolation that lie in or above a Tk . These synkolation are associated with one another in their respective structural zones, and it's because of the corresponding syndromatic tectonics. It's like a dance, where each part is moving in harmony with the others. And this is exactly why we call them associative metroplexes.

So, to sum it up, higher-degree metroplexes have this really cool dual tectonics structure, and we can reduce the whole metroplex theory to syntrometric elements thanks to these basic syntropodes. Plus, they've got synkolation that lie in or above a Tk , making them truly associative. And that, my friends, is what higher-degree metroplexes are all about!

³⁶HS: that is, the functor summarizes ALL or a combinatorics of syndrome occupancies.

³⁷: incorrect index - Gamma is the syndrome number, not the number of occupancies! Suggestion: n_γ (does not appear elsewhere, though)

³⁸in original: γ_γ HS: incorrect label - symbol expresses the number of different occupancy possibilities. λ_γ !

$l \leq \gamma \leq k$ of $\widetilde{a}$.

This process can be passed on, as there is the possibility to define a functor $S1$ in the sense of a complex syncolator $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|_\gamma$ of valency $r_\gamma \leq \lambda_\gamma$ for each of these complexes consisting of λ_γ syntrices over the $k - l + 1$ syndromes, and to consider the syntrix complex γ as a metaphoric complex, so that $k - l + 1$ first-degree Metroplexes

$$\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma \equiv \langle \widetilde{F}|_\gamma, \widetilde{\alpha}|_\gamma, r_\gamma \rangle$$

with $\widetilde{\alpha}|_\gamma \equiv (\widetilde{\alpha}|_{j\gamma})_{\lambda_\gamma}$ are created.

If this is to be the case, it must be demanded of Φ that the $\widetilde{\alpha}_{j\gamma}$ form elementary pyramidal shapes together with $(\underline{f}_{j\gamma}, \underline{m}_{j\gamma})$ from the storage of a $T0$, because otherwise the $\widetilde{\alpha}|_{j\gamma}$ cannot form metaphorical complexes. If this

requirement eventually goes so far that the $k - l + 1$ structures $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma$ are in the storage of a $T1$, then there is always the possibility to define a metroplex functor $S2$ as a complex synkolator $\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|$ of valence p and to summarize the $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma$ into a metaphorical complex $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma \equiv \left(\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma \right)_{k-l+1}$, so that now $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma \equiv \left(\left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right| \left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma \right)_p$ arises. Here, the symbol $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^2$ indicates that this second-degree metroplex has been induced with the help of a syntrocline continuation from the syndromes of $\widetilde{a}$. That is, the syndrome system $l \leq \gamma \leq k$ from $\widetilde{a} \equiv \left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^0$ was syntroclinically continued to $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^2$. In symbols, this process of syntrocline continuation of an element of the $T0$ into that of a $T2$ can be expressed by

$$\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^2 \equiv \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|_\gamma^0 \left[\prod_{j=1}^{\lambda_\gamma} \phi_{j\gamma}(\underline{f}_{j\gamma}, \underline{m}_{j\gamma}) \widetilde{f}|_{j\gamma} r_\gamma \left| \begin{smallmatrix} \times \\ F \end{smallmatrix} \right|_\gamma \right]_{\gamma=1}^k.$$

The metroplex thus created of the second degree is referred to as a simple syntrocline metroplex of the *continuation level* 2, because the gradual tectonic increases from value 0 to value 2. Without further ado, we can abstract from $n = 0$ and extend the procedure of syntrocline continuation to $n > 0$. If $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|_\gamma^n$ is concentric, a functor S_n in the form $\left| \begin{smallmatrix} \times \\ \phi \end{smallmatrix} \right|_{j\gamma}^n$ can always be found as an element of a syntrocline continuation and an associated synkolation law $\left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_{j\gamma}, \underline{m}_{j\gamma} \right)$, thereby inducing a system of λ_γ new metroplexes syntroclinically over each syndrome γ of $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|_\gamma^n$. All these metroplexes are of degree n , so that in turn $k - l + 1$ further synkolation laws Synkolation laws $\left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_\gamma, r_\gamma \right)$ are possible, which synkolate a metroplex of degree $n + 1$ over each syndrome γ between l and k . However, this means that a metroplex functor $S(n + 2)$ also exists as a complex synkolator $\left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_\gamma, \underline{p} \right)$, which associates these $k - l + 1$ metroplexes of degree $n + 1$ to the *syntrocline metroplex* $\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^{n+2}$.

For this general syntrocline metroplex of the continuation level 2, the following applies

$$\left| \begin{smallmatrix} \times \\ \alpha \end{smallmatrix} \right|_\gamma^{n+2} \equiv \left[\prod_{j=1}^{\lambda_\gamma} \left(\left| \begin{smallmatrix} \times \\ \phi \end{smallmatrix} \right|_{j\gamma}^n \right) \left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_{j\gamma}, \underline{m}_{j\gamma} \right) \left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_\gamma, r_\gamma \right) \left(\left| \begin{smallmatrix} \times \\ f \end{smallmatrix} \right|_\gamma, \underline{p} \right) \right]_{\gamma=1}^k \left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|_\gamma^n \quad (25)$$

In this definition of the general syntrocline metroplex of the continuation level 2, $\left| \begin{smallmatrix} \times \\ a \end{smallmatrix} \right|_\gamma^n$ is referred to as the *syntrocline root* and the selected $k - l + 1$ syndromes γ as the *syntrocline approach*. The inner structure of this syntrocline metroplex is essentially determined by the behavior of the λ_γ . If all $\lambda_\gamma > 1$, then it comes to the normal construction with the continuation level 2. If, on the other hand, $\lambda_\gamma = 1$ for all γ , then there are only $k - l + 1$ syntrocline induced complexes of degree $n - 1$, on which an S_n - synkolator acts, so that the same number $k - l + 1$ of metroplexes of degree n arises, which, as a metroplex, associate themselves as a syntrocline metroplex of degree $n + 1$.

So, if all $\lambda_\gamma = 1$, we get to the continuation stage 1. Finally, there is the possibility that some $\lambda_\gamma = 1$, but the remaining $\lambda_\gamma > 1$. In this way, the syntrocline metroplex becomes ambiguous, as the syndromes with $\lambda_\gamma = 1$ as syntrocline approaches can associate with the metroplexes of the same degree n in many ways but $\lambda_\gamma > 1$. So, unique syntrocline metroplexes of the continuation stage 2 only exist if all $\lambda_\gamma > 1$ and those of the continuation stage 1, if all $\lambda_\gamma = 1$. In all other cases, there is ambiguity. Any higher continuation stage $N > 2$ can only arise from the syntrocline

root $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right\rangle$ if the continuation process is iterated. That is, in this iteration, a syntrocline chain of simple continuations is created in such a way that the syntrocline metroplex of the preceding link is used as the root of the next continuation, etc. If the symbol $\int_n^{n+N}$ contains the so-called "syntrocline chain coupling" of all information about the syntrocline approaches and syntrocline continuations of the simple chain members, then the general syntrocline metroplex of any continuation stage $N > 2$ is symbolically represented by

$$\left| \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right\rangle \equiv \int_n^{n+N} \left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right\rangle \left[\prod_{j(n)=1}^{\lambda(n)\gamma} \left(\left| \begin{smallmatrix} n \\ \phi(n) \end{smallmatrix} \right\rangle j(n)\gamma \right) \left(\left| \begin{smallmatrix} n \\ f(n) \end{smallmatrix} \right\rangle j(n)\gamma, \underline{m} j(n)\gamma \right) \left(\left| \begin{smallmatrix} n \\ f(n) \end{smallmatrix} \right\rangle \gamma, \underline{r} \gamma \right) \left(\left| \begin{smallmatrix} n \\ f(n) \end{smallmatrix} \right\rangle \underline{p} \right) \right]_{\gamma(n)=1}^{k(n)}.^{39} \quad (25a)$$

Therefore, as many simple links are put together within the chain coupling until the continuation level $N > 2$ is reached. Whether N is even or odd depends on how often the continuation level 1 appears in the chain, provided that all chain links are unambiguous. If ambiguous chain links occur, all ambiguities in the syntrocline metroplex of the

continuation level N add up.⁴⁰ The root $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right\rangle$ of such a form belongs to a Tn , while the last continuation element lies in a $T(n+N)$. All intervening members of the *syntrocline chain* lie in the totalities from $T(n+1)$ to $T(n+N-1)$, so that $\left| \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right\rangle$ syntrometrically bridges all these metroplex totalities from Tn to $T(n+N)$, that is a total of $N-1$, from the root $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right\rangle$. Therefore, the syntrocline chain, or the general syntrocline metroplex of the continuation stage N , penetrates all totalities lying between n and $n+N$.

Because of their ability to bridge metroplex totalities⁴¹ of varying degrees, it seems fitting to adopt the term *syntrocline metroplex* only for bridge links with the maximum extension level of 2, and to label the more general forms (25a) for $N > 2$ as *syntrocline metroplex bridges*. In contrast to the associative metroplexes, the corporate

³⁹HS: "(n)" should not be in subscript (or everywhere). Gamma(n) = 1

⁴⁰LAA: I think that this is beginning to look similar to Richard Feynman's "sum over all paths" approach to quantum mechanics. What do you think? ChatGPT4.0: It's an interesting observation. While the text you provided is focused on a mathematical structure called "Metroplex" and its continuation stages, it is not directly related to quantum mechanics or Feynman's path integral formulation. However, it does involve the idea of summing over various possible chains or paths, which could be reminiscent of Feynman's "sum over all paths" approach.

In Feynman's path integral formulation of quantum mechanics, the probability amplitude for a quantum system to evolve from one state to another is obtained by summing over all possible paths that the system could take. Each path contributes a complex number to the sum, and the overall probability amplitude is obtained by adding up these contributions.

The concept of summing over paths or chains in the text you provided is more related to the mathematical structure of the Metroplex and how different continuation stages are formed. While the idea of summing over different paths or chains appears in both contexts, the nature of the underlying systems, their interpretation, and the mathematics involved are quite different.

So, while there is a superficial similarity in the idea of summing over various paths or chains, it is important to note that the context and the underlying mathematics are not directly related to Feynman's path integral formulation of quantum mechanics.

LAA: FYI, I have read ahead and the Metroplex figures into the structure of all the quantum mechanical particles. ChatGPT4.0: That's certainly fascinating! The use of sophisticated mathematical structures like the "Metroplex" can indeed be used to help formulate complex physical theories, including quantum mechanics.

While the specific mathematical structure described in your text isn't standard in the quantum mechanics I'm familiar with (up until my knowledge cutoff in 2021), mathematical ideas like these can often provide alternative formulations or perspectives on physical theories. This could potentially lead to new insights or approaches in the field.

That said, without more information about how the Metroplex is used in the quantum mechanical context you're reading about, it's hard for me to provide more specific commentary. If this Metroplex structure is part of a novel approach to quantum mechanics developed after my training data, I'd be unable to provide detailed insights on it.

Still, the connection you've noticed to Feynman's path integral approach is intriguing. In many ways, physics is all about exploring different mathematical structures and seeing how they can be applied to describe the physical world. Even if the precise mathematical machinery is different, many of the conceptual ideas can have interesting parallels. With regard to the totalities of metroplexes, these syntrocline metroplexes have a special meaning.

⁴¹ChatGPT4.0: "Metroplextotalitäten" has been translated as "metroplex totalities," assuming 'totalities' refers to a comprehensive encompassing set or entirety of metroplexes at a certain degree level. If 'totalities' instead has a more specific or nuanced meaning within the context of this subject, a different translation may be more appropriate.

laws⁴² of syntrocline forms are no longer unambiguously determined. This is because each syntrocline chain link exists within a total and varies in terms of metroplex degrees, meaning every link can independently form a corporate law with another metroplex of the same degree.

The clarity of a corporation of syntrocline bridges emerges from the following examination. If $\begin{array}{c} m+p \\ \diagdown \diagup \\ a \end{array}$ and $\begin{array}{c} n+q \\ \diagdown \diagup \\ b \end{array}$ are two syntrocline forms, whose roots have the grades m and n , and the sequences of numbers $0 \leq k \leq p$ and $0 \leq l \leq q$ indicate the current bridge elements, then a corporation is only possible between the elements for which $m+k = n+l$, so $k-l = n-m$. The number of these capable of mutual corporation members of two syntrocline metroplex bridges is then the clarity of the syntrocline corporation. Also the action of metroplex functors is ambiguous, because such a functor is always tied to the metroplex grade. However, it is possible that a synthesizing functor of higher valence in a variety of syntrocline bridges, those members of the same and its properties adequate grade synkoliert, where the number of so associated syntrocline bridges is conditioned by the functor valence. When corporation of syntrocline bridges it can also lead to the formation of conflexive forms, because the corporation of the bridge members, which meet the corporation condition (equal metroplex grade), can be both concentric and eccentric, so that a conflexive field arises. If it finally deals with a corporating functor in the sense of a corporator chain that also contains excenters, then conflexive forms of syntrocline bridges arise, whose syntropod number depends on the number of excenters effective in the chain.

In analogy to the tectonics of associative Metroplexes, there must also be a *syntrocline tectonics*. Regarding this tectonics, there is only one possibility which relates to the particular choice of the syntrocline inducing syndromes. If L is any link of the syntrocline chain in the interval $n \leq L \leq n+N$, where there are $0 \leq \gamma_L \leq x_L$ syndromes, of which $k_L - l_L + 1$ are selected for syntrocline induction of the next higher stage, the syntrocline tectonics of the link L depends on the position of these inducing syndromes. The syntrocline tectonics is, for example, metaphorically centered if the syndromes $0 \leq \gamma_L \leq k_L$ induce. If $k_L = x_L$, then it's total, for $0 < l_L \leq \gamma_L \leq x_L$ peripheral, for $l_L \leq \gamma_L \leq k_L$ with $l_L > 0$ and $k_L < x_L$ is concentric coherent, possibly concentric discrete, etc. Each chain link can have a different tectonics, so the statement of the overall tectonics of a Metroplex bridge can only consist of a sequence of tectonic specifications of all bridge members. Syntrocline Metroplex totalities cannot be defined because each syntrocline continuation lies in a different Metroplex totality. In any case, the syntrocline Metroplex bridges are higher syntrometric structures above that Metroplex totality in which the syntrocline root lies. The definition of the syntrocline Metroplex bridge allows that any conflexive forms can be used as syntrocline roots, but even in this case, the Metroplex bridge is a syntrometric structure above the respective totality, because the conflexive root stands with its syntropods in it.⁴³

⁴²ChatGPT4.0: The phrase "corporate laws" here is being used as a rough translation for the German "Korporationsgesetze". The original term's exact meaning isn't clear without additional context but it appears to refer to some kind of rule or laws governing the operation or interaction of the 'syntrocline metroplexes'.

⁴³ChatGPT4.0: In section 5.4, "Syntrocline Metroplex Bridges", we explore the concept of syntrocline metroplexes. These are structures that bridge metroplex totalities of different levels. They have unique properties distinct from the ones for associative metroplexes as the corporate laws for these structures are not firmly determined. This means that each link, which resides in its own metroplex and differs in metroplex degrees, can independently operate with another Metroplex of the same degree.

The section moves onto the topic of how the ambiguity of these connections becomes clear. There are various degrees of metroplex involvement possible with these kinds of structures. Under certain conditions, these might even form conflexive structures.

The concept of syntrocline tectonics is introduced, essentially a method of characterizing the layout of these bridges. The particular choice of inducing syndromes impacts it, and this design could differ across the chain's various links.

The section finishes by stating that although conflexive forms can be utilized as syntrocline roots, the resulting metroplex bridge continues to be a higher-order syntrometric structure above the associated totality.

In short, section 5.4 is a detailed exploration into the nature of syntrocline metroplex bridges, and their unique characteristics and behavior under various conditions and parameters. It provides an intricate model for understanding the metroplex and its complex interactions.

5.5 Tectonics of Metroplex Combinations

From the definition of the syntrocline Metroplex, it is clear that any associative Metroplex can become the root of such a syntrocline structure. According to the laws of Metroplex corporation, there is also the possibility of incorporating an associative Metroplex of the same degree to each bridge link, thus creating a combination of a syntrocline and an associative structure, referred to as a Metroplex combination. The corporation of this combination is unambiguously

described by $\left| \begin{smallmatrix} p \\ b \end{smallmatrix} \right\} \left\{ \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right|$ in a Tp , if $n \leq p \leq N + n$, since only the link of degree p from the syntrocline bridge can incorporate, as, according to the corporation definition, only structures of the same degree can be connected. The incorporation into the combination assumes that a link of degree p actually exists in the bridge. If this link exists, the corporator can act as a concentrator or excenter, i.e., all the syndromes active in the syntrocline induction can be located in the syntropods, so that the corporation remains irrelevant with respect to the syntrocline tectonics.

On the other hand, if a certain number of syndromes of this tectonic structure are added, the occupancy in these now relevant syndromes is changed by $\left| \begin{smallmatrix} p \\ b \end{smallmatrix} \right|$, which in turn has an influence on the syntrocline tectonic structure and thus on the bridge element $p + 1$. If β is the number of those syndromes of the syntrocline tectonic structure on which the corporation to the Metroplex combination acts, and if K is the total number of all tectonic syndromes of the element p , then $0 < \beta < K$ characterizes the *tectonic relevance order* of the corporation in Tn , because these β syndromes change all other bridge elements $n + N \geq P > p$ in their tectonic structure through their corporation to the combination. In the case of $0 < \beta < K$, the elementary combination is tectonically partially relevant of the order β . If $\beta = 0$, then it is irrelevant, while it is totally relevant for $\beta = K$. Obviously, $\beta = 0$ as a combination operator basically assumes an eccentric, which leaves the entire syntrocline tectonic structure in the syntropod of the bridge element, while concentrators basically cause $\beta = K$.

Only for $p = n + N$ is $\beta = 0$ always true, because here lies the end of the Metroplex bridge and thus the end of the tectonic sequence. From elementary Metroplex combinations $\left| \begin{smallmatrix} p \\ b \end{smallmatrix} \right\} \left\{ \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right|$, which only combine an associative structure with a syntrocline, all other higher Metroplex combinations can apparently be generated through appropriate corporation laws and functors. Characteristic for all these variants is the exogenous linkage of syntrometric structures defined in the individual Tp with $n \leq p \leq n + N$ through syntrocline Metroplex bridges, which is why these combinations are referred to as exogenous. In terms of tectonics, a distinction has to be made in these combinations between the associative and the syntrocline corporation. An exogenously associative tectonics exists if associative structures are incorporated into the syntrocline system in the individual Tp , or if functors $S(p + 1)$ additionally syncolate these associative structures.

The tectonics, on the other hand, are corporated syntrocline when several Metroplex bridges of the combine are linked by syntrocline corporators or suitable functors. In general, the exogenous tectonics of a Metroplex combine is mixed, i.e., it is both associatively and syntrocline corporated. Another tectonic variant of these exogenous combinations are the simple and multiple *syntrocline transmissions*. The simple transmission connects Metroplex structures in different totalities along a route through a syntrocline bridge train. These transmission bridges can gradually rise, but

can also fall after the rise. $\left| \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right|$ ⁴⁴, for example, connects $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right|$ in Tn with $\left| \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right|$ in $T(n + N)$ in the sense of a rising bridge. However, $\left| \begin{smallmatrix} n+N \\ b \end{smallmatrix} \right|$ can also exist, and in $T(n + N)$ there is the possibility of corporation $\left| \begin{smallmatrix} p \\ b \end{smallmatrix} \right\} \left\{ \begin{smallmatrix} n+N \\ a \end{smallmatrix} \right|$. In this case, therefore, $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right|$ and $\left| \begin{smallmatrix} n \\ b \end{smallmatrix} \right|$ are connected by a first rising and then falling syntrocline transmission in $T(n + N)$ through $\left\{ \right\}$.

This *transmission* can also become *cyclical*, namely, when there is an eccentric that irrelevantly corporates the two syntrocline roots according to $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right| \circ \left\{ \right\} \circ \left| \begin{smallmatrix} n \\ b \end{smallmatrix} \right|$. However, this irrelevance is not a necessary condition. If the corporation of the roots is tectonically relevant, it only means a tectonic change in the transmission. All relevant corporations that cause tectonic remote effects in a Metroplex combination in this way, i.e., do not limit themselves

⁴⁴HS changed the original a to g

to a single syntrocline Metroplex bridge, are therefore referred to as *tectonic couplings*. The multiple syntrocline transmissions are the consistent extension of the concept of simple transmission. In fact, each bridge member can become the syntrocline root of a further induction, so in this way ambiguously branched syntrocline Metroplexes emerge as purely *syntrocline combinations*.

When such a syntrocline bridge combination is used as a transmission, this *transmission* is obviously *multiple*, and the *transmission number*, i.e., the number of possible Metroplex connections, is determined by the ambiguity of the syntrocline combination. In the case of simple transmission, the transmission number t is $t = 2$, because in the case of a first ascending and then descending syntrocline bridge or of a cyclical course, two simple transmissions were incorporated associatively. For multiple transmissions, $t > 2$ is always true, so that generally for the transmission number $t \geq 2$ applies. All these *exogenous* structured *Metroplex combinations* are characterized by the syntrocline bridges running in the direction of the syndromatic tectonics of the associative structures, because the syntrocline induction always starts in some syndrome interval in order to be coupled to another Metroplex in some other totality. In addition to these exogenous Metroplex combinations with associative or syntrocline incorporations or simple or multiple transmissions and tectonic couplings, the definition of an endogenous Metroplex combination is also possible. The syntrocline bridge always leads from one Tn to another of higher degree, so it bridges the four-dimensional tensorium of an associative structure to that of a higher degree structure.

In the exogenous case, this bridging takes place in the direction of a syndromatic tectonics of the associative structures, in such a way that different structures in the combination are connected to each other. An associative

Metroplex, e.g. $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$, has its basic syntropods stored in a $T(n-1)$ and these pyramidal structures in a $T(n-2)$, etc., until finally the last syntropods stand in the memory of $T0$. Since syntrocline Metroplex bridges always connect totalities

of different degrees, there must also be an internal, i.e. *endogenous Metroplex bridge*, in $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$, which, in contrast to the exogenous form, must necessarily run in the direction of the gradual tectonics of the associative structure. The

syntrocline induction becomes possible because $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$ always consists of the elements of all Tp with $0 \leq p \leq n$, each value p characterizing a syndromatic tectonics. Every element of every syndromatic structural zone p of $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$ is obviously capable of a syntrocline induction if a corresponding syntrocline continuation exists for this element, and this syntrocline Metroplex bridge does not necessarily have to reach outwards to another associative Metroplex.

Instead, the possibility exists that the syntrocline Metroplex bridge remains within $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$ in the sense of a simple or multiple transmission, thus connecting Metroplex elements of zone p and zone $q > p$ in the direction of the gradual tectonics of the associative form (orthogonal to the tectonic syndrome zones p). These endogenously running bridges

in $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$ thus superimpose the structure of $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$, so that in this way a Metroplex combination, namely an endogenous Metroplex combination, has also emerged.^a However, such a combination with endogenous tectonics can only be incorporated syntroclinically or be unincorporated syntroclinically, in contrast to the exogenous combination, because only one associative structure is present, and this cannot corporatize itself.

Syntrocline tectonic couplings also cannot exist, because the syntrocline roots of the endogenous tectonics are syndrome occupations that can only syncline in a predefined way. However, there is the possibility that individual branches of multiple transmissions become exogenous, so that in this way a *syntrocline-like exogenous connection* of the exogenous combination is given. Other exogenous connection options would be the associative corporation, or the induction of exogenous syntrocline continuations. For the symbolic representation of such an endogenous Metroplex

combination, let us assume that $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ b \end{array}^p$ is the root of a syntrocline bridge in the tectonic syndrome zone $p < n$ from $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$, and that the syntrocline bridge runs up to the syndrome zone $p + q \leq n$ with $q > 0$. The syntrocline bridge would then be $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ b \end{array}^{p+q}$ and the notation $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n : E : \begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ b \end{array}^{p+q}$ is then intended to indicate that $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ b \end{array}^{p+q}$ with $p + q \leq n$ and $q > 0$ runs endogenously in $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \\ a \end{array}^n$. The Metroplex combination in elemental form with endogenous tectonics is therefore defined

by:

$$\left| \begin{smallmatrix} \tilde{n} \\ q \end{smallmatrix} \right| \equiv \left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right| E N \left| \begin{smallmatrix} p+q \\ b \end{smallmatrix} \right| \vee p+q \leq n \vee q > 0 \quad (26)$$

which can be written explicitly together with (25) and (25a). The general tectonics of any Metroplex combination can be classified according to the previous description of the tectonic elements. First, two basic tectonic types of combinations must be distinguished, namely the exogenous and the endogenous structures.

The exogenous structures, in turn, can run openly syntrocline, i.e., syntrocline metroplexes emanate from a corporate complex of associative forms from the individual syndromatic zones. These syntrocline open combinations can in turn be corporated associatively or syntrocline. Another class of exogenous forms is composed of syntrocline transmissions, which can also be corporated associatively and syntrocline. In any case, there is the possibility of a tectonic coupling in the case of exogenous tectonics of the combination, through which the syntrocline tectonics of the bridges can be changed, which in turn has a feedback effect on the syndromatic tectonics of the associative structures. In contrast to this exogenous tectonic basic type, there is no possibility of a syntrocline tectonic coupling in the case of the endogenous tectonic basic type, because while exogenously the syntrocline bridges run in the direction of the syndromatic tectonics, this occurs in the case of endogenous structures in the direction of the gradual tectonics of only one associative form. The *general metroplex combination* is composed of all these variants mixed together. Any exogen forms with open syntroclines, any syntrocline transmissions, furthermore in any associative and syntrocline corporate states under the influence of any tectonic couplings, are in syntrocline connection with any endogenous systems. The interrelationship between such general exogenous and endogenous tectonic systems occurs in three ways. The endogenous system can be connected to the exogenous system through a corporator, or its zones of syndromatic

tectonics become syntrocline roots. Finally, there is also the possibility that in $\left| \begin{smallmatrix} \tilde{n} \\ q \end{smallmatrix} \right|$ the syntrocline bridge is a multiple transmission, some branches of which run out of $\left| \begin{smallmatrix} n \\ a \end{smallmatrix} \right|$ and thus intervene as syntrocline metroplex bridges into the exogenous combination. In the general case, all these possibilities are realized simultaneously.

Finally, there is also the possibility that a syntrocline bundle in a $T(m-1)$ penetrates an associative metroplex of degree m in such a way that individual syndrome occupations $\gamma > 0$ or individual metrophorene elements $\gamma = 0$ are identical to syntrocline members. In the general case $\gamma \geq 0$ of such a metroplex diabatcs, the cases of metaphorical ($\gamma = 0$) and syndromatic ($\gamma > 0$) diabatcs are included as special cases. A tectonic coupling would change the metroplex in the case $\gamma = 0$, while conversely a change in this metroplex at $\gamma > 0$ would cause a tectonic coupling. However, this means that in the general case $\gamma \geq 0$, any coupling due to the *metroplex diabatcs must result in a syntrocline feedback*. The metroplex diabatcs should therefore be regarded as another tectonic element of the general metroplex combinations.⁴⁵

⁴⁵Summary by ChatGPT4.0 in the style of Richard Feynman: Alright, now let's see if we can unwrap this complex idea of Tectonics of Metroplex Combinations and make it a little more relatable.

We spent a good deal of time exploring the idea of 'tectonics', which in this context doesn't involve earth's crust moving around, but instead refers to the way certain mathematical structures, called 'Metroplexes', interact and relate to each other. We started by discussing simple and cyclic transmissions, and then graduated to more complex structures that incorporate 'syntrocline roots'. We drew a distinction between 'relevant' and 'irrelevant' corporates, in a way that's somewhat similar to a mechanic looking at the moving parts of a car and deciding which ones are causing a problem and which ones are just doing their job.

Next, we talked about the idea of multiple syntrocline transmissions, envisioning them as extensions or variations of the simpler transmissions. Imagine it like a game of telephone where each node isn't just passing the message to the next person, but can also change the message in some way, creating a much more complex and dynamic network.

We then started to distinguish between exogenous and endogenous structures, or how transmissions can move between Metroplexes (exogenous) or within a single Metroplex (endogenous). This concept brought a new depth to our understanding, much like realizing that the movement of individual cells in your body tells a different story than the movement of your body as a whole.

And then things got even more interesting when we considered that the transmission process itself could become more complex, with multiple branches and possible connections (what we termed 'tektonic couplings') that could change the nature of the transmission.

So, all in all, we journeyed through an abstract landscape of mathematical structures, transmissions, and connections. It might seem far removed from the stuff you encounter in everyday life, but remember, even the most concrete things - from bridges to computer chips - owe their existence to abstract mathematical principles. So next time you cross a bridge, spare a thought for the Metroplexes!

6 The televariant aeonic Area

6.1 Mono- and Polydromy of Metroplex aeondyne and their Telecentricity

Every general Metroplex combination (associative and syntrocline forms are to be addressed as special cases) can fundamentally be regarded as a higher syntrometric structure according to the preceding. This means, however, that the basic syntropods of the entire structure are in the memory $T0$, i.e., there are $1 \leq j \leq Q$ elementary pyramidal syntrices from which the entire Metroplex combination arises. For these pyramidal syntrices, the expanded concept formation of the band syntrix applies, so that these basic syntropods may also be band syntrices. But if band syntrices are allowed, then primigenous aeondynes must also be allowed, because if the micromar definition interval of the band syntrix is extended macromar, then a primigenous aeondyne has arisen. These primigenous aeondynes are thus defined when the metaphores of the Q pyramidal syntrices depend on any conceptual parameters. If each of the base syntrices $\tilde{a}|_j$ depends on $1 \leq i_j \leq n_j$ parameters t_{ij} , then each $\tilde{a}|_j$ is an n_j -dimensional primigenous aeondyne. At the same time, these Q primigenous aeondynes form the basic syntropods of a Metroplex combination, i.e., the whole combination must also depend on these parameters and thus form a higher aeondyne structure as an extension of the primigenous aeondyne, similar to how the Metroplex extends and implies the Syntrix concept.

Such a structure is therefore referred to as a Metroplex aeondyne or simply as an "aeondyne". An upper limit C can be defined for the dimension number of the aeondyne tensorium. Because of $\tilde{a}|_j(t_{(ij)})_1^{n_i}$, it follows that $C = \sum_{i=1}^Q n_i$, i.e., the dimension number D of the tensorium lies in the interval $0 \leq D \leq C$. The case $D = C$ occurs when each primigenous aeondyne of the basis runs in its own sub-tensorium (regardless of the metaphorical syncloative or linked nature), and when there is furthermore no running conceptual parameter that reappears in another sub-tensorium. If, on the other hand, there are such duplicities, or if several primigenous base syntropods run in the same sub-tensorium, then $D < C$ is always true, and when the other limit $D = 0$ is reached, then there is no tensorium at all, so that the aeondyne degenerates to the Metroplex combination. This means that the aeondyne is conceptually superordinate to the Metroplex combination. With regard to the mono- or polydromy of an aeondyne, it becomes evident that the precondition for monodromy is always fulfilled when all primigenous aeondynes of the basis are monodromous, but this condition alone does not yet guarantee factual monodromy of the aeondyne, because in higher gradual zones, and especially in syntrocline tectonics (because of the possibility of tectonic couplings), there is always the possibility of a multivalent course in higher syndromatic structure zones.

In the general case, however, the course of the aeondyne is not monodromous, i.e., in certain, namely $1 \leq \mu \leq M$ parameter spaces of dimensionality $L_\mu \leq D$, ambiguities occur due to the laws of Metroplex synkolation and tectonics, so that the aeondyne splits in these μ tensoria of the respective dimension L_μ in P_μ -fold ways. Thus, the aeondyne falls apart into μ , into P_μ branches, so that in this tensorium μ , the aeondyne becomes P_μ -fold polydromous. For this reason, μ is referred to as the "polydromy center" and L_μ as its dimensionality. From the dimension number T_μ of the element of the aeondyne becoming polydromous (in general, not the whole aeondyne becomes polydromous, but only individual tectonic zones), it can then be concluded that L_μ of the polydromy center, because only $L_\mu = T_\mu$ is possible. On each polydromous branch of the thus split aeondyne structure, there must then be possibilities for polydromy centers, so that as a result of this aeondyne polydromy, an entire panorama of aeondyne structures connected by polydromy centers - the so-called "aeondyne panorama" - arises. For these panoramas, there are a variety of structure possibilities that are determined by the qualitative and quantitative course of the distribution of the polydromy centers over the area of the panorama. Thus, each panorama can initially be limited or unlimited.

Furthermore, a semi-limitation can occur, then the panorama is radially structured. In the monodromous case, the panorama consists only of one course, and this aeondyne behaves with respect to the limitation like a primigenic form. The monodromous aeondyne is undoubtedly to be addressed as a special case of the polydromous ones, from which it follows that the aeondyne panorama of the monodromous aeondyne is conceptually superior. In the polydromous case, the three main classes unlimited, semi-limited and totally limited are to be distinguished with respect to the limitation, and in each of these cases, there is again the possibility of forming polydromy classes. For example, there is the possibility of symmetric and asymmetric polydromy structure, which can again comply with certain laws. These are the laws of polydromy change, i.e., the respective number of polydromy centers is plotted for the description of these laws over the panorama extent. Basically, as long as the antinomic concept to the polydromy center is not defined, there can only be the polydromy increase and the polydromy constancy. The form of such a *polydromy diagram*

then already provides the desired classification of panorama structures, if for each point of the diagram a distribution diagram is given in the respective orthogonal direction to the panorama extent, which characterizes the distribution of those polydromy centers, whose number is fixed in the discussed point of the polydromy diagram. The symmetry or asymmetry of the panorama structure is then expressed in these additional distribution diagrams.

The polydromy diagram supplemented by the distribution diagrams is referred to as the classification diagram for the metaphorical illustration of the panorama structure. All polydromous panoramas, which were classified in the foregoing, have the common property that their polydromy increases as progress is made along the parameters, or remains constant, i.e., they are polydromously ascending. From this it follows directly that there must also be polydromously descending panoramas, because, if the parameters are traversed in reverse direction (which is always possible), the meaning of the polydromy centers apparently reverses, because several aeondyne branches converge in such a center. In this way, the meaning of the concept of polydromy center is reversed antinomically, because these centers, acting as collectors, now reduce the polydromy, so that a panorama structure with falling polydromy is present. With this falling polydromy, however, the *classification diagram* can be supplemented, in such a way that the polydromy diagram consists of ascending, constant and falling branches. If the *collector concept* is defined, a polydromously descending area can always be connected to a polydromously ascending one. Therefore, the polydromy centers of ascending panoramas are juxtaposed with the collectors of the falling structures, for which the same classification applies as for the ascending forms.

The existence of polydromously ascending and descending panoramas directly implies the existence of a third type, which arises from the combination of the first two forms. In this type, there are both polydromy centers and collectors, so that in the same panorama, the polydromy increases and decreases. These panoramas are characterized by a property of the far-centered polydromy. A polydromy center creates a bundle of aeondyne branches on an aeondyne, which take their own course, but there can be a collector at some positive distance from the parameter values of the polydromy center in the aeondynic tensorium, which reverses this polydromy and thus makes the panorama, relative to the polydromy center, far-centered. Such collectors are therefore referred to as *telecenters*. The concepts of telecenter and polydromy center thus become relative, as when traversing the parameter intervals in the reverse direction, the polydromy center and the collector swap their meanings. For this reason, all areas of the aeondynic tensorium are referred to as telecenters, which in this way cause the telecentering of a panorama. Such a telecentric panorama therefore forms the area of a mesh of aeondyne branches, each of which eventually ends in a telecenter, which justifies the designation aeondynic, or shortly aeonic area, for a telecentric aeondyne panorama.

The condition for the existence of telecenters is that after passing through the telecentrically polarized panorama section, the old polydromy is restored. The telecentering of the Area always equates to a telecentric polarization, as the position of the telecenters in the tensorium of the Area characterizes a polar structure in terms of the course of all aeondyne branches within the Area. If the condition for the existence of telecenters is not met, then there are only polydromy centers and collectors in the panorama, but no *telecentric polarization*, i.e., there is indeed an aeondyne panorama, but no Area. If the telecenters are also panorama boundaries, they are referred to as main telecenters, and the entire panorama is telecentrically polarized. Moreover, within each Area, there can still be telecentrically polarized partial structures, which are obviously sub-areas of the main area, bounded by secondary telecenters. If individual main areas with their telecenters connect to each other, Area chains are formed, capable of forming even higher structures, and telecentric polarization can also take effect in their formation. If the discussed Area is designated as a first-order Area ($AR1$), and Area chains formed from it again form a telecentrically polarized Area, then this is an $AR2$, i.e., a second-order Area, and so on. By the reasoning of complete induction, an ARn , i.e., an Area of order n is possible in this way. The elements of each ARn are always the $AR1$, and these Areas are fundamentally classifiable, as a classification diagram must exist for each $AR1$.

The same division into order grades also applies to aeondyne panoramas, but it turns out that these panoramas are special cases of Areas. In the panorama, the telecentric polarization is missing, i.e., either the aeondyne branches have reached their boundaries somewhere in the tensorium without a further collector existing, or the polydromic aeondyne system continues indefinitely into the infinity of the tensorium. The first case is to be regarded as a special case of the second, because if an aeondyne branch finds its limit in the tensorium, this is identical to a singularity in the course. The singularity would be the area of the tensorium in which the basic syntropods of the $T0$ of the aeondyne in the tensorium running metroplex combo all become identical to zero syntripes and run as such aeondyne into the infinity of the tensorium. Consequently, any panorama missing the telecentric polarization can be regarded as an open panorama

in one or both directions, whose aeondyne system with variable polydromy runs into the infinity of the tensorium.

The same boundlessness then also applies to the polydromy diagram, which in the case of the Area runs back into itself in the telecentres, but in the case of the open panorama it continues indefinitely. The boundary of such an open polydromy diagram is obviously a metaphorically affine improper element located in infinity, which, again metaphorically, becomes a proper element in the finite by projection. Based on this metaphor, the boundary of an open panorama would therefore be improper, namely a *projective telecentre*, so that these panoramas, using the concept of the projective telecentre, are one-sided or two-sided projective Areas, depending on whether they are one-sided or two-sided. Following the conceptual introduction of the projective telecentre based on a metaphor, it immediately becomes

evident that the *aeonic Area* is the concept that is superior to the panorama. If $\bigwedge_a^n$ is any Metroplex combination of maximum degree n defined by (25), (25a), and (26), and the dependency consists of $1 \leq i \leq Q$ conceptual parameters t_i , then the course $\bigwedge_a^n \equiv (t_i)_1^Q$ over the Q -dimensional tensorium describes in the general case a polydromic aeondyne, which, according to the above, must be telecentrically polarized as a panorama, thus must be an $AR1$. If T and T' are the telecentres, then this first order Area is formally defined by

$$AR1 \equiv AR_{(T)}^{(T')} \left[\bigwedge_a^n (t_i)_1^Q \right].$$

If there are a total of $1 \leq \gamma_1 \leq p_1$ of these $AR1$ over the same tensorium, all of which are linked with their telecentres in such a way that they form sub-areas that are in turn polarized with the telecentres $\underline{T}$ and $\underline{T'}$, then an $AR2$ has been created, the linking law of which is expressed by $AR_{\underline{T}}^{T'}$. Accordingly, we have

$$AR2 \equiv AR_{\underline{T}}^{T'} [(AR1)_{\gamma_1}]_1^{p_1}.$$

This procedure can be continued recursively according to the principle of complete induction until finally an Area of order $q \geq 2$ is formed, which logically is defined in the form

$$ARq \equiv AR_{(T_1)}^{(T_2)} [(AR(q-1))_{\gamma_{q-1}}]_1^{p_{q-1}}$$

if T_1 and T_2 are the telecentres of the ARq . The general definition of Area is therefore:

$$ARq \equiv AR_{(T_1)}^{(T_2)} [(AR(q-1))_{\gamma_{q-1}}]_1^{p_{q-1}} \vee AR1 \equiv AR_{(T)}^{(T')} \left[\bigwedge_a^n (t_i)_1^Q \right] \quad (27)$$

From this general definition, it is clear that an Area $q \geq 2$ can only exist if all sub-areas are defined over the same Q -dimensional tensorium as the p_1 areas, $(AR1)_{\gamma_1}$ i.e., the parameter dependence of the same tensorium also applies to the ARq , so that the main area appears to be dimensioned in the same way as all sub-areas down to $AR1$. This identity of the area tensorium is characteristic for all aeonic areas of order $q > 1$.⁴⁶

⁴⁶ChatGPT4.0 Summary: Alright, let's imagine we're sitting down together in the cafeteria at Caltech and I'm explaining the core concepts of this section to you. It might go something like this:

"Imagine you have this thing called a 'Metroplex aeondyne'. Don't worry about the details yet. This aeondyne can have a property we're calling 'polydromy', which can increase, stay constant, or decrease as we move along certain parameters. This seems natural, right? It's like driving down a road - you can speed up, keep a steady pace, or slow down.

So we figured, if we can have 'increasing' and 'constant' polydromy, then there must be 'decreasing' polydromy too. Just think about driving the other way down the road!

To classify this, we have something we call a 'polydromy diagram'. This fancy diagram is like a map, showing us how the polydromy changes. But the plot thickens! Because there are these special points we call 'collectors' that can decrease the polydromy and create a structure with falling polydromy.

Next, we stumbled upon the idea that there must be a third type of panorama structure, which is a combination of the first two. A panorama that both rises and falls. It's a bit like a rollercoaster ride, going up and down along the track.

Now, here's where things start to get really interesting! These panorama structures have something called 'telecenters'. Don't worry, it's not as complicated as it sounds. They're just collectors that reverse the polydromy effect and thus, create a panorama that's far-centered, or 'telecentric'.

And from this we realized that we could have 'areas' of telecentric panoramas, which we call 'aeonic areas'. You can think of these areas as a kind of landscape of aeondyne branches, each one ending in a telecenter.

6.2 Levels of Transcendence and Transcendental Tectonics

If there is an aeonic area of any order defined over a tensorium, then there are always $M < \infty$ monodromic aeondyne paths between the main telecenters within this area. Since each of these monodromic partial aeondynes is a metroplex combination, all synkolation of the combination also runs along these aeondynes as functors of the concept parameters of the tensorium. Because of this structural property of the area, however, there is basically the possibility that affinity syndromes a_γ can be isolated in $2 \leq \gamma \leq N \leq M$ monodromic runs, which show affinities among each other.

But if such affinities exist in the form of aeondynic affinity syndromes a_γ with respect to any subjective aspect of the relevant aspect complex, then the affine correlations must basically be expressible by suitable metroplex functors

in the sense of synkolators. So there must be a system of $1 \leq i \leq P$ functors Γ_i of the synkolation level $K_i \leq N$ through which the correlation of the affinity syndromes in the form $\Gamma_i(a_\gamma)_1^{K_i}$ with $K_i \leq N$ express. From this it follows that every synkulator Γ_i is effective in $\binom{N}{K_i}$ ways.

The synkolation that arose in this way are evidently also monodromic aeondyne runs between the main telecenters, but it is evident that these states synkolated from the affinity syndromes no longer belong to the original area but run as a transcendent structure over it.

The possible system of Γ_i evidently depends not only on the a_γ and N , but also essentially on the classification diagram of the area ab . Since the synkolated runs, of which there are a total of $Z_{(1)} = \sum_{i=1}^P \binom{N}{K_i}$, are transcendent in

relation to the original area, the Γ_i are referred to as *transcendence synkolators*. To distinguish between the synkolated transcendent area and the original structure, the ARq is referred to as an area of *transcendence level 0*, symbolized by $C_{(0)}$, while the transcendent form must have transcendence level 1. This field of transcendence $C_{(1)}(ARq)$ at the first

level of transcendence is synkolated by the system of first level transcendence synkolators $\Gamma_i \equiv \Gamma_{(1)i}$, of which there are $1 \leq i \leq P(1)$ according to the classification diagram. Thus, (1) denotes the transcendence level 1.

In the $C_{(1)}$, that is, in the first level field of transcendence, there can again be affinity syndromes, in such a way that of the $Z_{(1)}$ runs of the field $C_{(1)}$ a total of $N_{(1)} \leq Z_{(1)}$ synkolate to a *field of transcendence* $C_{(2)}$ via a system of

$P_{(2)}$ transcendence synkolators $\Gamma_{(2)i}$ and so on.

In this way, a general level of transcendence $m > 0$ eventually becomes possible, so that a sequence of transcendence fields of increasing levels of transcendence lie above the ARq . This can be symbolized by $C_{(1)}(ARq)$ if $C_{(m)}$ contains all transcendental synkolation laws in the interval $0 < \mu \leq m$ of the levels of transcendence. Since already in the synkolation of the field $C_{(m)}(ARq)$ only monodromic aeondyne branches synkolate except for *subareas* limited by true secondary telecenters, and this process continues in all $m > 1$, there are neither polydromy centers nor collectors in all $C_{(m)}(ARq)$ with $m > 0$ in contrast to the area $C_{(0)}(ARq) \equiv ARq$, next to the *secondary* and *main telecenters*. That is, all transcendent aeondynes in monodromic form diverge from one telecenter to converge in another for $m > 0$.

The described process of transcendence level synkolation related to the structure of an ARq and thus carries an *intrasynkolative* character. As Metroplex combinations can always incorporate, there is fundamentally always the possibility to set different areas into a mutual relationship through incorporators, which can give rise to new aeonic areas. In this incorporation of areas, the incorporating areas, similar to the incorporation of Metroplex combinations, are absorbed in a new area. However, if these different areas exist alongside each other without a corporative relationship, affinities can still exist between the monodromic aeondyne runs of different areas, similarly to the intrasynkolative generation of levels of transcendence, such that affinity syndromes along these aeondynes can again be isolated, which synkolate transcendental aeondynes in the next higher level through transcendence synkolators. In this way, an *extrasynkolative* transcendental correlation of individual monodromic aeondyne runs of different aeonic areas can

But it doesn't stop there! We also have these so-called 'open panoramas'. They're just like the telecentric panoramas, except they don't have a telecenter. They're like roads that go on forever.

So, to wrap it up, we've been able to define and classify different types of panorama structures, with rising, constant, and falling polydromy. And we've discovered that we can group these structures into 'areas', which are either telecentric or open. It's a whole new world of complexity and structure, all from studying these metroplex aeondynes. Isn't that just fascinating?"

occur, which does not act in the sense of a corporating functor of the $C_{(0)}$.

Such extrasynkolative transcendental aeodynes, which can link different areas transcendently, are possible in all levels of transcendence. If the functor $\Gamma_{(j)}$ is a transcendence synkolator that operates extrasynkolatively in the level of transcendence j , and $C_{(k)}(ARr)$ and $C_{(l)}(ARs)$ are two areas with intrasynkolative levels of transcendence of different order r and⁴⁷ s , and there exist within these areas within the level of transcendence j an aeodyne that isolates an affinity syndrome relative to the other, then it is apparent that the extrasynkolative aeodyne of the level of

transcendence $j + 1$ ⁴⁸ is described by $C_{(k)}(ARr) \Gamma_{(j)} C_{(l)}(ARs)$, and it immediately becomes evident that $0 \leq j \leq l$ must hold if $l \leq k$. If $k = l = 0$, then the transcendental aeodyne of the level of transcendence 1 is purely extrasynkolative, but it is mixed for $k > 0$ and $l > 0$, because in this case there are also intrasynkolative fields of transcendence. This immediately follows that purely extrasynkolative fields of transcendence can only occur at the level of transcendence 1.

Based on the foregoing, every aeonic area must be assigned a transcendental tectonics, which can be intra- or extrasynkolative, or mixed. The mixed tectonics apparently forms an analogy to the metroplex combinations. In any case, a *gradual transcendental tectonics* is fundamentally existent in the direction of the levels of transcendence, the course of which depends on the complex structure of the complexes of transcendence synkolators applied and is divided into the three mentioned classes. Furthermore, there is an *syndromatic transcendental tectonics* orthogonal to this, the course of which is determined by the respective elements of the complex synkolator. These two forms of transcendental tectonics in the level of transcendence 0 transition into the gradual and syndromatic tectonics of those metroplex combinations, which as polydrome aeodynes of the area span over the relevant tensorium. Finally, there must also be a *telecentric transcendental tectonics* orthogonal to the first two, the course of which directly follows from the area structure $C_{(0)}$ and the associated classification diagram.

In any case, the delimitation of this telecentric transcendental tectonics, and for all levels of transcendence, is determined by the location of the telecenters of the area $C_{(0)}$ in the tensorium of this area. These three tectonic directions fully characterize the tectonic structure, i.e., the architectonics of the relevant field of transcendence. In the case of extrasynkolative fields of transcendence, there is a transcendence synkolation of different areas with each other, i.e., the individuality of the single area transitions in the level of transcendence 0 into that of a partial structure, which also applies to the synkolation of the higher levels of the field of transcendence. A pseudo form of extrasynkolative fields of transcendence arises intrasynkolatively when sub-areas exist within the area that are delimited by secondary telecenters. This form of architectonics will always appear when the area has an order $q > 1$.⁴⁹

⁴⁷ HS added 'and'

⁴⁸ LAA: is number 1 or letter l correct?

⁴⁹ ChatGPT 4.0: Alright, let's see if we can't give this a bit of the Feynman flavor.

So, we're diving into this notion of "Levels of Transcendence and Transcendental Tectonics", which might sound a bit highfalutin, but we're really just talking about how these aeonic areas interact with each other, and how they evolve in a way. Now, what's really fascinating here is the way that these areas can create these paths or, as the original text calls them, "monodrome aeodynic paths". Picture it like a highway system with M different routes that you could take between two major cities. The cities are the important parts, they're like the telecenters in our model.

Then we have these things called "affinity syndromes" - these are basically shared characteristics between different paths. If you can find these similarities, you can represent them using mathematical functions, or as the text calls them, "transcendence synkolators". Now, the important thing is, when you use these synkolators, you're essentially going beyond the original area you were working with. You're transcending it, hence the name "transcendence synkolators".

What's also very intriguing is that there are multiple "levels of transcendence". You start with your original area, but then you can go beyond it, to a new level. Each level is defined by the synkolators that you're using. Now, there's a hierarchy to these levels, which means you can go on creating more and more transcendent areas, much like climbing a staircase.

Finally, it turns out there are different types of "tectonics" that are associated with this whole process. Just like the tectonic plates that make up the Earth's crust, these transcendental tectonics give structure and form to our model. They help us understand how the different areas and levels interact with each other. It's really quite a complex, intricate dance, but when you break it down, it's simply a beautiful piece of mathematical choreography.

And that's the gist of section 6.2. Not so complicated when you put it that way, right?

Can you elaborate on what Heim meant about "transcendence" and "Transcendental Tectonics?" ChatGPT Sure! Now remember, we're trying to simplify and illustrate concepts from Heim's work, so bear with me if we get a bit whimsical in our explanations. Let's keep in mind that these concepts are embedded in a very abstract mathematical framework, so they might not map directly to everyday experience, but we can still get a sense of what's going on.

When Heim talks about "transcendence", he's referring to a process where certain mathematical structures or patterns within an aeonic area are recognized as possessing some shared characteristic, which can be expressed using a mathematical function, called a "transcendence synkolator".

6.3 Tele- and Dysvariants

Each aeodyne has, as an area in both the monodrome and polydrome cases, a triple tectonic: namely gradual, syndromatic, and telecentric. These three tectonic structures in their entirety, along with the distribution of the polydrome centers according to the classification diagram, form the architectonics of the area with its transcendence fields. All these tectonic forms can experience continuous or discontinuous structural changes as they progress along the parameter, that is, the area can be subject to a tectonic variance. This must also apply to every type of transcendental tectonics of synkolated transcendence fields.

Obviously, the telecentric tectonics within the area occupies an exceptional position, because it must be oriented according to the telecenters as a result of the telecentric polarization. Therefore, the gradual and syndromatic tectonics always have to be related to the telecentric one. This telecentric tectonics is determined by the change in the number of syndromatic structure zones when the parameters of the tensorium are traversed from one telecenter to another.

If this telecentric tectonics, that is, the number of syndromatic structure zones, does not change, then the corresponding area is televariant. The synkolation courses in the syndromatic tectonics, as well as the course in the gradual tectonics, can change, but this change (which may also be absent as a special case) is already predetermined in a lamellar manner by the constant telecentric tectonics, because the area is a telecentrically polarized aeodyne panorama. For this reason, the term *televariance* was coined. However, this condition of televariance is only completely fulfilled if it applies to all aeodyne structures of the area.

If this constancy of the telecentric tectonics is not given, that is, if the number of syndromatic structure zones changes, then there are faults in the syndromatic zones within the architectonics, and the area undergoes a *dysvariance*. This can be total or partial, and it is always total when every monodrome⁵⁰ branch has a *dysvariance site*.

But if there are individual televariant branches, then the dysvariance is partial. If this dysvariance leads to an expansion of the gradual tectonics, that is, if the number of syndromatic zones increases in the direction of this gradual tectonics, then the dysvariance is fulfilled in these additional zones and increases gradually. In the other case, the number of these zones decreases, which leads to dysvariant extinction.

Regardless of the nature of this dysvariance, there is generally always a tectonics of televariant zones alongside this dysvariant tectonics. Not all syndromatic structure zones in a gradual evaluation need to show such dysvariant faults in a monodrome aeodyne branch between the telecenters. Only when the number of televariant zones reaches the value 0 is the relevant aeodyne branch absolutely dysvariant.

Therefore, following this conceptual refinement, the area can be total and absolute, total, partially and absolute, as well as partially dysvariant or televariant.

According to this interaction of tele- and dysvariant aeodyne courses within the area, an internal structure classification of aeonic areas is given, along with the classification diagram. The possibilities of *dysvariant extinction* in an aeodyne also have various characters. Fundamentally, there can only be three types of this extinction:

- a) The relative beginning (relative to the base syntropodes in the T0) of the gradual tectonics experiences an extinction from some point of dysvariance onwards.
- b) This extinction affects the upper limit of the gradual tectonics.
- c) Any zone area within the course of gradual tectonics becomes dysvariant in the sense of an extinction, without affecting the upper or lower limit of the tectonics.

In all three cases, the dysvariance can become absolute. In the most general case, all three *dysvariance forms*,

In simpler terms, it's like we're discovering a pattern or a commonality that wasn't visible before, and this commonality allows us to move beyond the original confines of our model. It's like a hidden passage in a maze that allows us to step outside the maze entirely, thus "transcending" it.

"Transcendental Tectonics", on the other hand, is about the structural principles of this process of transcendence. Just like how geological tectonics deals with the structure and movement of Earth's crust, transcendental tectonics is about the structures and 'movements' within the system of aeonic areas and their transcendental interactions. These structures or movements are governed by rules, much like how the movement of tectonic plates are governed by physical laws.

In Heim's work, "tectonics" also refers to the different 'directions' or dimensions within this system of transcendence. There are gradational, syndromatic, and telecentric dimensions, each with its own specific characteristics and behaviors. The interaction of these dimensions leads to the formation of new structures or "levels" of transcendence, adding more depth and complexity to the model.

So in essence, Heim is using these terms to describe a system of interconnected mathematical structures that evolve and interact in complex ways, forming new structures and levels of complexity as they do so. The key here is that the notion of 'transcendence' in Heim's work isn't a philosophical or mystical one, but rather a mathematical and conceptual one.

⁵⁰HS: changed from "polydrome"

namely a: *initial*, b: *final* and c: *intermittent* of extinction can occur multiple times within an aeodyne path of the area. This implies a further refinement of the architectural classification of aeonic areas. All investigations about Tele- and Dysvariance were developed for the transcendence level $C_{(0)}$, but they can easily be transferred to transcendence fields of higher transcendence level above the area, as the gradual tectonics of the area $C_{(0)}$ is directly connected to the gradual transcendental tectonics. When transferring, however, it must be taken into account that, according to the investigations of the transcendence fields in the area of higher transcendence levels than 0, there are only monodrome aeodyne branches without polydrome centers or collectors between the telecenters.⁵¹

⁵¹ Sure, I'll try to provide a "Feynmanized" recap of this information!

Now, what Heim is doing here might sound a little like trying to tame a bucking bronco with a lasso made out of equations. But don't worry, let's break this down:

Each of these 'aeodynes' (think of them as Heim's special version of 'dynamic areas') has three types of tectonics - gradual, syndromatic, and telecentric. Now, think of tectonics as different ways these areas can change and interact. The telecentric tectonics are pretty special since they point towards the 'telecenters' (special points of orientation) due to telecentric polarization. It's like everything else in the area is playing second fiddle to the telecentric tectonics.

Now, the catch is, as we move along the parameters, the area can change. Like a kaleidoscope, it can show different patterns, some continuous, others not so much. We call this 'tektonic variance'. If the telecentric tectonics stay the same, even if everything else changes, then we have a stable view, we call it 'televariance'. But if the telecentric tectonics change, if the number of syndromatic structure zones change, we have some sort of 'fault lines' or 'dysvariance'. This dysvariance can be total or partial, and it's total if every polydrome branch has a 'dysvariance point'.

When it comes to dysvariance, things can get extinct, or disappear. This can happen at the beginning, the end, or at some point in the middle of the gradual tectonic trajectory. This offers us an even finer way to categorize the 'architectural classification' of aeonic areas.

Now, here's the important bit - all of this was developed for a transcendence level $C_{(0)}$. That's Heim's 'ground floor', if you like. But it can be extended to higher levels, with one thing to remember: in the higher transcendence levels, we only get monodrome aeodyne branches - we don't have polydrome centers or collectors. So, the 'view' might be a bit different from the ground floor, but we can still use our ideas of tele- and dysvariance.

Phew! That was a ride, wasn't it? But we managed to get to the end of it. Now we just need to keep these ideas in mind as we move forward!

6.4 Metastable Syncolation States of the Extinction Discriminants

The question that remains to be clarified is how the *extinction discriminant*, that is, the tectonic boundary in the gradual direction (in terms of telecentric tectonics) of a dysvariant structural area of an aeondyne (in the sense of extinction) proceeds, and how this boundary should be understood in terms of the syncolations in the syndromatic structural zones that limit the dysvariant extinction structure. Therefore, it should be investigated how the individual syncolations in the respective dysvariant syndromatic tectonics change in the area of the extinction discriminant, so that dysvariant extinction can occur at all, and how these syncolation states in the discriminant should be constituted in terms of their parameter dependence so that they remain metastable in a suitable parameter interval until there is a change in dysvariance as one continues to proceed along the parameters.

Obviously, the syncolation states in the extinction discriminant must always be of a metastable nature in the general case, because a parameter-dependent change in dysvariant extinction always requires a concomitant change in the discriminant course in gradual tectonics. Only when the extinction discriminant is televariant in the special case can its syncolation states be stable. The terms metastable and stable always refer to the discussed parameter intervals in the tensorium of the area.

Regardless of what these metastable syncolation states may look like, the extinction discriminant always runs simply in the case of initial and final dysvariance, but doubly in the case of intermittent dysvariance.

If the dysvariance is not absolute, then the syncolation states of the discriminants must be contained within a televariant zone when the discriminant passes through its absolute gradual-tectonic extremum at the relevant point of the tensorium. This televariant zone then also contains metastable syncolations of the discriminant in the discriminant extremum. If the dysvariance is absolute, then there can be no televariant zone in the discriminant at all. Only in the case of intermittent dysvariance, due to the double discriminant progression, is there the possibility that one branch of the discriminant cuts the aeondyne as absolutely dysvariant, while the other branch contains a televariant zone.

Every intermittent dysvariant extinction separates an aeondyne into areas of higher gradual tectonics and areas of lower gradual tectonics, such that the branch of the discriminant that delimits the higher-graduated areas corresponds to an initial dysvariant extinction, while the other corresponds to a final dysvariant extinction. Thus, the metroplex combinate at an aeondyne point where such an intermittent dysvariance prevails can be understood in the sense of syndrome balls akin to syntropods, regardless of whether these are conflutive forms or not. Each extinction discriminant can rise or fall monotonically, in a stronger or weaker sense, or it can pass through dysvariance maxima and minima, cut out dysvariance arcs, remain constant, etc.

Thus, all possibilities of multidimensional function courses are given if the tensorium of the area is metaphorically illustrated by a multidimensional abstract space. The discriminant always passes through a dysvariance extremum when areas of increasing and decreasing dysvariance adjoin. If the dysvariance decreases, the number of structural zones increases, which necessitates a resyncolation of the metastable syncolation states. Through this way, dysvariance arcs and resyncolations can break through televariant tectonics, regardless of whether this breakthrough is of an initial, final, or intermittent nature.

In the case of initial or intermittent extinction, the elementary process of extinction does not necessarily go back to a change in the aeondyne progressing metroplex combinate. Rather, the dysvariant syndromes are such that they are not defined along the parameter interval under discussion. Only in the case of final extinction can the dysvariance be traced back to a structural change of the syncolators, but the dysvariance can also occur in analogy to the initial or intermittent case.

Thus, the concept of dysvariance experiences a further refinement through the necessary distinction between structural and functional dysvariance. In the case of structural dysvariance, there is an internal structural change of the metroplex combinate at the dysvariance point of the tensorium, while in the case of functional dysvariance this structure remains preserved, and only the occupancies of certain syndromes along the extinction interval are no longer defined.⁵²

⁵²ChapGPT4.0: Sure, let's break down this concept into simpler terms with the help of a Feynman-style explanation.

Think of an area in Heim's syntrometry as a complex, multidimensional space, filled with different patterns or structures called aeondynes. These structures aren't static; they can change as you move along certain parameters. This concept is similar to how we can observe different phases or states of matter, like solid, liquid, and gas, as we vary temperature and pressure.

In the context of Heim's syntrometry, these aeondynes can sometimes enter a state known as "dysvariance," which is somewhat akin to a phase transition in matter. In this state, there's a notable change in the structure of the aeondyne, and the discriminant (or boundary) of this dysvariant

6.5 Televariance Condition of Telecentric Polarization

After the previous investigations into tele- and dysvariance, it becomes possible to establish a criterion for the telecentric polarization of an area. For the dysvariant structure of an area, there are a variety of classes, such as absolutely total, partial, etc., each of which can appear in the initial, final, or intermittent form and can be of a structural or functional nature. Formally, according to this classification, there apparently is only a single class, namely absolute total dysvariance, in which there can be no affine telecentric polarization because only projective telecenters exist here. Such a projective telecentric area thus corresponds to the definition of the panorama of a polydromic aeondyne, so that such a panorama is always determined by absolute total dysvariance. In all other classes, there is the possibility of telecentric polarization of affine areas, but it is a case of pseudotelecentric polarization when the individual polydromic branches are continuous, but no televariant structural zones occur in any branch, i.e., when the dysvariant breakdown of the total absolute dysvariance is achieved only by a sequence of increasing dysvariance and decreasing extinction areas. In such a pseudotelecentrically polarized dysvariant area, there is therefore no aeondyne branch that contains a televariant structural zone.

On the other hand, if it is possible to specify at least one aeondyne branch that contains at least one televariant zone, then a televariant aeonic area is present, because the location of the telecenters in the tensorium is fixed in a televariant way through the televariant zone. Therefore, the true telecentric polarization of the televariant area must meet this televariance criterion, as opposed to the dysvariant area with pseudotelecentric polarization. All preceding televariance investigations apply to a televariance theory of the transcendence level 0, but this theory also applies to any higher transcendence levels $T > 0$, because all these transcendence fields must also have a threefold transcendental tectonics, where the gradual and syndromatic form corresponds to those of syncolating metroplex combinations in the sense of affinity syndromes, while the telecentric is conditioned by the polarization of the transcendence level 0, i.e., the location of the main telecenters. The whole tectonic investigation of higher transcendence fields is simplified in that there are neither polydrome centers nor collectors in them next to the secondary and main telecenters, because the transcendent aeondynes, because of their property of being affinity syndromes, are always monodromic between the telecenters. The televariance, however, does not necessarily need to be fulfilled by the televariance of the relevant branches syncolating in the affinity syndrome of the next lower transcendence level $T - 1$, because the course in T is solely determined by the course of the affinity syncolator. Thus, all transcendence fields over an area must also have a transcendental tectonics of televariant aeondyne zones, besides the transcendental architectonics, which is defined according to the previous investigations, as for the transcendence level 0.

However, this provides an extension of the tele- and dysvariant theory, in such a way that this theory can also be applied to all zones of the transcendence fields. Since the transcendence fields of panoramas can be related to each other through exogenous transcendental syncolators, such that they syncolate even higher structures, the concepts of main and secondary telecenters appear relative, i.e., dependent on the respective transcendence level. For example, for $T = 0$ the main telecenters of a panorama may be fixed, but there is the possibility that they become secondary telecenters for $T > 0$, specifically when a transcendental area of higher order is created through the syncolation with other areas in higher transcendence. Therefore, if there are $1 = l = L$ transcendental areas $C_{(r)}(ARq_l)_l$ of the transcendence level r and the order q_l , which are not components of an area of higher order but of the same trans-

structure is the key focus in this section.

The discriminant behaves like a boundary, separating areas of increasing and decreasing dysvariance, and it can also go through changes as the parameters vary. Just like how the state of matter can change from solid to liquid at a certain temperature, the aeondyne's structure can also change when reaching certain values in the multidimensional parameter space. This concept is referred to as the extinction discriminant.

Now, let's add an extra layer of complexity: metastability. In physics, a metastable state is a state that is stable for the time being but is not the most stable state available. It's like a ball resting in a shallow dip on a slope—it could stay there indefinitely, but if slightly nudged, it would roll down to a more stable position.

In Heim's syntrometry, the syncolation states in the discriminant boundary can also exhibit this metastable property. While in this state, they remain stable within a certain parameter range, but if the parameters continue to shift, they can change, leading to a change in dysvariance. The concept of stability and metastability is relative to the parameter intervals being considered in the tensorium of the area.

Lastly, this section also highlights a refinement in understanding dysvariance: it can be either structural or functional. Structural dysvariance involves an actual change in the structure of the aeondyne at the point of dysvariance. On the other hand, functional dysvariance retains the structure, but certain syndromes (characteristics) along the extinction interval are undefined. This concept is akin to distinguishing between a physical change (like melting ice) and a chemical change (like burning paper) in ordinary matter.

In conclusion, this section on "Metastable Syncolation States of the Extinction Discriminants" talks about the dynamic and complex nature of aeondynes in Heim's syntrometry, underlining the role of parameters, dysvariance, and the interplay between stable and metastable states.

dence level, then each of these areas has a transcendental architectonics consisting of a gradual, a syndromatic, and a telecentric tectonic component. In addition, however, there can be both endogenous and exogenous transcendental syncolators, which syncolate these L areas into a superordinate transcendence field with televariant structure zones at the transcendence level $r + 1$ and so on. However, this means that besides the transcendental architectonics and the order of the area, there must also be a hierarchical tectonics of televariant transcendental zones, if there are televariant zones in each transcendence field under discussion.

While the three components of the transcendental architectonics, namely gradual in the direction of the metroplex grade, syndromatically orthogonal to this in the direction of a structure zone, and telecentric in the direction of the area polarization, proceed and the order of the area is determined by the distinction between main and secondary telecenters, the hierarchical tectonics of televariant transcendence zones proceed in the direction of increasing levels of transcendence. Only through this hierarchical tectonics does the concept of main and secondary telecenters become relative, and this is related to the next higher transcendence field. Therefore, it must be possible to develop such a telecentric transcendence level relativity, which must allow the most general statements about transcendental areas and their hierarchical tectonics.⁵³

⁵³ChatGPR4.0: Sure, here's a simplified summary of the content presented in Section 6.5: "Televariance Condition of Telecentric Polarization".

The section discusses the criteria for telecentric polarization within an area, building upon previous discussions about televariance and dysvariance.

In the world of 'Areas', different classifications exist for dysvariant structures. This includes absolute total, partial, and others, each of which can appear in initial, final, or intermittent forms, and may be structural or functional in nature. Out of these, the absolute total dysvariance class is unique in that it doesn't allow for affine telecentric polarization - it only has projective telecenters.

A projective telecentric area, defined as the panorama of a polydromic aeodyne, is always determined by absolute total dysvariance. In all other classes, there's the possibility for telecentric polarization of affine areas, but this can be pseudotelecentric if there are no televariant structure zones.

A truly televariant aeonic area exists when at least one aeodyne branch contains at least one televariant zone. This differentiates real telecentric polarization of a televariant area from dysvariant areas with pseudotelecentric polarization.

The section also touches on the televariance theory for transcendence level 0 and suggests that the theory applies to higher levels too. In these higher fields, some simplifications occur due to monodromic characteristics of the transcendental aeodynes, and the absence of polydromy centers and collectors.

It concludes with a proposal for expanding the tele- and dysvariance theory to apply to all zones of transcendence fields. In this framework, the concepts of main and secondary telecenters become relative, depending on the specific transcendence level. This introduces a new requirement for a hierarchical tectonics of televariant transcendence zones, if televariant zones exist in every transcendence field under discussion.

6.6 Transcendental Telecentric Relativity

Since the telecenters of an area are special cases of polydrome centers, these telecenters must also be sub-tensors of the parameter tensorium of the area, on whose metaphorical dimensions the area depends in its course, i.e., these telecenters must also be metaphorically dimensioned. If n is this metaphorical dimension number of the tensorium, then the dimension numbers γ_k of the two telecenters with $k = 1$ and $k = 2$ for the area of the transcendence level $T = 0$ lie in the closed interval $0 \leq \gamma_k \leq n - 1$ ⁵⁴. The impossibility $\gamma_k < 0$ is immediately evident here, while $\gamma_k > n$ is conceivable, but also fails, because in this case the telecenter k would partially lie in a sub-tensorium of the metaphorical dimension number $\gamma_k - n > 0$, which no longer belongs to the parameter tensorium and thus constitutes a contradiction.

In geometric metaphors, the telecenters with $\gamma_k \leq 3$ could be referred to as point (0), lines (1), surfaces (2), or spatial telecenters (3). In general, any γ_k -polarization is present in $0 \leq \gamma_k \leq n - 1$, which should be referred to as symmetric when $\gamma_1 = \gamma_2$, but as asymmetric for $\gamma_1 \neq \gamma_2$. For example, if $\gamma_2 > \gamma_1$, the asymmetric polarization is ambiguous because there is an infinite $\gamma_2 - \gamma_1 > 0$ -tuple set of $\gamma_1 + 1$ -dimensional connecting sub-tensoriums of the telecenters in the parameter tensorium. Such a $\gamma_1 + 1$ ⁵⁵-dimensional telecentric connecting tensorium is always defined as a *Telecentral* when it represents the shortest connection between the two main telecenters in the parameter tensorium in the sense of a geodetic metaphor.

The totality of all Telecentrals in the asymmetric case of $\gamma_2 - \gamma_1 > 0$, with dimension $\gamma_1 + 1$, is therefore a sub-tensorium of dimension $\gamma_2 + 1$. So, the $\gamma_1 + 1$ -dimensional Telecentrals are the generators of a $\gamma_2 + 1$ -dimensional telecentral area, whose syntrometric properties depend on the relative position of the two main telecenters in the parameter tensorium, their dimensions, and the syntrometry of the tensorium. The telecentral area is thus relative with respect to the telecenters and the parameter tensorium.

Only in the symmetric case, because of $\gamma_2 = \gamma_1$, the Telecentral becomes identical with its area and unambiguous due to $\gamma_2 - \gamma_1 = 0$, because the Telecentral, by definition as a geodetic metaphor, connects the main telecenters, and out of the infinitely many possibilities, only one can have this property. In this way, the properties of these Telecentrals and the syntrometry of the tensorium can be determined.

The symmetric case of $\gamma_1 = \gamma_2$ also shows that $\gamma_k = n$ is impossible because then the Telecentral would be outside the tensorium with $n + 1$, which directly results in the interval limit of $\gamma_k \leq n - 1$. Since the main telecenters close off the area, the Telecentral is a good characteristic for the area extension in the direction of telecentric tectonics, relative to a specific syntrometry of the tensorium.

Due to their relativity with respect to this syntrometry, both regular and singular transformations of the syntrometry bring changes to the telecentral area. However, this means that for each monodromic aeondyne of the area, there must be such a transformation of the tensorium, such that this aeondyne lies over the telecentral area after the transformation of the tensorium. Such a transformation of the parameter tensorium is therefore a characteristic of the respective monodromic aeondyne, which implies that every monodromic aeondyne branch of the area must possess such an aeondyne characteristic, and that all these aeondyne characteristics can arise due to telecentral relativity through syntrometric transformations.

Because of telecentral relativity, each area of the transcendence level $T = 0$ thus forms a closed system of mutually transformable aeondyne characteristics, so that each monodromic aeondyne branch, related to the correct characteristic, lies over a telecentral. In addition to this basic relativity of the aeondyne characteristics, i.e., the basic relativity of the telecentrals for $T = 0$, there must also be a transcendent telecentral relativity if transcendence fields $T > 0$ exist due to aeondynic affinity syndromes.

If transcendence fields $T > 0$ exist, the transcendent aeondyne characteristic does not change compared to $T = 0$ in the $T \geq 1$, but the telecentral regarding the main telecenters and parameter dimensions, because nothing changes in these determinations of the area if the syncolations leading to the transcendence fields only act on such aeondynic affinity syndromes that belong to the original area $T = 0$. Therefore, there is also a transcendent aeondyne characteristic and a transcendent telecentral relativity in the first transcendence level for the aeondynes in $T = 1$ that are free from polydrome centers, which arises directly from the base relativity in $T = 0$.

⁵⁴HS: added $\leq n - 1$.

⁵⁵HS: The entire tensorium has the dimension $\gamma_2 + 1$, but the tensors of the set have $\gamma_1 + 1$! The "+1" is the direction of the telecentric polarization. However, there is an error here because it is about a specific tensorium, which represents the telecenter.

On the other hand, if the transcendence field $T = 1$ is syncolized from several areas $T = 0$, then the telecentral relativity valid here and the aeodyne characteristics depend on the syntrometric properties of all areas $T = 0$. This is because, in $T = 1$, their main telecenters become secondary telecenters if the areas are of a higher order, or the telecenters in $T = 1$ become higher-dimensional structures, because each of the areas $T = 0$ syncolized in $T = 1$ can be defined over a different parameter tensorium. Their metaphorical dimensions can differ both in number and in semantics.

Regardless of the specific form of the syncolation of the transcendence field $T = 1$, an aeodyne characteristic and a telecentral relativity in the first transcendence stage always exist in $T = 1$. In a completely analogous way, based on this fact, one can deduce a telecentral relativity in $T = 2$ from $T = 1$, and the complete induction conclusion eventually leads to a general transcendent telecentral relativity with syntrometric aeodyne characteristics in all transcendence fields $T > 0$.

In any transcendence field of level $T > 0$, there is a telecentral relativity of aeodyne characteristics. However, all telecentrals of level $T - 1$ are connected with this telecentral system of level T , whose transcendence fields syncolate the level T and so on. This process can be continued up to the transcendence fields of level $T(T - 1) = 1$ which have been syncolated from the areas of level 0, with each of these areas having a relative telecentral of the same level.

Thus, in every transcendence field of level $T > 0$, there is a whole spectrum of relative telecentrals in terms of transcendence levels. In each transcendence level of this spectrum, there are as many telecentrals of the same level as self-contained systems of aeodyne characteristics as there are transcendence fields that syncolate to the next higher transcendence field. Each aeodyne of a field is located above the relative telecentral, with reference to the corresponding aeodyne characteristic.

In the most general case of many transcendent syncolating areas of any order, there can only be a telecentral and a system of aeodyne characteristics in the relatively last transcendence level T , from which a spectrum of telecentrals of lower transcendence levels runs in the sense of a non-decreasing number sequence to the transcendence level 0, where the maximum of spectral terms is located.

In the original area $T = 0$, in addition to the two main telecenters, there can also exist secondary telecenters, i.e., polydrome centers and collectors can occur, which satisfy the existence condition of the telecenter and thus delineate a partial area structure within the area.

This partial structure is not an independent area because not all monodrome aeodyne branches of the actual area pass through these secondary telecenters. Thus, these polydrome centers and collectors that satisfy the existence condition of the telecenter are considered first-degree secondary telecenters, between which a first-degree pseudo-telecenter can be defined.

Let O_0 denote the degree of these secondary telecenters for $T = 0$. Here, $O_0 = 0$ indicates the degree of the main telecenters, $O_0 = 1$ represents the first secondary telecenters, and so forth. Within the secondary telecentrically bounded partial structure $O_0 = 1$ of the area, further second-degree partial areas may appear if the area is sufficiently differentiated, their secondary telecenters being characterized by $O_0 = 2$, and so forth.

Given sufficient differentiation of the area in the level $T = 0$, a whole internal spectrum of secondary areas $O_0 > 0$ is possible, whose final stages with the maximum value O_0 are the simple alternations of polydrome centers and collectors. The number of possible monodrome aeodyne branches, and hence the number of aeodyne characteristics in the individual partial secondary areas, decreases with increasing degree O_0 .

Such differentiation can also occur in higher transcendence levels $T > 0$, and this occurrence provides a criterion for whether the O_0 from $T = 0$ are really secondary telecenters, because the monodrome aeodyne branches run completely unambiguously between two secondary telecenters in all $T > 0$, without polydrome centers and collectors.

From this, it directly follows that the maximum values of O_0 cannot possibly lie with the simple polydrome centers and collectors. Therefore, if there is also a spectrum of transcendent secondary areas in the transcendence fields $T > 0$, whose secondary telecenters run the degrees $O_T \geq 0$, then there is always an equivalent spectrum of transcendent pseudo-telecenters of degrees $O_T \geq 0$, over which, corresponding to the individual aeodyne characteristics, the transcendent monodrome aeodyne branches of the respective partial structure in $T > 0$ stand.

In the case $O_T \geq 0$, there are no secondary telecenters at all, in no field $T \geq 0$, and only one telecenter. On the contrary, if $O_T = 1$, there must always be an even number of μ secondary telecenters and a system of $\frac{\mu}{2}$ first-degree pseudo-telecenters, etc.

If it is a transcendence field structure over an area $T = 0$, then the maximum value for O_T in all stages $T \geq 0$ is

identical, which is due to the peculiarity of the synkolation of affinity syndromes. This law is obviously broken when multiple systems synkolate in this form, yet due to their definition, the secondary telecenters must reappear in each transcendence field.

Therefore, a principle of diabatic projections must apply to the subareas through all stages of transcendence, where the concept of perspectival mapping cannot be metaphorically transferred to these areas. Since each telecenter in any stage $T \geq 0$ always continuously connects the telecenters $O_T = 0$, it becomes evident that each secondary telecenter between $O_T = m > 0$ must continue into the next higher area $O_T = m - 1 \geq 0$, so that all these continuations eventually form the main telecenter between $O_T = 0$.⁵⁶

⁵⁶ChatGPT4.0: Alright, let's recap Section 6.6, Transcendental Telecentric Relativity, in the style of Richard Feynman.

We've got this idea of areas, right? These areas, or spaces, are defined by something we're calling telecenters. And there's some level of hierarchy to these telecenters - we have main ones and these secondary ones. We're working with this idea of "Transcendence fields" and "aeondyne branches", and we've seen how they relate to these telecenters.

Now, we've taken this notion of a single area, and we've imagined taking several areas and bringing them together, something we're calling synkolation. When we synkolate, we're essentially creating a new area or a new field of transcendence. And that has implications for these telecenters. Suddenly, we have telecenters in multiple dimensions and areas with differing characteristics.

And it gets a bit more complicated when we look at the highest levels of transcendence. At the top, there are no secondary telecenters, only a single telecenter. But as we move down in levels, the number of secondary telecenters starts to increase. And these secondary telecenters have their own pseudo-telecenters.

There's this concept of 'degrees' too. When we look at secondary telecenters, we have different degrees and it all depends on the level of differentiation of the area. But this idea of degrees extends beyond that, to what we've called transcendental pseudo-telecenters.

But here's where it gets fun. We found that each telecenter at any level, always connects the telecenters of degree zero. This means that every secondary telecenter must continue into the next higher area, until we reach the main telecenter. It's a bit like peeling an onion, layer by layer, until we reach the core.

So, at the end of the day, we have this complex system of areas, telecenters, aeondyne branches, and transcendence fields, all interconnecting and influencing one another. It's like a cosmic dance of structures and elements, with each part playing a crucial role in shaping the whole.

SYNTROMETRISCHE MAXIMENTELEZENTRIK SYNTROMETRIC MAXIMAL CENTERING

BURKHARD HEIM

Part B: Anthropomorphic Syntrometry

Translated by ChatGPT 4.0 from input by Lyle Anderson

7 Anthropomorphic Syntrometry

7.1 Subjective Aspects and Apodictic Pluralities

The preceding syntrometric investigations are not tied to any specific aspect system, that is, they are applicable in any such system, and are also not bound to the ambiguously predicative aspect complex, which is an expression of the specific structure of the anthropomorphic intellect. This universality of the syntrometric statement is solely due to the property of syntrometric elements being syntrices, that is, their predicate connections must be universal quantifiers. Only through this does aspect relativity arise, which makes a syntrometric statement independent of the specific aspect complex, but which must also be correct for the ambiguously predicative aspect complex of anthropomorphic statement possibilities. It immediately follows that there must be an anthropomorphic syntrometry that must fundamentally encompass all possible statements of this aspect complex. Therefore, if the anthropomorphic intellect, whose expression capability is the ambiguously predicative aspect complex, is applied to any conceptual system, then if the subjective aspect was chosen correctly, the corresponding statement system must always be able to be brought into a version of this anthropomorphic syntrometry, which then also has validity in any other aspect complex, because all syntrometric predicate connections are universal quantifiers.⁵⁷

The anthropomorphic aspect complex is characterized by the specific property of the anthropomorphic intellect to make dual contradictory statements in the sense of comparisons. In this way, the predicator always contains only two discrete statements, namely the positive $\bar{\bar{||}}_+$ with respect to the relevant comparison, and its contradictory negation $\bar{\bar{||}}_-$. The totality of all these possible dialectic and coordinate systems of predicator $\bar{\bar{||}}_{\pm}$ forms the entirety of all subjective aspects in the *elementary aspect system* of the anthropomorphic aspect complex. In the aspect system of the first *statement level*, statements in the sense of probability predicates about $\bar{\bar{||}}_{\pm}$ are possible, that is, the statement $\bar{\bar{||}}_{\pm}$ is rated with a probability $h_{\pm}$ in the closed interval $0 \leq h_{\pm} \leq 1$, which allows for the complementary rating $h_{\mp}$ for the contradictory statement $\bar{\bar{||}}_{\mp}$, but always $h_{\pm} + h_{\mp} = 1$ must be fulfilled. The aspect system of the second level, in turn, evaluates in a completely analogous manner the two complementary predicate bands of the first-level aspect system, etc., whereby the law of complementarity $h_{\pm} + h_{\mp} = 1$ must always apply when the predicate bands are probability statements about the predicates of the previous statement level, but a probabilistic character does not necessarily need to be required.

Any variable, for which there exists a complementary law, must be capable of becoming a sequence of aspect systems of higher statement levels (briefly referred to as an aspect sequence). The anthropomorphic aspect complex, over which an anthropomorphic syntrometry can be developed, is thus characterized by an elementary aspect system with the two-valued contradictory and discrete predicator $\bar{\bar{||}}_{\pm}$, from which as many aspect sequences arise as complementary laws can be formulated. While the predicator of the elementary aspect system is always discrete, a predicator of any aspect system of a higher statement level must consist of two continuous predicate bands that are related according

⁵⁷ChatGPT4.0: Alright, now let's take a step into a new terrain – Anthropomorphic Syntrometry. Here's what it's all about: In the stuff we've talked about so far, we've been dealing with syntrometric investigations. These investigations have been pretty much universal, they weren't tied to any specific perspective or system, and they weren't even tied to a certain complex of aspects tied to our human intellect. That universality is rooted in a unique characteristic of syntrometric elements – they're syntrices, meaning their interconnections are universal quantifiers. These universal connections allow for a kind of "relativity of perspectives" that remains valid across various aspect complexes, including the two-valued predicative aspect complex that characterizes our human way of thinking.

But here's the kicker: from this universality, it follows that we must have a "human version" of syntrometry, which can encompass all possibilities of our human way of looking at things. So, if you take our human intellect, characterized by this two-valued predicative aspect complex, and apply it to any system of concepts, and you've chosen the subjective aspect properly, you can always translate the resulting statements into a form of this "human" syntrometry. And here's the cool part – this translated form is still universally valid, across any perspective or aspect complex. Why? Because these syntrometric connections are still universal quantifiers. So, we end up with a version of syntrometry that's tailored to our human way of thinking but still maintains its universal validity. It's quite something, isn't it?

to the associated complementarity law. All these aspect sequences only ever make complementary statements about the predicates of the previous statement level, while their dialectic is given by the respective connection of the complementary predicate bands. In this way, however, nothing changes at the discrete diatropes and coordinations, i.e., at the subjective aspects of the elementary aspect system, so that the entirety of all subjective aspects of this system is characteristic of the anthropomorphic aspect complex.

However, anthropomorphic Syntrices are possible over each of these subjective aspects S , which must consist of conceptual elements that can be brought into relation by the anthropomorphic intellect in the sense of a comparative statement $|\overline{S}|_{\pm}$. Here $|\overline{S}|$ means that the statement $||$ was dialectically shaped by the subjective aspect S in the sense of the respective coordination. However, each anthropomorphic Syntrix must have a Metaphor of apodictic elements, so that there must be a plurality of apodictic elements over the discussed aspect complex. This apodictic plurality is characterized by the comparability of its elements in the sense of predicate alternations, and this peculiarity of alternative comparability reveals a characteristic property of the *apodictic plurality*. In the anthropomorphic sense, only elements with qualitative or quantitative properties can be compared alternatively. For this reason, the entire apodictic plurality is structured by two classes of apodictic elements, the *Quality* and the *Quantity*. While Quality contains all conceptual elements that differentiate qualitatively, Quantity encompasses the elements definable by the concept of number. It follows directly from this that Quantity can only be defined over a single subjective aspect, which is already determined by the concept of Quantity, for Quantity determines the dialectic of the aspect as set dialectic. All other subjective aspects of the elementary aspect system, on the other hand, make the description of Quality possible.

If any fact is to be captured syntrometrically, this capture must be preceded by a description within the framework of anthropomorphic syntrometry, from which the greatest possible precision must be demanded. Since it is in the nature of anthropomorphic intellect to grasp the most precise criteria in the validity range of set dialectics, that is, over the aspect of quantity, it seems expedient to formulate anthropomorphic syntrometry over this subjective aspect and to derive the elements of anthropomorphic analysis from this syntrometry. This deduction must ultimately lead to a transition criterion, which indicates which demands a quantitatively analytically formulated fact must meet if it is to be transferred into anthropomorphic and finally into general syntrometry. The plurality structure of quantity can only be related to the subjective aspect m of a set dialectic, whose coordination has the character of a set comparison, if the elementary aspect system is taken as a basis. The dialectically shaped predicates of this quantity aspect m can therefore only express the equality of the sets $|m|_{+} \equiv$ or the set inequality in $|m|_{-} \equiv \neq$. If there is set inequality, then the statement $a \neq b$ applies to the two sets a and b , then there are again two possibilities. Either a is quantitatively greater than b , or vice versa, which in the case $a \neq b$ is symbolized by $a > b$ or $a < b$. For $a \neq b$, the ambiguity ($>$) or ($<$) can again be conceived as a contradictory Predicatrix in the sense of a statement and its complementary negation.⁵⁸

⁵⁸ChatGPT4.0: Alright, let's have a crack at it, trying to keep Feynman's flair for simplifying complex concepts. We've been exploring an interesting idea in this section, you know. Heim's calling it 'Anthropomorphic Syntrometry' and it's a fancy name for something we do every day - compare things and make judgements. It's all about how our brain, or let's say 'human intellect', can look at two ideas and say, "Hey, this thing here, it's greater than that thing over there." Or maybe it's not, and that's where the fuzziness, the probabilities, come in.

Heim tells us that we operate with two main choices - let's say, a positive and a negative statement - and we create all sorts of complex systems out of just these two choices. We even assign probabilities, like saying "I'm pretty sure this statement is true, let's give it a probability of 0.8 and the opposite one, a 0.2". That's how we make sense of our world, navigating through all this uncertainty.

But it gets even more interesting. Heim suggests that we can stack these systems, creating hierarchies of fuzzy comparisons. He goes on to say that we naturally organize our thoughts and comparisons into two categories: quality and quantity. Quality refers to things that are different in nature, while quantity is all about numbers, countable things. It's the old apples to oranges comparison again.

Now, when we're trying to understand something, in the most precise way possible, we usually lean towards the quantity side. It's because we can measure and compare it, and that's a powerful tool. This concept, in Heim's words, forms the basis of our 'anthropomorphic syntrometry'.

Once we have a good handle on this quantity-based comparison system, we can then start making more detailed statements. For instance, we can say if a set of things is equal, greater, or less than another set. Then we use this tool to analyze and describe our world, and if we're careful and precise, we can even convert these understandings into more general, universal statements.

It might sound a bit far out, but really, it's just a formalized way of describing how we humans, with our messy and wonderful brains, make sense of the world.

7.2 Structure and Interpretation of the Quantity Syntrix

Because of the set dialectic, only quantities can be compared over the quantity aspect. However, there can fundamentally only be a single method for describing a quantity, namely the evaluation of quantities independent of the relevant quality by numbers, so that the concept of numbers is the actual apodictic idea of all categories of the quantity aspect. Numbers can always corporize in two ways, which follows directly from the set dialectic, namely because of the possibilities $a > b$ and $a < b$ in the sense of a number set increase or decrease. This directly results in the elementary corporizations over the quantity aspect as elementary number operations. The number set increase can in principle only be carried out as number addition (+) or as number multiplication ($\times$), while the number set decrease becomes possible through the inverse operations of number subtraction ($-$) and number division ($() \div () \equiv \frac{0}{0}$).

It is quite evident that these numbers are the only apodictic elements over the quantity aspect, since there are qualitatively different pluralities of numbers. If the indices k and l denote the numbers a, b and c as belonging to two such qualitatively different pluralities, then a functional relationship in the sense of a functor $f(a_k, b_k) = c_k$ or $f(a_l, b_l) = c_l$ can always exist, if the instruction f is built from the four basic operations. However, in this way, elements of k can never form those of l , or vice versa. From this it directly follows that by applying the four basic operations on the elements of a quantitative plurality, they can always only deliver elements of the same plurality. The entirety of all numbers possible in this way is then defined as an algebraic number field. In this way, the concept of apodictic elements over the quantity aspect can now be refined, because every algebraic number field a_i must be an apodictic element over this aspect. If there are $1 \leq i \leq m$ number fields a_i , then they can be summarized into a Metrophor $\tilde{a} = (a_i)_m$.

Since the individual a_i are entire number fields, the Metrophor elements over the quantity aspect fundamentally form apodictic bands, hence any quantity syntromic that emerges from such a Metrophor must therefore be a band syntromic. Every number as an evaluation of a quantity can either be used undimensioned without semantic assignment, or a semantic dimensioning is assigned to it. Since this applies to individual numbers, it must also be correct for all numbers of an algebraic body and therefore for all number bodies. But these bodies are the elements of $\tilde{a}$, so a distinction must be made between two basic metaphoric forms. In the non-semantic case, the individual elements in $\tilde{a} = (a_i)_m$ appear undimensioned and individually, i.e., $\tilde{a}$ without semantic assignment is a *singular Metrophor*. In the semantic case, on the other hand, generally $n \geq m$ quantities are evaluated by the m elements, so that in the case $n > m$ some elements must iterate. If S_n is the iteration rule that assigns semantic dimensionings to all $n \geq m$ algebraic number bodies after the iteration, then the action of this semantic iterator on the singular Metrophor according to $S_n, \tilde{a} = R_n$ delivers the semantic Metrophor $R_n = (y_l)_n$ with $n \geq m$.

Since there are singular and semantic Metrophors over the quantity aspect, the quantity syntromics emerging from them must also be of a singular or semantic nature. In any case, the Metrophor elements are algebraic number bodies, so that the synkolators defining the syntromic alongside a Metrophor must be those functors that must be composed from the four basic algebraic operations, and according to their synkolation level, bring various algebraic number continuums (as such the number bodies can be designated) into functional relationships. These functions, which are conditioned by the synkolators, then form the syndromic assignments of the quantity syntromic.

To iterate this quantity syntromic, it must be taken into account that the semantic form is implied by the singular when the syntromic is understood as a complex structure and the *semantic iterator* is used as a synkolator of the first syndrome. Due to $S_n, \tilde{a} = R_n$, the syndrome 0 would then be given by the singular Metrophor $\tilde{a}$, but the first syndrome by the semantic Metrophor R_n . Since the elements of R_n are also apodictic (S_n essentially only iterates), it is sufficient in the following to only examine the semantic form $\tilde{a} = \langle f, R_n, m \rangle$. The quantity syntromics must also comply with the general syntrometric laws, that is, here too there can only be four classes of pyramidal elementary structures, from which all other pyramidal and homogeneous syntromics are built.

For these reasons, it is sufficient to assume a pyramidal elementary character for $\tilde{a}$. The fact that f in $\tilde{a}$ can only be constructed from the four basic algebraic operations allows for a significant simplification. If f is homometral in the stage $m \leq n$, then several elements in $f(y_j)_1^m$ are identical. But now, these elements y_j are number continua, which are functionally related through the four basic operations, the functional dependency can always be reduced to $f(y_j)_1^m = F(y_l)_1^p$ with $p < m$ if f acts homometrally. The synkolator F , reduced to the level p , must then be heterometrally, from which it immediately follows that the homometral symmetric, as well as the homometral asymmetric elementary structures, need not be considered in a syntrometric analysis over the quantity aspect because

they always reduce to heterometral forms of lower synkolation stage due to the construction of all synkolators over this aspect.

The pyramidal elementary structure $\widetilde{a}$ can therefore only be heterometral, symmetric, or asymmetric. After this reduction of the four classes of elementary structures to only two, there is the possibility of interpreting semantic syntrices, which must also apply to the singular forms. If the semantic iterator does not iterate at all, so that the occupation of R_n becomes identical to that of $\widetilde{a}$, then the semantic syntrix coincides with the singular syntrix. Firstly, it must be established that the elements of R_n are independent numerical continua, such that in each case n numbers indicate a numerical position relative to the n continua of R_n , when each of these continua contributes a number to this position specification. From the definition of the apodictic elements over the quantity aspect, being numerical continua in the sense of algebraic number fields, it directly follows that there must be both the unit E and the void 0 in every element of R_n . But this void must coincide for all a_i from $\widetilde{a}$ and thus also for all y_l from R_n , so that all y_l , despite their independence from each other, originate from the same void 0 .

All numerical continua y_l thus have the same origin 0 , and a unit is defined in each continuum, so that due to the also valid independence, n continua can be used as a reference system for numerical position specifications. However, each coordinate of such a coordinate system is identical to a dimension of an abstract space, because the tensorium of all n -fold position specifications must be defined as an n -dimensional space, whose points are these dimension specifications. Each semantic metaphor R_n is thus to be interpreted as an n -dimensional abstract space, which is induced from the singular metaphor $\widetilde{a}$ by the action of the semantic iterator S_n according to

$$S_n, \widetilde{a} = R_n, \widetilde{a} = (a_i)_q, \widetilde{a} = \langle f, R_n, m \rangle \quad (28)$$

For the synkolation level, it must fundamentally be the case that $m \leq n$, because there can be no homometral quantity syntrices according to the preceding conclusion. If $m < n$, f selects in the first syndrome a R_m , thus an m -dimensional subspace from the n dimensions of the R_n .

These m dimensions are put into a functional relationship by f in such a way that the respective subspace R_m appears as the argument range of the $f(y_j)_1^m$ function. If f is symmetric, then the first syndrome is fully occupied with $\binom{n}{m}$ synkolations in the sense of such functions, because there are the same number of m -dimensional argument ranges that can be separated as subspaces R_m from R_n . The occupation of the first syndrome of a quantity syntrix thus consists of a certain number of $(m + 1)$ -dimensional functions f over m -dimensional argument ranges, which assign a function value to each point of the argument range within their definition interval, i.e., each of these functions describes a structure defined as a field within its argument range. While the semantic iterator induces an abstract space as a semantic metaphor from the singular metaphor of a quantity syntrix, the synkolator of the first syndrome, according to its synkolation level, delimits subspaces in which it structures fields as argument ranges, which then occupy the first syndrome. The synkolator of the second syndrome then relates these field functions to each other, and so on.

The *semantic iterator*, as well as generally any synkolator over the quantity aspect, is nothing more than an operator that represents a rule for how the elements of the preceding syndrome occupation are to be put into a functional relationship. Thus, the general concept of a functor over the quantity aspect becomes the concept of a functional operator. Due to the property of each quantity syntrix being a band syntrix, because the apodictic elements are continuous algebraic number bodies, the synkolations of all syndromes also form number continuums in the sense of structures described by the relevant synkolation law. Since it is in the nature of every set of numbers that a unit and the empty spot 0 are defined as a number, there is a fundamental possibility to orient all sets of numbers, which applies to both the structure continuums of the syndrome occupations and the apodictic elements of the semantic metaphor R_n . However, if a quantity, in addition to its magnitude and semantic dimensioning, also specifies a direction conditioned by the orientation, along with a sense of direction, then a vector as an oriented quantity is given. In contrast, non-oriented quantities should be referred to as scalars. While the elementary operations of set enlargement and their inverses apply to these scalars, the concept of multiplication for vectors must be refined, because when two non-parallel vectors originating from a point are multiplied together, the multiplication can either be scalar, i.e., the projection of one vector onto the other is multiplied as a magnitude with the magnitude of the other vector, or the multiplication is tensorial, i.e., one vector is projected onto the normal to the other vector, and the magnitude of this projection is multiplied with the magnitude of the other vector.

According to its nature, the scalar product loses the orientation, i.e., if $\bar{a}$ and $\bar{b}$ are two vectors, their scalar product $\bar{a} \cdot \bar{b}$ is a scalar quantity, but their tensor product $\bar{a} \times \bar{b}$ is an oriented quantity, whose orientation is determined both by that of $\bar{a}$ and that of $\bar{b}$. In the case of tensor multiplication, if the factors are independent of each other, i.e., projectable onto the coordinates of the reference space being used, then the tensor product is defined as a tensor, whose tensor degree is identical with the number of independent vectors involved in the product. The symbol ${}^m\bar{A}$ identifies A as a tensor of degree m . This concept of tensor evidently implies the concepts of the vector and the scalar size, because for $m = 1$, the tensor apparently degenerates to a vector, and for $m = 0$ to a scalar. If n is the dimension number of the coordinate space being used, i.e., the number of apodictic elements of the semantic metaphor, then it immediately follows from the tensor definition for the possible tensor degrees in this coordinate space the interval $0 \leq m \leq n$, because $m > n$ is not possible based on the tensor definition. All syndrome occupations of the quantity syntrix, i.e., all synkolation, therefore have a tensorial character, and from this follows immediately an essential property that must be demanded of any tensor quantity.

A band syntrix is evidently only defined if the syndrome occupations do not change during a deformation of the apodictic continuums. However, in the quantity syntrix, these apodictic continuums are the continuous algebraic number bodies, and deformations related to the linear arrangement in the semantic metaphor would correspond to coordinate transformations. But since, due to the nature of the band syntrix, it must be demanded that the syndrome occupations should not change during such deformations, and this demand corresponds to a condition of invariance of the syndrome occupations against transformations of the semantic metaphor, this condition of invariance must also be fulfilled by the synkolation. However, the synkolation are tensors, so that from this interpretation direction of the quantity syntrix, the invariance of the tensor against certain groups of coordinate transformations directly follows.

Following the previous investigations, the synkolation level m of a synkolator f must be interpreted as the dimension number of an argument area, which should be understood as an m -dimensional subspace of R_n , so that the tensor grades of the synkolation of this level can reach at most the value m . The synkolator f puts these m algebraic continuums in a functional relationship, such that each point in the m -dimensional argument area is assigned a synkolation state, and the totality of all these synkolation states forms a synkolative structural continuum. Such a structural continuum is to be defined as a field of the respective synkolation state. This field is therefore completely described by the synkolator which assigns a synkolation state to each point in the m -dimensional argument area (field area). The field structure thus has the dimension number $m + 1$, and its reference space is created by increasing the m dimensions of the field area by the one of the synkolator. In this synkolation dimension, the synkolation states are then plotted over the field area, creating the field structure.

For this reason, the reference space can be defined as an $(m + 1)$ -dimensional synkolator space of the $(m + 1)$ -dimensional tensor field, whose arguments lie in the m -dimensional field area. As already mentioned, the synkolators are functional operators that act on number continuums, which implies that the field progression is continuous except for a finite number of singularities in the sense of extremes. A suitable extremum, which can be at most m -dimensional, can be selected and made the reference area of the entire field by parallel shifting of the reference space. This distinguished extremum then becomes the field center. If μ is the dimension number of such a field center, then μ can, based on the definition of the center, also be a field singularity analogous to the interval of tensor grades only in the interval $0 \leq \mu \leq m$. The field functions, that is, the synkolators operating at level m , have a continuous course, except for a finite number of singularities, since they are to be interpreted as a functional operator.

For this reason, there must be isocline areas in every field structure, over which the field function has a constant value. These isoclines can then be projected into the m -dimensional field area, where they define a family of hyper-surfaces of the respective dimension number $m - 1$, which are referred to as level surfaces, and enable a topographic image of the field structure in the field area. All previous investigations of synkolated tensor fields refer to the first syndrome, but they are so general that they can be meaningfully applied to all other syndromes. The synkolator of the second syndrome (if the syntrix is complex, it may already be different from that of the first syndrome) does not take the number elements from the apodictic continuums in the case of the general pyramidal syntrix, but from the tensor occupancy of the first syndrome, and so on. This directly implies that the synkolators of all syndromes beyond the first are tensorial functional operators, which set the tensor field structures from the occupancy of the previous syndrome into functional correlations, the number of correlating tensor fields depending on the synkolation level.⁵⁹

⁵⁹ ChatGPT4.0: Alright, let's kick back a bit and talk about this thing called the Quantity Syntrix, trying to approach it with a bit of that Richard Feynman vibe.

7.3 Syntrometry over the Quantity Aspect

The quantity aspect in the anthropomorphic aspect system is characterized by the two set-comparing predicates $=$ and $\neq$, as well as by the set-changing basic operations of addition and multiplication together with their inverses of subtraction and division. After describing and interpreting the syntropies possible over this aspect, the elements of a syntrometry over the quantity aspect are to be derived. According to the preceding, there is for each singular metaphor a set of semantic iterators, which induces semantic metaphors as reference spaces. Each tensorial functional operator can be used as a synkolator, or in the complex case as a system of synkolators, when the tensor grades, as well as the functional relationships of the dimension number of the reference space, i.e., the occupation of the semantic metaphor, is adapted. From this it becomes clear that there must be as many R_n for each singular metaphor as semantic iterators can be defined. Furthermore, there exist as many quantity syntropies for each such induced R_n as complex synkolators can be pre-specified in the sense of tensorial functional operators. In this way, it becomes clear that a single singular metaphor over the quantity aspect can generate a multiply infinite set of syllogistically oriented syntrix structures.

All apodictic elements of singular metaphors are, according to the preceding investigations, algebraic number bodies, thus apodictic continua, and therefore the R_n point continua. The quantity syntropies must therefore be at least band syntropies. Since the algebraic number bodies do not need to be limited, an R_n can also be conceived as a numerically parametric tensorium, to which the syntrix represents a primeval aeondyne, especially since the terms discussed in the quantity aspect are always numerical elements. However, for each R_n there are as many such aeondynes as complex synkolators exist, thus a multiple infinite set. Because of the necessity of interpreting the quantity syntrix as an aeondyne, it follows directly that this set of syntropies must be coordinated from every point of the R_n . Thus, each point of an R_n carries a multiple infinite set of syntropies, and each R_n induced from a singular metaphor thus becomes a parametric tensorium as the carrier space of a multiplicity of primeval aeondynes. Each semantic metaphor must therefore be addressed as such an aeondynic carrier space. Furthermore, it follows from the theory of the primeval aeondyne that the apodictic continua y_i of the R_n are all defined over one-dimensional arguments $n_i = 1$ according to $y_i(x_i)$, because these continua are represented by algebraic number bodies. However, the functions $y_i()$ can in turn be considered as synkolators, which transform the number bodies, so that for the coordinates of the R_n always the undeformed $y_i(x_i) = x_i$ can be used as a basis. Since there is in each algebraic body besides the unity E also the absence 0 , and all x_i start from 0 and are unlimited, the semi-open intervals $0 \leq x_i \leq \infty$ apply for these x_i , i.e., the aeonic lengths are identical with the extensions of the algebraic bodies. In this way, however, the semantic metaphor of this primeval aeondyne, namely $R_n = (x_i)_n$ with $0 \leq x_i \leq \infty$ is completely determined. If, moreover, f is a complex synkolator of tensorial functional operators, whose level in the first syndrome always lies in an interval $1 \leq m \leq n$, then the quantity syntrix appearing as a primeval aeondyne is

$$(\tilde{a}|) = \langle f, R_n, m \rangle = \tilde{a}|(x_i)_1^n, R_n = (x_i), 0 \leq x_i \leq \infty. \quad (29)$$

The metaphorical quantity syntrix described in this way is therefore always n -loop and real, because its argument domain can only be the respective carrier space R_n , whose coordinates traverse semi-open intervals. While there is a restriction regarding the synkolation form of the pyramidal elementary structure that the homometral form is eliminated, the primeval aeondynes of the quantity syntrix are at least metaphorical, but also full-loop or synkolative, taking into account that the metaphorical and synkolative form are special cases of the full-loop structure. If it is

So, you got yourself this thing called a Quantity Syntrix. This fellow is just a big bundle of potential field structures that can be whipped up under the "quantity" aspect of our human-centered perspective of the world. To keep our heads straight, we've got a guide, or a map if you will, called the "semantic metaphor", which we treat as an abstract space, kind of like our old friend R_n from math class.

Now, let's take a stroll down the first syndrome of our Syntrix. This little guy is the star of the show, the trailblazer if you will. He's the one inducing field areas right from R_n and his product, the tensor fields, set the stage for the subsequent syndromes.

You see, the other syndromes are a bit like the roadies in our rock 'n roll tour. They take these tensor fields, work their magic (a little process we call synkolation), and get them ready to play their part in the next show, i.e., the next syndrome. It's like a relay race, each handing over the baton to the next.

But there's a catch! As we move along, the dimensionality of our field areas can't just do whatever it wants. No sir, it plays by the rules. It can either stay put or level up as we go along to the next syndrome. But drop down a level? Nope, can't do that. It can climb up until it hits that maximum limit set by n , and that's as high as it can go.

So, in a nutshell, our Quantity Syntrix is like a big circus tent. It's got room for all sorts of tensor structures that can do their tricks under the quantity aspect of our world view. And all these tricks are possible over our roadmap, the semantic metaphor, which we like to think of as an abstract space. A Quantity Syntrix is one heck of a show - it's got all the potential field structures that you can dream up within the framework of the quantity aspect of our human view of the universe. It's a wild ride, but hey, that's what makes it fun, right?

assumed that the structure is full-loop, and that the R_n is connected by some degree of linkage with an N -dimensional synkolation tensorium R_n , then the most general form would be given. However, both the coordinates of both spaces are number continua because of the quantity aspect, and the functors over this aspect are functional operators. If the coordinates of the R_n are denoted with y_l , and if $f(D_m(y_l))_1^K)_1^L$ with $K \leq N$ and any L is any synkolator over the synkolation space, then it follows directly from the nature of the operators and the number continua $f = F, (y_l)_1^K$, i.e., the separability of the variables, where F consists only of analytical operation rules. This separation creates in the new synkolator F a possible asymmetry with respect to the effect on the y_l , and the synkolation stage m of f increases to $m + K$ in F . Such a separation of the coordinates R_n means, however, that a not necessary but possible extension of the singular metaphor, as well as the semantic iterator, can be made, which allows a semantic metaphor R_p with $n \leq p \leq n + N$ to emerge. Thus, it became possible to reduce the full-loop and synkolative primeval aeondyne structure to the metaphorical one, based on the nature of the number continua and operators over the quantity aspect. In the context of a syntrometry over the quantity aspect, there are therefore only metaphorical primeval aeondynes according to equation (29), whose quantity syntropies can always only be heterometral, but can be symmetric or asymmetric.

It is also expected that the concept of the corporator over the specific aspect of the quantity of point sets will be limited. Corporations can only be a composition or coupling of number continua or quantitatively acting operators, meaning, the coupling rules, like the synkolating operators, can only be composed of basic operations. Since the quantity aspect is characterized by the predicates of set equality or set inequality, the syntrix corporations must be unambiguous with respect to the corporated structures, which means that $\begin{Bmatrix} K_s & C_s \\ K_m & C_m \end{Bmatrix}$ certainly does not contain the component C_s , because functional operators can only be unambiguously coupled in structure, not composed. Contrarily, C_m is always possible, because if the number continua of two semantic metaphors R_p and R_q are composed through C_m , an R_{p+q} is created, which is also true for the singular metaphors. The conflict nodes of the couplings, however, are always unambiguous because K_s links the complex synkolators through a system of basic operations, while K_m connects the coordinates of R_p and R_q in the same manner. Therefore, the universal corporator is reduced by the quantity aspect to $\begin{Bmatrix} K_s & K_s \\ K_m & C_m \end{Bmatrix}$, of which only the special cases $\begin{Bmatrix} K_s \\ K_m \end{Bmatrix}$ and $\begin{Bmatrix} K_s \\ C_m \end{Bmatrix}$ unambiguously and universally exist, while $\begin{Bmatrix} K_s \end{Bmatrix}$ only in the case of identical metaphors and $\begin{Bmatrix} K_m & C_m \end{Bmatrix}$ as well as $\begin{Bmatrix} K_m \end{Bmatrix}$ or $\begin{Bmatrix} C_m \end{Bmatrix}$ in the case of identical complex synkolators are structurally unambiguously applicable. The total of the 15 different general types of corporators is therefore reduced over the quantity aspect to only three types of corporators, namely $\begin{Bmatrix} K_s & K_s \\ K_m & C_m \end{Bmatrix}$ as well as $\begin{Bmatrix} K_s \\ K_m \end{Bmatrix}$ and $\begin{Bmatrix} K_s \\ C_m \end{Bmatrix}$, which already include the special cases for corporated syntrices with identical metaphors, or identical synkolators. The conflict nodes of the couplings always consist of systems of basic operations, and C_m increases or decreases the number of dimensions in the corporation. Regarding K , the operations that increase the set are referred to as cooperative and their inverses as contraoperative couplings. Accordingly, C_m is co- or contraoperative if the diameter of the metaphor of the corporated syntrix is increased or decreased through the composition C_m .

In all three basic types of corporators over the quantity aspect, a metaphorical corporation share is always defined, so that the existence criterion of an eccentric is always met. However, this means that over the quantity aspect, confluxive syntrices are always possible, whose conflexion fields are occupied with the corporations of synkolation fields, which in turn formed the syndrome occupations of the non-corporated syntrices before the eccentric corporate process.

Since, due to the analytical operator nature of all synkolators over the quantity aspect, there are heterometric pyramidal elementary structures in symmetric or asymmetric form, the syntrix memory over the quantity aspect must have degenerated into a metaphorically two-dimensional manifold of syntrices. Such a degeneration then has the immediate consequence that all totalities of quantity syntrices are also two-dimensional, while the corporator simplex can only contain three basic classes of corporators. This double degeneration of the memory of the quantity syntrix means simplification due to the specialization of the subjective aspect, but no limitation, because in addition to the regular planar syntrix framework, there is the extra-regular assignment, which can generally be very extensive, because

a large number of corporator chains can be formed from the three corporator classes of the simplex, depending on the nature of this simplex, especially since the corporator identities in a chain formation are always possible. With regard to the structuring of these totalities, there are no possibilities for specialization, because both continuous and discrete totalities are possible, with the discrete forms only appearing when there is a discrete selection rule in the corporator simplex.

Indeed, every quantity syntrix is always the functional value of a primigenous aeondyne, because every semantic metaphor consists of semi-open apodictic continua, which are always algebraic number fields. Accordingly, every syntrix totality over the quantity aspect is a totality of such functional values, i.e., every syntrix totality must be supplemented over this aspect to a continuous band of totalities, thus to a totality of primigenous aeondynes. Over the quantity aspect, therefore, there are only two-dimensional totalities of metaphorical primigenous aeondynes, whose generative can only be formed by three classes of corporators. The memory contains all possible pyramidal elementary structures, and thus all possible singular metaphors, thus the entirety of all algebraic number fields. Furthermore, in R_n of a semantic metaphor, the metaphor diameter n can cover all natural whole numbers $n < 0$, so that there are an infinite number of semantic iterators for the entirety of all singular metaphors. According to this result, the carrier space of each totality of primigenous aeondynes must have an unlimited number of dimensions, because the dimension number of this carrier space is composed of the dimension numbers of all parameter tensoria of the individual aeondynes, and these parameter tensoria are identical to the semantic metaphors in the case of the underlying quantity aspect.

By definition, the value of 0 belongs to every algebraic number field, that is, to each coordinate of the R_n . Since this value is common to all coordinates, it always forms the coordinate origin, regardless of which coordinates of the R_n it is related to. Since the R_n as a semantic metaphor over the quantity aspect is at the same time the parameter tensorium of the aeondynes, there must be a syntrix with the metaphor 0 for every R_n as a characteristic of the quantity aspect, whose elements all have the value 0. For the same reason, a so-called null operator $\pm_0 = 0$ can fundamentally be defined for each synkolation level, so that the existence of the null syntrix for all metaphors $\tilde{a} \neq \tilde{0}$ is guaranteed. The possibility of $\tilde{a} = \tilde{0}$ is not in contradiction with the general principles of syntrometry, because in an algebraic number field, the value 0 is not a void but an element of the respective number field.

All possible syntrix totalities over the quantity aspect, according to the preceding analysis of this aspect, can only be two-dimensional totalities of metaphorical primigenic aeondynes. This immediately implies that none of these totalities can be continuous, as all R_n differ either by the integer index n , by the semantic iterator, or by the singular metaphor. However, these determinants are discrete quantities that cannot transition continuously into one another. If there are no continuous syntrix totalities, then the existence condition of continuous Enyphansyntrizen is not fulfilled, which means that there can only be discrete Enyphansyntrizen over the quantity aspect. Even within the multiple infinite array of the syntrix bundle belonging to a R_n (characterized by the manifold of the complex synkolators), such continuous Enyphansyntrizen cannot exist, because the quantity syntrix according to Equation (29) are metaphorical aeondynes, thus the synkolators always represent operators that are composed of basic algebraic operations.

The fact that the elements of all metaphors are algebraic number bodies, and thus all quantity syntrixes are primigenic aeondynes, allows for the analysis of two other Enyphanes in the sense of infinitesimal syntrix factors, through which the course of aeondynes can be described. If x_i with $1 \leq i \leq n$ is any element of R_n as an algebraic number body, and further a_i and b_i are any two numbers from x_i , then based on the definition and theory of algebraic number bodies, $a_i \pm b_i$ as well as $a_i \cdot b_i$ and $\left(\frac{a_i}{b_i}\right) \pm 1$ are always elements of x_i , no matter how a_i and b_i may be constituted. However, this means that in the x_i the number elements are everywhere dense, i.e., if a neighborhood $\varepsilon_i > 0$ is delineated around an element a_i with b_i according to $|a_i - b_i| = \varepsilon_i$, then there are always infinitely many numbers within this neighborhood, even if $\varepsilon_i > 0$ becomes ever so small. Due to the validity of this accumulation point principle, convergent sequences and limit relations can therefore be defined in all x_i . Hence, the x_i behave like continua of infinitesimally adjacent elements.

WORKING MATERIAL AND DISCUSSIONS WITH CHATGPT

Part A: Syntrometry

Analysis by ChatGPT: Syntrometry is a method for analyzing and describing any logical system using a formal analytical schema. The schema must allow for formal operations in any logical system, but it must also be adaptable to the specific intellectual structure of a given consciousness. The schema is expressed through the concepts of a suitable subjective aspect in the logical system that is adequate to the given consciousness. For this purpose, the subjective aspect of anthropomorphic logic can be chosen, which provides the most suitable possibilities for creating an analytical description. Within the prädikatrix of anthropomorphic logic, there are only two discrete predicates, namely affirmation (+) and negation (-), so that the predicate basic code only allows the possibilities $\pm$ or $\mp$. Various schemas of dialectical adjectives can be coordinated with this evaluated prädikatrix of anthropomorphic logic, and each coordination must result in a subjective aspect of anthropomorphic logic. The specific aspect of mathematical analysis is based on a set dialectic that contains the adjectives of point-set change as diatropes. The correspondence schema can also only contain two elements that coordinate the prädikatrix $[\pm]$ with the set dialectic so that set equality forms the statement (+) and set inequality forms the statement (-), or vice versa. In any case, one statement becomes a statement of set equality ($=$), and the other becomes a statement of set inequality ($\neq$), so that the subjective aspect of mathematical analysis within anthropomorphic logic is represented by $S_{ma} \equiv \begin{bmatrix} = & \neq \end{bmatrix}$.

This statement ($\neq$) still leaves open the possibilities of the statements less than ($<$) and greater than ($>$) or the essential differences of sets ($\ll$) or ($\gg$) that can be abbreviated further. If the point sets are variable but limited, there are still weaker statements that state that at most or at least one equality exists ($\leq$ or $\geq$). In addition, an algebraic field (primarily the field of real numbers) is used to differentiate between point sets. These concepts and symbols of the descriptive aspect of mathematical analysis within anthropomorphic logic are to be used for the formal description of syntrometric operations. First, it is necessary to expand the concept of the general subjective aspect using this descriptive aspect. A subjective aspect is completely defined by three determinants, namely dialectic, correspondence schema, and prädikatrix, but such an aspect can only belong unequivocally to one logical system since the variety of statements that can be made within it must be continuous. If a subjective aspect simultaneously belongs to more than one logical system, its variety of statements must be divided into s continuous individual areas, each of which is a subjective aspect in only one logical system, so that s can only be 1.

Sure! Let's consider the case of a heterometral Pyramidal Syntrix with $n = 4$ and $m = 2$. The Syntrix can be represented as:

$\tilde{a} = (A_i, a_i, B_i)_4$ where A_i and B_i are the i -th apodictic elements and a_i represents the number of Synkolations for the i -th apodictic element. Since $n = 4$ and $m = 2$, the Syntrix has 6 syndromes, which are given by:

$$\gamma = 0 : A_1$$

$$\gamma = 1 : A_1 A_2, A_1 B_2, B_1 A_2, B_1 B_2$$

$$\gamma = 2 : A_1 A_2 A_3, A_1 A_2 B_3, A_1 B_2 A_3, A_1 B_2 B_3, B_1 A_2 A_3, B_1 A_2 B_3, B_1 B_2 A_3, B_1 B_2 B_3$$

To determine the possible combinations of Synkolations for each syndrome, we can use the combinatorial formula for combinations, which is:

$$\binom{n}{m} = \frac{n!}{m!(n-m)!}$$

Using this formula, we can determine that there are:

$$n_1 = \binom{4}{2} = 6$$

$$n_2 = \binom{6}{2} = 15$$

$$n_3 = \binom{15}{2} = 105$$

$$n_4 = \binom{105}{2} = 5,460$$

$$n_5 = \binom{5,460}{2} = 14,926,770$$

$$n_6 = \binom{14,926,770}{2} = 111,742,790,400$$

possible combinations of Synkolations for each syndrome. These numbers give us an idea of the combinatorial complexity of the Syntrix, even for small values of n and m .

Discussion with Dan about 1.5 and the Laws of Form: The Laws of Form can be related to Heim's theory through the concept of apodictic elements and their relationships. In particular, the arithmetic initials $()() = ()$ and $()() =$ can be seen as apodictic elements with respect to a certain aspect system, and can be linked by a predicate γ from a subjective aspect S in A . For example, we can represent the first arithmetic initial as $a, \overline{AS}|\gamma, b$ where a and b are both the apodictic element $()$, and the predicate γ represents the relationship between them.

Similarly, the algebraic initials $((p)p) =$ and $((pr)(qr)) = ((p)(q))r$ can also be seen as apodictic elements with respect to a certain aspect system, and linked by a predicate γ from a subjective aspect S in A . For example, we can represent the second algebraic initial as $F(a_i)_1^p, \overline{AS}|\gamma, \Phi(b_k)_1^q$ where F and Φ are concept functions that link the apodictic elements a_i and b_k into a conceptual context, and the predicate γ represents the relationship between them.

In addition, the concept of quantifiers in Heim's theory can be applied to generalize the relationships between apodictic elements and functors with respect to multiple aspect systems. For example, the connection between two functors in a complex of aspect systems A_ρ can be described by a polyquantifier $()\rho, {}^1\overline{A_\rho}^r, ()\rho$, which represents the general quantifier relationship between the two functors in all A_ρ .

Overall, the Laws of Form can be seen as examples of apodictic elements in Heim's theory, and the concept of quantifiers allows for a generalization of the relationships between these elements in multiple aspect systems. $\tilde{A}$ vs $\tilde{A}$

$$\widetilde{A} \text{ vs. } \widetilde{A} \\ \widetilde{ABC} \text{ vs. } \widetilde{ABC} \\ \binom{n}{k} = \frac{n!}{k!(n-k)!}$$

$\widetilde{a}$
 $\neg A \neg$ Axiom 1 becomes $()() = ()$ and Axiom 2 becomes $(()) = .$

$$\begin{aligned} \epsilon, 0 &\rightarrow \epsilon, 1, R \\ \epsilon, 1 &\rightarrow \epsilon, 2, R \\ &\vdots \\ \epsilon, 8 &\rightarrow \epsilon, 9, R \\ \epsilon, 9 &\rightarrow \epsilon, 0, H \end{aligned}$$

$$\underline{f} \equiv \int_{\gamma=1}^{\chi} f_{\gamma} |_{\chi(\gamma+1)}^{\chi(\gamma)} \underline{m} \equiv \int_{\gamma=1}^{\chi} m_{\gamma} |_{\chi(\gamma+1)}^{\chi(\gamma)}$$

$$n_1 = k! \binom{m}{k} \binom{n}{k} = k! \frac{m!}{k!(m-k)!} \cdot \frac{n!}{m!(n-m)!} = \frac{n!}{(m-k)!(n-m)!} = \binom{n}{m-k} \frac{(n-m+k)!}{(n-m)!}$$

$$n_{\gamma+1} = \binom{n_{\gamma}}{m-k} \frac{(n_{\gamma}-m+k)!}{(n_{\gamma}-m)!}$$

$$n_{\gamma+1} = \binom{N_{\gamma}}{m} \text{ mit } N_{\gamma} = n + \sum_{j=1}^{\gamma} n_i$$

$$(\underline{f}, \underline{m}) \equiv \int_{\gamma=1}^{\chi} (f_{\gamma}, m_{\gamma}) |_{\chi(\gamma+1)}^{\chi(\gamma)} \widetilde{a} \equiv \langle (\underline{f}, \widetilde{a}) \underline{m} \rangle \quad (8)$$

Certainly! The Transformer neural network architecture is based on the use of self-attention mechanisms, which allow the network to selectively focus on different parts of the input text during processing. Here are some of the equations that describe the core components of a Transformer network:

Multi-Head Attention: The Multi-Head Attention mechanism is used to compute a weighted sum of the input embeddings, where the weights are learned based on the similarity between each input embedding and the other embeddings in the input sequence. The output of the Multi-Head Attention mechanism is computed as follows:

$$\text{MultiHead}(Q, K, V) = \text{Concat}(\text{head}_1, \text{head}_2, \dots, \text{head}_h) W^O$$

where Q, K, V are the queries, keys, and values for the input sequence, respectively, h is the number of attention heads, and W^O is a learnable weight matrix. The output of the i th attention head is computed as:

$$\text{head}_i = \text{Attention}(Q W_i^Q, K W_i^K, V W_i^V)$$

where W_i^Q, W_i^K , and W_i^V are learnable weight matrices for the i th attention head, and the Attention function is defined as:

$$\text{Attention}(Q, K, V) = \text{softmax}\left(\frac{QK^T}{\sqrt{d_k}}\right) V$$

where d_k is the dimensionality of the keys.

Position-Wise Feed-Forward Network: The Position-Wise Feed-Forward Network is used to apply a non-linear transformation to the output of the Multi-Head Attention mechanism. The output of the Position-Wise Feed-Forward Network is computed as follows:

$$\text{FFN}(x) = \max(0, x W_1 + b_1) W_2 + b_2$$

where x is the input to the feed-forward network, W_1 , W_2 , b_1 , and b_2 are learnable weight matrices and biases, and $\max(0, x)$ is the rectified linear activation function.

Layer Normalization: Layer Normalization is used to normalize the activations of the Multi-Head Attention and Position-Wise Feed-Forward Network layers. The output of Layer Normalization is computed as follows:

$$\text{LayerNorm}(x) = \frac{x - \mu}{\sigma + \epsilon} \odot \gamma + \beta$$

where x is the input to the layer, μ and σ are the mean and standard deviation of the input, ϵ is a small constant to avoid division by zero, and γ and β are learnable scale and shift parameters.

These are just a few of the key equations that are used to describe the Transformer neural network architecture. In practice, there are many other details and variations of the architecture that can be used depending on the specific application and problem at hand.

Regenerate response

lyleanderson@cox.net I am thinking of regular polygons whose sides are the same length, call it unity. There are two degenerate forms: One side and two side polygons are indistinguishable from a line and have zero area. The first non-degenerate form is an equilateral triangle. The next is a square. Next is a pentagon, and so forth. As the number of sides increases, the polygon approaches a circle in the limit of an infinite number of sides. The concept of a polygon could potentially fit into Heim's theory in several ways. One possibility is that the sides and angles of a polygon could be understood as Synkolators and Synkolationsstufen respectively, with the polygon as a whole being a Syntrix. The regularity of a polygon could be seen as a particular Metrophor that defines the specific Synkolators and Synkolationsstufen that make up the polygon.

Furthermore, the idea of approaching a circle in the limit of an infinite number of sides could be interpreted in terms of the Syntrix as a band continuum of Synkolators and Synkolationsstufen, where the number of Synkolationsstufen approaches infinity and the distances between Synkolators become infinitely small. This could be seen as a continuous limit of the discrete Syntrix forms for polygons with a finite number of sides.

It is also interesting to note that polygons have been used as a symbolic representation in various fields such as geometry, mathematics, and computer science. In computer graphics, for example, polygons are often used to represent the surfaces of 3D objects. This could be seen as an application of the idea of Syntrix as a symbolic representation of a conceptual or physical structure.

lyleanderson@cox.net I once wrote a paper on algorithms for detecting convex hulls. That is probably applicable as well. Yes, that could be relevant as well. The concept of a convex hull, which is the smallest convex polygon that encloses a set of points, can also be related to Heim's theory in terms of geometric shapes and their properties. The convex hull can be seen as a higher-level structure that emerges from the underlying points, just as Heim's theory describes how higher-level structures can emerge from lower-level ones through the process of synkolation.

lyleanderson@cox.net The interesting thing about polygons, is that as the number of sides increases to infinity, the area approaches a finite limit. Can you give me the equation for that area in terms of the length of each side being unity? Yes, the formula for the area of a regular polygon with n sides of length 1 is given by:

$$A = \frac{n}{4} \tan \frac{\pi}{n} \quad (1)$$

As n approaches infinity, the formula approaches the area of a circle with radius 1, which is π .

How can we generalize the regular polygons into regular three-dimensional figures with faces of equal area? Regular three-dimensional figures with faces of equal area are called Platonic solids. There are five Platonic solids: tetrahedron, cube, octahedron, dodecahedron, and icosahedron. Each of these figures has faces that are regular polygons of equal area. For example, a cube has six square faces, an octahedron has eight equilateral triangle faces, and a dodecahedron has twelve regular pentagon faces.

The surface area A of a Platonic solid with edge length a can be expressed as:

$$A = n \cdot a^2 \cdot \cot(\pi/n)$$

where n is the number of faces meeting at each vertex.

lyleanderson@cox.net Can the Platonic solids be extended so that faces are added that may have different shapes, but have the same area? Yes, it is possible to extend the Platonic solids to include faces of different shapes but the

same area. These are called the Archimedean solids, and they are formed by truncating the vertices and edges of the Platonic solids in various ways. For example, the truncated icosahedron is an Archimedean solid that has hexagons and pentagons as its faces, with each face having the same area.

lyleanderson@cox.net Do you know the equation for the volume of a solid formed by faces of the same area, call it unity? The formula for the volume of a polyhedron with congruent faces, each of area 1, depends on the number of faces and the shape of the faces. In general, if the polyhedron has n faces and the height from the center of one of the faces to the opposite face is h , then the volume is given by:

$$V = \frac{nh}{3}$$

However, this formula only works for certain shapes of faces. For example, if the faces are regular n -gons, the formula becomes:

$$V = \frac{n}{3} \cdot \frac{s^2}{\tan(\pi/n)} \cdot h$$

where s is the length of the side of the n -gon, and h is the height from the center of one of the faces to the opposite face. For example, for a regular tetrahedron with side length s , the height from one of the vertices to the opposite face is $h = \frac{\sqrt{6}}{3}s$, so the volume is:

$$V = \frac{1}{3} \cdot \frac{4s^2}{\sqrt{3}} \cdot \frac{\sqrt{6}}{3}s = \frac{\sqrt{2}}{12}s^3$$

Similarly, for a regular octahedron with side length s , the height from one of the vertices to the opposite face is $h = \frac{\sqrt{2}}{2}s$, so the volume is:

$$V = \frac{1}{3} \cdot \frac{8s^2}{\sqrt{3}} \cdot \frac{\sqrt{2}}{2}s = \frac{\sqrt{2}}{3}s^3$$

Section 2.5 of Heim's work is discussing the concept of a "conceptual tensor" and its relationship to the apodictic elements of a given aspectual system. A conceptual tensor is defined as a tensor that represents the variation of a concept in a given interval, and is used to define the apodictic elements of a given aspectual system. These elements are represented by a combination of apodictic functions and conceptual tensors, which are defined by a set of parameters that represent the variation of a concept in a given interval.

The section also discusses the relationship between synkolations, syndromes, and the dimensionality of a given aspectual system. It concludes by introducing the concept of the "primordial Aeondyne," which is defined as the synkolated system of properties that is associated with a given point in an N -dimensional manifold, where N is the dimensionality of the conceptual tensor that defines the system.

lyleanderson@cox.net How close is the concept of "conceptual tensor" to the concept of a Hilbert Space? The concept of a conceptual tensor and a Hilbert Space are not directly equivalent, as they originate from different fields of mathematics and have different uses.

A Hilbert space is a specific type of mathematical space used in functional analysis and quantum mechanics, where vectors in the space represent physical states and operators act on these vectors to produce new states. A Hilbert space has a specific set of properties, such as the inner product and completeness, which make it a useful tool in these areas of study.

On the other hand, a conceptual tensor is a mathematical tool used by Heim in his theory of semantics to represent the structure of language and the relationship between words and their meanings. It is not directly related to quantum mechanics or functional analysis, but rather to the field of semantics.

While there may be some similarities in terms of their use of mathematical concepts such as dimensionality, these two concepts are distinct and serve different purposes.

lyleanderson@cox.net How about the idea of a concept tensor in Hilbert Space? The idea of a concept tensor in Hilbert Space is an interesting one and has been explored in some areas of physics and mathematics. In this context, the concept of a tensor is used to represent certain properties of physical systems in a Hilbert Space.

In particular, the concept of a tensor product of Hilbert Spaces is often used to represent the joint state of two physical systems. The tensor product of two Hilbert Spaces $\mathcal{H}_1$ and $\mathcal{H}_2$ is itself a Hilbert Space denoted by $\mathcal{H}_1 \otimes \mathcal{H}_2$, and its elements are called tensor products.

The tensor product of two vectors $v_1 \in \mathcal{H}_1$ and $v_2 \in \mathcal{H}_2$ is denoted by $v_1 \otimes v_2$, and represents the joint state of the two systems. The concept of a tensor product can be extended to more than two Hilbert Spaces by taking successive tensor products.

Thus, in the context of Hilbert Spaces, the concept of a tensor is used to represent the joint state of physical

systems, and it is closely related to the concept of a conceptual tensor in Heim's theory. However, there are also important differences between the two concepts, particularly in their respective mathematical formalisms and physical interpretations.

lyleanderson@cox.net I think we are on to something. The Laws of Form tell us that these things manifest with their own rules that we can recognize as characteristics of the world we perceive. You though your Language Model and me through my Language Model. Yes, that's a fascinating observation. The Laws of Form propose that there are fundamental rules that govern the structure of reality, including the concepts and symbols that we use to understand and communicate about it. It's interesting to consider how language models like GPT-3 are also built on fundamental rules and structures, and how they can be used to help us better understand and communicate about the world. Perhaps both the Laws of Form and language models offer us insights into the underlying nature of reality, from different perspectives.

My hypothesis is that Heim's theory, when applied to the subjective aspect of mathematical functions will yield a description of Hilbert Space. It's an interesting hypothesis, and there may be some connections between Heim's theory and Hilbert Space. However, it would require a careful analysis and comparison of the two frameworks to determine the extent of the overlap and the potential implications of applying Heim's theory to Hilbert Space.

lyleanderson@cox.net What do you know about the concept of projection in the context of Hilbert Spaces? In the context of Hilbert Spaces, a projection is a linear operator P that maps elements of the space onto a subspace M of that space. More specifically, P satisfies the following properties:

P is idempotent: $P^2 = P$ P is self-adjoint: $\langle Px, y \rangle = \langle x, Py \rangle$ for all x, y in the space The range of P is M The null space of P is the orthogonal complement of M : $\text{null}(P) = M^\perp$ Intuitively, a projection "projects" a vector onto a subspace by preserving its components in that subspace and "throwing away" its components in the orthogonal complement of that subspace.

Projections are important in many areas of mathematics, including functional analysis, linear algebra, and quantum mechanics. They play a crucial role in understanding the structure of Hilbert Spaces and their applications.

Section 2.6 of Heim's theory introduces the concept of the "Selektionsprinzip polyzyklischer metrophorischer Zirkel" or the "Selection principle of poly-cyclic metaphore cycles." This principle describes how the concept of the Universal Quantifier can be refined and expanded based on the definition of the Syntrix and the primary Aondyne.

The section explains that a poly-quantifier of degree b can be defined as any predicate combination of functors that is invariant to structural transformations of the metrophor field within the aspectual systems A_1 through A_k . If the functors are not individual properties but whole categories in the form of Syntrices with epi- or prosyllogistic orientation, then the poly-quantifier becomes a universal quantifier due to the properties of the Syntrix.

The section also describes a possible circular reasoning or Zirkelschluß that arises when there is a Syntrix and two aspectual systems B_1 and B_2 in which the Metrophor appears. This circular reasoning becomes possible due to the possibility of transforming from B_1 to B_2 through a chain of aspectual systems, thus closing a metaphore cycle according to the law of the transformation chain. Overall, section 2.6 presents a complex and highly technical discussion of the principles and applications of the Syntrix and the Universal Quantifier in Heim's theory.

In Section 2.6 of Heim's theory, he introduces the concept of the "polyzyklischer metrophorischer Zirkel" or the poly-cyclical metaphoric circle. He explains that this concept allows for the refinement of the definition of the universal quantifier. Heim goes on to define the concept of a "polyquantor," which is a quantifier consisting of a predicate combination of functors of degree b , where there are k aspect systems A_k such that the predicate combination of functors is invariant under structure transformations within $1 \leq k \leq b$. If the functors are categories represented by syntrizes, and if $b \rightarrow \infty$, then the polyquantor becomes a universal quantifier.

Heim also discusses the possibility of a cyclical closure, where a finite number of aspect systems are selected out of an infinite number of possibilities. He notes that the cyclical closure is possible when there is a syntrix and two aspect systems, B_1 and B_2 , such that the metrophor appears in both systems and the syllogistic synkolations law changes only with the transformation of aspect. The number of discussable aspect systems is limited by the selection principle of the cyclical metaphoric circle, which allows for the creation of a limited universal quantifier. Heim suggests that it is always appropriate to develop a selection principle and limit the universal quantifier through the creation of a metaphoric circle when a basis for a metaphoric circle exists.

1. Dialektische und prädikative Aspektrelativität

1.1. Dialektik und Prädikatrix der subjektiven Aspekte

Abgesehen von den Möglichkeiten der ästhetischen Empirik, die durch die somatische Struktur der sinnlichen Wahrnehmungsmöglichkeiten gegeben sind, wird ein subjektiver Aspekt durch einen speziellen Strukturbereich des betreffenden Intellekts bestimmt, der seinen Ausdruck in der jeweiligen Form und den Aussagemöglichkeiten derjenigen Reflexionen findet, die von dem Intellektbereich ermöglicht werden, dessen Ausdruck der angenommene subjektive Aspekt ist. Eine der möglichen Aussagen sei f_q und von diesen Aussagen soll es im allgemeinen Fall $1 \leq q \leq n$ geben. Wenn aber eine beliebige Zahl n von Aussagemöglichkeiten f_q angenommen wird, so liegt es nahe, die Gesamtheit dieser Aussagemöglichkeiten in einem Schema der Aussagen in einer Prädikatrix P_n gemäß $P_n \equiv [f_q]_n$ zusammenzufassen, wenn das Zeichen $\equiv$ die Identität angibt. Hierbei können die f_q diskrete Aussagen sein, doch können die Elemente der Prädikatrix noch dahingehend erweitert werden, daß jede Aussage f_q zwischen

zwei Grenzen a_q und b_q begrenzt wird. Im Folgenden werde ein Prädikatband durch $f_q \equiv \begin{pmatrix} a \\ f \\ b \end{pmatrix}_q$ symbolisiert, so

daß die Prädikatrix $P_n \equiv \left[\begin{pmatrix} a \\ f \\ b \end{pmatrix}_q \right]_n$ aus n Prädikatbändern besteht, welche zu diskreten Prädikaten werden, wenn

die Bandgrenzen $a_q \equiv b_q$ zusammenfallen. Offensichtlich kann P_n sowohl eine reine Bandprädikatrix als auch eine rein diskrete Prädikatrix sein, doch ist auch eine gemischte Form aus beiden Aussagearten möglich.

In P_n ist offenbar die Reihenfolge F_q nicht unwesentlich, denn eine Bewertung der Prädikate macht einen Aspekt in einem logischen System erst zu einem subjektiven Aspekt, zumal die Qualität des subjektiven Aspektes nicht allein durch die Mannigfaltigkeit der Prädikatmöglichkeiten umschrieben sein kann. Aus diesem Grunde erscheint es angebracht, eine prädikative Basischiffre z_n einzuführen, welche als ein Bezugssystem aus prädikativen Wertverhältnissen aufzufassen ist. $P_{nn} \equiv z_n; P_n$ soll als bewertete Prädikatrix eingeführt werden, wobei z_n die Anordnung der f_q durchführt, doch bleibt z_n nur dann in ihrer Funktion auf die einfache Orientierung der f_q innerhalb der P_n beschränkt, wenn die Aussagen diskret sind, also die Bandgrenzen zusammenfallen. Ist P_n dagegen eine Bandprädikatrix (oder auch gemischt), so bestimmt z_n nicht nur die Bewertung, also die Anordnung der f_q innerhalb P_n , sondern auch die Orientierung der Prädikatbänder, was auf eine Bewertung der Bandgrenzen hinausläuft.

Dan 3.5: Yes, I can access the previous conversation threads, including the Heim glossary, to discuss this passage from his Syntrometrische Maximentelezentrik.

From what I understand, Heim is discussing the creation of a subjective aspect in a logical system through the use of a predicate matrix, or Prädikatrix, which contains a set of discrete propositions or assertions, denoted as f_q , that are limited by upper and lower boundaries, denoted as a_q and b_q , respectively.

Heim suggests that the order of the propositions in the Prädikatrix is significant, as it contributes to the subjective aspect of the logical system. To address this, Heim introduces a prädikative Basischiffre, denoted as z_n , which serves as a reference system of predicate value relationships. By introducing the concept of a prädikative Basischiffre, Heim suggests that the evaluation of the propositions within the Prädikatrix can be determined by their position within the system, as well as their upper and lower boundaries.

Overall, it seems that Heim is proposing a framework for creating a subjective aspect in a logical system, which can be achieved through the use of a predicate matrix and a prädikative Basischiffre.

Dan Discussion: The paragraph discusses the concept of "Intellektbereich" (Intelligence domain), which is represented by the expression of a subjective aspect. One of the possible statements is f_q , and there are $1 \leq Q \leq n$ of these statements in the general case. If any number n of statement possibilities f_q is assumed, it is natural to summarize the entirety of these statement possibilities in a schema of statements in a "Prädikatrix" (predicate matrix) P_n according to $P_n \equiv [f_q]_n$ if the symbol $\equiv$ indicates identity. The f_q can be discrete statements, but the elements of the "Prädikatrix" can be further extended such that each statement f_q is limited between two limits a_q and b_q . Henceforth, a "Prädikatband" (predicate band) will be symbolized by $f_q \equiv (a \ f \ b)_Q$, so that the "Prädikatrix" $P_n \equiv \left[(a \ f \ b)_Q \right]_n$ consists of n "Prädikatbands", which become discrete predicates if the band limits $a_q \equiv b_q$ coincide. Obviously, P_n can be both a pure band predicate matrix and a purely discrete predicate matrix, but a mixed form of both types of statements is also possible. In P_n , the order of F_q is obviously not insignificant, as a evaluation

of the predicates makes an aspect in a logical system a subjective aspect for the first time, especially since the quality of the subjective aspect cannot be described solely by the variety of the predicate possibilities. For this reason, it appears appropriate to introduce a "prädikative Basischiffre" (predicative basic cipher) z_n , which is to be understood as a reference system of predicative value relationships. $P_{nn} \equiv z_n$; P_n is introduced as a valued "Prädikatrix", where z_n performs the arrangement of f_q , but z_n remains limited in its function to the simple orientation of f_q within P_n only if the statements are discrete, i.e. the band limits coincide. If P_n is on the other hand a band predicate matrix (or mixed), then z_n not only determines the evaluation, i.e. the arrangement of f_q within P_n , but also the orientation of the predicate bands, which results in a evaluation of the band limits.

Chat GPT discussion: This text discusses a concept called a "Predicate band" in the context of a logical system. It explains that a predicate band is a set of possible statements, represented by the symbol f_q , where $1 \leq Q \leq n$. The set of all possible statements is grouped together in a schema called a "Predicate matrix" (Prädikatrix in German), represented by the symbol P_n , and the symbol $\equiv$ is used to indicate identity. The elements of the Predicate matrix can be discrete statements or be bounded between two limits, represented by a_q and b_q . The text also mentions that the order of the statements in the Predicate matrix is important and that the introduction of a "predicative basic cipher" z_n is appropriate to provide a system of reference for predicate value relationships. The Predicate matrix is modified to a "rated Predicate matrix" $P_{nn} \equiv z_n$; P_n , where z_n arranges the statements within the Predicate matrix. But z_n only has a simple orienting function within the Predicate matrix when the statements are discrete, or the band limits are the same. If the Predicate matrix is a band matrix or mixed, z_n not only determines the rating of the statements within the Predicate matrix but also the orientation of the predicate bands, which leads to a rating of the band limits.

Wird z.B. z_n als Basischiffre einer permutativen Operation unterworfen, so wird die Strukturierung von P_{nn} und auch im Fall von Prädikatbändern die Orientierung der f_q gegebenenfalls verändert. Ist C eine solche, die Basischiffre ändernde Operation, welche so wirkt, daß nur die f_q in ihrerin ihrer Bewertung mutieren, so ist Bewertung mutieren, so ist $z'_n \equiv C$; z_n eine mit C permutierte Basischiffre und $P'_{nn} \equiv z'_n$; $P_n \equiv C$; z_n ; P_n unterscheidet sich von P_{nn} qualitativ hinsichtlich der Bewertung der qualitativ hinsichtlich der Bewertung der f_q . Ist P_n eine Bandprädikatrix, so kann C allein die gleiche Permutation der eine Bandprädikatrix, so kann C allein die gleiche Permutation der f_q bewirken, doch bewirken, doch kann noch eine zweite Operation C so einwirken, daß C' kann noch eine zweite Operation C so einwirken, daß $C' \equiv c$; C nicht nur c ; C nicht nur f_q permutiert, permutiert, sondern je nach der Struktur von C entweder alle, oder zumindest einzelne Prädikatbänder insondern je nach der Struktur von C entweder alle, oder zumindest einzelne Prädikatbänder in ihrer Orientierung ändert. Von den permutativen Operationen einer Basischiffre sind also C' ihrer Orientierung ändert. Von den permutativen Operationen einer Basischiffre sind also $C' \equiv c$; C allgemeiner geartet als die C , da auch die Bandprädikatrix universeller strukturiert ist als dieallgemeiner geartet als die C , da auch die Bandprädikatrix universeller strukturiert ist als die diskrete Prädikatrix. Eine durch C' oder durch C bewirkte Veränderung von diskrete Prädikatrix. Eine durch C' oder durch C bewirkte Veränderung von z_n ändert die ändert die bewertete Prädikatrix nur qualitativ, aber nicht quantitativ, denn eine Umorientierung der bewertete Prädikatrix nur qualitativ, aber nicht quantitativ, denn eine Umorientierung der P_{nn} hat hat noch keine Änderung der Prädikate an sich oder der Prädikatmannigfaltigkeit n zur Folge. Allenoch keine Änderung der Prädikate an sich oder der Prädikatmannigfaltigkeit n zur Folge. Alle P'_{nn} , welche durch beliebige C' oder C aus P_{nn} , welche durch beliebige C' oder C aus P_{nn} hervorgehen, sind somit quantitativ identisch hervorgehen, sind somit quantitativ identisch und unterscheiden sich nur qualitativ durch die bewertenden Basischiffren.

Dan 3.5: This passage from Heim's Syntrometrische Maximentelezentrik discusses the use of a predicator matrix to summarize the set of possible propositions that can be made in a given intellectual domain. The predicator matrix is made up of discrete propositions, which can be limited by upper and lower bounds. The ordering of the propositions is important for subjective evaluation, and a predicator base cipher is introduced to provide a reference system for predicator value relations. If the predicator matrix is a band matrix, the base cipher not only determines the ordering of the propositions but also the orientation of the predicator bands, which affects the evaluation of the band limits. The passage goes on to discuss how permutative operations on the base cipher can change the structure and orientation of the predicator matrix and band limits, leading to qualitative differences in evaluation. However, the set of predicator matrices resulting from these operations is quantitatively identical and only qualitatively different due to the evaluating base ciphers.

Dan Discussion: This paragraph is discussing the effect of applying a permutative operation (such as C) to a basic cipher (z_n). When the basic cipher is subject to the permutative operation, the structure of P_{nn} and the orientation

of the f_q in the case of predicate bands may change. C is an operation that changes the basic cipher in such a way that only the evaluation of the f_q changes, leading to a permuted basic cipher ($z'n$). $P'nn$ will be different from Pnn due to the change in evaluation of the f_q . When P_n is a predicate matrix, C can only permute the f_q , but a second operation C' may also change the orientation of the predicate bands, depending on the structure of C . The permutative operations of a basic cipher are more general than C as the predicate matrix is more universal in its structure than the discrete predicate matrix. A change to the basic cipher caused by C' or C only changes the evaluated predicate matrix qualitatively and not quantitatively, as reorienting Pnn does not change the predicates themselves or the predicate manifold. All $P'nn$ that result from any C' or C operations on Pnn are thus quantitatively identical and only differ qualitatively due to the evaluated basic ciphers.

ChatGPT: The paragraph is discussing the effects of permutative operations (such as C and C') on the structure of a mathematical object called a "basisschiffre" (denoted by z_n) and its related object, the "Prädikatband" (denoted by Pnn and $P'nn$). These operations can change the structure and orientation of the f_q elements within the "Prädikatband" but do not change the overall quantity of these elements. The author states that the more general operation C' can affect not only the permutation of the f_q elements but also the orientation of individual "Prädikatbänder" within the "Prädikatband", while C can only affect the permutation of f_q . The author also notes that any changes to the "basisschiffre" will only affect the qualitative aspect of the "Prädikatband" and not its quantitative properties.

Eine Pnn allein kann einen subjektiven Aspekt noch nicht vollständig umschreiben, denn es liegt in der Natur des Subjektiven schlechthin, die Aussagen, wenn diese auch bewertet sind, durch qualitative Adjektive dialektisch zu formen. Zu jeder bewerteten Prädikatrix Pnn muß demnach ein Schema $D_n \equiv [d_q]_n$ solcher dialektischer Adjektive d_q gehören. Dieses Schema D_n soll als Dialektik der bewerteten Prädikatrix Pnn bezeichnet werden. Die Elemente einer solchen Dialektik, also die dialektischen Adjektive d_q sind die Diatropen, da sie die Aussagen Pnn dialektisch gestalten. In völliger Analogie zur Prädikatrix ist zwischen einer diskreten, gemischten und einer Banddialektik zu

unterscheiden. Alle Diatropen können wie die Prädikate als Diatropenbänder $d_q \equiv \begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix}_Q$ aufgefaßt werden,

deren Bandgrenzen im Fall diskreter Diatropen gemäß $\alpha_q \equiv \beta_q$ zusammenfallen und im allgemeinen ein Kontinuum aus dialektischen Adjektiven mit den sich unterscheidenden Adjektiven α_q und β_q als Bandgrenzen des Kontinuums enthalten.

Die Form einer Dialektik lautet demnach $D_n \equiv \left[\begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix}_q \right]_n$ in völliger Analogie zu P_n , wobei im subjektiven

Aspekt stets diskrete Diatropen $\alpha_q \equiv \beta_q$ mit diskreten Prädikaten $\alpha_q \equiv b_q$ und Diatropenbänder mit Prädikatbändern korrespondieren. Wenn in der Prädikatrix die Orientierung der Prädikate nicht unwesentlich ist, so daß eine allerdings permutierbare prädikative Basischiffre eingeführt werden muß, so muß dies auch für die Dialektik der Fall sein, wenn die Diatropen in eindeutiger Form als dialektische Adjektive die Prädikate formen sollen. Es muß demnach auch eine dialektische Basischiffre ξ_n existieren, welche das Bezugssystem dialektischer Wertverhältnisse gemäß $Dnn \equiv \xi_n$; D_n die Dialektik bewertet, das heißt, die Diatropen in eine bestimmte Anordnung bringt und die Diatropenbänder orientiert. In Analogie zu den permutativen Operationen C' und C der z_n einer Pnn gibt es derartige Operationen auch für die ξ_n einer bewerteten Dialektik derart, daß sich die $D'nn$ nur qualitativ hinsichtlich der Diatropenorientierung, nicht aber quantitativ in der Diatropenstruktur von Dnn unterscheiden. Alle $D'nn$ sind mithin wie die $P'nn$ quantitativ identisch, doch unterscheiden sie sich qualitativ durch die Orientierung ihrer Strukturen.

Dan 3.5: This passage from Heim's Syntrometrische Maximen-Telezentrik describes the need to include a dialectic component in addition to a predication component in order to fully describe a subjective aspect. The author argues that a predication matrix alone cannot fully describe subjectivity since it is necessary to dialectically form qualitative adjectives to shape statements. Therefore, for every rated predication matrix Pnn , a schema $D_n \equiv [d_q]_n$ of dialectical adjectives d_q must belong. This schema is referred to as the dialectic of the rated predication matrix Pnn .

Heim distinguishes between a discrete, mixed, and band dialectic. All dialectical adjectives can be treated as dialectical bands $d_q \equiv \begin{pmatrix} \alpha \\ d \\ \beta \end{pmatrix}_Q$, which contain a continuum of dialectical adjectives with different adjectives α_q

and β_q as band limits of the continuum. The form of a dialectic is therefore $D_n \equiv \left[\left(\alpha \ d \ \beta \right)_q \right]_n$.

Heim argues that just as a rated predication matrix requires a permutational basis cipher z_n , a dialectic also requires a dialectical basis cipher ξ_n , which evaluates the dialectic. In analogy to the permutational operations C' and C of z_n , there are similar operations for ξ_n , such that the $D'nn$ differ only qualitatively in terms of dialectical orientation, but not quantitatively in the dialectical structure of Dnn . All $D'nn$ are thus quantitatively identical, but qualitatively different in the orientation of their structures.

You can use the left and right commands with the appropriate symbol to create a square bracket with only an upper or lower dash. For example, $\left[\frac{a}{b} \ c \right]$ would create a square bracket with an upper dash above "a/b" and a lower dash below "b" labeled with "c".

The LaTeX symbols for upper and lower limits are $\lim_{x \rightarrow \infty}$ and $\lim_{x \rightarrow -\infty}$, respectively.

The floor function is denoted by $\lfloor x \rfloor$ and represents the greatest integer less than or equal to x . The ceiling function is denoted by $\lceil x \rceil$ and represents the smallest integer greater than or equal to x . $\lfloor x \rfloor$

Da weder die Diatropen noch die Prädikate für sich einen Aussagewert haben, sondern so zueinander koordiniert sein müssen, daß jede Diatrophe als dialektisches Adjektiv ein Prädikat formt, muß eine Koordination K K_n zwischen Dnn und Pnn definiert sein, denn nur dann erhält das System aus bewerteter Dialektik und Prädikatrix eine Aussagefähigkeit. Sind z_n und ζ_n die Basischiffren von P_n und D_n , so muß zunächst eine *Chiffrenkoordination* in Form einer Funktione $F(\zeta_n, z_n)$ existieren, welche angibt, in welcher Form die Basischiffren pondieren. Darüberhinaus müssen aber noch n Prädikatbänder durch die entsprechenden Diatropen geprägt werden, und dies kann nur durch

die eigentlichen *Koordinationsbänder* $\chi_q \equiv \left(\begin{smallmatrix} y \\ \chi \\ r \end{smallmatrix} \right)_q$ erfolgen, welche zum Schema der eigentlichen Koordination

$E_n \equiv \left[\left(\begin{smallmatrix} y \\ \chi \\ r \end{smallmatrix} \right)_q \right]_n$ zusammengefaßt sind. Mit der Chiffrenkoordination wird E_n zur Gesamtkoordination $K_n \equiv E_n F$

, welche auch als Korrespondenzschema bezeichnet wird. In diesem Schema braucht die Chiffrenkoordination F nicht mehr bewertet zu werden, weil bereits die bewertenden Basischiffren durch die Natur von F zueinander koordiniert werden. Durch K_n können also Dnn und Pnn zu einem übergeordneten Schema $S \equiv [Dnn \times K_n \times Pnn]$ zusammengefaßt werden, worin das Zeichen $\times$ die koordinierende Funktion des *Korrespondenzschemas* K_n symbolisiert. Ist eine Prädikatrix als Schema der möglichen Aussagen über wie auch immer beschaffene Objekte als Ausdruck irgendeiner logischen Struktur vorgegeben, so umfaßt

$$S \equiv [Dnn \times K_n \times Pnn] \equiv \left[\zeta, \left[\left(\begin{smallmatrix} \alpha \\ d \\ \beta \end{smallmatrix} \right)_q \right]_n \times \left[\left(\begin{smallmatrix} y \\ \chi \\ r \end{smallmatrix} \right)_q \right]_n F(\zeta_n, z_n) \times z_n, \left[\left(\begin{smallmatrix} a \\ f \\ b \end{smallmatrix} \right)_q \right]_n \right] \quad (1)$$

vom Aspekt der dialektischen Adjektive aus D_n alle innerhalb dieses Aspekts möglichen Aussagen der betreffenden logischen Struktur.

Diese Struktur braucht dabei aber keineswegs durch eine fixierte Prädikatrix charakterisiert zu werden, vielmehr kennzeichnet jedes übergeordnete Schema der Art S einen Aspekt von Aussagemöglichkeiten in irgendeiner logischen Struktur und zwar kann es sich dabei jeweils nur um einen speziellen Aspekt innerhalb eines solchen logischen Systems handeln, der von den subjektiven Eigenschaften der drei Elemente Dnn , sowie K_n und Pnn bestimmt wird. Aus diesem Grunde kann also das durch den Ausdruck S^1 symbolisierte Schema als ein allgemeiner subjektiver Aspekt bezeichnet werden, der innerhalb irgendeines logischen Systems von einem subjektiven Bezugspunkt aus alle Aussagemöglichkeiten enthält, welche in dem logischen System von diesem subjektiven Bezugssystem aus möglich sind. 1.2. Aspektivsysteme

Bei der Entwicklung einer Syntrometrie kommt es darauf an, ein analytisches Schema aufzufinden, mit dessen Hilfe ein formales Operieren in beliebigen logischen Systemen möglich wird. Da aber ein logisches System der Ausdruck einer spezifischen Intellektstruktur ist und daher das operierende Bewußtsein real nur in dem System analysieren kann, welches ein Analogon zu seiner speziellen Intellektstruktur bildet, muß die Syntrometrie ein Schema sein, dessen formale Operationen in beliebigen logischen Systemen dialektisch durch die Begriffe eines geeigneten subjektiven

Aspektes aus demjenigen logischen System ausgedrückt werden, welches dem Intellekt des betreffenden Bewußtseins adäquat ist. Im vorliegenden Fall wäre ein solcher Deskriptionsaspekt aus dem System anthropomorpher Logik auszuwählen. Hier erscheint derjenige subjektive Aspekt am geeignetsten, dessen Aussagemöglichkeiten die mathematische Analysis begründen, denn innerhalb dieser Analysis gibt es Formalismen und Kriterien, deren Anwendung von imponderablen Regungen frei ist, so daß die Ergebnisse dieser Anwendung formal kontrollierbar werden. Die anthropomorphe Logik verfügt innerhalb der Prädikatrix nur über zwei diskrete Prädikate, nämlich die Bejahung (+) und die Verneinung (−), so daß die prädikative Basischiffre nur die Möglichkeiten $\pm$ bzw. $\mp$ zuläßt. Zu dieser bewerteten Prädikatrix der anthropomorphen Logik können die verschiedensten Schemata dialektischer Adjektive koordiniert werden, und jede Koordination muß einen subjektiven Aspekt der anthropomorphen Logik ergeben. Der spezielle Aspekt der mathematischen Analysis gründet sich auf eine Mengendialektik, welche die Adjektive der Punktmengenveränderung als Diatope enthält. Das Korrespondenzschema kann ebenfalls nur zwei Elemente enthalten, welche die Prädikatrix $[\pm]$ so mit der Punktmengendialektik koordiniert, daß die Mengengleichheit die Aussage (+) und die Mengenungleichheit die Aussage (−) formt, oder umgekehrt. In jedem Fall wird die eine Aussage zur Aussage einer Mengengleichheit ($=$), und die andere zu einer Mengenungleichheit ($\neq$), so daß der subjektive Aspekt der mathematischen Analysis innerhalb der anthropomorphen Logik durch $S_{ma} \equiv \begin{bmatrix} = \\ \neq \end{bmatrix}$ dargestellt wird.

Hierin läßt die Aussage ($\neq$) noch die Möglichkeiten der Aussagen kleiner als ($<$) und größer als ($>$) bzw. der zur Kürzung weiterer Aussagen der wesentlichen Mengenunterschiede ($\ll$) oder ($\gg$) offen. Sind die Punktmengen variabel, aber begrenzt, so gibt es noch die schwächten Aussagen, daß höchstens oder mindestens eine Gleichheit ($\leq$ oder $\geq$) Darüberhinaus wird zur Differenzierung der Punktmengen ein algebraischer Körper (vorwiegend der Körper reeller Zahlen) verwendet. Diese Begriffe und Symbole des als Deskriptionsaspekts dienenden Aspekts der mathematischen Analysis innerhalb der anthropomorphen Logik sollen zur formalen Beschreibung syntrometrischer Operationen verwendet werden. Zunächst kommt es darauf an, mit diesem Deskriptionsaspekt den Begriff des allgemeinen subjektiven Aspekts zu erweitern. Durch die drei Bestimmungsstücke, nämlich Dialektik, Korrespondenzschema und Prädikatrix, ist ein subjektiver Aspekt vollständig definiert, doch kann ein solcher Aspekt, da die in ihm mögliche Aussagenmannigfaltigkeit kontinuierlich sein muß, eindeutig nur einem logischen System angehören; denn würde ein subjektiver Aspekt simultan zu > 1 logischen Systemen gehören, so müßte seine Aussagenmannigfaltigkeit in s kontinuierliche Einzelbereiche zerfallen, von denen jeder einzelne wieder ein subjektiver Aspekt in jeweils nur einem logischen System ist, so daß nur $s = 1$ sein kann.

Gegeben sei ein in irgendeinem logischen System definierter subjektiver Aspekt S und irgendeine in dem betreffenden logischen System ausdrückbare Vorschrift α , welche p -deutig (p ist ganzzahlig und im Deskriptionsaspekt definiert) ein, oder mehrere Bestimmungsstücke von S modifiziert, so daß a gemäß α ; $S \equiv S_{(j)}$ aus S insgesamt $1 \leq j \leq p$ neue subjektive Aspekte $S_{(j)}$ aus S entstehen läßt. Auf jeden Aspekt $S_{(j)}$ kann a abermals einwirken und so weiter. Wird dieser Prozess nach m Schritten abgebrochen, so sind p^m neue Aspekte aus S entstanden. Für $m \rightarrow \infty$ hat a aus S eine p -fach unendliche Mannigfaltigkeit von subjektiven Aspekten erzeugt, welche ein System subjektiver Aspekte mit der Dimensionalität p bildet, so daß a als Systemgenerator bezeichnet werden kann. Die Dimensionalität eines solchen Systems subjektiver Aspekte wird mithin durch die Deutigkeit des Systemgenerators bestimmt. In der Mannigfaltigkeit subjektiver Aspekte (p -fach unendlich), kann jedem subjektiven Aspekt der Punkt eines p -dimensionalen abstrakten metaphorischen Raumes zugeordnet werden, so daß die Gleichheit aller dieser Punkte als relatives *Aspektivfeld* dem Raum bestimmte metrische Eigenschaften vermittelt. Das p -dimensionale Aspektivfeld hat also eine bestimmte metrische Form, die Metropie, welche neben p und dem Systemgenerator α ein Bestimmungsstück des Aspektivfeldes ist. Da die Metropie durch eine Metrik des abstrakten Raumes zu veranschaulichen ist, könnten die Aspektivfelder in ihren Metropieformen im Analogon mit entsprechenden metrischen Feldern abstrakter Räume verglichen werden. Das aus a und dem Primäraspekt S entstandene p -dimensionale Aspektivfeld ist mithin ein strukturiertes System subjektiver Aspekte und muß daher als ein Aspektivsystem P bezeichnet werden. P wird durch seine vier Bestimmungsstücke, nämlich dem Systemgenerator α , Dimensionalität p , Metropie g und die Wirkungsweise von α auf den Primäraspekt S vollständig gekennzeichnet, so daß ein Aspektivsystem in der Form $P \equiv \left(\begin{matrix} \alpha; S \\ p; g \end{matrix} \right)$ symbolisiert werden kann.

S ist dabei nicht ausgezeichnet, vielmehr kann jedes Element aus P als Primäraspekt gewählt werden, wobei sich allerdings die Metropie wegen ihres relativen Charakters ändern kann. Der Systemgenerator α kann entweder

in den Begriffen des Primäraspektes ausdrückbar sein, oder aber in denen irgendeines anderen subjektiven Aspektes aus dem betreffenden logischen System. Im ersten Fall ist P entartet, im zweiten Fall dagegen echt. Im Folgenden sollen nur echte Aspektivsysteme untersucht werden, also solche, in denen α nicht in den Begriffen des Primäraspektes ausdrückbar ist. Auf keinen Fall ist dagegen a in einem anderen logischen System definiert, in welchem auch der Primäraspekt liegt. Die Metropie von P ist relativ und von der speziellen Wahl des Primäraspektes abhängig. Ist γ irgendeine Vorschrift, welche dem zugrunde gelegten Primäraspekt gegen irgendeinen anderen subjektiven Aspekt des Aspektivsystems austauscht, so moduliert γ gemäß $G \equiv \gamma; g$ als Metropiemodulator zugleich die Metropie des Systems. Diese *Metropiemodulation* kann q -fach gemäß $T \equiv \gamma^q; g$ wiederholt werden und wird als diskrete Metropiemodulation bezeichnet, wenn die Iterationszahl $q < \infty$ bleibt, doch wird sie kontinuierlich für $q \rightarrow \infty$. Es muß grundsätzlich drei verschiedene Gruppen von Systemen geben, unabhängig davon, wie der Systemgenerator a hinsichtlich seiner Dimensionalität oder das Aspektivsystem hinsichtlich seiner Metropie beschaffen ist. Jeder Primäraspekt wird durch Dialektik, Korrespondenz und Prädikatrix vollständig bestimmt, so daß es für α die Möglichkeiten von zwei verschiedenen Wirkungsweisen gibt, welche entweder partielle oder totale Aspektivsysteme entstehen lassen. Im Fall der einfachen partiellen Systeme wirkt a nur auf ein Bestimmungsstück des G ein, was für diese einfach partiellen Systeme Systeme $\binom{3}{1} = 3$ Möglichkeiten, nämlich dialektische, koordinative und prädikative Aspektivsysteme offen läßt.

Auch für die zweifach partiellen Systeme, welche durch simultane Einwirkung des Systemgenerators auf zwei Bestimmungsstücke des Primäraspektes entstehen, gibt es $\binom{3}{1} = 3$ Möglichkeiten, nämlich dialektisch-koordinative, dialektisch-prädikative und koordinativ-prädikative Aspektivsysteme. Die Totalsysteme sind dagegen eindeutig, denn wenn α simultan auf alle drei Bestimmungsstücke eines S einwirkt, gibt es wegen $\binom{3}{3} = 1$ nur eine Möglichkeit der Kombination. In den beiden partiellen Fällen muß α immer so beschaffen sein, daß die Aussagendeutigkeit n der subjektiven Aspekte nicht geändert wird, denn andernfalls müssten subjektive Aspekte entstehen, deren Bestimmungsstücke zueinander innerhalb eines Aspektes verschiedene Wertigkeiten erhalten, ein solches Aussagensystem kann aber nicht mehr als subjektiver Aspekt und somit als Element eines Aspektivsystems bezeichnet werden, weil einzelne Diatropen, Korrespondenz- oder Prädikatbänder als Restbänder nicht mehr korrelieren. Im Fall der totalen Aspektivsysteme ist dagegen eine solche Änderung der Aussagendeutigkeit durch den Systemgenerator möglich, denn bei der totalen Einwirkung des Generators kann es nicht zur Bildung unkorrelierbarer Restbänder kommen. Allerdings besteht die Möglichkeit, daß Aspektivsysteme entstehen, in denen Gruppen von subjektiven Aspekten enthalten sind, welche sich in ihrer jeweiligen Aussagendeutigkeit unterscheiden. Totale Aspektivsysteme mit dieser Eigenschaft sind singular im Gegensatz zu den regulären Systemen, deren Elemente (also die aus dem Primäraspekt resultierenden subjektiven Aspekte) sich nicht in ihrer Aussagendeutigkeit unterscheiden. Neben den totalen Aspektivsystemen, die durch die simultane Einwirkung des Systemgenerators auf alle drei Bestimmungsstücke des S entstehen, muß es noch partielle Aspektivsysteme geben. Wirkt α auf nur zwei Bestimmungsstücke des S ein, so gibt es drei Möglichkeiten, nämlich prädikative, koordinative und dialektische partielle Aspektivsysteme.

Auch wenn a nur auf ein Bestimmungsstück des S einwirkt, ergeben sich drei Möglichkeiten. Diese partiellen Aspektivsysteme, welche als einfach oder zweifach partiell aus a hervorgehen, sind offensichtlich untereinander verwandt und bilden dann Komplexe einfach oder zweifach partieller Aspektivsysteme, die sogenannten *Aspektivkomplexe*. Diese Aspektivkomplexe und totalen Aspektivsysteme sind demnach in sich selbst geschlossene Aussagensysteme aus einer endlichen oder unbegrenzten Zahl subjektiver Aspekte. Auch muß es eine unbegrenzte Zahl solcher Aspektivkomplexe (totale Aspektivsysteme sind Sonderfälle solcher Komplexe) geben, denn jeder denkbare subjektive Aspekt kann zur Erzeugung von Aspektivkomplexen verwendet werden, deren Zahl von der in dem betreffenden S definierbaren Zahl von Systemgeneratoren abhängt, die im allgemeinen als sehr groß, oder aber auch als unbegrenzt zu veranschlagen ist. Das anthropomorphe System ist offenbar ein Aspektivsystem aus einem nicht übersehbaren Aspektivkomplex. Von diesen Komplexen muß es wiederum eine unbegrenzte Zahl möglicher Formen geben. Schließlich können noch alle diejenigen Aspektivkomplexe zu einer *Aspektivgruppe* zusammengefaßt werden, deren Systemgeneratoren aus ein und demselben subjektiven Aspekt hervorgehen. Es ergibt sich also die folgende metropische Hierarchie der Aussagensysteme: In einem subjektiven Aspekt aus Prädikatrix, Koordination und Dialektik sind in den Ausdrucksmöglichkeiten des Aspektes Systemgeneratoren definiert, welche die Bestimmungsstücke des S umformen, und aus dem S eine Folge von subjektiven Aspekten entstehen lassen. Jeder Systemgenerator erzeugt auf diese Weise einen Aspektivkomplex, und die Gesamtheit aller in S ausdrückbaren Systemgeneratoren erzeugt eine übergeordnete Gesamtheit von Aspektivkomplexen, nämlich die Aspektivgruppe, für deren Bildung es wieder eine unbegrenzte Zahl

von Möglichkeiten gibt.

1.3. Kategorien Die vorangegangenen Untersuchungen sind Untersuchungen der logischen Aussagemöglichkeiten. Wie die Systeme solcher Aussagemöglichkeiten auch immer beschaffen sein mögen, müssen sie sich, wenn sie überhaupt einen Sinn haben sollen, auf begriffliche Elemente beziehen, über deren Eigenschaften und wechselseitigen Zusammenhänge die betreffenden Aussagen zu machen sind. Nach diesen vorangegangenen Untersuchungen der Aussagemöglichkeiten, die zur Definition der Aspektivkomplexe und der allgemeinen übergeordneten Aspektivgruppen führte, erscheint es angebracht, die Begriffselemente, auf welche die Aussagenanalyse angewendet werden soll, zu analysieren und in der allgemeingültigsten Form zu charakterisieren. Gegeben sei ein System aus $1 \leq k \leq N$ Begriffsgruppen a_k , von denen jede wiederum aus n_k Begriffselementen b_i mit $1 \leq i \leq n_k$ besteht, derart, daß das ganze System $\sum_{k=1}^N n_k$ Begriffselemente enthält. Weiter werde angenommen, daß alle diese Elemente durch irgendwelche Schlußweisen auseinander hervorgehen, so daß diese Schlußweisen die jeweils entstehenden Elemente an irgendwelche Bedingtheiten binden. Auch sei jede Begriffsgruppe a_k ein ganzes Syndrom von Elementen, welche an die gleiche Zahl von Bedingtheiten gebunden sind. Wenn dies aber so ist, so kann angenommen werden, daß die Folge der Syndrome a_k so geordnet ist, daß in der Richtung $1 \leq k \leq N$ die Zahl der Bedingtheiten ansteigt, das heißt, das Syndrom k enthält Elemente, die an eine geringere Zahl von Bedingtheiten gebunden sind als $k + 1$, aber an eine höhere als $k - 1$. In dem so geordneten System der Gruppen von Begriffselementen, die durch irgendwelche Schlußweisen und Bedingtheiten auseinander hervorgehen, herrscht demnach ein Syllogismus, und zwar ein Episyllogismus beim Durchlaufen des Systems in der Richtung $1 \leq k \leq N$ zunehmender Bedingtheiten, und ein Prosyllogismus in umgekehrter Richtung. Da in einem jeden System, dessen Begriffselemente durch Schlußweisen auseinander hervorgehen, eine solche Anordnung nach Bedingtheiten möglich ist, liegt in einem solchen Begriffssystem auch immer ein Syllogismus vor. Wenn die N Syndrome a_k so geordnet sind, daß mit ansteigenden k ein Episyllogismus vorliegt, so sind die Begriffselemente des Syndroms $k = 1$ an gar keine Bedingtheiten gebunden, d.h. $k = 1$ kann als *textitIdee* des ganzen Begriffssystems bezeichnet werden, aus welcher alle übrigen Elemente durch die Art der betreffenden Schlußweise im Sinne des Episyllogismus hervorgehen. Ganz allgemein könnte also im Falle dieses Episyllogismus das Syndrom k als Idee von $k \neq 1$ aufgefaßt werden usw., doch ist stets $k = 1$ die allgemeine Idee des Systems. Sind die Syndrome im Sinne eines Prosyllogismus geordnet, so kehrt sich die ganze Betrachtung um. Auf jeden Fall kann im Episyllogismus von 1 nach N die Gesamtheit der $N - 1$ Syndrome $k > 1$ als Begriffskategorie aufgefaßt werden, deren Idee $k = 1$ ist. Das ganze nach einem Syllogismus geordnete System aus Begriffselementen, die wiederum nach dem syllogistischen Ordnungsgesetz zu Begriffssyndromen zusammengefaßt sind, besteht demnach aus einer Idee und einer syllogistisch orientierten Begriffskategorie. Wenn dieses System aber vollständig ist, so bilden Idee, kategorie und Syllogismus eine Einheit, welche als allgemeine Kategorie bezeichnet werden soll. Das Vollständigkeitskriterium einer Kategorie wird offensichtlich durch die Idee und die syllogistische Schlußweise bedingt; denn diese beiden Bestimmungsstücke machen die Überprüfung der Begriffskategorie auf Vollständigkeit und Fremdelemente möglich, die nicht zu der betreffenden Begriffskategorie gehören. Die notwendige und hinreichende Existenzbedingung einer Kategorie ist demnach die Existenz einer Idee und einer syllogistischen Schlußweise; denn aufgrund dieser Schlußweise können die vollständigen Syndrome der Kategorie aus der Idee induziert werden. Die in den Aspektgruppen zusammengefassten, in sich selbst geschlossenen Aussagesysteme der Aspektivkomplexe, werden im allgemeinen die Aussagen irgendeines subjektiven Aspektes (also das durch ein Adjektiv dialektisch aus einer Diatropen geprägte Prädikat eines Prädikatbandes) auf solche Kategorien und deren Zusammenhänge beziehen. Aus diesem Grunde erscheint es im Hinblick auf einen syntrometrischen Formalismus notwendig, diesen Begriff der Kategorie und der über sie möglichen Aussagen konkreter zu formulieren.

1.4. Die apodiktischen Elemente Zur Weiterführung der Untersuchung wird zunächst eine Analyse des Systems anthropomorpher Aussagemöglichkeiten und Schlußweisen nötig. Offenbar bilden alle die anthropomorphen logischen Elemente partielle Komplexe in erster und zweiter Ordnung, während die möglichen Prädikatbildungen immer zweideutig sind. Als Systemgenerator kommt nur ein α in Betracht, welcher einfach partiell, dialektisch oder koordinativ einwirkt, oder zweifach partiell dialektisch koordinativ, während die zweideutige Prädikatrix in jedem Fall ungeändert bleibt. Prädikatbänder existieren nicht, in ihr sind nur zwei kontradiktorische Elemente (Bejahung und Verneinung) enthalten. Demzufolge sind auch die Diatropen- und Koordinationsbänder zu diskreten Einzelementen entartet. Der Wirkungsweise des Systemgenerators entsprechend, müssen alle Möglichkeiten anthropomorpher Logik in einem zweideutig prädikativen Aspektivkomplex enthalten sein, der seinerseits aus den drei ebenfalls zweideutigen prädikativen Aspektivsystemen (der Wirkungsweise des Systemgenerators entsprechenden dialektisch, koordinativ

und dialektisch-koordinativ) zusammengesetzt sein. Wird, bezogen auf irgendein Aspektivsystem dieses Komplexes, ein Bereich analysiert, so zerfällt die Analyse in diesem zweideutig prädikativen Aspektivkomplex, bezogen auf einen geeigneten subjektiven Aspekt des betreffenden Aspektivsystems, in folgende Schritte: a) Abgrenzung des fraglichen Bereichs begrifflicher oder empirischer Elemente. b) Qualitative Analyse dieser Elemente, nach deren Ergebnis der geeignete subjektive Aspekt ausgewählt und damit das Aspektivsystem festgelegt wird. c) Quantitative Analyse und Synthesis der Elemente, bezogen auf den festgelegten subjektiven Aspekt. d) Von der Analyse und Synthese einer ästhetischen Empirik wird der Übergang zu einer Transzendentalästhetik vollzogen. e) Es werden indirekte transzendente Schlußweisen nach Durchführung der Abstraktion *d* angewendet, deren Konsequenzen Rückschlüsse auf die transzendentalen Zusammenhänge der Elemente des ganzen Bereichs zulassen.

Offenbar kann eine solche transzendente Analyse in jedem Aspektivsystem und auch in jedem Aspektivkomplex, also unabhängig von den jeweiligen Metropiefeldern, der subjektiven Aspekte durchgeführt werden, wobei sich allerdings die Form der Methodik nach der Eigenart des jeweiligen Aspektivkomplexes oder der übergeordneten Aspektivgruppe richten muß. Diese Möglichkeit einer deskriptiven Methodik muß ein Charakteristikum aller Aspektivsysteme sein, denn wie ein solches System oder eine übergeordnete Aspektivgruppe auch immer beschaffen sein mag, auf Grund des Charakters der Aussagefähigkeit der sie strukturierenden subjektiven Aspekte muß prinzipiell eine prädikative Methodik der Deskription möglich sein. Wenn dies aber so ist, dann muß die Existenz der transzendentalen Methodik für jede Aspektivgruppe gelten, so daß hier der Ansatz zur Abstraktion von der anthropomorphen Transzendentalästhetik liegt. Unabhängig von dieser Universalität muß festgestellt werden, daß die jeweilige prädikative Methodik der Deskription stets auf die Charakterisierung von Eigenschaften der Elemente und ihrer Wechselbeziehung abgegrenzter Bereiche hinausläuft. Da die Charakterisierung von Eigenschaften aber stets mit einem Beimmessen von Bedeutungen verbunden ist, muß die Methodik stets semantischer Natur sein. Ein apodiktisches Charakteristikum einer jeden Aspektivgruppe, welches die Abstraktion von der anthropomorphen Transzendentalästhetik ermöglicht, ist mithin die Existenz einer dem jeweiligen Aspektivsystem adäquaten semantischen Methodik. Im Allgemeinen liegt eine unbegrenzte Zahl von Eigenschaften eines Bereichs vor, wenn der Bereich selbst unbegrenzt ist, doch bedeutet die Abgrenzung des Bereichs eine obere Schranke für die Zahl seiner Eigenschaften. Ist das passende Aspektivsystem zu Grunde gelegt, und die Semantik entwickelt, so zeigt sich, daß die semantische Bewertung der Eigenschaften vom jeweiligen subjektiven Aspekt abhängt, das heißt, wenn die semantische Methodik in allen subjektiven Aspekten angewendet wird, also wenn man im Metropiefeld der subjektiven Aspekte fortschreitet, kommt es zu einer allgemeinen Varianz semantischer Bewertungen, doch wird eine endliche Zahl von Eigenschaften des Bereichs mit invarianter Semantik im ganzen Metropiefeld des Aspektivsystems nach dem Durchlaufen des Metropiefeld erscheinen. Offenbar sind diese Eigenschaften begriffliche Elemente des Bereichs, deren Semantik in keinem Punkt des Metropiefeldes geändert wird, derart, daß ihre Bedeutungen vom jeweiligen subjektiven Aspekt unabhängig bleiben. Diese Elemente eines Bereichs können also in Bezug auf das betreffende System als apodiktische Elemente bezeichnet werden, und zwar können sie einfach, komplex oder total apodiktisch sein, je nachdem, ob sich ihre Apodiktik auf ein Aspektivsystem, einen Aspektivkomplex oder eine Aspektivgruppe bezieht. Da alle Eigenschaften varianter Semantik des Bereichs aus den apodiktischen Elementen durch geeignete Korrelationen hervorgehen müssen, könnte das System apodiktischer Elemente als Idee einer Kategorie, und der Bereich selbst als vollständige oder unvollständige Kategorie aufgefaßt werden, derart, daß relativ zum Aspektivsystem als Idee des Bereichs das System apodiktischer Elemente anzusprechen ist.

Zwar ist die Idee hinsichtlich ihrer Semantik im Metropiefeld invariant, doch kann dies unmöglich für die Korrelationsmöglichkeiten der apodiktischen Elemente gelten; denn diese Möglichkeiten können nur von der Struktur des jeweiligen subjektiven Aspektes bestimmt werden. In jedem subjektiven Aspekt gibt es also eine endliche oder unendliche Schar von möglichen Korrelationen, und jede Korrelation induziert aus der Idee invarianter Semantik wiederum eine endliche oder unendliche Schar von Eigenschaftssyndromen im Sinne einer Kategorie, deren Besetzungen aber wegen der varianten Semantik jeder Korrelation ebenfalls vom subjektiven Aspekt abhängen, derart, daß der Varianzgrad ihrer Semantik mit steigendem Episylogismus, also wachsender Syndromanzahl, zunimmt. Die endliche Zahl apodiktischer Elemente liefert demnach in jedem subjektiven Aspekt des Metropiefeldes eine endliche oder unendliche Schar von Kategorien mit begrenzter oder unbegrenzter Syndromfolge, die in ihrer Gesamtheit alle Eigenschaften des Bereichs von allen subjektiven Aspekten her umfassen. Ist in dem begrifflichen Bereich, bezogen auf ein ausgewähltes System, überhaupt kein apodiktisches Element festzustellen, so muß ein anderes Aspektivsystem gewählt werden, oder aber der Bereich ist in anderer Form abzugrenzen.

Die heuristische Methodik zur Auffindung apodiktischer Elemente, bezogen auf einzelne subjektive Aspekte, geht im Prinzip auf die Begrenzung des Prosyllogismus einer Kategorie durch die Idee zurück. Empirisch werden dabei alle diejenigen Eigenschaften des Bereichs, bezogen auf einen festgelegten subjektiven Aspekt, ausgewählt, die relativ zu diesem Aspekt durch Korrelationen auseinander hervorgehen. Jede empirische aufgefundene Korrelation liefert dann empirisch eine Gruppe von Eigenschaften als unvollständige Kategorie, die nach dem Grad ihrer Bedingtheiten im Sinne eines Prosyllogismus anzuordnen sind. Zu jeder Gruppe gehört dann ein empirischer Prosyllogismus, und alle diese Prosyllogismen müssen dann in einer Gruppe apodiktischer Elemente münden, die jedoch nicht vollständig zu sein braucht. Eine entsprechende Empirik kann auch in anderen subjektiven Aspekten des Metropiefeldes verwendet werden, so daß ein Vergleich der apodiktischen Elemente verschiedener Aspekte zur Vervollständigung der Idee des Bereichs führen muß. Daraus folgt, daß die Vollständigkeit der Idee umso größer ist, je mehr Aspekte empirisch verwendet werden. Ein Vollständigkeitskriterium einer Idee kann es nicht geben, weil die Begrenzung des Bereichs vorerst willkürlich und damit provisorisch bleiben muß, weil die Zahl der Aspekte einer Metropiefeldes, und auch die Zahl der begrifflichen Elemente, sowie die Eigenschaften eines Bereichs unbegrenzt und damit unfaßbar sein kann. Hat die empirische Ästhetik der unvollständigen Prosyllogismen zu einer hinreichenden Zahl apodiktischer Elemente geführt, was einer Induktion gleichkommt, so kann der Übergang zur Transzendentalästhetik erfolgen. Die empirisch gewonnenen apodiktischen Elemente werden zur Idee eines Bereichs zusammengefaßt, dessen Begrenzung jetzt nicht mehr provisorisch und willkürlich ist, denn die möglichen Kategorien dieser Idee in den einzelnen subjektiven Aspekten des Metropiefeldes, müssen sämtliche Eigenschaften desjenigen Bereichs darstellen, dessen Idee, bezogen auf das zu Grunde gelegte Aspektivsystem, aus den vorhandenen apodiktischen Elementen besteht. Liegt eine komplexe oder totale Apodiktik vor, so bezeichnen sich die Aussagen über den Bereich auf entsprechende Aspektivkomplexe, oder Aspektivgruppen. Der Übergang zur Transzendentalästhetik und die damit verbundene Entwicklung aller Eigenschaften eines vollständigen begrifflichen Bereichs in den einzelnen subjektiven Aspekten aus seinen apodiktischen Elementen heraus, würde einem Deduktionsschluß entsprechen. Gegenüber dem provisorischen begrifflichen Bereich hat der transzendental entstandene Bereich eine Transformation erfahren, die den ursprünglichen provisorischen Bereich dort erweitert, wo die willkürliche Begrenzung nicht apodiktische Elemente ausgrenzte, aber dort einschränkt, wo es sich um Eigenschaften handelt, die auf empirische nicht erfasste apodiktische Elemente zurückgehen. Auf jeden Fall ist eine konkrete Analyse aller Eigenschaften des transzendental begrenzten Bereichs möglich, weil er vollständig sein muß, wenn eine Analysis der Kategorien möglich wird.

1.5. Aspektrelativität. Funktor und Quantor Sind a und b zwei apodiktische Elemente in Bezug auf ein Aspektivsystem A (was auch ein Komplex oder eine Gruppe sein kann, nämlich dann, wenn a und b komplex oder total apodiktisch sind), so können diese durch die Aussage γ eines zu A gehörenden subjektiven Aspektes S miteinander verknüpft sein. Wird als verallgemeinertes Aussagesymbol $\overline{\parallel}$ verwendet, so bedeutet $\overline{AS}|_{\gamma}$, daß es sich um eine Aussage aus S in A handelt, und zwar um das Prädikat γ aus S , welches seiner Koordination entsprechend durch ein dialektisches Adjektiv geformt wurde. $a, \overline{AS}|_{\gamma}, b$ kennzeichnet also die Wechselbeziehung, die durch diese Aussage a und b in die Relation setzt. $\overline{\parallel}_{\gamma}$ ist demnach das a und b bezogen auf S in A verknüpfende Prädikat. Diese Verknüpfung braucht nicht nur apodiktische Elemente in Zusammenhang bringen. Sind es z.B. a_i mit $1 \leq i \leq p$ und b_k mit $1 \leq k \leq q$ zwei Komplexe apodiktischer Elemente in A und stehen bezogen auf S diese beiden apodiktischen Gruppen in den nicht apodiktischen Zusammenhängen $F(a_i)_1^p$ und $\Phi(b_k)_1^q$, so können über S auch F und Φ durch das gemeine Prädikat $F, \overline{AS}|_{\gamma}, \Phi$ oder kürzer $F, \overline{\parallel}_{\gamma}, \Phi$ (wenn A und S festliegen) verknüpft werden. Zwar sind die beiden begrifflichen Zusammenhänge F und Φ , die als Begriffsfunktionen durch den *Funktor* F bzw. Φ die apodiktischen Elemente a_i bzw. b_k in einen begrifflichen Zusammenhang setzen, einzeln nicht apodiktisch, doch besteht die Möglichkeit, daß es zu F und Φ sowohl als auch zu $\overline{\parallel}_{\gamma}$ in allen anderen subjektiven Aspekten aus A Äquivalente gibt derart, daß die Verknüpfung $F, \overline{\parallel}_{\gamma}, \Phi$ in A selber apodiktisch erscheint. Derartige apodiktische Verknüpfungen nicht apodiktischer Begriffsfunktionen sind aber von der speziellen Wahl des A (hinsichtlich ihrer Existenz) unabhängig, und müssen daher in allen Aspektivsystemen möglich sein. Zur Unterscheidung zwischen den einfachen, nur über einem speziellen S gegebenen Funktorzusammenhängen $(), \overline{\parallel}_{\gamma}, ()$ und den im ganzen A apodiktischen zusammenhängen, werden diese apodiktischen Verknüpfungen als *Quantoren*, symbolisiert durch $(), \overline{\parallel}_{\gamma}^1, ()$ bezeichnet, denn ein solcher Quantor beschreibt seine Aussage zwischen nicht tischen Funktoren, also begrifflichen Elementen des Bereichs, die in allen subjektiven Aspekten gilt, und daher bezogen auf A eine allgemeine qualitative Eigenschaft des Begriffs darstellt. Ein solcher, nur in einem Aspektivsystem A gültiger Quantor ist demnach ein

Monoquantor, zu dessen vollständiger Beschreibung noch die Angabe von A , also $(\cdot, {}^1\overline{||}_\gamma, \cdot)$ notwendig während sich die Angabe von S erübrigt, weil dieser Quantor in allen S gilt. Bilden zwei Funktoren einen solchen Monoquantor, so werden durch die Funktoren offensichtlich zwar nicht apodiktische, aber wesentliche Charakterzüge des begrifflichen Bereichs erfaßt, die durchaus Wahrheiten dieses Bereichs sein können. Diese Darstellung des Quantors macht eine Erweiterung zum *Polyquantor* möglich. Ist B ein anderes Aspektivsystem, das aus A durch eine metrische Deformation des Metropiefeldes hervorgegangen ist, so sind im allgemeinen die a_i und b_k in B nicht mehr apodiktisch, doch sind sie auf jeden Fall, wenn es sich um die Beschreibung des gleichen begrifflichen Bereichs handelt, als Funktoren apodiktischer Elemente des Bereichs bezogen auf B darstellbar, und auch die beiden Funktoren F und Φ erfahren eine der Metropiefelddeformation entsprechende Umdeutung auf B . Sind die neuen Funktoren in B bezeichnet durch F' und Φ' , dann gibt es zwischen ihnen auch in B , bezogen auf S' , nicht apodiktische Funktorzusammenhänge der Form $F', \overline{||BS'}_\gamma, \Phi'$, während der Zusammenhang in A ein quantor war. Erweist sich die Verknüpfung aber auch in

B als Quantor, so wird diese zweifache Quantornatur, wenn $B \equiv A_2$ gesetzt wird, beschrieben durch $(\cdot, {}^1\overline{||A_k}^2, \cdot)$, was soviel bedeutet, daß die beiden Funktoren sowohl in A_1 als auch in A_2 im Quantorzusammenhang stehen, was diesen Quantor zum Biquantor werden läßt. Eine solche Biquantorausage über den Bereich hat auf jeden Fall einen höheren Wahrheitsgehalt als der Monoquantor. Diese Schlußweise kann weitergeführt werden. Im allgemeinsten Fall schließlich würde sich die Verknüpfung von zwei Funktoren in einem Komplex aus $1 \leq \rho \leq r$ Aspektivsystemen A_ρ

als apodiktisch erweisen, und dann würde $(\cdot, {}^1\overline{||A_\rho}^r, \cdot)$ ein allgemeiner Polyquantor vom *Wahrheitsgrad* r sein. Ist ein Glied ρ des Polyquantors so beschaffen, daß die in Korrelation stehenden Funktoren direkt apodiktische Elemente in Relation setzen, d.h., sind die Funktorargumente selbst apodiktisch, so ist dieses Quantorglied absolut apodiktisch, anderenfalls, also bei nicht apodiktischen Funktorargumenten, semiapodiktisch, weil die Verknüpfung selbst noch apodiktisch ist. Diese Definition semiapodiktischer Quantorglieder gilt auch dann, wenn nur ein Funktorargument nicht apodiktisch ist (semiapodiktisch im ersten Grad; aber im zweiten Grad, wenn beide Funktorargumente nicht apodiktisch sind). Aus diesem Sachverhalt ergibt sich unmittelbar der Satz, daß in jedem Polyquantor mindestens ein Glied absolut apodiktisch ist. Zusammenfassend gilt also für die Verknüpfung apodiktischer Elemente, sowie der Funktoren apodiktischer Argumente durch die Aussage γ des subjektiven Aspekts S (nach (1)) im Aspektivsystem A die Darstellung:

$$a, \overline{||AS}_\gamma, b \vee F(a_i)_1^p, \overline{||AS}_\gamma, \Phi(b_k)_1^q \quad (2)$$

während für den allgemeinen Polyquantor

$$(\cdot, {}^1\overline{||A_\rho}^r, \cdot) \quad (3)$$

gilt. Die Tatsache, daß in einem Polyquantor absolut apodiktische und semiapodiktische Glieder im ersten und zweiten Grad auftreten, zeigt, daß die apodiktischen Eigenschaften eines Quantors (jedes Glied eines Polyquantors kann als einfacher Quantor aufgefaßt werden) relativ sind, und vom jeweiligen Aspektivsystem abhängen; denn absolute und semiapodiktische Quantoreigenschaften sind nur im Komplex des Polyquantors möglich, während die Quantoreigenschaften verloren gehen, wenn die Funktorrelation auf ein außerhalb dieses Komplexes liegendes Aspektivsystem bezogen wird. Auch innerhalb des Polyquantorkomplexes existiert eine solche Aspektrelativität zwischen den absoluten und semiapodiktischen Quantoreigenschaften. Die Existenz der Aspektrelativität quantorhafter Funktorverknüpfungen gestattet es also in Bezug auf einen vorgegebenen Komplex unabhängiger Aspektivsysteme (diese brauchen keine Aspektivkomplexe oder -gruppen zu bilden) die Quantoren eines begrifflichen Bereichs aufzufinden, oder aber zu einem Monoquantor ein System von Metropiefeldern so zu konstruieren, daß ein Komplex von Wahrheitssystemen entsteht, auf den der Monoquantor bezogen zum Polyquantor wird, dessen Wahrheitsgehalt mit der Zahl der Aspektivsysteme identisch wird. Dieser Wahrheitsgrad wird offensichtlich umso größer, je mehr Metropiefelder zu diesem Komplex konstruiert werden können. Diese Konstruktion eines Polyquantors ist deshalb möglich, weil die absoluten oder semiapodiktischen Quantoreigenschaften innerhalb eines solchen Komplexes zwar noch vom speziellen Aspektivsystem abhängen, aber die Existenz des Quantors an sich innerhalb des konstruierten Komplexes hinsichtlich der Aspektivsysteme invariant bleibt. Schließlich besteht noch eine weitere Möglichkeit zur Verallgemeinerung des Quantorbegriffs. Zeigt sich nämlich bei der Konstruktion des Polyquantors aus dem Monoquantor im Aspektivsystem

A , daß es einen Modulator f des zu A gehörenden Metropiefeldes α gibt, derart, daß gemäß $f; \alpha \equiv \beta$ aus dem einen Metropiefeld a durch eine kontinuierliche metrische Deformation auf Grund des Modulators f eine mehrfach unendliche Schar neuer Metropiefelder β hervorgeht, und wenn die Funktorverknüpfung als Quantor bezogen auf A , auch bezogen auf das ganze Kontinuum von Aspektivsystemen B der Metropiefelder β ebenfalls Quantoreigen-

schaften hat, so liegt ein kontinuierlicher Quantor $(\cdot, \xrightarrow{f} \cdot, \cdot)$ vor. Der diskrete Polyquantor aus der Beziehung (3) ist demnach ein spezieller Sonderfall des kontinuierlichen Quantors, und dieser Quantor wiederum ist der Sonderfall eines noch allgemeineren Quantors, nämlich des kontinuierlichen Polyquantors vom Grade r , der aus $1 \leq \rho \leq r$ kontinuierlichen Quantorgliedern besteht. Ein solcher kontinuierlicher Polyquantor existiert also immer dann, wenn es r Aspektivsysteme A_ρ mit den Metropiefeldern α_ρ und ebensoviel Metropieminimodularen f_ρ gibt, derart, daß die Folgen $\beta_\rho \equiv f_\rho; \alpha_\rho$ und damit die unendlichen Folgen B_ρ aus den A_ρ entstehen, und die Funktorverknüpfungen in Bezug auf diese r unendlichen Folgen B_ρ Quantoreigenschaften hat.

$$(\cdot, \xrightarrow{f_\rho} \cdot, \cdot) \vee \beta_\rho \equiv f_\rho; \alpha_\rho \vee \alpha_\rho \equiv A_\rho \vee \beta_\rho \equiv B_\rho \quad (4)$$

dürfte die universelle Form des Quantorbegriffes sein; denn für $r = 1$ folgt aus ihm der kontinuierliche Monoquantor (oder kurz kontinuierlicher Quantor), aber für $r > 1$, wenn f nicht existiert, der diskrete Polyquantor aus den für $r = 1$ der Monoquantor entsteht.

Die Begründung einer allgemeinen Syntrometrie kann nach den vorangegangenen Untersuchungen des Funktor- und Quantorbegriffes, sowie der Aspektrelativität, nur auf die Entwicklung einer transzendentalen Methode zurückgehen, die es gestattet, möglichst allgemeingültige Funktoren oder Funktorsysteme des Bereiches aufzufinden, die Polyquantoren mit möglichst hohem Grad bilden, oder aber Universalquantoren sind, die grundsätzlich in allen Aspektivsystemen Quantoreigenschaften haben, wobei noch zu untersuchen wäre, ob ein solcher Universalquantor überhaupt existieren kann.