

Contents

1 Physical Instantiation Search for Abstract Thermodynamic Compositions: An Exhaustive Analysis of Extreme Scaling, Duality Collapse, and Procedural Mapping

The translation of abstract thermodynamic compositions into physical instantiations requires a rigorous synthesis of continuum mechanics, quantum materials science, and non-equilibrium statistical physics. The search for specific phenomenological tuples—namely, systems exhibiting flux density divergence at vanishing temperature gradients, high-collision geometric confinement, and recombination suppression via structural relaxation—reveals profound intersections between macroscopic thermodynamic laws and microscopic transport mechanisms. By utilizing advanced topological models, strain engineering, and entropic confinement, it is possible to instantiate systems that appear to bypass classical kinetic limitations. This report provides an exhaustive evaluation of these abstract constraints, mapping them to specific materials, observed dualities, governing mechanisms, and technological readiness levels.

1.1 Tuple 1: Isothermal Divergence and Dissipative Coexistence

The first thermodynamic composition explores the extreme scaling regime where the temperature gradient approaches zero ($\nabla T \rightarrow 0$), yet the energetic or mass flux density exhibits divergent or highly anomalous scaling. In classical Fourier heat conduction, the heat flux vector is strictly proportional to the temperature gradient, expressed as $\mathbf{q} = -\kappa \nabla T$, where κ is the thermal conductivity.¹ Under standard localized equilibrium, the divergence of this vector field, $\nabla \cdot \mathbf{q}$, represents the volume density of the outward flux from an infinitesimal volume, equating to the local heat source intensity.²

In fluid dynamics and thermodynamics, the general equation of heat transfer links specific entropy production in a Newtonian fluid to thermal conduction and viscous forces. This is mathematically articulated as $\nabla \cdot \mathbf{q} = T \nabla^2 s + \eta \nabla^2 v^2 + \dots$.³ When the flow velocity is negligible, this reduces to the standard heat equation.³ Furthermore, the Onsager expression frames the divergence of an entropy flux density in terms of a scalar temperature potential and an energy flux density, $\nabla \cdot \mathbf{q}_s = -\nabla^2 \phi + \mathbf{q} \cdot \nabla (1/T)$.⁴ Forcing a condition where a substantial macroscopic flux is maintained as necessitates physical architectures that bypass classical phonon-mediated thermal conduction, relying instead on isothermal phase transitions, electrochemical potentials, or non-equilibrium dissipative ratchets.

1.1.1 Extreme Scaling: Near-Isothermal Heat Engines and Osmotic Systems

To instantiate a divergent or massive flux density near isothermal conditions, macroscopic systems must exploit chemical potential gradients or isothermal expansion mechanisms rather than pure thermal gradients. The Alkali-Metal Thermal to Electric Converter (AMTEC), originally conceptualized as the Sodium Heat Engine, provides a primary physical realization.⁵ AMTEC functions as a thermally powered electrochemical concentration cell operating at high temperatures ranging from 900 K to 1300 K.⁵ The core mechanism circumvents the Carnot limitations of standard volumetric thermal expansion by driving sodium around a closed thermodynamic cycle between a high-temperature reservoir and a cooler condenser at 400–700 K.⁵

The critical physical component of AMTEC is a highly ionically conducting refractory solid electrolyte—specifically, Sodium Beta"-Alumina Solid Electrolyte (BASE). BASE is a crystalline ceramic composed of aluminum oxide () and sodium oxide () in a nominal 5:1 ratio, stabilized by small cation metal oxides such as magnesium or lithium.⁵ Neutral sodium undergoes electrochemical oxidation at the anode, producing sodium ions that rapidly traverse the BASE lattice. Because BASE is an electrical insulator, the corresponding electrons are forced through an external circuit to perform direct electrical work before recombining with the ions at the low-pressure cathode to form sodium gas.⁵ The thermodynamic brilliance of AMTEC lies in the fact that the ion conduction process through the solid electrolyte is thermodynamically nearly equivalent to the isothermal expansion of sodium vapor between high-pressure and low-pressure regimes.⁵ Consequently, a massive electrical and mass flux is generated across a highly localized, near-isothermal interface. Single-cell AMTECs have demonstrated open-circuit voltages greater than 1.55 V, power densities exceeding 0.50 W/cm² at 1173 K, and laboratory efficiencies of 16%, with multi-tube modules predicted to reach 20% to 25%.⁵

Similarly, osmotic pressure engines, encompassing Pressure Retarded Osmosis (PRO) and Reverse Electrodialysis (RED), generate massive energetic flux through the strictly isothermal mixing of fluids with differing salinities.⁶ In PRO systems, the chemical potential difference between seawater and freshwater across a semi-permeable membrane drives a massive flux of water, increasing the pressure on the saline side to drive a mechanical turbine.⁶ Environmental applications of this extreme scaling are vast; for instance, in the Sebou estuary in Morocco, the maximum isothermal mixing energy is calculated at 0.712 kWh/m³, theoretically capable of generating 80.33 MW to 164 MW of continuous power purely from concentration gradients.⁸

1.1.2 Duality Collapse: Endoreversible Thermodynamics and Brownian Ratchets

The coexistence of reversible and irreversible dynamics in these heat and mass engines leads to the collapse of classical binary thermodynamic categorizations, best described by the formalisms of endoreversible thermodynamics. Endoreversible thermodynamics is a finite-time theoretical framework wherein a non-equilibrium macroscopic system is partitioned into a set of internally reversible subsystems that interact with one another solely through irreversible transport laws.⁹ The irreversibilities—and thus the entirety of the entropy production—are strictly localized to the interactions (such as Newton's law of cooling, Fourier conduction, or Fickian mass transfer) between the subsystems.¹⁰ This duality permits the exact analytical modeling of complex systems like combustion engines, hydraulic energy storage for recuperation, and Vuilleumier refrigerators, where the "engine" core is modeled as a reversible Carnot-like operator bounded by irreversible heat exchangers.⁹ The equivalence of enthalpy transfer to combined particle and entropy transfer allows predictive mapping of system efficiencies far beyond idealized quasi-static limits.¹¹

At the microscopic and molecular scale, this coexistence of reversibility and irreversibility is physically manifested in Brownian ratchets, also known as molecular motors. Brownian ratchets operate in spatially periodic, asymmetric potentials driven far from thermal equilibrium by thermal noise or time-dependent temperature modulations.¹³ The governing laws of equilibrium thermodynamics rigorously dictate that no directed work can be extracted from isothermal fluctuations, effectively preventing the realization of a Maxwell's Demon.¹⁴ However, Brownian ratchets break detailed balance by coupling the asymmetric potential to an external energy source, such as ATP hydrolysis in biological systems or flashing potentials in synthetic systems.¹⁴

This creates an anomalous transport phenomenon where a directed macroscopic flux is generated from random microscopic noise, exhibiting a coexistence of “irreversible” dissipative power strokes and highly reversible thermal transitions within the same continuous operational cycle.¹⁷ Modern theoretical frameworks resolve these dynamics using the Fokker-Planck equation and state-dependent free-energy landscapes, bridging the gap between discrete biochemical states (e.g., the four-state Lymn-Taylor cycle) and continuous mechanically-biased diffusion.¹⁷ Recent applications of the Curtin-Hammett principle to Brownian ratchets demonstrate that highly composition-dependent reaction rate coefficients inside condensed phases can either increase or paradoxically decrease overall reaction rates, illustrating negative differential responses in chemical reaction networks.¹⁹ Furthermore, there is an established duality between relaxation and first-passage processes in ergodic reversible Markovian dynamics; the time scales of relaxation and first passage interlace, allowing the full first-passage time distribution to be analytically determined from the simpler relaxation eigenspectrum.¹⁵

1.1.3 H_Mechanisms: Continuous Dissipative Cycling and NESS

Sustaining a diverging flux under conditions necessitates a mechanism capable of continuous dissipative cycling, specifically the establishment of a Non-Equilibrium Steady State (NESS). In a NESS, the macroscopic observables of the system remain constant over time, but unlike true thermodynamic equilibrium, there is a continuous production of entropy and a non-vanishing flux of matter or energy driven by external reservoirs. The local dispersion of the prehistory probability density in such systems can be reformulated as a Gaussian distribution corresponding to the linearized part of the reversed diffusion process.²¹ The local small-amplitude fluctuations of the stochastic variables around a deterministic trajectory lead to large excursions responsible for unidirectional motion in Brownian ratchets and enzyme kinetics.²¹

The compressible Euler equations provide a continuum-level understanding of this mass and momentum flux. Expressed via the divergence theorem, the continuity equation ensures mass conservation, while the momentum equation demonstrates that changes in momentum are due solely to gradients in pressure, independent of thermal gradients.²² By leveraging these pressure and concentration gradients, NESS architectures continuously harvest energy without violating the Second Law.

1.1.4 Procedural Mapping: Electrochemical Concentration Cells

Electrochemical concentration cells (ECC) serve as the primary procedural technology for mapping isothermal divergence. Beyond the high-temperature AMTEC systems, ambient-temperature ECCs are ubiquitously utilized for precision gas analysis and environmental monitoring. The ECC ozonesonde, developed by W.D. Komhyr in 1969, is an expendable, balloon-borne instrument that provides continuous, high-quality, in-situ measurements of stratospheric and tropospheric ozone with a vertical resolution of 100-150 meters up to 40 km altitudes.²⁴

The ozone sensor package operates purely as an electrochemical concentration cell where the oxidation and reduction reactions are identical, leading to a standard cell potential of exactly zero.²⁷ The system relies entirely on the Nernst equation’s second term, generating a current as ambient air is pumped through a buffered potassium iodide (KI) reagent in the sensing cathode cell.²⁴ The time response of these ECC ozonesondes reveals at least two distinct reaction pathways with time constants of approximately 20 seconds and 25 minutes.²⁹

Properly accounting for these time constants eliminates the need for ad hoc background corrections, drastically reducing uncertainty in regions of low ozone and strong concentration gradients.²⁹ These systems are highly mature and have been deployed globally, from the Sebou estuary trials to the NASA GSFC Southern Hemisphere ADditional OZonesondes (SHADOZ) project, which tracks stratospheric and tropospheric exchanges with relative precisions of (1-2 ppb).³⁰

System Component	Thermodynamic Operating Regime	Primary Flux Mechanism	Associated Analytical Framework
AMTEC (Sodium Heat Engine)	900–1300 K (Isothermal Expansion)	Na ⁺ ion conduction across BASE lattice	Endoreversible Thermodynamics
PRO / RED Membranes	Ambient (Isothermal Mixing)	Osmotic / Salinity concentration gradients	Dissipative NESS Scaling
Brownian Ratchets	Ambient / Biological	Noise-driven spatial asymmetry	Fokker-Planck / Lymn-Taylor
ECC Ozonesondes	Ambient / Stratospheric	KI reagent redox driven by Nernst potential	Compressible Euler Fluid Dynamics

1.1.5 TUPLE_1_PHYSICS:

- : Alkali-Metal Thermal to Electric Converters (AMTEC) utilizing Sodium Beta"-Alumina Solid Electrolytes (BASE); Pressure Retarded Osmosis (PRO) and Reverse Electrodialysis (RED) membranes; Molecular motor proteins operating as Brownian ratchets.
- : Isothermal expansion equivalent to ion conduction across ceramic interfaces; Directed macroscopic transport generated from isotropic thermal noise in asymmetric potentials; Complete localization of entropy production to the interfaces of internally reversible subsystems.
- : Electrochemical oxidation of neutral sodium with decoupled electron/ion transport via external circuits and solid electrolytes; Spatially periodic asymmetric potentials breaking detailed balance via external dissipative driving; Continuous Non-Equilibrium Steady State (NESS) cycling.
- : AMTEC systems are currently at TRL 3–5, having demonstrated 16% efficiency and power densities exceeding 0.50 W/cm² in laboratory environments, with multi-tube modules predicted to reach 25% efficiency.⁵ Osmotic PRO systems have reached TRL 6–7, with world-first grid-connected prototypes demonstrated by Statkraft.⁶ ECC Ozonesondes are fully mature at TRL 9, deployed globally for decades.²⁶

1.2 —

Tuple 2: High Collision Regimes and Geometric Confinement

The second thermodynamic composition necessitates a rigorous analysis of physical systems operating under severe geometric confinement, where the structural boundaries dictate the kinetic, entropic, and thermodynamic behavior of the enclosed particles. As the characteristic

dimensions of a channel or pore approach the atomic or molecular scale, bulk continuum descriptions universally fail. In these regimes, boundary-dominated phenomena such as the Knudsen transition, single-file diffusion, and entropic trapping supersede classical Fickian laws.

1.2.1 Extreme Scaling: Knudsen Regime and Geometric Confinement

In classical fluid mechanics, transport is governed primarily by intermolecular collisions. However, in highly confined nanoporous materials such as zeolite catalysts, single-walled carbon nanotubes (SWNTs), and synthetic supramolecular architectures, the system transitions abruptly into the Knudsen diffusion regime.³² The Knudsen number (Kn) is formally defined as the ratio of the mean free path of the bulk gas (λ) to the mean flight path constrained by the pore geometry (L).³² For an ideal gas, the bulk mean free path is expressed as $\lambda = \frac{k_B T}{\sqrt{2} n \sigma^2 P}$, where k_B is the Boltzmann constant, σ is the collision diameter, and P is pressure.³²

When $Kn \gg 1$, intermolecular collisions become statistically negligible, and gas transport is entirely dominated by high-frequency collisions between the molecules and the pore walls.³² The Knudsen diffusion coefficient is calculated as $D_K = \frac{1}{3} \lambda v$, where v is the mean molecular velocity.³⁴ In typical Fluid Catalytic Cracking (FCC) units, Knudsen diffusion occurs extensively within the meso- and macropores of the catalyst particles, dictating the catalytic conversion rates.³³ Zeolites provide the ultimate realization of this geometric confinement. The shape selectivity of zeolites—such as reactant selectivity, transition state selectivity, and product selectivity—allows for “molecular engineering” in catalyst design.³⁵ For example, in the toluene disproportionation process over ZSM-5 zeolites, the diffusivity of p-xylene is approximately 1000 times faster than its ortho- and meta- isomers. Because the reaction is diffusion-limited, the geometric confinement forces the product selectivity of the highly valuable p-xylene to exceed 97%.³⁵ Mesoscopic kinetic Monte Carlo simulations and pulsed field gradient (PFG) NMR tracking in NaX crystals reveal that the Knudsen tortuosity factor is highly dependent on the non-exponential distribution of flight path lengths traversing between solid collisions.³⁶

1.2.2 Duality Collapse: Confinement and Single-File Accessibility

As geometric confinement reaches the sub-nanometer scale, particles are restricted such that they can no longer physically pass one another, precipitating a thermodynamic phase transition from normal Fickian diffusion to single-file diffusion (SFD).³⁷ In Fickian diffusion, the mean square displacement (MSD) scales linearly with time ($\langle x^2 \rangle \propto t$, where $\langle x^2 \rangle$ is the MSD). In the SFD regime, the collective many-body interactions and absolute interparticle exclusion force the sequence of the particles to be strictly conserved.³⁷ Consequently, the MSD exhibits an anomalous time dependence, scaling as $\langle x^2 \rangle \propto t^{1/2}$.³⁸

This duality of simultaneous high mobility and restricted accessibility has been extensively verified experimentally. Hyperpolarized NMR spectroscopy in self-assembled bis-urea macrocycles demonstrates that phenyl ether bis-urea macrocycles—which form zigzag nanochannels with cross-sectional dimensions of ~ 0.5 nm—enforce persistent single-file behavior over tens of seconds.³⁷ Conversely, expanding the channel diameter to utilizing phenylethynylene bis-urea macrocycles completely restores normal Fickian diffusion, demonstrating absolute synthetic control over the thermodynamic transport state.³⁷

Similar duality collapses are observed in fluidic nanotechnology. Continuous-flow ^1H NMR studies of water confined within 1D single-walled carbon nanotubes (SWNTs) reveal that the

confined water completely resists freezing down to 223 K, maintaining highly mobile, chain-like single-file transport dynamics characterized by a Hz NMR signal.³⁸ Molecular dynamics simulations of KCl solutions in sub-nanometer SWNTs indicate the transition parameter drops from 0.95 in 1.5-nm tubes to 0.8 in 1.1-nm tubes.⁴³ In 2D membranes such as graphene oxide (GO) sheets, Fickian diffusion () dictates parallel interfacial transport, while single-file constraints govern vertical permeation.⁴¹ Interestingly, 2D hexagonal boron nitride (h-BN) and molybdenum disulfide (MoS₂) nanopores can exhibit liquid flow velocities exceeding 1 m/s in channels accommodating only a few layers of water, driven by surface wiggling and increased structural order.⁴⁴

1.2.3 Entatic States in Metalloenzymes

In the realm of bioinorganic chemistry, this duality of confinement and functional accessibility is perfectly encapsulated by the “entatic state” (or rack-induced state) in metalloenzymes.⁴⁵ First proposed by Vallee and Williams, the entatic state occurs when the rigid tertiary structure of a protein scaffold forces a coordinated transition metal ion and its first coordination sphere into an energized, highly strained geometric and electronic configuration.⁴⁵ The biological matrix acts as a sub-nanometer clamp, placing structural restrictions on the metal that approximate the geometry of the catalytic transition state.⁴⁶

For instance, in the blue-copper protein azurin, the Cu²⁺ ion is held in a distorted configuration by three strong ligands (one thiolate sulfur and two imidazole nitrogens), establishing a geometry intermediate between the preferred states of Cu²⁺ and Cu^I [*image38* *lambda*) required for electron transfer is drastically minimized.⁴⁵ According to Marcus theory, lowering this reorganization energy results in an exponential increase in the electron transfer rate.⁴⁵ Recent quantifications in Cu(I)-based photosensitizers reveal entatic states reaching kcal/mol of strain energy relative to conformationally flexible benchmarks, allowing engineered synthetic models (like guanidine quinolinyl systems) to achieve phenomenal electron transfer rates on the order of .⁴⁵ Steady-state spectroscopic studies of mutant cytochrome structures confirm that relieving this steric confinement immediately removes the entatic imposition, reverting the kinetics to slow, bulk-like behavior.⁴⁸

1.2.4 H_Mechanisms: Geometric Confinement without Energy (Entropic Trapping)

When a flexible, highly extended macromolecule—such as double-stranded DNA or large polyelectrolytes—is forced into a pore structure smaller than its bulk radius of gyration (), it undergoes purely geometric entropic trapping without the need for external energetic input.⁴⁹ Transporting the molecule through a nanoconstriction requires the polymer chain to stretch and elongate, which imposes a severe thermodynamic penalty due to the catastrophic loss of conformational degrees of freedom (rotational and translational entropy).⁴⁹

The molecule instead becomes temporarily trapped at the interface of nanopores or within larger interstitial voids in the porous matrix (such as within agarose gels or artificial hydrogels), where it can expand and maximize its conformational entropy.⁴⁹ This creates a unique transport mechanism intermediate between Ogston sieving and reptation.⁴⁹ Experimental implementations utilizing ultrathin nanoporous silicon nitride (NPN) membranes as capping nanofilters over hexagonal microwell grids demonstrate an exceptionally high entropic barrier to escape.⁵¹ In the entropic trapping regime, DNA electrophoretic mobility scales anomalously as , where the exponent ranges from 1.5 to 2.5 depending on the DNA size, fundamentally deviating from hydrodynamic chromatography principles.⁴⁹ Advances in two-photon fabrication have yielded nanofluidic traps containing 50 nm slit channels

intersected by nm transverse grooves, allowing DNA to uncoil and map physically without surface immobilization, increasing residence times by 3- to 5-fold purely via local steric confinement.⁵²

Transport Regime	Dimensional Scaling	Primary Governing Interaction	Observable Metric
Fickian Diffusion	Bulk/Macroporous	Intermolecular collisions	
Knudsen Diffusion		Molecule-wall collisions	
Single-File Diffusion	Sub-nanometer	Interparticle strict exclusion	
Entropic Trapping	Pore Size	Conformational entropy loss	

1.2.5 Procedural Mapping: Nanoporous Membrane Catalysis

The macroscopic manifestation of these nanoscale confinement effects is realized in industrial nanoporous membrane catalysis and gas separation. As global demands for greenhouse gas mitigation accelerate, Mixed Matrix Membranes (MMMs) leveraging precisely engineered filler materials offer superior cost-to-efficiency ratios compared to traditional chemical absorption.⁵⁵

Techno-economic evaluations categorize carbon dioxide separation technologies into distinct tiers. Physical absorption currently stands as a Tier 1 performer (93% efficiency, 1.03 GJ/tCO₂).⁵⁷ However, advanced membrane filtration technologies (Tier 2) are rapidly closing the gap, achieving up to 88% and capture costs of \$2.57. The HyCaps membrane module, currently at Technology Readiness Level (TRL) 6, has successfully undergone pilot testing in the UK and is scaling to a commercial 10,000 tonnes-per-annum (tpa) plant at Holcim’s Höver cement facility, with a path to a 800,000 tpa industrial scaleout by 2028.⁵⁸

1.2.6 TUPLE 2 PHYSICS:

- : Hyperpolarized in bis-urea macrocycles; Sub-nanometer single-walled carbon nanotubes (SWNTs); Nanoporous silicon nitride (NPN) membranes for DNA sieving; Blue-copper metalloenzymes (azurin) and de novo artificial metalloproteins (ArM MPP1).
- : Transition from Fickian () to single-file () transport dynamics; Anomalous electrophoretic mobility scaling due to conformational entropy gradients; Ligand-field-destabilized high reduction potentials and energized structural geometries reaching 20 kcal/mol of strain energy relative to relaxed states.
- : Interparticle exclusion interactions in 1D sub-nanometer lattices preventing sequence exchange; Energetic barriers induced by the requirement to compress the polymer radius of gyration during pore translocation; Protein-matrix “rack” forcing intermediate coordination geometries to exponentially lower electron-transfer reorganization barriers.
- : Zeolite catalysis and standard Knudsen-regime filtration membranes are highly mature (TRL 8–9) and operate at massive industrial scales.³³ Advanced CO₂ capture

membranes (HyCaps) are at TRL 6.58 Engineered synthetic entatic state models and nanofluidic entropic trapping arrays remain largely in the laboratory validation phase (TRL 3-4).⁴⁵

1.3 —

Tuple 3: Structural Relaxation and Topological Protection

The third thermodynamic composition targets the absolute suppression of charge recombination through structural relaxation, non-equilibrium symmetry breaking, and topological protection. In conventional semiconductor optoelectronics and photovoltaics, electron-hole pairs generated by incident photoexcitation are highly susceptible to rapid radiative or non-radiative recombination, fundamentally limiting energy conversion efficiency to the Shockley-Queisser limit.⁶⁰ Bypassing this limitation requires decoupling the spatial, energetic, or topological overlap of the charge carriers.

1.3.1 Extreme Scaling: Ferroelectric Photovoltaics and Structural Relaxation

Ferroelectric photovoltaics (FPVs) operate on physical principles entirely distinct from the conventional p-n junctions that dictate classical solar cells.⁶¹ In FPVs, the spontaneous inversion symmetry breaking intrinsic to the non-centrosymmetric ferroelectric lattice generates a permanent, bulk depolarization field.⁶² This macroscopic spontaneous polarization drives a continuous, ballistic separation of photoexcited charge carriers, resulting in anomalous above-bandgap photovoltages completely untethered from the material's fundamental bandgap limits.⁶²

To further suppress recombination, advanced FPV materials—such as Bismuth Ferrite (or BFO) and Metal Halide Perovskites (MHPs)—utilize targeted structural relaxation and strain engineering.⁶² In highly crystalline, pristine BFO, the bandgap is nearly direct, which mathematically necessitates a high rate of radiative recombination. However, polar order engineering via A-site substitution (e.g., replacing with Lanthanum) deliberately destabilizes the local ferroelectric order, inducing a direct-to-indirect bandgap transition.⁶²

At the quantum electronic level, the valence band maximum (VBM) in BFO is comprised of strongly hybridized Bi, Fe, and O antibonding states.⁶² The substitution of La removes the highly stereochemically active Bi lone-pair electrons, lowering the energy of these states and introducing prominent indirect band features.⁶² Consequently, photoexcited electrons rapidly thermalize to the conduction band minimum, resulting in a profound momentum-space mismatch with the localized holes.⁶² This mechanism, coupled with local strain and off-center displacement variations (smaller displacement for La dopants), substantially suppresses radiative recombination.⁶² In MHPs like $\text{CH}_3\text{NH}_3\text{PbBr}_3$, the bandgap reduction is driven by the reorientation of polar organic cations that distort the inorganic framework, weakening the overlap between I and Pb orbitals and extending charge carrier lifetimes to several nanoseconds.⁶⁶

Furthermore, structural phase transitions and the existence of ferroelectric domain walls act as highly efficient, distinct transport channels. At these domain boundaries, abrupt changes in polarization orientation establish massive local electrostatic potential gradients. Theoretical and experimental analyses demonstrate that this localized charge separation mechanism plummets the overall carrier recombination rate by an astonishing 3 to 4 orders of magnitude relative to single-domain bulk perovskites.⁶¹ Enrichment of specific defects or

impurities at the domain wall further alters the electronic structure, establishing a Schottky junction effect that definitively isolates the carriers.⁶⁸

1.3.2 Duality Collapse: Hysteretic Memory and Topological Protection

The coupling between structural configurations and charge separation inherently yields highly non-linear, non-ergodic macroscopic behaviors, classified mathematically as hysteresis. Hysteretic materials, ranging from magnetic alloys to complex relaxor ferroelectrics like Potassium-Tantalate-Niobate (KTN) crystals, exhibit a profound history-dependence.⁶⁹ The macroscopic response of these materials to an external field is dictated by the slow structural relaxation of internal domains transitioning through a rugged, high-dimensional free-energy landscape.⁷⁰

In KTN crystals, intricate metastable domain patterns emerge near the cubic-to-tetragonal structural phase transition.⁷⁰ The macroscopic layout of these ferroelectric superlattices is governed by the absolute thermodynamic necessity to minimize volume polarization charge, dictating that the divergence of the polarization field must locally vanish ($\nabla \cdot \mathbf{P} = 0$).²⁹ This stringent charge-screening requirement enforces continuous “flux-closure” conditions, creating vortex-like domain arrangements.²⁹

Because these vortices require the simultaneous, coordinated realignment of all surrounding domains to be unwound, the individual domains acquire “topological protection”.²⁹ They are effectively locked into localized potential minima and shielded from isolated thermal fluctuations.²⁹ The Topological Breakdown Model (TBM) demonstrates that frustration in these materials arises precisely from the geometric conflict between standard collinear dipole-dipole interactions at domain walls and non-collinear polarization-charge screening interactions at walls.²⁹ When the applied electric field or temperature overcomes this protective threshold, the system undergoes abrupt percolative transitions, releasing the pinned domains in discrete Barkhausen jumps and yielding characteristic slim-loop hysteresis cycles.²⁹

This mechanism of topological protection against recombination and signal decay is currently being mapped onto advanced data storage and memory technologies. In spintronics, magnetic skyrmions—localized, stable topological magnetic textures—move and interact like ordinary particles under spin-polarized currents.⁷³ Direct observations in Co/Ni multilayers reveal dipolar skyrmions possessing arbitrary topological charges up to 10.⁷⁴ Their topological protection prevents the spin texture from continuously transforming into a saturated ferromagnetic state, unless annihilated through a structural Bloch point due to extreme thermal fluctuations.⁷³ Similarly, in photonics, dynamically tunable Chern numbers encode multi-bit memory states protected from environmental and fabrication defects, bypassing the need for dissipative light-trapping elements.⁷⁵

Protection Mechanism	Carrier / Quasiparticle	Functional Device Application	Recombination Reduction
Indirect Bandgap Mismatch	Electron-Hole Pairs	BFO / MHP Photovoltaics	Thermalization mismatch
Domain Wall Segregation	Polarized Free Charges	Ferroelectric Solar Cells	3 to 4 orders of magnitude
Flux-Closure Vortices	Ferroelectric Dipoles	KTN Non-Volatile Memory	Pinned by

Protection Mechanism	Carrier / Quasiparticle	Functional Device Application	Recombination Reduction
Topological Winding	Magnetic Skyrmions	Spintronic Racetrack Memory	Protected by integer

1.3.3 Procedural Mapping: Ferroelectric and Electret Charge Separation

Macroscopic implementations of these structurally relaxed, charge-separated states are realized in bulk electrets and advanced thin films. Electret polymers, such as uniaxially stretched polyvinylidene fluoride (PVDF), maintain long-term macroscopic charge separation by trapping injected charges in deep structural potential wells.⁷⁶ When poled under intense external electric fields (e.g., 300 kV/cm), the resulting sensors display massive piezoelectric coefficients of approximately 200 pC/N, far exceeding standard PVDF-TR films.⁷⁶ These electroresponsive polymers operate effectively at high Technology Readiness Levels (TRL 8-9) in robust commercial applications, including Emfit medical respiration and epileptic seizure monitors.⁷⁶ In parallel, Metal Halide Perovskite (MHP) solar cells—capitalizing on the precise strain engineering and domain manipulation outlined above—have skyrocketed in viability, with single-junction architectures achieving certified power conversion efficiencies of 25.5%, and silicon-MHP tandem cells pushing beyond 29.5%.⁶⁵

1.3.4 TUPLE_3_PHYSICS:

- : Bismuth Ferrite () bulk samples and thin films; Metal Halide Perovskites (MHPs) such as ; Potassium-Tantalate-Niobate (KTN) relaxor ferroelectric crystals; Electret polymer films (PVDF); Co/Ni multilayer magnetic skyrmions.
- : Direct-to-indirect bandgap transitions resulting from -site substitution and structural lattice strain; 3 to 4 orders of magnitude reduction in electron-hole recombination rates at ferroelectric domain walls; The emergence of Barkhausen jumps and non-ergodic slim-loop hysteresis corresponding to the breakdown of flux-closure topologies.
- : Spontaneous inversion symmetry breaking driving a bulk depolarization field for continuous charge separation; Removal of stereochemical lone-pair electrons altering valence band hybridization and carrier momentum matching; Geometric frustration between collinear dipole-dipole interactions and the strict non-collinear charge-screening conditions.
- : Fundamental ferroelectric photovoltaics utilizing complex oxides like BFO remain in the early research phase (TRL 1–3) due to low overall efficiencies.⁶² Metal Halide Perovskite solar cells are rapidly maturing through industrial prototyping (TRL 4–6).⁶⁵ Electret-based charge separation devices (PVDF) for piezoelectric sensing are fully commercialized and globally deployed (TRL 8–9).⁷⁶

1.4 —

FEASIBILITY_ASSESSMENT:

The theoretical viability of synthesizing these exact abstract thermodynamic compositions into physical engineering space demands rigorous scrutiny, specifically regarding limits

imposed by the Second Law of Thermodynamics and the constraints of physical continuum scaling.

Tuple 1 Feasibility (Divergent Flux at): The mathematical stipulation of a diverging heat flux density strictly as presents an apparent physical singularity. Under classical Fourier conduction, . Maintaining a finite or diverging with a vanishing thermal gradient requires the thermal conductivity to mathematically approach infinity, which is an inherently unphysical proposition for any macroscopic solid.¹ However, the composition is completely physically sound and legally evades Second Law violations when transitioning the fundamental driving force from a pure thermal gradient to an electrochemical gradient, as explicitly observed in AMTEC systems, or an osmotic concentration gradient, as seen in RED/PRO engines.⁵ The divergence theorem () merely dictates the continuity relationship between the boundary flux and the internal volume source.² In AMTEC, the isothermal expansion of the sodium vapor is powered entirely by the localized electrochemical differential across the BASE electrolyte, perfectly sidestepping the Fourier limitation.⁵ Similarly, Brownian ratchets operate legally within the Second Law solely because they are coupled to an external driving source (e.g., flashing potentials or chemical hydrolysis) that fundamentally breaks detailed balance; without this continuous external energy input, extracting directed work from isothermal thermal fluctuations would constitute an impossible, entropy-destroying Maxwell’s Demon.¹⁴ Therefore, Tuple 1 does not violate the Second Law, provided the system is chemically or electrically driven into a Non-Equilibrium Steady State (NESS).

Tuple 2 Feasibility (High-Collision Confinement): The physics of Knudsen diffusion and single-file diffusion are mathematically coherent and experimentally verified up to the limits of physical channel persistence.³² However, the theoretical scaling of the mean square displacement () in perfect SFD mathematically assumes an infinitely long, perfectly defect-free uniform channel. In any physical realization, finite channel lengths, thermal wall fluctuations, and lattice defects allow for minute molecular bypass probabilities, which eventually forces the system to revert to normal Fickian diffusion at extremely long macroscopic timescales.³⁷ Furthermore, the “entatic state” rack mechanism in metalloenzymes—while remarkably effective at lowering reorganization energy—requires a massive expenditure of evolutionary folding energy to permanently hold the protein matrix in a strained geometry (up to 70 kJ/mol).⁴⁵ The thermodynamic penalty is paid by the global stability of the tertiary protein structure, meaning such active sites possess highly restricted operational windows and are highly vulnerable to denaturation if removed from strictly controlled physiological temperatures and pH levels.

Tuple 3 Feasibility (Recombination Suppression via Structural Relaxation): Ferro-electric photovoltaics theoretically bypass the classical Shockley-Queisser limit by relying on spontaneous bulk polarization rather than bandgap-limited p-n junctions.⁶⁰ The proposition of dropping recombination rates by 3-4 orders of magnitude via domain wall engineering is computationally sound and experimentally validated.⁶⁷ However, the direct-to-indirect bandgap transition achieved via -site substitution (e.g., La into BFO) highlights a severe, fundamental engineering trade-off. While the momentum mismatch of an indirect bandgap radically extends the charge carrier lifetime by suppressing radiative recombination, the laws of quantum mechanics dictate that it simultaneously reduces the material’s initial photon absorption coefficient.⁶² The system is forced to trade photon generation efficiency for carrier longevity. Additionally, the topological protection observed in systems like KTN crystals or magnetic skyrmions is completely dependent on maintaining the delicate balance of continuous field configurations (e.g.,). This protection is not infinite; high external electric fields or elevated temperatures easily strip the system of this nonergodic stability, collapsing the topological protection through a Bloch point annihilation and irrevocably resetting

the memory states.²⁹ The tuple requires no unphysical conditions such as negative heat capacities or infinite Q-factors, provided the boundaries of thermal stability are explicitly parameterized.

1.4.0.1 Works cited

1. Lesson 10. Applications of Divergence Theorem Consider incompressible fluid of constant density ρ in a region T with the boundary - Purdue Math, accessed April 4, 2026, <https://www.math.purdue.edu/~gabriea/MA528/lecture10.pdf>
2. What does divergence of heat flux mean? - Physics Stack Exchange, accessed April 4, 2026, <https://physics.stackexchange.com/questions/697548/what-does-divergence-of-heat-flux-mean>
3. General equation of heat transfer - Wikipedia, accessed April 4, 2026, https://en.wikipedia.org/wiki/General_equation_of_heat_transfer
4. Steady-State Thermodynamics Thermal Dissipation - Quondam's Notebook, accessed April 4, 2026, https://www.pdquondam.net/Thermal_Dissipation_III.pdf
5. Report on nuclear batteries | DOCX - Slideshare, accessed April 4, 2026, <https://www.slideshare.net/slideshow/report-on-nuclear-batteries/33799341>
6. Deployable Membrane-Based Energy Technologies: the Ethiopian Prospect - MDPI, accessed April 4, 2026, <https://www.mdpi.com/2071-1050/12/21/8792>
7. Pathways to a Net-Zero- Carbon Water Sector Through Energy-Extracting Wastewater Technologies - INL Research Library Digital Repository - Idaho National Laboratory, accessed April 4, 2026, https://inldigitallibrary.inl.gov/sites/sti/sti/Sort_63191.pdf
8. Pressure Retarded Osmosis (PRO): Past experiences, current developments, and future prospects | Request PDF - ResearchGate, accessed April 4, 2026, https://www.researchgate.net/publication/290789305_Pressure_Retarded_Osmosis_PRO_Past_experiences_current_developments_and_future_prospects
9. Publications | Research | Computational Physics - TU Chemnitz, accessed April 4, 2026, <https://www.tu-chemnitz.de/physik/CPHYS/Publications/>
10. Dynamics and Statistical Physics Division Fachverband Dynamik und Statistische Physik (DY) - DPG-Verhandlungen, accessed April 4, 2026, <https://www.dpg-verhandlungen.de/2014/dresden/dy.pdf>
11. Endoreversible Thermodynamics of a Hydraulic Recuperation System - Monarch: Qucosa, accessed April 4, 2026, <https://monarch.qucosa.de/api/qucosa%3A34015/attachment/ATT-0/>
12. Thermodynamics 2.0, accessed April 4, 2026, https://iaisae.org/wp-content/uploads/BoA-Final_v2.pdf
13. Brownian motors: noisy transport far from equilibrium - University of Utah Math Dept.,

accessed April 4, 2026, <http://www.math.utah.edu/~bresslof/ratchets>

14. Introduction to the physics of Brownian motors - ResearchGate, accessed April 4, 2026, https://www.researchgate.net/publication/225392185_Introduction_to_the_physics_of_Brownian_motors
15. Stochastic Thermodynamics: Experiment and Theory - Max Planck Institute for the Physics of Complex Systems, accessed April 4, 2026, <https://www.pks.mpg.de/stochastic-thermodynamics-experiment-and-theory/poster-contributions>
16. Collective transient ratchet transport induced by many elastically interacting particles - PMC, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC8355274/>
17. Cellular motions and thermal fluctuations: The Brownian Ratchet - ResearchGate, accessed April 4, 2026, https://www.researchgate.net/publication/14830744_Cellular_motions_and_thermal_fluctuations_The_Brownian_Ratchet
18. Allocating and Splitting Free Energy to Maximize Molecular Machine Flux | The Journal of Physical Chemistry B - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.jpccb.7b10621>
19. Analysis of kinetic asymmetry in a multi-cycle chemical reaction network establishes the principles for autonomous compartmental - ChemRxiv, accessed April 4, 2026, <https://chemrxiv.org/doi/pdf/10.26434/chemrxiv-2024-pzhcs>
20. Chemical Kinetics and Mass Action in Coexisting Phases - Max Planck Institute for the Physics of Complex Systems, accessed April 4, 2026, https://www.pks.mpg.de/fileadmin/user_upload/MPIPKS/group_pages/BiologicalPhysics/juelicher/Highlights_2022/CKaMAiCP2022.pdf
21. A Prehistorical Approach to Optimal Fluctuations for General Langevin Dynamics with Weak Gaussian White Noises - MDPI, accessed April 4, 2026, <https://www.mdpi.com/2076-3417/12/17/8569>
22. 11-Applications of the Divergence Theorem: The Compressible Euler Equations - UC Davis Mathematics, accessed April 4, 2026, <https://www.math.ucdavis.edu/~temple/MAT22C/LecturesMat22CW15/11-ApplsDivThm-22C-W15.pdf>
23. Untitled - Yale University, accessed April 4, 2026, http://www.astro.yale.edu/vdbosch/astro595_notes.pdf
24. Formation And Transport Of Oxidants Along Gulf Coast And In Northern Us - epa nepis, accessed April 4, 2026, <https://nepis.epa.gov/Exe/ZyPURL.cgi?Dockkey=2000XRK3.TXT>
25. Summary Report 1990 - NOAA Global Monitoring Laboratory, accessed April 4, 2026, https://gml.noaa.gov/publications/summary_reports/summary_report_19.pdf
26. STCE Annual Report 2021, accessed April 4, 2026, https://www.stce.be/movies/STCE%20AR/STCE%20Annual%20Report%202021_FINAL.pdf

27. *****A***t*****2 - ERIC, accessed April 4, 2026, <https://files.eric.ed.gov/fulltext/ED324245.pdf>
28. Lidar profiling of ozone above the central San Joaquin Valley during - the California Base-line Ozone Transport Study (CABOTS) - NOAA Chemical Sciences Laboratory, accessed April 4, 2026, <https://csl.noaa.gov/projects/cabots/CABOTSfinalreport15R012.pdf>
29. Stratospheric ozone loss in the Arctic winters between 2005 and 2013 derived with ACE-FTS measurements - Copernicus ACP, accessed April 4, 2026, <https://acp.copernicus.org/articles/19/577/2019/acp-19-577-2019-relations.html>
30. Feasibility of aerial measurements of methane emissions from landfills - GOV.UK, accessed April 4, 2026, https://assets.publishing.service.gov.uk/media/5a7f0195ed915d74e6227db6/SC130034_Report.pdf
31. NASA Scientific and Technical Aerospace Reports - AMiner, accessed April 4, 2026, https://static.aminer.org/pdf/PDF/000/589/450/microsourcing_using_information_technology_to_create_unexpected_work_relationships_and.pdf
32. Solving the inverse Knudsen problem: Gas diffusion in random fibrous media - TU Delft Repository, accessed April 4, 2026, https://repository.tudelft.nl/file/File_275e5245-9be0-4ae9-b7df-79cb4b269b2c?preview=1
33. Diffusion in Fluid Catalytic Cracking Catalysts on Various Displacement Scales and Its Role in Catalytic Performance | Chemistry of Materials - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/cm050031z>
34. Investigation of Molecular Mean Free Path, Molecular Kinetic Energy, and Molecular Polarity Affecting Knudsen Diffusivity along Pore Channels - MDPI, accessed April 4, 2026, <https://www.mdpi.com/2297-8739/9/5/130>
35. THE DIFFUSION MECHANISM OF HYDROCARBONS IN ZEOLITES Jirong Xiao B.S., East China Institute of Chemical Technology Submitted to t - DSpace@MIT, accessed April 4, 2026, <https://dspace.mit.edu/bitstream/handle/1721.1/33810/22929092-MIT.pdf?sequence=2&isAllowed=y>
36. Mesoscopic simulations of the diffusivity of ethane in beds of NaX zeolite crystals - AIP Publishing, accessed April 4, 2026, https://pubs.aip.org/aip/jcp/article-pdf/doi/10.1063/1.2567129/13381173/094702_1_online.pdf
37. Crystalline Bis-urea Nanochannel Architectures Tailored for Single-File Diffusion Studies | ACS Nano - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acsnano.5b01895>
38. Single-File Diffusion of Confined Water Inside SWNTs: An NMR Study | ACS Nano, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/nn901554h>
39. Diffusion in Confined Geometries, accessed April 4, 2026, <https://d-nb.info/1255616113/34>

40. The Physics Institutes - Report 2005 - Fakultät für Physik und Erdsystemwissenschaften - Universität Leipzig, accessed April 4, 2026, https://www.physes.uni-leipzig.de/fileadmin/Fakult%C3%A4t_PhysGeo/Dokumente/Forschung/2005_Forschungsbericht_Physik.pdf
41. Molecular Insight into the Adsorption Thermodynamics and Interfacial Behavior of GOs at the Liquid-Liquid Interface - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.jpcb.0c10118>
42. Exploring Single-File Diffusion in One-Dimensional Nanochannels by Laser-Polarized ¹²⁹Xe NMR Spectroscopy | The Journal of Physical Chemistry A - ACS Publications - American Chemical Society, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/jp002322v>
43. Fluids and Electrolytes under Confinement in Single-Digit Nanopores - PMC - NIH, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC10037271/>
44. Transport Phenomena in Nano/Molecular Confinements - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acsnano.0c07372>
45. Quantifying Entatic States in Photophysical Processes: Applications to Copper Photosensitizers | Inorganic Chemistry - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.inorgchem.9b02976>
46. Direct Evidence for a Geometrically Constrained “Entatic State” Effect on Copper(II/I) Electron-Transfer Kinetics As Manifested in Metastable Intermediates | Journal of the American Chemical Society, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/ja002995r>
47. Artificial Metalloproteins: At the Interface between Biology and Chemistry | JACS Au, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/jacsau.2c00102>
48. Disabling the Entatic Control of Methionine Ligation through Additive Destabilization of Ferric Cytochrome c - PMC, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12188557/>
49. Molecular sieving using nanofilters: Past, present and future - Mechanical Engineering, accessed April 4, 2026, https://me-web.engin.umich.edu/ibbl/pdf/2007_LabonChip_Han.pdf
50. macromolecules: Topics by Science.gov, accessed April 4, 2026, <https://www.science.gov/topicpages/m/macromolecules>
51. Entropic Trapping of DNA with a Nanofiltered Nanopore | ACS Applied Nano Materials, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acsanm.9b00606>
52. Nanofluidic Traps by Two-Photon Fabrication for Extended Detection of Single Macromolecules and Colloids in Solution - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acsanm.1c03691>
53. 2009 APS March Meeting, accessed April 4, 2026, <https://higherlogicdownload.s3.ama>

zonaws.com/APS/22d184d0-d6d2-4493-97da-847d341d3d82/UploadedImages/Documents/Unit_MAR09_DFD.pdf

54. Nanofluidic Traps by Two-Photon Fabrication for Extended Detection of Single Macromolecules and Colloids in Solution | Request PDF - ResearchGate, accessed April 4, 2026, https://www.researchgate.net/publication/358869209_Nanofluidic_Traps_by_Two-Photon_Fabrication_for_Extended_Detection_of_Single_Macromolecules_and_Colloids_in_Solution
55. Review on Synthesis and Characterization of Advanced Nanomaterials-based Mixed Matrix Membranes (MMMs) for CO₂ Capture: Progress, Challenges, and Prospects | Energy & Fuels - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.energyfuels.4c03305>
56. Membrane-Based CO₂ Capture Across Industrial Sectors: Process Conditions, Case Studies, and Implementation Insights - PMC, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC12298987/>
57. Integrated CO₂ capture and conversion: dual-functional materials, mechanisms, and pathways to industrial decarbonization - EES Catalysis (RSC Publishing) DOI:10.1039/D5EY00322A, accessed April 4, 2026, <https://pubs.rsc.org/en/content/articlehtml/2026/ey/d5ey00322a>
58. STATE OF THE ART: CCS TECHNOLOGIES 2025 - Global CCS Institute, accessed April 4, 2026, <https://www.globalccsinstitute.com/wp-content/uploads/2025/08/State-of-the-Art-CCS-Technologies-2025-Global-CCS-Institute.pdf>
59. Biomolecule Separation by Steric Hindrance using Nanofluidic Filters - Mechanical Engineering, accessed April 4, 2026, https://me-web.engin.umich.edu/ibbl/pdf/Proceeding_2004_IEEE_EMBS_Han.pdf
60. Unleashing the Potential of Strain-Engineered Ferroelectric Oxynitrides for Optimal Photovoltaic Performance | The Journal of Physical Chemistry C - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.jpcc.3c03824>
61. Efficient charge separation at localized 2D ferroelectric domains in perovskite solar cells, accessed April 4, 2026, <https://pubs.rsc.org/en/content/articlehtml/2025/ee/d5ee00640f>
62. Enhancing ferroelectric photovoltaic effect by polar order engineering - PMC - NIH, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC6035034/>
63. Unveiling charge utilization mechanisms in ferroelectric for water splitting - PMC - NIH, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11810992/>
64. 2. Supplementary Information: - OSTI, accessed April 4, 2026, <https://www.osti.gov/servlets/purl/1991698>
65. Ferroic Halide Perovskite Optoelectronics - OSTI, accessed April 4, 2026, <https://www.osti.gov/pages/servlets/purl/1819578>

66. Ferroelectric Polarization Suppresses Nonradiative Electron-Hole Recombination in CH₃NH₃PbI₃ Perovskites: A Time-Domain ab Initio Study | The Journal of Physical Chemistry Letters - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.jpclett.9b02931>
67. Enhanced Photovoltaic Properties Induced by Ferroelectric Domain Structures in Organometallic Halide Perovskites | The Journal of Physical Chemistry C - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.jpcc.7b03091>
68. Ferroelectric Photovoltaic Pyroelectric Coupling Effect: Mechanism and Applications, accessed April 4, 2026, https://www.researchgate.net/publication/393859671_Ferroelectric_Photovoltaic_Pyroelectric_Coupling_Effect_Mechanism_and_Applications
69. Time and temperature dependent magnetic viscosity experiments on Sr/Co nanoferrite particles, accessed April 4, 2026, <https://uu.diva-portal.org/smash/get/diva2:1776852/FULLTEXT01.pdf>
70. Topological protection breakdown: A route to frustrated ferroelectricity - DIGIBUG Principal, accessed April 4, 2026, <https://digibug.ugr.es/bitstream/handle/10481/107233/h6j7-cgwz.pdf?sequence=1&isAllowed=y>
71. New modalities of strain-control of ferroelectric thin films | Request PDF - ResearchGate, accessed April 4, 2026, https://www.researchgate.net/publication/303326920_New_modalities_of_strain-control_of_ferroelectric_thin_films
72. Topological Protection Can Arise from Thermal Fluctuations and Interactions - DSpace@MIT, accessed April 4, 2026, <https://dspace.mit.edu/bitstream/handle/1721.1/121190/PhysRevLett.122.118001.pdf?sequence=1&isAllowed=y>
73. Skyrmionics—Computing and memory technologies based on topological excitations in magnets | Journal of Applied Physics | AIP Publishing, accessed April 4, 2026, <https://pubs.aip.org/aip/jap/article/130/7/070908/158211/Skyrmionics-Computing-and-memory-technologies>
74. Dipolar skyrmions and antiskyrmions of arbitrary topological charge at room temperature - PMC, accessed April 4, 2026, <https://pmc.ncbi.nlm.nih.gov/articles/PMC11021192/>
75. Nontrapping Tunable Topological Photonic Memory - arXiv.org, accessed April 4, 2026, <https://arxiv.org/html/2502.19398v1>
76. DETECTION OF HUMAN VITAL SIGNS BY SEAT-EMBEDDED FERROELECTRIC FILM SENSORS AND - Research - Department of Transportation, accessed April 4, 2026, <https://www-nrd.nhtsa.dot.gov/departments/esv/22nd/files/Session%203%20Written.pdf>
77. Electromagnetic Machines Which Utilize Microgeometry Field Structures - DTIC, accessed April 4, 2026, <https://apps.dtic.mil/sti/tr/pdf/ADA219828.pdf>
78. Ultra Wide-Bandwidth Micro Energy Harvester - DSpace@MIT, accessed April 4, 2026, <https://dspace.mit.edu/bitstream/handle/1721.1/63072/725893053-MIT.pdf?sequence=2&isAllowed=y>

79. Photoelectrocatalytic CO₂ Reduction Using g-C₃N₄ Hybrids: Review and Perspectives | Energy & Fuels - ACS Publications, accessed April 4, 2026, <https://pubs.acs.org/doi/10.1021/acs.energyfuels.5c01356>