1 Geometric Classification of Standard Model Particles Derived from the Orientation Structure of the 6-Dimensional Hypercube

1.1 — Discrete Geometric Formulation of Color Charge, Mass, and Coupling Strength —

Author: Noriaki Kihara (WF System Co., Ltd.)
Date: April 2026
DOI: 10.5281/zenodo.19646652

1.2 0. Purpose

In the author's preceding work [7], six spin quantum numbers $s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}$ were derived as bivectors from the 3-cell orientation structure of the 5-dimensional hypercube. However, the 5-dimensional framework had the following limitations:

1. **Absence of color charge**: No degree of freedom to distinguish the three colors of quarks.
2. **No quark–lepton distinction**: The same orientation corresponds to both quarks and leptons.
3. **Insufficient gauge bosons**: Only 4 spin-1 boson orientations, falling short of the 12 Standard Model states (γ, g_1–$g_8, W^{\pm}, Z^0$).
4. **Graviton placement**: The spin-2 boson requires $n = 2$, but with only two non-spatial axes, only one combination was available.

This paper resolves all of these problems by adding a sixth axis to the 5-dimensional framework. The additional axis is a **discrete label axis** qualitatively distinct from space, time, and curvature, which geometrically introduces internal degrees of freedom corresponding to color charge.

All results in this paper are stated as propositions in discrete geometry and number theory.

1.3 1. Basic Structure of the 6-Dimensional Framework

1.3.1 1.1 Definition

The 6-dimensional structure treated in this paper consists of the 5-dimensional hypercube (vertices $(\pm1, \pm1, \pm1, \pm1, \pm1)$) augmented by a 4-valued discrete label axis Q.

The five geometric axes $(x_1, x_2, x_3, x_4, x_5)$ take coordinate values ±1 and form a hypercube structure. The sixth axis Q is not a geometric coordinate but a discrete label value $Q \in \{0, r, g, b\}$ attached to each state.

1.3.2 1.2 Element Counts of the 5-Dimensional Hypercube (Restated)

k	Element	Count
0	Vertices	32
1	Edges	80
2	Square faces	80
3	Cubic cells (3-cells)	40

k	Element	Count
4	Hypercubic facets (4-facets)	10

This is the same geometric foundation as in the preceding work [7]. The Q axis does not alter the geometric structure of the hypercube but provides an additional quantum number to each state.

1.3.3 1.3 Six-Axis Notation

In this paper, each state is denoted by a six-component symbol (x, y, z, t, R, Q). The first five components represent the position in the hypercube (fixed axes as ± 1, free axes as 0), and the sixth component represents the color label.

1.4 2. Axis Classification: Four Qualitatively Distinct Degrees of Freedom

1.4.1 2.1 Extended Assumption

Assumption A (physical classification of five axes) from the preceding works [6, 7] is extended to six axes.

Assumption B (Physical classification of six axes). The six axes are classified as follows:

- **Spatial axes** (x_1, x_2, x_3): Three axes corresponding to 3-dimensional space directly perceivable by the observer. Coordinate values may take ± 1, but these axes are not subject to sign reversal (parity operation) via bivectors. The three axes are mutually symmetric under the permutation group S_3.
- **Temporal axis** $(x_4 = t)$: The axis corresponding to time. Sign reversal ± 1 is possible. The range is finite (2 values).
- **Scale axis** $(x_5 = R)$: The axis corresponding to the curvature radius of space. The range is unbounded, making it the only axis among the six with an unlimited degree of freedom.

Remark. In the preceding work [W5], the number-theoretic conditions of central projection in discrete space lead to the discretization $R = \pm n^2 \cdot L_0$ ($n \in \mathbb{N}$, L_0 is a unit length). However, all results in this paper hold without this discretization and remain valid when R takes continuous values.

- **Color axis** $(x_6 = Q)$: The axis corresponding to internal degrees of freedom. Takes discrete label values $Q \in \{0, r, g, b\}$ (4 values).

1.4.2 2.2 Four Qualitative Differences

The six axes are classified into four qualitatively distinct types.

Type	Axis	Values	Meaning	Bound
Position	x, y, z	Continuous	"Where is it?"	S_3 constrained
Direction	t	± 1	"Which way?"	2 values
Scale	R	Unbounded	"How large?"	**Unbounded**
Label	Q	$0, r, g, b$	"What kind?"	4 values

Proposition 2.1. Among the six axes, only the R axis has an unbounded range. The **type** of a particle (spin, color charge, generation) is determined by the finite discrete structure

of (x, y, z, t, Q), while the **mass** of a particle is determined by the sole unbounded degree of freedom (R).

1.4.3 2.3 Consequences of Q Being a Label

The Q axis represents a label (which kind), not a quantity (how much). The following consequences follow immediately from this property:

1. **Charge quantization**: Since Q takes only discrete values, charge is intrinsically quantized – not as a consequence of continuous U(1) symmetry, but as an inherent property of the axis discreteness.
2. **Physical determination of label count**: The number of values of Q is not constrained by hypercube geometry. The number of values is determined by what the physics requires – namely, the number of color degrees of freedom.
3. **Coupling strength and channel count**: The coupling strength of a force is proportional to the number of interaction channels provided by Q labels (detailed in Section 7).

1.5 3. Stability of the Spin Classification

1.5.1 3.1 Spin Derivation (Restated)

As in the preceding work [7], spin is determined by the number n of non-spatial axes contained in the normal bivector.

Since a bivector is composed of two axes, the number of non-spatial axes is limited to $n = 0, 1, 2$. This holds regardless of whether there are two or three non-spatial axes.

n	Structure of C	Boson spin	Fermion spin
0	No non-spatial reversal	0	1/2
1	One non-spatial reversal	1	3/2
2	Two non-spatial reversals	2	5/2

Proposition 3.1 (Spin stability). The set of spin quantum numbers $s \in \{0, \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}\}$ remains unchanged when the number of non-spatial axes increases from two (5-dimensional) to three (6-dimensional).

Proof. A bivector is composed of two axes. Since n is the number of non-spatial axes contained in the bivector, $n \leq 2$. The maximum value of n cannot exceed 2 regardless of how many non-spatial axes exist. □

1.5.2 3.2 Increase in the Number of Types

While the set of spin values is invariant, the number of types belonging to each spin increases.

Spin	n	5D types	6D types	Source of increase
$s = 0, \frac{1}{2}$	0	3	3	None
$s = 1, \frac{3}{2}$	1	6	**9**	3 types $e_i \wedge e_Q$
$s = 2, \frac{5}{2}$	2	1	**3**	2 types $e_t \wedge e_Q$, $e_R \wedge e_Q$
Total		**10**	**15**	+5

1.6 4. Geometric Derivation of Color Charge

1.6.1 4.1 Four Values of the Q Axis and Color

The four values $Q \in \{0, r, g, b\}$ have the following physical correspondence:

Q value	Color charge	Particle type
0	Colorless (color-neutral)	Lepton
r	Red	Quark
g	Green	Quark
b	Blue	Quark

Proposition 4.1 (Quark–lepton theorem). States with $Q = 0$ correspond to leptons (no color charge), while states with $Q \in \{r, g, b\}$ correspond to quarks (with color charge). For a given geometric orientation (x, y, z, t, R), the four values of Q cause each generation of fermions to split into one lepton type and three quark colors.

Remark (On the degrees of freedom for electric and color charge). In the Standard Model, electric charge and color charge are orthogonal, independent quantum numbers. In the present framework, color charge is given by the Q-axis value (0 vs $\{r, g, b\}$), while the sign of electric charge is distinguished by the orientation signs of the spatial axes (see Table A in Section 8: within each generation, spatial signs $+$ and $-$ separate charged and neutral particles). Furthermore, the denominator 3 in quark fractional charges $(\pm 1/3, \pm 2/3)$ coincides with the number of nonzero Q-axis values ($|\{r, g, b\}| = 3$). That is, leptons ($Q = 0$, colorless) carry electric charge in units of 1, while quarks ($Q \neq 0$, 3 colors) carry electric charge in units of 1/3. The complete derivation of electric charge (in particular, the asymmetry between $+2/3$ and $-1/3$) remains a task for future work (see Section 9).

We also considered adding electric charge as an independent seventh axis. However, increasing the number of non-spatial axes to four would raise the number of $n = 2$ (spin-2) types to $\binom{4}{2} = 6$ and expand the spin-3/2 fermion families to four, thereby producing a particle spectrum similar to that of supersymmetry theories. Given the current experimental situation (non-observation of supersymmetric particles), this paper adopts the approach of reading electric charge from the Q-axis structure and spatial axis signs. This design allows natural extension by adding a seventh axis should supersymmetric particles be discovered in the future.

1.6.2 4.2 Derivation of Eight Gluon States

Gluons are spin-1 gauge bosons that mediate color charge transitions. For the three nonzero values $\{r, g, b\}$ of the Q axis, gluons are represented as color–anticolor pairs.

Proposition 4.2 (Gluon theorem). The number of independent gauge boson states mediating transitions among color labels $\{r, g, b\}$ is

$$3 \times 3 - 1 = 8$$

The -1 arises because the color singlet (the state proportional to $r\bar{r} + g\bar{g} + b\bar{b}$) is not included among the physical gluons.

Proof. Color transitions are described by pairs $Q_{\text{initial}} \to Q_{\text{final}}$. There are $3^2 = 9$ transitions from three colors to three colors. However, one linear combination of the diagonal components (the color singlet) acts identically on all colors and does not mediate color transitions. Therefore, the number of independent states is $9 - 1 = 8$. □

Remark. In the Standard Model, the eight gluon states are derived as the 8-dimensional adjoint representation of the Lie algebra $\mathfrak{su}(3)$ of SU(3). In this paper, the same number 8 is derived purely combinatorially from the 4-label structure of the Q axis. No continuous group structure of SU(3) needs to be assumed.

1.6.3 4.3 Complete Classification of Gauge Bosons

According to the type of non-spatial axis, spin-1 gauge bosons are classified as follows:

Fixed axis	$n = 1$ type	Boson orientation	SM correspondence	Count
t axis	Temporal	$(0, 0, 0, \pm, 0, Q)$	W^+, W^-	2
R axis	Curvature	$(0, 0, 0, 0, \pm, Q)$	γ, Z^0	2
Q axis	Color	$(0, 0, 0, 0, 0, Q_i \to Q_j)$	Gluons g_1–g_8	8
Total				**12**

The two W boson states ($t = +1, -1$) correspond to positive and negative charges. γ and Z^0 correspond to the sign of the R axis, with the distinction between massless ($M(\gamma) = 0$) and massive ($M(Z) \neq 0$). The eight gluon states follow from Proposition 4.2.

The total of 12 agrees with the number of gauge boson states in the Standard Model ($\gamma + g \times 8 + W^+ + W^- + Z^0 = 12$).

1.7 5. Geometric Placement of the Graviton and Spin-2 Bosons

1.7.1 5.1 Resolution by Three Non-Spatial Axes

In five dimensions (two non-spatial axes), the only $n = 2$ bivector was $e_4 \wedge e_5$ ($t \wedge R$) – **one combination**. In six dimensions (three non-spatial axes: t, R, Q), $n = 2$ increases to **three combinations**:

Remark (On Q-axis participation in bivectors). The Q axis is a label axis and does not participate in the geometric faces of the hypercube (the 80 faces of Section 6.3). However, in the classification of bivectors that determine spin, Q participates as one of the non-spatial axes in the type enumeration. That is, $e_t \wedge e_Q$ and $e_R \wedge e_Q$ are not geometric 2-faces but **abstract bivector types** defined as combinations of non-spatial axes.

Normal bivector	Fixed axes	C matrix
$e_t \wedge e_R$	t, R	$\mathrm{diag}(+, +, +, -, -, Q)$
$e_t \wedge e_Q$	t, Q	$\mathrm{diag}(+, +, +, -, R, \bar{Q})$
$e_R \wedge e_Q$	R, Q	$\mathrm{diag}(+, +, +, t, -, \bar{Q})$

Remark (C matrix notation for the Q component). In the C matrix, the Q component is written as Q when it is preserved. When the Q component is "reversed," it is written as $\bar{Q}$, meaning anticolor substitution ($r \leftrightarrow \bar{r}$, $g \leftrightarrow \bar{g}$, $b \leftrightarrow \bar{b}$, $0 \leftrightarrow 0$).

Proposition 5.1. In the 6-dimensional framework, spin-2 bosons come in three types.

Type	Fixed axes	Force property	SM correspondence
tR type	t, R	Attractive only ($n = 2$, even)	Graviton
tQ type	t, Q	Attractive only	**Prediction**
RQ type	R, Q	Attractive only	**Prediction**

The graviton naturally corresponds to the tR type. The tQ and RQ types are predictions of spin-2 bosons with no counterpart in the Standard Model.

1.7.2 5.2 Resolution of the 5-Dimensional Problem

The problem in five dimensions – that "the graviton does not fit into a 4-facet center" – was due to dimensional insufficiency. In six dimensions, with three non-spatial axes, multiple $n = 2$ bivectors exist, providing the graviton with a natural geometric position.

1.8 6. Geometric Derivation of Mass via the Fifth Axis (R Axis)

1.8.1 6.1 Uniqueness of the R Axis: The Sole Unbounded Degree of Freedom

The R axis is the only one among the six axes with an unbounded range.

Axis	Number of possible values
x, y, z	S_3 constrained (3 generations)
t	2
Q	4
R	∞

1.8.2 6.2 Mass and Area

The R axis corresponds to the curvature radius and has dimension L (length). The area of square faces containing the R axis is proportional to the value of R, so mass is determined by the area structure of R-containing faces.

Definition 6.1. The **signed area** $M(\sigma)$ of an orientation structure σ is defined as the sum of the products of orientation signs and areas over all square faces containing the R axis:

$$M(\sigma) = \sum_{\text{square face } f \text{ containing } R} \operatorname{sgn}(f) \cdot \operatorname{Area}(f)$$

1.8.3 6.3 Classification of R-Containing Faces

Among the 80 square faces, those containing the R ($= x_5$) axis are classified as follows:

Face type	Bivector	Type count	Face count	Area scale
Spatial–R	$e_i \wedge e_5$	3	$3 \times 8 = 24$	$L \cdot R$
t–R	$e_4 \wedge e_5$	1	8	$T \cdot R$
Total		**4**	**32**	

Faces containing the R axis account for 32 out of 80 total faces, or $\frac{2}{5}$ of the total.

1.8.4 6.4 Masslessness Condition

Proposition 6.1. When the signed area $M(\sigma) = 0$ (complete cancellation of signs among all R-containing faces), the orientation corresponds to a massless particle.

The masslessness of the photon and gluons is explained as a consequence of the vanishing signed area of the corresponding orientations.

1.9 7. Determination of Coupling Strength by Channel Count

1.9.1 7.1 Channel Count and Coupling Constants

Proposition 7.1 (Qualitative indication). The coupling strength of a force may be qualitatively related to the type of axis involved in the interaction and the number of channels.

Force	Mediator	Axis involved	Coupling characteristic
Strong	g_1–g_8	Q axis (8 color-transition channels)	Between color-charged particles only
Electromagnetic	γ	R axis	Between particles with nonzero electric charge (see Section 4 remark)
Weak	$W^{\pm}, Z^0$	t axis	All fermions
Gravity	G	$t \wedge R$ ($n = 2$)	All particles (Q-independent)

Remark. In the Standard Model, coupling constants near the energy scale M_Z are $\alpha_s \approx 0.1$, $\alpha \approx 1/137$, $\alpha_W \approx 1/30$, $\alpha_G \approx 10^{-39}$; the strong, electromagnetic, and weak forces are of comparable order, while gravity alone is extremely weak. Quantitative derivation of this hierarchy structure remains a task for future work.

1.10 8. Complete Correspondence Table with Standard Model Particles

1.10.1 Table A: Confirmed Particles ↔ 6-Dimensional Orientations

Bosons

Particle	Symbol	s	Orientation (x, y, z, t, R, Q)	n	Mass (MeV)	Note
Higgs	H^0	0	$(\pm, 0, 0, 0, 0, 0)$ etc.	0	125,250	$Q = 0$, color-neutral
W boson	W^+	1	$(0, 0, 0, +, 0, 0)$	1	80,379	t type
W boson	W^-	1	$(0, 0, 0, -, 0, 0)$	1	80,379	t type
Photon	γ	1	$(0, 0, 0, 0, +, 0)$	1	0	R type, $M = 0$
Z boson	Z^0	1	$(0, 0, 0, 0, -, 0)$	1	91,188	R type, $M \neq 0$
Gluon	g_1–g_8	1	$(0, 0, 0, 0, 0, Q_i \to Q_j)$	1	0	Q type, 8 color transitions
Graviton	G	2	$(0, 0, 0, \pm, \pm, 0)$	2	0	tR type

Fermions: Leptons ($Q = 0$, color-neutral)

Particle	Symbol	s	Gen.	Orientation (x, y, z, t, R, Q)	Mass (MeV)
Electron	e^-	1/2	1	$(+, +, 0, 0, 0, 0)$	0.511
Electron ν	ν_e	1/2	1	$(+, -, 0, 0, 0, 0)$	$< 10^{-6}$
Muon	μ^-	1/2	2	$(+, 0, +, 0, 0, 0)$	105.66
Muon ν	ν_μ	1/2	2	$(+, 0, -, 0, 0, 0)$	< 0.17
Tau	τ^-	1/2	3	$(0, +, +, 0, 0, 0)$	1,776.9
Tau ν	ν_τ	1/2	3	$(0, +, -, 0, 0, 0)$	< 15.5

Antileptons correspond to orientations with reversed spatial axis signs.

Fermions: Quarks ($Q \in \{r, g, b\}$, color-charged)

Particle	Symbol	s	Gen.	Orientation (x, y, z, t, R, Q)	Mass (MeV)
Up	u	1/2	1	$(+, +, 0, 0, 0, r/g/b)$	2.16
Down	d	1/2	1	$(+, -, 0, 0, 0, r/g/b)$	4.67
Charm	c	1/2	2	$(+, 0, +, 0, 0, r/g/b)$	1,270
Strange	s_q	1/2	2	$(+, 0, -, 0, 0, r/g/b)$	93.4
Top	t_q	1/2	3	$(0, +, +, 0, 0, r/g/b)$	172,760
Bottom	b	1/2	3	$(0, +, -, 0, 0, r/g/b)$	4,180

Each quark has three color states $Q = r, g, b$. Antiquarks correspond to reversed spatial signs and anticolors.

Summary

Category	s	Q	Types	States	SM states
Higgs	0	0	1	1	1
Gauge boson (t type)	1	0	2	2	2 ($W^\pm$)
Gauge boson (R type)	1	0	2	2	2 (γ, Z)
Gluon (Q type)	1	transition	8	8	8
Graviton (tR type)	2	0	1	1	1
Lepton	1/2	0	6	6	6
Antilepton	1/2	0	6	6	6
Quark	1/2	r,g,b	6	18	18
Antiquark	1/2	$\bar{r}, \bar{g}, \bar{b}$	6	18	18
Total				**62**	**61 + G**

Remark. The Standard Model comprises 61 states (excluding the graviton). The graviton G is not an element of the Standard Model but a hypothetical particle of quantum gravity. The 62 states of this paper provide a unified classification of the 61 Standard Model states plus one graviton state ("SM + gravity").

1.10.2 Table B: Predictions of Undiscovered Particles

Prediction	s	B/F	n	Orientation feature	Gen.	States
tQ type boson	2	B	2	t, Q fixed	–	Predicted
RQ type boson	2	B	2	R, Q fixed	–	Predicted
t type fermion	3/2	F	1	Spatial–t normal	3 gen.	12 x 4(Q)
R type fermion	3/2	F	1	Spatial–R normal	3 gen.	12 x 4(Q)
Q type fermion	3/2	F	1	Spatial–Q normal	3 gen.	12 x 4(Q)
tR type fermion	5/2	F	2	t–R normal	None	4 x 4(Q)
tQ type fermion	5/2	F	2	t–Q normal	None	4 x 4(Q)
RQ type fermion	5/2	F	2	R–Q normal	None	4 x 4(Q)

Remark. The $s = 3/2$ fermions split into three families (t type, R type, Q type), each possessing a three-generation structure. Compared with the gravitino (one species) of supersymmetry theory, this predicts a far richer structure.

1.10.3 Summary of Improvements from Five Dimensions

Problem	5D (W7)	6D (this paper)	Status
Color charge	Absent	$Q \in \{0, r, g, b\}$	**Resolved**
Quark–lepton distinction	Impossible	$Q = 0$ vs $Q \neq 0$	**Resolved**
Gluon state count	Insufficient (4 vs 12)	$3 \times 3 - 1 = 8$	**Resolved**
Gauge boson total	4	12 ($W^{\pm} + \gamma + Z + g \times 8$)	**Resolved**
Graviton	Position unclear	tR type, $n = 2$	**Resolved**
Spin types	6	6 (stable)	**Confirmed**

1.11 9. Open Problems and Future Work

1.11.1 9.1 Quantitative Computation of Signed Area

Explicitly computing the signed area $M(\sigma)$ of Definition 6.1 for each orientation structure and comparing with known particle mass ratios is the most direct test of this theory. Particularly important are:

- $m_\mu/m_e \approx 206.8$: inter-generation mass ratio
- $m_W/m_Z \approx 0.881$: relation to the Weinberg angle
- $m_t/m_e \approx 3.4 \times 10^5$: maximum mass ratio

1.11.2 9.2 Derivation of the Gauge Group

There is a suggestive correspondence between the 4-label structure of the Q axis and SU(3), the 2 values of the t axis and SU(2), and the continuous scale of the R axis and U(1). A rigorous proof of the mechanism by which continuous gauge groups emerge from discrete structures remains open.

1.11.3 9.3 CKM and PMNS Matrices

How inter-generation mixing (CKM matrix, PMNS matrix) is derived as a breaking of S_3 symmetry remains unresolved.

1.11.4 9.4 Higgs Mechanism

The Higgs boson corresponds to $n = 0$ (spatial axes only), but how spontaneous symmetry breaking by the Higgs field is understood within this framework is a task for future work.

1.11.5 9.5 Additional Spin-2 Boson Predictions

If tQ-type and RQ-type spin-2 bosons (other than the graviton) exist, their experimental consequences must be clarified.

1.11.6 9.6 Geometric Derivation of Electric Charge Values

This paper adopted the approach of reading the sign of electric charge from the orientation signs of spatial axes, and the unit of the charge magnitude (1 vs 1/3) from the Q-axis sector ($Q = 0$ vs $Q \neq 0$) (see Section 4.1 remark). However, the specific charge assignment rules of Table A — namely, that for the same spatial orientation pattern, $Q = 0$ (lepton) yields charge -1 while $Q \neq 0$ (quark) yields charge $+2/3$, and for another spatial orientation pattern, $Q = 0$ yields charge 0 while $Q \neq 0$ yields charge $-1/3$ — have not been derived from a deeper geometric principle. In particular, the asymmetry between $+2/3$ and $-1/3$ is the most prominent open problem of this paper.

1.12 10. Conclusion

From the 6-dimensional framework obtained by adding a sixth axis (Q axis: 4-valued discrete label) to the 5-dimensional hypercube, the following have been derived:

1. **Stability of spin classification**: The set of spin types is the same six as in five dimensions ($s = 0, \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}$) and is stable under dimensional increase.

2. **Geometric derivation of color charge**: The four labels $\{0, r, g, b\}$ of the Q axis provide color charge and distinguish $Q = 0$ (leptons) from $Q \neq 0$ (quarks).

3. **Derivation of eight gluon states**: The number is determined as $3 \times 3 - 1 = 8$ color transitions without assuming SU(3).

4. **Complete correspondence of 12 gauge boson states**: The three families – t type ($W^{\pm}$), R type (γ, Z^0), and Q type (g_1–g_8) – reproduce all 12 Standard Model gauge boson states exactly.

5. **Natural placement of the graviton**: With three non-spatial axes, the $n = 2$ (spin-2 boson) naturally corresponds to the graviton as the tR type. Additionally, tQ-type and RQ-type spin-2 bosons are newly predicted.

6. **Areal origin of mass**: The signed area of faces containing the R axis determines mass. The R axis is the sole unbounded degree of freedom among the six axes, providing an unbounded range for mass.

7. **Qualitative classification of forces**: A qualitative classification has been given in which different types of axes are involved in different forces (strong $\leftrightarrow Q$ axis, electromagnetic $\leftrightarrow R$ axis orientation, weak $\leftrightarrow t$ axis, gravity $\leftrightarrow t \wedge R$). Quantitative derivation of the coupling constant hierarchy remains a task for future work.

8. **Reproduction of 62 SM + gravity states**: A correspondence has been given for 14 boson states ($H + W^{\pm} + \gamma + Z + g \times 8 + G$) and 48 fermion states (12 leptons + 36 quarks). Of these, 61 states correspond to the Standard Model and the graviton state corresponds to quantum gravity.

The structure in which six coordinate axes divide into four qualitatively distinct types – **position** (3 spatial axes), **direction** (1 temporal axis), **scale** (1 curvature axis), and **label** (1 color axis) – suggests that many of the free parameters of the Standard Model are geometric consequences.

1.13 References

[1] Finkelstein, D. & Rubinstein, J. "Connection between Spin, Statistics, and Kinks." *J. Math. Phys.* 9, 1762 (1968).

[2] Atiyah, M.F., Manton, N.S. & Schroers, B.J. "Geometric Models of Matter." *Proc. R. Soc. A* 468, 1252 (2012). arXiv: 1108.5151.

[3] Furey, C. "Standard Model Physics from an Algebra?" arXiv: 1611.09182 (2016).

[4] Kaluza, Th. "Zum Unitatsproblem der Physik." *Sitzungsber. Preuss. Akad. Wiss. Berlin* 966-972 (1921).

[5] Coxeter, H.S.M. *Regular Polytopes.* Dover (1973).

[6] Kihara, N. "Geometric Classification of Spin Derived from the Orientation Structure of the 5-Dimensional Orthoplex." DOI: 10.5281/zenodo.19630972 (2026).

[7] Kihara, N. "Bivectorial Classification of Spin Derived from the Orientation Structure of the 5-Dimensional Hypercube." DOI: 10.5281/zenodo.19643358 (2026).

[W5] Kihara, N. "Complete Spherical Coverage of Central Projection in Discrete Space and the Number-Theoretic Necessity of the 5-Dimensional Background Space." DOI: 10.5281/zenodo.19624957 (2026).