

Week 2-4: Propositional Logic (Tutorial starts from Page 8)

Wang Xiyu

Contents

1 Syntax	2
1.1 Well-formed Formula (WFF)	2
1.2 Closed Set	2
1.2.1 All Closed Set	2
1.3 FORM is the minimum of the set of Closed	3
1.4 Subformula	4
2 Semantics	4
2.1 Truth Assignment/Proposition evaluation	4
2.2 Entail/Model	4
2.2.1 Valuation and Model	5
2.2.2 Logical Equivalence	5
2.3 Satisfiability of Formula	5
2.3.1 Valid formula(tautology)	5
2.4 Satisfiability of a set of formula	5
2.4.1 Relevance	6
2.4.2 Logical consequence	6
2.5 Modelling Computations as Propositional Logic	7
3 Compactness Theorem	8

1 Syntax

Definition 1.1. *DSyntax decides what are allowed to be written*

1.1 Well-formed Formula (WFF)

A formula ψ is obtained by composing $p \in P$ with $c \in C$, where P is the set of propositions and C is the set of connectors. here we use any set of complete logic, like $C = \{\neg, \vee\}$. This is not a precise definition as counter examples that do not form a valid formula can be formulated based on this, for example $\neg \vee p$ means nothing. To formalize the definition we may attempt to define a language with the following grammar:

$$\Sigma = P \cup C \cup \{(\cdot)\}$$

$$FORM : \{\psi = p \in P \mid \neg\psi \mid \psi \vee \psi \mid (\psi)\}$$

However this is a recursive definition as the definition of the a formula ψ involves itself. We need to find a declarative way to define this recursive/inductive construction.

Definition 1.2. *FORM is the smallest set of strings over Σ where*

1. $P \subseteq FORM$
2. $\forall \psi \in FORM, \neg\psi \in FORM$
3. $\forall \psi_1, \psi_2 \in FORM, \psi_1 \vee \psi_2 \in FORM$

We can see as a construction rule, we are admitting strings into the set of formula, but we need to somewhat impose restriction on what is not acceptable.

1.2 Closed Set

Let's remove the attributive **the smallest** first and define another set that contains all the set that satisfy these rules listed above:

Definition 1.3. *Closed is a set of strings over Σ where*

1. $P \subseteq Closed$
2. $\forall \psi \in Closed, \neg\psi \in Closed$
3. $\forall \psi_1, \psi_2 \in Closed, \psi_1 \vee \psi_2 \in Closed$

1.2.1 All Closed Set

And we define a set ACS (All-closed sets). We can define FORM via the following constriction:

$$FORM = \bigcap ACS$$

Suppose $BCS = \emptyset$ and from definition

$$U_0 = \bigcap BCS = \{w \in \Sigma^* \mid \forall S \in BCS, w \in S\}$$

However, $BCS = \emptyset$, The statement $\forall S \in BCS, w \in S$ is vacuously true $\forall w \in \Sigma^*$, which means that

$$U_0 = \Sigma^* = \bigcap BCS$$

A contradiction. Therefore we have proved that ACS cannot be empty, thus $\forall P, \exists S$ such that S is a closed set.

1.3 FORM is the minimum of the set of Closed

First let's prove that FORM is closed. It seems to be trivial that the intersection of closed sets is closed but formal proof is still needed. We denote a set from the intersection construction as T_0 .

Lemma 1.4. T_0 is closed

We are proving that the set T_0 we get from the intersection construction still satisfy the rules that define what is closed.

Proof. From rule 1,

$$\begin{aligned} \forall S \in ACS, P \subseteq S \\ \rightarrow \forall p \in P, \forall S \in ACS, p \in S \\ \rightarrow \forall p \in P, p \in \bigcap ACS \\ \rightarrow P \in T_0 \end{aligned}$$

From rule 2,

$$\begin{aligned} \forall S \in ACS, T_0 \subseteq S \\ \text{let } w \in T_0, \forall S \in ACS, w \in S, \neg w \in S \\ \rightarrow \forall w \in T_0, \neg w \in \bigcap ACS \\ \rightarrow \forall w \in T_0, \neg w \in T_0 \end{aligned}$$

From rule 3,

$$\begin{aligned} \forall S \in ACS, T_0 \subseteq S \\ \text{let } w_1, w_2 \in T_0, \forall S \in ACS, w_1 \vee w_2 \in S \\ \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in \bigcap ACS \\ \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in T_0 \end{aligned}$$

□

Lemma 1.5. T_0 is **a** smallest Closed set

Proof. The proof of minimality is trivial. $T_0 = \bigcap ACS \rightarrow \forall S \in ACS, T_0 \subseteq S$

□

Lemma 1.6. T_0 is **the** smallest Closed set (unique). In other word,

$$\forall T'_0 \in \{T \mid \forall S \in ACS, T \subseteq S\}, T'_0 = T_0$$

Proof. Suppose otherwise, $\exists T'_0 \in ACS \neq T_0$.

$$\begin{aligned} T_0, T'_0 \in ACS \rightarrow \forall S \in ACS, T_0 \subseteq S, T'_0 \subseteq S \\ \rightarrow T_0 \subseteq T'_0 \wedge T'_0 \subseteq T_0 \\ \rightarrow T_0 = T'_0 \end{aligned}$$

A contradiction.

□

Now we can use the generative rule to define formula, which is called **Backus-Naur form**. Note that a context-free grammar is a type of formal language. Backus Naur form is a specification language for this type of grammar. It is used to describe language syntax.

1.4 Subformula

Definition 1.7. ϕ_1 is a subformula of ϕ_2 , denoted by

$$\phi_1 \preceq \phi_2$$

as a relation is

- *reflexive:* $\forall \phi \in FORM, \phi \preceq \phi$
- *transitive:* $\forall \phi_1, \phi_2, \phi_3 \in FORM, (\phi_1 \preceq \phi_2) \wedge (\phi_2 \preceq \phi_3) \rightarrow \phi_1 \preceq \phi_3$

Inductive definition:

$$\begin{aligned} \forall \phi \in FORM, \phi \preceq (\neg \phi) \\ \forall \phi_1, \phi_2 \in FORM, \phi_1 \preceq (\phi_1 \vee \phi_2) \\ \phi_2 \preceq (\phi_1 \vee \phi_2) \end{aligned}$$

We might intuitively think that subformula can be defined using the notion of substring, since we have been considering formula as concatenation of proposition and connector symbols. However based on our inductive definition, we can come up with the following counter example to a definition using substring: A formula ϕ_1 is a subformula of ϕ_2 if and only if ϕ_1 is a well-formed formula, and is a substring to ϕ_2

$$p \vee q \not\preceq \neg p \vee q$$

What if we tighten the syntax by strictly enforce the use of parentheses whenever we can

2 Semantics

Definition 2.1. *Semantics describe meanings.*

2.1 Truth Assignment/Proposition evaluation

A truth assignment or a valuation of propositions is a function that maps from the set of propositions to true and false.

$$\begin{aligned} v : P \rightarrow \{T, F\} \\ v \in [P \rightarrow \{T, F\}] \end{aligned}$$

Example 2.2. Consider $v = \{p_1 \mapsto T, p_2 \mapsto F\}$, we can derive

$$\begin{aligned} v(p_1) &= T \\ v(p_2) &= F \\ v(\neg p_2) &= T \\ v(p_1 \vee p_2) &= T \text{ etc.} \end{aligned}$$

2.2 Entail/Model

We denote entailment of an evaluation function to formula as

$$v \models \psi$$

We also inductively define the following rule

$$\begin{aligned} \forall p \in P, v \models p &\iff v(p) \\ v \models \psi &\iff v \not\models \neg \psi \\ v \models \psi_1 \vee \psi_2 &\iff (v \models \psi_1) \vee (v \models \psi_2) \end{aligned}$$

2.2.1 Valuation and Model

A valuation v is a model of a formula ϕ , if and only if $v(\phi) = T$.

- $\forall \phi$ over P , $\exists 2^{|P|}$ valuations.
- A formula can have many or one or none models

2.2.2 Logical Equivalence

Definition 2.3. *Syntax definition of logical equivalence is as follows:*

$$\phi_1 \equiv \phi_2 \iff \forall v : P \mapsto \{T, F\} (v \models \phi_1), v \models \phi_2$$

Example 2.4.

$$p \vee (\neg p) \equiv q \vee (\neg q)$$

$$\begin{aligned} \forall v &\models (p \vee (\neg p)), \\ v &\models p \text{ or } v \models (\neg p) \\ v &\models p \text{ or } v \not\models p \\ \forall v : P &\mapsto \{T, F\}, \end{aligned}$$

TODO

2.3 Satisfiability of Formula

Definition 2.5. *Let $\phi \in FORM$, ϕ is satisfiable*

$$\iff \exists v : P \mapsto \{T, F\}, v \models \phi$$

2.3.1 Valid formula(tautology)

Definition 2.6. *Tautology: ϕ is a tautology $\iff \forall v, v \models \phi$*

Example 2.7.

$$\phi = \neg(p \vee (\neg p))$$

From example 2.4, we have

$$\begin{aligned} \forall v, v &\models (p \vee (\neg p)) \\ \rightarrow \forall v, v &\not\models \neg(p \vee (\neg p)) \end{aligned}$$

Thus ϕ is NOT satisfiable.

2.4 Satisfiability of a set of formula

Set Satisfication

Definition 2.8. *Let $\Gamma \subseteq FORM$,*

$$v \models \Gamma \iff \forall \phi \in \Gamma, v \models \phi$$

Set Satisfiability

Definition 2.9. Let $\Gamma \subseteq FORM$, Γ is satisfiable

$$\iff \exists v, \forall \phi \in \Gamma, v \models \phi$$

Example 2.10. $\Gamma = \emptyset$, Γ is satisfiable.

Proof. Consider $v : P \mapsto \{T, F\}$, $v(p) = T \forall p \in P$, $\forall \phi \in \Gamma, v \models \phi$ is vacuously true. $\square$

Example 2.11. $\Gamma = FORM$, Γ is NOT satisfiable.

Proof. Suppose otherwise,

$$\begin{aligned} \exists v, \forall \phi \in \Gamma, v \models \phi \\ \forall \phi \in \Gamma, (\neg \phi) \in \Gamma \\ v \models (\neg \phi) \end{aligned}$$

A contradiction. $\square$

2.4.1 Relevance

Definition 2.12. Define set $OCCUR(\phi) \subseteq P$, which contains propositions appear in ϕ . Inductively defined as

$$\begin{aligned} OCCUR(\phi) &= \{p\} \iff \phi \equiv p, p \in P \\ OCCUR(\neg \phi) &= OCCUR(\phi) \\ OCCUR(\phi_1 \vee \phi_2) &= OCCUR(\phi_1) \cup OCCUR(\phi_2) \end{aligned}$$

Lemma 2.13. Let $\phi \in FORM$, let v_1, v_2 be valuations over P .

$$\forall p \in OCCUR(\phi), v_1(p) = v_2(p) \rightarrow (v_1 \models \phi \rightarrow v_2 \models \phi)$$

v_1, v_2 can be different valuations over $p' \notin OCCUR(\phi)$

2.4.2 Logical consequence

Semantic logical consequence:

Definition 2.14.

$$\Gamma \models \phi \iff \forall v \models \Gamma, v \models \phi$$

Syntactic logical consequence: (logical derivability)

Definition 2.15. A formula ϕ is a syntactic consequence of Γ if there exists a formal proof (a finite sequence of formulas) ending in ϕ , where each step is either a premise from Γ , an axiom, or follows from previous steps via an inference rule (like Modus Ponens).

Material Implication ($\rightarrow$) vs. Logical Consequence ($\models$)

- Material implication is a connector within a formula, $p \rightarrow q \equiv \neg(p \vee q)$, and can therefore be eval to true or false in a single model.
- Logical consequence is a relation between formulas (meta-logic), it is a statement about all models. $\{p, p \rightarrow q\} \models q$

2.5 Modelling Computations as Propositional Logic

Computation problems can be considered as membership check of an encoded problem instance to a language.

$$L = \{p \mid \langle M, p \rangle\}$$

Here instead of arbitrary encoding, we consider problem encoding with propositional logics only, and solve with SAT solvers by deciding whether the set of formula we form from any instance of the problem is satisfiable. That is

$$\text{Encode} : p \mapsto \text{FORM}, \forall p, \exists v \models \text{Encode}(p) \iff p \in L$$

The rules of such modelling include:

- Propositional logics only
- The final expression must one well formed fomula
- for problem of size n , the size of the formula ϕ must be in $O(\text{poly}(n))$

Note that there can be different definition on the size of the

- problem: in unary, binary etc;
- Formula: Whether by the number of nodes in the syntax tree or length of the string **Anything else?**

Example 2.16. *Graph coloring problem: given a undirected graph $G = (V, E)$, whether there exists a k coloring of all vertices such that no neighbouring vertices have the same color. Consider the following construction of a proposition set P :*

$$P = \{p_{v,i} \mid v \in V, i \in [1, k]\}$$

The valuation of proposition $p_{v,i}$ conceptually represents vertex v is assigned color i .

We can formulate constriants as well formed formula, and use conjunction to connect the formula to model satisfaction of multiple constraints.

Corollary 2.17. $\forall \phi_1, \phi_2 \in \text{FORM}, (\phi_1 \wedge \phi_2) \in \text{FORM}$

First let us list down the constraints of this problems, including implicit ones in plain words

1. *Neighbouring vertices have different colors.*
2. *All vertices as assigned with exactly one color.*

1. ϕ_{nbr}

First lets formula the constraint between any 2 vertices,

$$\forall (\alpha, \beta) \in E, c(v_\alpha) \neq c(v_\beta)$$

And from our proposition set, $c(v_\gamma) = i \iff \exists v, v \models p_{\gamma,i}$. As such we have

$$\phi_{\alpha,\beta,i} = \neg(p_{\alpha,i} \wedge p_{\beta,i})$$

This models vertex α and vertex β are not colored with color i at the same time. For every color this needs to hold, thus we have

$$\phi_{nbr(\alpha,\beta)} = \bigwedge_{i \in [1,k]} \phi_{\alpha,\beta,i}$$

For every pair of neighbouring vertices this also needs to hold. Therefore we have

$$\phi_{nbr} = \bigwedge_{(\alpha,\beta) \in E} \phi_{nbr(\alpha,\beta)}$$

2. $\phi_{\text{exactly-1}}$

To model exactly 1 coloring on each vertex, we can formulate $\phi_{\geq 1}$ and $\phi_{\leq 1}$ first and use conjunction to connect them. Given our propositions, to make sure vertex v is at least assign a color ($\phi_{v,\geq 1}$),

$$\phi_{v,\geq 1} = \bigvee_{i \in [1,k]} p_{v,i}$$

This needs to hold for every vertex, thus we have

$$\phi_{\geq 1} = \bigwedge_{v \in V} \phi_{v,\geq 1}$$

Then we need to model at most 1, (or less than 2) coloring on each vertex. Vertex v is not assignment with both color i and j can be formulated as

$$\phi_{v,i,j,\leq 1} = \neg(p_{v,i} \wedge p_{v,j})$$

This needs to hold for every distinct pair of colors, thus

$$\phi_{v,\leq 1} = \bigwedge_{i \neq j \in [1,k]} \phi_{v,i,j,\leq 1}$$

For all vertices

$$\phi_{\leq 1} = \bigwedge_{v \in V} \phi_{v,\leq 1}$$

Finally we have

$$\phi_{\text{exactly-1}} = \phi_{\leq 1} \wedge \phi_{\geq 1}$$

And the overall formula for an instance of $G = (V, E)$ becomes

$$\phi = \phi_{\text{nbr}} \wedge \phi_{\text{exactly-1}}$$

Similar to reduction, we need also now to show that

$$\exists v : P \mapsto \{T, F\}, v \models \phi \iff \langle G = (V, E), k \rangle \in L_{\text{color}}$$

TODO

3 Compactness Theorem

Definition 3.1. *Axiom of Choice: formally*

$$\forall X, \emptyset \in X \vee \exists f, \text{dom}(f) = X \wedge \forall A \in X, f(A) \in A$$

A choice function (or selection) f defined on a collection of nonempty sets X , such that for every set A in the collection X , $f(A)$ is an element in A . That is, Axiom—For any set X of nonempty sets, there exists a choice function f that is defined on X and maps each set of X to an element of that set.


Theorem 3.2. *If P is finite, FORM over P is finite.*

Theorem 3.3. *If P is countable, FORM over P is countable.*


Proof. Given $\forall \phi \in \text{FORM}$, ϕ is finitely long, this theorem is another way of stating: The language formed from countable alphabet $P \cup C$ with the inductive construction rules, is countable. We can prove a stronger theorem first. $\square$

Theorem 3.4. *All languages formed from a countable alphabet is countable.*

Rational Numbers ($\mathbb{Q}^+$)



Countable Language (Σ^2)



Countable Language (Σ^*)




Figure 1: Comparison of diagonal enumeration for numbers vs. strings.

Proof. Consider $L = \Sigma^*$, we will prove that Σ is countable $\rightarrow L$ is countable. We use the same diagonal proof for the countability of $\mathbb{Q}$ to show an enumeration of all strings formable from a countable alphabet. To formalize, we use induction to prove that $\forall i \geq 0, \Sigma^i$ is countable

Base case: $\Sigma^1 = \Sigma$, countable as defined.

Inductive hypothesis: Σ^n is countable: we enumerate all strings in Σ^n , and use the same diagonal proof to show the concatenation of one symbol from the countable alphabet to every string Σ^{n+1} is also countable. Thus we can give an enumeration of all Σ^i and use the diagonal argument again to show the existence of an enumeration of Σ^* .

Alternatively, Given

$$\Sigma^* = \bigcup_{i \geq 0} \Sigma^i$$

From the Axiom of Choice, which implies the union of countably many countable set is countable, Σ^* is countable. Since $\forall L$ over $\Sigma, L \subseteq \Sigma^*, \Sigma^*$ is countable, all languages over countable Σ is countable. $\square$

Definition 3.5 (Finitely Satisfiable). *A set of formulas Γ is said to be **finitely satisfiable** if and only if for all finite subsets $\Gamma_0 \subseteq \Gamma$, Γ_0 is satisfiable.*

Theorem 3.6 (Compactness Theorem).

$$\Gamma \text{ is satisfiable} \iff \Gamma \text{ is finitely satisfiable}$$

Proof. Consider a special case where Γ is “ T -good”, defined as:

$$\forall p \in P, (p \in \Gamma \vee \neg p \in \Gamma)$$

Claim: If Γ is T -good, then Γ is finitely satisfiable $\implies \Gamma$ is satisfiable.

Suppose Γ is T -good and finitely satisfiable. We define a candidate valuation v_L such that for all $p \in P$:

$$v_L(p) = \begin{cases} T & \text{if } p \in \Gamma \\ F & \text{if } \neg p \in \Gamma \end{cases}$$

First, we prove that for every $p \in P$, **exactly one** of $\{p, \neg p\}$ is in Γ . By the T -good property, at least one is in Γ . Suppose both $\{p, \neg p\} \subseteq \Gamma$. Consider the finite subset:

$$\Gamma' = \{p, \neg p\}$$

Since Γ' is a finite subset of a finitely satisfiable set, it must be satisfiable. However, $\neg \exists v$ such that $v \models \{p, \neg p\}$, which is a contradiction. Thus, v_L is well-defined.

Now we show $v_L \models \Gamma$. Suppose $\exists \phi \in \Gamma$ such that $v_L \not\models \phi$. Define the following finite sets based on the atoms occurring in ϕ :

$$\begin{aligned}\Gamma_\phi^+ &= \{p \mid p \in \text{OCCUR}(\phi) \wedge v_L(p) = T\} \\ \Gamma_\phi^- &= \{\neg p \mid p \in \text{OCCUR}(\phi) \wedge v_L(p) = F\} \\ \Gamma_\phi &= \{\phi\} \cup \Gamma_\phi^+ \cup \Gamma_\phi^-\end{aligned}$$

Since Γ_ϕ is a finite subset of Γ , it must be satisfiable by some valuation v . For this v :

- $\forall p$ such that $p \in \Gamma_\phi^+$, $v(p) = T$.
- $\forall \neg p$ such that $\neg p \in \Gamma_\phi^-$, $v(\neg p) = T \implies v(p) = F$.

By construction, v and v_L agree on all atoms in $\text{OCCUR}(\phi)$. By the **Relevance Lemma**, $v(\phi) = v_L(\phi)$. Since $v \models \Gamma_\phi$, we have $v \models \phi$. This implies $v_L \models \phi$, contradicting our assumption $v_L \not\models \phi$. Therefore, $\forall \phi \in \Gamma, v_L \models \phi$, which implies $v_L \models \Gamma$, Γ is satisfiable.

Now we will generalize this to all Γ which may not be ‘ T -good’, by showing that we can construct a ‘ T -good’ finitely satisfiable set Δ , from any finitely satisfiable Γ , and therefore show that $\Gamma SAT \iff \Delta SAT$.

We will only prove the case for countable P now. Let an enumeration of $p \in P$ be $\{p_1, p_2, \dots\}$

Define finite subset Γ_i of Γ as such:

$$\begin{aligned}\Gamma_0 &= \Gamma \\ \Gamma_{i+1} &= \begin{cases} \Gamma_i \cup \{p_i\} & \text{if } \Gamma_i \cup \{p_i\} \text{ is finitely satisfiable} \\ \Gamma_i \cup \{\neg(p_i)\} & \text{Otherwise} \end{cases}\end{aligned}$$

Here we need to prove that $\forall i$, between $\Gamma_i \cup \{p_i\}$ and $\Gamma_i \cup \{\neg(p_i)\}$ at least one case is also finitely satisfiable.

Theorem 3.7. *Let Γ be finitely satisfiable set, and ϕ be a formula, then $\Gamma \cup \{\phi\}$ is finitely satisfiable or $\Gamma \cup \{\neg(\phi)\}$ is finitely satisfiable.*

Proof. We prove the contrapositive of this statement: if $\Gamma \cup \{\phi\}$ and $\Gamma \cup \{\neg(\phi)\}$ are both not finitely satisfiable, Γ is not finitely satisfiable. Given both are not finitely satisfiable, there exist finite subsets $\Gamma_1, \Gamma_2 \subseteq \Gamma$ such that $\Gamma_1 \cup \{\phi\}$ and $\Gamma_2 \cup \{\neg(\phi)\}$ are not satisfiable. Therefore for all valuations, it must not model either Γ_1 or $\{\phi\}$, and not model either Γ_2 or $\{\neg(\phi)\}$. Denote the set of valuations that models formula ψ as $\text{models}(\psi)$, the set of valuations that models set of formula Σ as $\text{models}(\Sigma)$

$$\begin{aligned}\text{models}(\Gamma_1) \cap \text{models}(\phi) &= \emptyset \\ &\rightarrow \forall v \in \text{models}(\Gamma_1), v \not\models \phi \rightarrow v \models (\neg\phi) \\ &\rightarrow \text{models}(\Gamma_1) \subseteq \text{models}(\neg\phi) \\ \text{models}(\Gamma_2) \cap \text{models}(\neg\phi) &= \emptyset \\ &\rightarrow \forall v \in \text{models}(\Gamma_2), v \not\models (\neg\phi) \rightarrow v \models (\phi) \\ &\rightarrow \text{models}(\Gamma_2) \subseteq \text{models}(\phi)\end{aligned}$$

Now consider the set of valuations that models $\Gamma_1 \cup \Gamma_2$

$$\begin{aligned} \text{models}(\Gamma_1 \cup \Gamma_2) &= \text{models}(\Gamma_1) \cap \text{models}(\Gamma_2) \\ (\text{models}(\Gamma_1) \cap \text{models}(\Gamma_2)) &\subseteq (\text{models}(\phi) \cap \text{models}(\neg\phi)) = \emptyset \end{aligned}$$

Since $\Gamma_1 \cup \Gamma_2$ is a finite subset of Γ that is not satisfiable, Γ is not finitely satisfiable. □

Now prove that $\forall i, \Gamma_i$ is finitely satisfiable. □

Tutorial

Definition 3.8 (Generic Closure). Let U be a universe of elements and let O be a set of functions whose domain is U and codomain is $\mathcal{P}(U)$. The arity of a function $f \in O$ is denoted by $\text{arity}(f)$. A function f of arity r has type

$$f : U^r \rightarrow \mathcal{P}(U).$$

Let $I \subseteq U$ be an initial set. A set $S \subseteq U$ is said to be (I, O) -closed if:

1. $I \subseteq S$.
2. For every $f \in O$ with $\text{arity}(f) = r$ and for all $e_1, \dots, e_r \in U$, if $e_1, \dots, e_r \in S$, then $f(e_1, \dots, e_r) \subseteq S$.

Lemma 3.9. Let U be a universe, O a set of operators, and $I \subseteq U$ an initial set. There exists a smallest (I, O) -closed set.

Define $\mathcal{P}(U)$ as a collection of all ordering of all elements in the power set $\mathcal{P}(U)$

Q1

Claim: The smallest (I, O) -closed set can be constructed by taking the intersection of all (I, O) -closed set, ie.

$$AIOCS = \{S \subseteq U \mid S \text{ is } (I, O)\text{-closed}\}$$

$$T_0 = \bigcap AIOCS$$

Existence

First we prove the existence of such a set.

Proof. Suppose otherwise,

$$T_0 = \bigcap AIOCS = \emptyset$$

$$T_0 = \{x \in \mathcal{P}(U) \mid \forall S \in AIOCS, x \in S\}$$

$$T_0 = \emptyset \rightarrow x \in S \text{ is vacuously true } \forall x \in \mathcal{P}(U) \rightarrow T_0 = \mathcal{P}(U)$$

contradiction. □

Closed

Now we prove that T_0 is (I, O) -closed.

Proof. Property 1:

$$\forall S \in AIOCS, I \subseteq S$$

$$\rightarrow \forall i \in I, \forall S \in AIOCS, i \in S$$

$$\rightarrow \forall i \in I, i \in \bigcap AIOCS$$

$$\rightarrow \forall i \in I, i \in T_0$$

Property 2:

$$\forall S \in AIOCS, T_0 \subseteq S$$

$$\text{Let } e_1, e_2, \dots, e_r \in T_0, \forall S \in AIOCS, e_1, e_2, \dots, e_r \in S$$

$$\rightarrow \exists f \in O, f(e_1, e_2, \dots, e_r) \in S$$

$$\rightarrow \forall e_1, e_2, \dots, e_r \in T_0, \exists f \in O, f(e_1, e_2, \dots, e_r) \in \bigcap AIOCS$$

$$\rightarrow f(e_1, e_2, \dots, e_r) \in T_0$$

□

Uniqueness

Claim:

$$\forall T'_0 \in \{T \in \mathcal{P}(U) \mid \forall S \in AIOCS, T \subseteq S\}, T'_0 = T_0$$

Proof. Suppose otherwise, $\exists T'_0 \in AIOCS, T'_0 \neq T_0$.

$$\begin{aligned} T_0, T'_0 \in AIOCS &\rightarrow \forall S \in AIOCS, T_0 \subseteq S, T'_0 \subseteq S \\ &\rightarrow T_0 \subseteq T'_0, T'_0 \subseteq T_0 \\ &\rightarrow T_0 = T'_0 \end{aligned}$$

□

Q2

For each of the following strings, prove or disprove that it is a well-formed formula:

1. $w_1 = \epsilon$ (the empty string).
2. $w_2 = (\neg p)$ where $p \in P$.
3. $w_3 = (p\neg)$ where $p \in P$.

2.1

$$w_1 = \epsilon \notin FORM$$

Claim: $\forall w \in FORM, |w| > 0$ I'd like to use definition 2.19 and lemma 2.21 which I proved later without using the conclusion from this question.

$$FORM = INDFORM = \bigcup_{i \geq 0} FORM_i$$

Proof. Base case:

$$\begin{aligned} FORM_0 &= P \\ \forall \phi \in FORM_0, \phi &= p \in P \\ &\rightarrow \forall \phi \in FORM_i, |\phi| > 0 \end{aligned}$$

Inductive Hypothesis: Suppose

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| > 0$$

$$\begin{aligned} FORM_{i+1} &= FORM_i \cup \{(\neg\phi) \mid \phi \in FORM_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in FORM_i\} \\ \forall \psi \in FORM_{i+1}, \exists \phi \in FORM_i, |\psi| &= |\phi| \text{ or } |\psi| = |\phi| + 3 \text{ or} \\ \exists \phi_1, \phi_2 \in FORM_i, |\psi| &= |\phi_1| + |\phi_2| + 3 \end{aligned}$$

□

2.2

$$p \in P \rightarrow p \in FORM \rightarrow (\neg p) \in FORM$$

2.3

$(p \neg) \notin FORM$

Proof. Suppose otherwise, $(p \neg \epsilon) \in FORM$

$$\begin{aligned} \forall \phi \in FORM, (\neg \phi) \in FORM \\ \rightarrow \epsilon \in FORM \end{aligned}$$

Contradiction from 2.1 □

Q3

Fix $U = \mathbb{R}$. For each of the following sets S , define a finite initial set $I \subseteq \mathbb{R}$ and a set of operators O with $|O| < 10$ such that the smallest (I, O) -closed set is S .

1. $S = \{0, 1, 2, 3, \dots\}$.
2. $S = \{1, 3, 5, 7, \dots\}$.
3. $S = \mathbb{Q}$.
4. $S = \{\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \frac{5}{3}, \frac{7}{3}, \dots\}$.

Answer:

1. $\mathbb{N} : I = \{0\}, O = \{f(x) = x + 1\}$
2. $Odd : I = \{1\}, O = \{f(x) = x + 2\}$
3. $\mathbb{Q} : I = \{0\}, O = \{f(x) = -x, f(x) = x + 1, f(x, y) = \begin{cases} \frac{x}{y} & \text{if } y \neq 0 \\ \emptyset & \text{if } y = 0 \end{cases}\}$
4. $\{\frac{x}{3} \mid x \in \mathbb{N}\} \setminus \mathbb{N} : I = \{\frac{1}{3}, \frac{2}{3}\}, O = \{f(x) = x + 1\}$

Finite Directed Graphs

Definition 3.10 (Graph Reachability). *Let $G = (V, E)$ be a directed graph. A path from s to t is a sequence of vertices $\pi = u_1, \dots, u_k$ such that:*

1. $u_1 = s$,
2. $(u_i, u_{i+1}) \in E$ for all $1 \leq i < k$,
3. $u_k = t$.

The length of π is $k - 1$. We say that t is reachable from s if $s = t$ or such a path exists. The set of ancestors of t is defined as

$$Reach_t = \{v \in V \mid t \text{ is reachable from } v\}.$$

Q4

Let $G = (V, E)$ be a directed graph and let $t \in V$. Fix the universe $U = V$.

1. Define an initial set I (with $|I| \leq 2$) and a unary operator f such that the smallest $(I, \{f\})$ -closed set is the set of vertices reachable from t .
2. Let $d \in \mathbb{N} \setminus \{0\}$ and $r < d$. Define I and a unary operator f such that the smallest $(I, \{f\})$ -closed set is the set of vertices reachable from s via paths of length ℓ with $\ell \equiv r \pmod{d}$.

4.1

$$I = \{t\}, O = \{f(v) = \text{succ}(v)\}$$

4.2

$$I = \{s\} \cup \{v \in V \mid v \in \text{succ}^r(s)\}, O = \{f(v) = \text{succ}^d(v)\}$$

Q5

Let $G = (V, E)$ be a directed graph and fix $U = V \times V$. Define an initial set I with $|I| \leq |V|$ and a binary operator f such that the smallest $(I, \{f\})$ -closed set is

$$\text{AllReachPairs} = \{(u, v) \mid v \text{ is reachable from } u\}.$$

5.1

$$I = \{(v, v) \mid v \in V\}, O = \{f((v_1, v_2), (v_3, v_4)) \mapsto \begin{cases} \{(v_1, v_4)\} & \text{if } v_2 = v_3 \\ \emptyset & \text{if } v_2 \neq v_3 \end{cases}\}$$

Inductive Definitions

Definition 3.11 (Inductive Well-Formed Formulae). *Let P be a set of propositions and $C = \{\neg, \vee\}$. Define sets $FORM_i$ as follows:*

$$\begin{aligned} FORM_0 &= P, \\ FORM_{i+1} &= FORM_i \cup \{(\neg w) \mid w \in FORM_i\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in FORM_i\}. \end{aligned}$$

Define

$$INDFORM = \bigcup_{i \geq 0} FORM_i.$$

Q6

1. Show that $FORM_i \subseteq FORM_{i+1}$ for all i .
2. Prove or disprove that $FORM_i = FORM_{i+1}$ for some i .
3. Prove or disprove that $FORM_i$ is (P, C) -closed for all i .
4. Prove or disprove that $FORM_i$ is (P, C) -closed for some i .

6.1

Base case: $FORM_0 \subseteq FORM_1$ as

$$FORM_1 = FORM_0 \cup \{(\neg w) \mid w \in FORM_0\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in FORM_0\}$$

Wait this is just trivial

6.2

Claim:

$$\exists i \geq 0, FORM_i = FORM_{i+1}$$

It is obvious that $\forall i \geq 0, FORM_i$ is finite. Suppose

6.3

Claim:

$$\forall i \geq 0, FORM_i \text{ is (P,C)-closed}$$

Proof. Suppose otherwise, consider $FORM_0$,

$$\begin{aligned} FORM_0 &= P \\ \forall p \in P, (\neg p) &\in FORM_0 \end{aligned}$$

Contradiction □

6.4

Claim:

$$\exists i \geq 0, FORM_i \text{ is (P,C)-closed}$$

Proof.

Lemma 3.12.

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

The longest formula in $FORM_i$ has size $2^{i+3} - 3$. Parentheses are counted in string length

Proof. Base case:

$$\begin{aligned} FORM_0 &= P \\ \forall \phi \in FORM_0, |\phi| &= 1 \end{aligned}$$

Inductive hypothesis:

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

Consider all new strings formed from the longest string $\phi \in FORM_i$,

$$\begin{aligned} |\phi| &= 2^{i+3} - 3 \\ |(\neg\phi)| &= 2^{i+3} \\ |(\phi \vee \phi)| &= 2 \times (2^{i+3} - 3) + 3 = 2^{(i+1)+3} - 3 \end{aligned}$$

□

Suppose $\exists i \geq 0, FORM_i$ is (P,C)-closed, and from lemma 2.20 the longest formula in $FORM_i$ has size $|\phi_{max}| = 2^{i+3} - 3$. $\rightarrow \forall \phi \in FORM_i, (\neg\phi) \in FORM_i \rightarrow \phi'_{max} = (\neg\phi_{max}) \in FORM_i, |\phi'_{max}| > 2^{i+3} - 3$, a contradiction.

□

Lemma 3.13. $FORM = INDFORM$.

Q7

Prove the lemma by showing:

1. $INDFORM$ is (P,C)-closed.
2. For every (P,C)-closed set S , $INDFORM \subseteq S$.

7.1

Claim: $INDFORM$ is (P,C)-closed

Proof. **Claim:** $P \subseteq \text{INDFORM}$

First condition of (P,C)-closed set:

$$\text{FORM}_0 \subseteq \text{INDFORM} \rightarrow P \subseteq \text{INDFORM}$$

Claim: $\forall w \in \text{INDFORM}, (\neg w) \in \text{INDFORM}$

$$\begin{aligned} \text{INDFORM} &= \bigcup_{i \geq 0} \text{FORM}_i \\ &\rightarrow \forall i \geq 0, \text{FORM}_i, \text{FORM}_{i+1} \in \text{INDFORM} \\ &\quad \forall w \in \text{INDFORM}, \exists j \geq 0, w \in \text{FORM}_j \\ &\rightarrow (\neg w) \in \text{FORM}_{j+1} \rightarrow (\neg w) \in \text{INDFORM} \end{aligned}$$

Claim: $\forall w_1, w_2 \in \text{INDFORM}, (w_1 \vee w_2) \in \text{INDFORM}$

Case 1: $\exists i \geq 0, w_1, w_2 \in \text{FORM}_i$

$$w_1, w_2 \in \text{FORM}_i \rightarrow (w_1 \vee w_2) \in \text{FORM}_{i+1} \rightarrow (w_1 \vee w_2) \in \text{INDFORM}$$

Case 2: $\nexists i \geq 0, w_1, w_2 \in \text{FORM}_i$. WLOG, suppose $w_1 \in \text{FORM}_i, w_2 \in \text{FORM}_j, i < j$

$$\forall w \in \text{FORM}_n, w \in \text{FORM}_{n+1} \text{ By induction } w_1 \in \text{FORM}_i \rightarrow w_1, w_2 \in \text{FORM}_j$$

Back to case 1. □

7.2

$$\forall (\text{P,C})\text{-closed set } S, \text{INDFORM} \subseteq S$$

To prove this, we only need to prove that $\text{INDFORM} \subseteq S_{\min} = \bigcap_{S \text{ is } (\text{P,C})\text{-closed}} S$.

Proof. Base case:

$$\bigcup_{i=0}^0 \text{FORM}_i = \text{FORM}_0 = P \subseteq S_{\min}$$

Inductive hypothesis: Suppose

$$\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min}$$

$$\begin{aligned} &\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min} \rightarrow \text{FORM}_n \in S_{\min} \\ &\quad S_{\min} \text{ is } (\text{P,C})\text{-closed} \rightarrow \forall i \in [1, n], \forall \phi \in \text{FORM}_n, (\neg \phi) \in S_{\min}, \\ &\quad \forall i \in [0, n], \forall \phi_1, \phi_2 \in \text{FORM}_i, (\phi_1 \vee \phi_2) \in S_{\min} \\ &\rightarrow \bigcup_{i=0}^n \text{FORM}_i \cup \{(\neg \phi) \mid \forall \phi \in \bigcup_{i=0}^n \text{FORM}_i\} \cup \{(\phi_1 \vee \phi_2) \mid \forall \phi_1, \phi_2 \in \bigcup_{i=0}^n \text{FORM}_i\} \subseteq S_{\min} \\ &\rightarrow \bigcup_{i=0}^{n+1} \text{FORM}_i \subseteq S_{\min} \end{aligned}$$

□

Definition 3.14 (Alternative Inductive Definition). Define $AltFORM_0 = P$ and for $i \geq 0$:

$$\begin{aligned} AltFORM_{i+1} = & \{(\neg w) \mid w \in AltFORM_i\} \\ & \cup \{(w_1 \vee w_2) \mid w_1 \in AltFORM_i, \exists j \leq i, w_2 \in AltFORM_j\} \\ & \cup \{(w_1 \vee w_2) \mid w_2 \in AltFORM_i, \exists j \leq i, w_1 \in AltFORM_j\}. \end{aligned}$$

Define

$$AltINDFORM = \bigcup_{i \geq 0} AltFORM_i.$$

Question 8

1. Show that $FORM_i = \bigcup_{j=0}^i AltFORM_j$ for all i .
2. Show that $INDFORM = AltINDFORM$.

8.1

Lemma 3.15.

Proof. Base case:

$$FORM_0 = \bigcup_{j=0}^0 AltFORM_j$$

$$FORM_0 = P = AltFORM_0$$

Inductive hypothesis: Suppose

$$\forall j \in [0, i], FORM_j = \bigcup_{k=0}^j AltFORM_k$$

$$\begin{aligned} FORM_{i+1} &= \bigcup_{k=0}^i AltFORM_k \cup \{(\neg \phi) \mid \phi \in \bigcup_{k=0}^i AltFORM_k\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i AltFORM_k\} \\ &\supseteq \bigcup_{k=0}^i AltFORM_k \cup \{(\neg \phi) \mid \phi \in AltFORM_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i AltFORM_k\} \end{aligned}$$

$$\bigcup_{k=0}^{i+1} AltFORM_k \subseteq FORM_{i+1}$$

Now we prove the other way:

$$\forall \phi \in FORM_{i+1}, \phi \in \bigcup_{k=0}^i AltFORM_k \text{ or } \phi = (\neg \psi), \psi \in \bigcup_{k=0}^i AltFORM_k \text{ or } \phi = (\psi_1 \vee \psi_2), \psi_1, \psi_2 \in \bigcup_{k=0}^i AltFORM_k$$

Case 1:

$$\phi \in \bigcup_{k=0}^i AltFORM_k \rightarrow \phi \in \bigcup_{k=0}^{i+1} AltFORM_k$$

Case 2:

$$\begin{aligned}
\psi &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j \leq i, \psi \in \text{AltFORM}_j \\
\phi &= (\neg\psi), \psi \in \text{AltFORM}_j, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

Case 3:

$$\begin{aligned}
\psi_1, \psi_2 &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j, l \leq i, \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l \\
\phi &= (\psi_1 \vee \psi_2), \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k \\
\text{FORM}_{i+1} &\subseteq \bigcup_{k=0}^{i+1} \text{AltFORM}_k \rightarrow \text{FORM}_{i+1} = \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

□

Formation Sequences

Definition 3.16 (Formation Sequence). A (P, C) -formation sequence is a nonempty finite sequence $\rho = w_1, \dots, w_k$ such that for each i :

- $w_i \in P$, or
- $w_i = (\neg w_j)$ for some $j < i$, or
- $w_i = (w_j \vee w_k)$ for some $j, k < i$.

The sequence belongs to w if $w_k = w$.

Q9

Give two distinct formation sequences that belong to

$$w = ((\neg p_1) \vee ((\neg p_3) \vee (\neg(p_2 \vee (\neg p_1)))))$$

Two distinct formation sequences ρ_1, ρ_2 are:

ρ_1 :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= (\neg w_1) \\
w_3 &= p_2 \\
w_4 &= (w_3 \vee w_2) \\
w_5 &= (\neg w_4) \\
w_6 &= (p_3) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_7 \vee w_5) \\
w_9 &= (w_2 \vee w_8)
\end{aligned}$$

ρ_2 :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= p_2 \\
w_3 &= p_3 \\
w_4 &= (\neg w_3) \\
w_5 &= (\neg w_1) \\
w_6 &= (w_2 \vee w_5) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_4 \vee w_7) \\
w_9 &= (w_5 \vee w_8)
\end{aligned}$$

These are distinct because the components at index $i = 2$ differ $((\neg p_1)$ vs p_2).

Lemma 3.17. $INDFORM = \{w \in \Sigma^* \mid w \text{ is formation-sequence-well-formed}\}$.

Q10

Prove the lemma by showing:

1. Every formation-sequence-well-formed string is in $INDFORM$.
2. Every $w \in INDFORM$ has a formation sequence.

Before attempting Q10 from the above definition of formation-sequence-well-formed, I will prove the following lemma using the definition below first:

Definition 3.18. Define $ExFORM_0 = P$

$$\begin{aligned}
ExFORM_{i+1} &= (\{(\neg w) \mid w \in ExFORM_i\} \\
&\cup \{(w_1 \vee w_2) \mid w_1 \in ExFORM_i, \exists j \leq i, w_2 \in ExFORM_j\} \\
&\cup \{(w_1 \vee w_2) \mid w_2 \in ExFORM_i, \exists j \leq i, w_1 \in ExFORM_j\} \\
&)\setminus \bigcup_{j=0}^i ExFORM_j
\end{aligned}$$

Lemma 3.19.

$$\bigcup_{j=0}^i ExFORM_j = FORM_i$$

Proof. Base case:

$$ExFORM_0 = P = FORM_0$$

Inductive Hypothesis: Suppose $\forall n \geq 0$,

$$\bigcup_{i=0}^n ExFORM_i = FORM_n$$

From definition 2.19,

$$\begin{aligned} FORM_{n+1} &= \bigcup_{i=0}^n ExFORM_i \cup \{(\neg w) \mid w \in \bigcup_{i=0}^n ExFORM_i\} \\ &\cup \{(w_1 \vee w_2) \mid w_1 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_2 \in \bigcup_{i=0}^n ExFORM_j\} \\ &\cup \{(w_1 \vee w_2) \mid w_2 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_1 \in \bigcup_{i=0}^n ExFORM_j\} \end{aligned}$$

Definition 3.20.

$$\begin{aligned} FromFORM : n &\mapsto \{(\neg w) \mid w \in \bigcup_{i=0}^n ExFORM_i\} \\ &\cup \{(w_1 \vee w_2) \mid w_1 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_2 \in \bigcup_{i=0}^n ExFORM_j\} \\ &\cup \{(w_1 \vee w_2) \mid w_2 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_1 \in \bigcup_{i=0}^n ExFORM_j\} \end{aligned}$$

Lemma 3.21.

$$FORM_{n+1} = \bigcup_{i=0}^n ExFORM_i \cup FromFORM(n)$$

Lemma 3.22.

$$\begin{aligned} ExFORM_{n+1} &= (\bigcup_{i=0}^n ExFORM_i \cup FromFORM(n)) \setminus \bigcup_{i=0}^n ExFORM_i \\ ExFORM_{n+1} &= FromFORM(n) \setminus \bigcup_{i=0}^n ExFORM_i \end{aligned}$$

From Lemma 3.21 and Lemma 3.22, we have

$$\begin{aligned} \bigcup_{i=0}^{n+1} ExFORM_i &= \bigcup_{i=0}^n ExFORM_i \cup ExFORM_{n+1} \\ &= \bigcup_{i=0}^n ExFORM_i \cup (FromFORM(n) \setminus \bigcup_{i=0}^n ExFORM_i) \\ &= \bigcup_{i=0}^n ExFORM_i \cup FromFORM(n) \\ &= FORM_{n+1} \end{aligned}$$

□

Lemma 3.23.

$$\bigcup_{i \geq 0} ExFORM_i = \bigcup_{i \geq 0} FORM_i$$

Proof. $\bigcup_{i \geq 0} ExFORM_i \subseteq \bigcup_{i \geq 0} FORM_i$

$\forall n \geq 0$, From Lemma 2.28

$$\phi \in \bigcup_{i=0}^n ExFORM_i \rightarrow \phi \in FORM_n \rightarrow \phi \in \bigcup_{i=0}^n FORM_i$$

$$\bigcup_{i \geq 0} ExFORM_i \supseteq \bigcup_{i \geq 0} FORM_i$$

Corollary from Lemma 2.28

□

Corollary 3.24.

$$\bigcup_{i \geq 0} ExFORM_i = INDFORM$$

$$\forall \phi \in ExFORM_i \rightarrow \phi \text{ is well-formed formula}$$

10.1, 2

Claim:

$$FORMSeq = \{w \in \Sigma^* \mid w \text{ is formation-sequence-well-formed}\} = INDFORM$$

Proof. Let ρ_i denotes the set of ρ that contains a sequence of i strings $w_1, w_2, \dots, w_i$

Define

$$\begin{aligned} str : \rho &\mapsto w, str(\rho = w_1, w_2, \dots, w_k) = w_k \\ str : \rho_i &\mapsto \{w\}, str(\rho = w_1, w_2, \dots, w_k) = \{w \mid w = str(\rho), \rho \in \rho_i\} \end{aligned}$$

Claim:

$$\bigcup_{k \geq 1} str(\rho_k) = \bigcup_{i \geq 0} ExFORM_i$$

Proof. $\bigcup_{k \geq 1} \rho_k \subseteq \bigcup_{i \geq 0} ExFORM_i$

$\forall w \in FORMSeq, \exists \rho = w_1, w_2, \dots, w_k, w = w_k$

Base case:

$$\bigcup_{k=1}^1 str(\rho_1) = \{w \mid w \in P\} = \bigcup_{i=0}^0 ExFORM_i$$

Inductive hypothesis: Suppose

$$\bigcup_{j=1}^{k+1} str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$$

$$\bigcup_{j=1}^{k+2} str(\rho_j) = \bigcup_{j=1}^{k+1} str(\rho_j) \cup \{w \mid \exists \rho \in \rho_{k+2}, w = str(\rho)\}$$

By definition, $w_{k+2} = str(\rho)$ for some $\rho \in \rho_{k+2}$, one of the 3 below holds:

1. $w_{k+2} \in P \rightarrow w_{k+2} \in \bigcup_{i \geq 0} ExFORM_i$
2. $w_{k+2} = (\neg w_n), n < k+2 \rightarrow w_n \in \bigcup_{j=1}^n str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$
3. $w_{k+2} = (w_m \vee w_n), m, n < k+2 \rightarrow w_m, w_n \in \bigcup_{j=1}^{\max(m,n)} str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$

$$\bigcup_{k \geq 1} \rho_k \supseteq \bigcup_{i \geq 0} ExFORM_i$$

Base case is the same as above.

Inductive hypothesis: (this will be used in Q11)

Lemma 3.25.

$$\bigcup_{i=0}^k ExFORM_i \subseteq \bigcup_{j=1}^{2(k+1)} str(\rho_j)$$

$\forall w \in ExFORM_{k+1}$, one of the 3 below holds: 1.

$$w \in P \rightarrow w \in ExFORM_0 \rightarrow w \in \bigcup_{i=0}^k ExFORM_i \subseteq \bigcup_{j=1}^{k+1} str(\rho_j)$$

2.

$$\begin{aligned} w = (\neg w_n), w_n \in ExFORM_k &\rightarrow \exists \rho = w_1, w_2, \dots, w_k \in \bigcup_{j=1}^{k+1} \rho_j, w_n = str(\rho) \\ &\rightarrow \exists \rho' \in \rho_{k+2}, \rho' = w_1, w_2, \dots, w_k, (\neg w_n) \\ &\rightarrow w \in str(\rho_{k+2}) \rightarrow w \in \bigcup_{j=1}^{k+2} str(\rho_j) \\ &\rightarrow w \in \bigcup_{j=1}^{2(k+1)} str(\rho_j) \end{aligned}$$

3. (Concat finite sequences that yield formula composing disjunctions gives finite sequence)

$$\begin{aligned} w = (w_m \vee w_n), WLOG, w_m \in ExFORM_k, \exists q \leq k, w_n \in ExFORM_q \\ &\rightarrow \exists \rho^m \in \rho_{k+1} = w_1^m, w_2^m, \dots, w_k^m, \\ &\quad \rho^n \in \rho_{q+1} = w_1^n, w_2^n, \dots, w_q^n \in \bigcup_{j=1}^{k+1} \rho_j, w_m = str(\rho^m), w_n = str(\rho^n) \\ &\rightarrow \exists \rho' = w_1^m, w_2^m, \dots, w_k^m, w_1^n, w_2^n, \dots, w_q^n, (w_k^m \vee w_q^n) \in \rho_{k+q+2} \\ &\rightarrow w \in \bigcup_{j=1}^{k+q+1} str(\rho_j) \\ &\rightarrow w \in \bigcup_{j=1}^{2(k+1)} str(\rho_j) \end{aligned}$$

Therefore,

$$\bigcup_{k \geq 1} str(\rho_k) = \bigcup_{i \geq 0} ExFORM_i$$

□

By Corollary 3.24,

$$\bigcup_{k \geq 1} \text{str}(\rho_k) = \text{INDFORM}$$

□

Q11

Lemma 3.26. *Every well-formed formula ϕ is of one of the following forms:*

1. $\phi \in P$,
2. $\phi = (\neg\psi)$,
3. $\phi = (\psi_1 \vee \psi_2)$.

Proof. From Corollary 3.24,

$$\forall \phi \in \text{INDFORM}, \exists k \geq 0, \phi \in \text{ExFORM}_k$$

Case $x = 0$:

$$x = 0 \rightarrow \phi \in \text{ExFORM}_0 = P$$

Case $x > 0$:

$$\begin{aligned} &\phi \in \text{ExFORM}_x \text{ and } \forall y \in [0, x), \phi \notin \text{ExFORM}_y \\ &\forall \phi \in \text{ExFORM}_x, \phi \in P \text{ or} \\ &\phi \in (\{(\neg\psi) \mid \psi \in \text{ExFORM}_{x-1}\} \\ &\quad \cup \{(\psi_1 \vee \psi_2) \mid \psi_1 \in \text{ExFORM}_{x-1}, \exists j \leq x-1, \psi_2 \in \text{ExFORM}_j\}) \end{aligned}$$

Since we define ExFORM_i to be pair wise disjoint, we guarantee that

$$\phi \in \text{ExFORM}_k \iff k \text{ is the minimum depth needed to form } \phi \text{ in the syntax tree}$$

Intuitively, x is the minimum depth required to formulate ϕ , thus with depth $d \in [0, x)$, ϕ cannot be formed. Suppose $\phi \in P$,

$$\phi \in \text{ExFORM}_0, \phi \in \text{ExFORM}_x, x > 0$$

Contradiction. Therefore

$$\phi \in (\{(\neg\psi) \mid \psi \in \text{ExFORM}_{x-1}\} \cup \{(\psi_1 \vee \psi_2) \mid \psi_1 \in \text{ExFORM}_{x-1}, \exists j \leq x-1, \psi_2 \in \text{ExFORM}_j\})$$

Now we will prove $\forall \phi \in \text{INDFORM}$

$$(\phi \in \{(\neg\psi) \mid \psi \in \text{ExFORM}_{x-1}\}) \oplus (\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in \text{ExFORM}_{x-1}, \exists j \leq x-1, \psi_2 \in \text{ExFORM}_j\})$$

Suppose we enforce the use of parentheses for all subformula, that is except $\phi \in P$, all formula starts with '('. If for some i , $\phi \in \{(\neg\psi) \mid \psi \in \text{ExFORM}_i\}$, ' $\neg$ ' must be at index 1. Likewise, if for some i , $\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in \text{ExFORM}_i, \exists j \leq i, \psi_2 \in \text{ExFORM}_j\}$, ' $\vee$ ' must not be at index 1. Thus $\nexists \phi \in \text{INDFORM}, \phi \in \{(\neg\psi) \mid \psi \in \text{ExFORM}_{x-1}\}$ and $\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in \text{ExFORM}_i, \exists j \leq i, \psi_2 \in \text{ExFORM}_j\}$ □

Lemma 3.27 (Characterization of Well-Formed Formulae). *Let P be a set of propositions and $C = \{\neg, \vee\}$. Every well-formed formula ϕ over (P, C) is of exactly one of the following forms:*

1. $\phi \in P$,
2. $\phi = (\neg\psi)$ for some well-formed formula ψ ,
3. $\phi = (\psi_1 \vee \psi_2)$ for some well-formed formulae ψ_1, ψ_2 .

Structural Induction Principle. A property $P(\phi)$ holds for all well-formed formulae if:

- $P(p)$ holds for all $p \in P$,
- $P(\psi)$ implies $P(\neg\psi)$,
- $P(\psi_1)$ and $P(\psi_2)$ imply $P(\psi_1 \vee \psi_2)$.

Q1

1.1

Theorem 3.28. *Let α be a well-formed formula. Let $c(\alpha)$ denote the number of occurrences of binary connective symbols ($\wedge, \vee, \rightarrow, \leftrightarrow$) in α , and let $s(\alpha)$ denote the number of occurrences of sentence symbols in α . Then*

$$s(\alpha) = c(\alpha) + 1.$$

1.2

Q2

Recall the following derived connectives:

- $\phi_1 \wedge \phi_2 \equiv \neg((\neg\phi_1) \vee (\neg\phi_2))$,
- $\perp \equiv \neg(p \vee \neg p)$ for some $p \in P$,
- $\phi_1 \rightarrow \phi_2 \equiv (\neg\phi_1) \vee \phi_2$.

Using structural induction, show that for every formula ϕ over $(P, \{\neg, \vee\})$ with $P \neq \emptyset$, there exists a formula ψ using only connectives $\{\perp, \rightarrow\}$ such that ϕ and ψ are logically equivalent.

Q3

Definition 3.29 (Conjunctive Normal Form). *A formula ϕ over $(P, \{\neg, \vee, \wedge\})$ is in conjunctive normal form (CNF) if*

$$\phi = C_1 \wedge C_2 \wedge \cdots \wedge C_m \quad (m \geq 1),$$

where each clause C_i has the form

$$C_i = \ell_{i,1} \vee \ell_{i,2} \vee \cdots \vee \ell_{i,k_i},$$

and each literal $\ell_{i,j}$ is either p or $\neg p$ for some $p \in P$.

Using structural induction, show that for every well-formed formula ϕ , there exists a logically equivalent formula ψ in CNF.

Q4

Definition 3.30 (Occurs). *Define $\text{occurs} : \text{FORM} \rightarrow \mathcal{P}(P)$ inductively by:*

[leftmargin=2em]

- If $\phi = p \in P$, then $\text{occurs}(\phi) = \{p\}$,
- If $\phi = \neg\psi$, then $\text{occurs}(\phi) = \text{occurs}(\psi)$,
- If $\phi = \psi_1 \oplus \psi_2$ with $\oplus \in \{\wedge, \vee\}$, then $\text{occurs}(\phi) = \text{occurs}(\psi_1) \cup \text{occurs}(\psi_2)$.

Lemma 3.31 (Relevance Lemma). *Let ϕ be a propositional formula and let $v_1, v_2 : P \rightarrow \{\text{true}, \text{false}\}$ be valuations such that $v_1(p) = v_2(p)$ for all $p \in \text{occurs}(\phi)$. Then $v_1 \models \phi$ if and only if $v_2 \models \phi$.*

Prove this lemma using structural induction.

Q5

Lemma 3.32 (Generalized Relevance Lemma). *Let ϕ be a propositional formula and let v_1, v_2 be valuations such that $v_1(p) = v_2(p)$ for all p in some set $S \supseteq \text{occurs}(\phi)$. Then $v_1 \models \phi$ if and only if $v_2 \models \phi$.*

Prove this lemma.

Q6

Lemma 3.33 (Relevance Lemma for Logical Equivalence). *Let ϕ_1, ϕ_2 be propositional formulae. The following are equivalent:*

[leftmargin=2em]

- *For all valuations v_1, v_2 agreeing on $\text{occurs}(\phi_1) \cup \text{occurs}(\phi_2)$, we have $v_1 \models \phi_1$ iff $v_2 \models \phi_2$.*
- *ϕ_1 and ϕ_2 are logically equivalent.*

Prove this lemma.

Q7

Let $G = (V, E)$ be an undirected graph and $k \in \mathbb{N}$. Construct a propositional formula ϕ_G such that:

1. $|\phi_G|$ is polynomial in $k + |V| + |E|$,
2. ϕ_G is satisfiable if and only if G has a vertex cover of size at most k .

Formally prove correctness of the encoding.

Q8

Given a partial $N^2 \times N^2$ Sudoku grid S , construct a propositional formula ϕ_S of size polynomial in N^2 such that ϕ_S is satisfiable if and only if S has a completion to a valid Sudoku grid. Prove correctness.

Q9

(Extra Credit) Encode the Independent Set problem as propositional logic.

1. Construct a formula $\phi_{G,k}$ satisfiable iff G has an independent set of size k .
2. Write $\phi_{G,3}$ explicitly for a given graph on five vertices.

Q10

(Extra Credit) Implement a solver for Independent Set using a SAT solver as an oracle. The program should read a graph and k , construct $\phi_{G,k}$, and determine satisfiability.

Background: Compactness

Definition 3.34 (Satisfiable). *A set Γ of propositional formulae is satisfiable if there exists a valuation v such that $v \models \Gamma$.*

Definition 3.35 (Finitely Satisfiable). *A set Γ is finitely satisfiable if every finite $\Gamma' \subseteq \Gamma$ is satisfiable.*

Theorem 3.36 (Compactness Theorem). *A set Γ of propositional formulae is satisfiable if and only if it is finitely satisfiable.*

Q11

Prove that if Γ is satisfiable, then Γ is finitely satisfiable.

Q12

Definition 3.37 (P -complete). *Γ is P -complete if for every $p \in P$, either $p \in \Gamma$ or $\neg p \in \Gamma$.*

Definition 3.38 (P -consistent). *Γ is P -consistent if for no $p \in P$ do both p and $\neg p$ belong to Γ .*

Lemma 3.39. *If Γ is finitely satisfiable and P -complete, then Γ is P -consistent.*

1. Prove the lemma.
2. Give examples as requested in the handout.

Q13

Let Γ be finitely satisfiable and let ϕ be a formula.

1. Prove that at least one of $\Gamma \cup \{\phi\}$ or $\Gamma \cup \{\neg\phi\}$ is finitely satisfiable.
2. Give examples as requested.

Q14

Let $P = \{p_1, p_2, \dots\}$ and define $\Gamma_0, \Gamma_1, \dots$ as in the handout. Prove that $\Delta = \bigcup_{i \geq 0} \Gamma_i$ is finitely satisfiable.

Q15

Prove that if Γ is finitely satisfiable, then Γ is satisfiable.

Q16

Solve Questions 10 and 11 on page 53 (end of Section 1.5) in *A Mathematical Introduction to Logic* by Herbert B. Enderton.