

# Week 2-4: Propositional Logic (HW2 start from page 29)

Wang Xiyu A0282500R

## Contents

|          |                                                               |           |
|----------|---------------------------------------------------------------|-----------|
| <b>1</b> | <b>Syntax</b>                                                 | <b>2</b>  |
| 1.1      | Well-formed Formula (WFF)                                     | 2         |
| 1.2      | Closed Set                                                    | 2         |
| 1.2.1    | All Closed Set                                                | 2         |
| 1.3      | FORM is the minimum of the set of Closed                      | 3         |
| 1.4      | Subformula                                                    | 3         |
| <b>2</b> | <b>Semantics</b>                                              | <b>4</b>  |
| 2.1      | Truth Assignment/Proposition evaluation                       | 4         |
| 2.2      | Entail/Model                                                  | 4         |
| 2.2.1    | Valuation and Model                                           | 4         |
| 2.2.2    | $\models$ as a relation                                       | 5         |
| 2.2.3    | Logical Equivalence                                           | 5         |
| 2.3      | Satisfiability of Formula                                     | 5         |
| 2.3.1    | Valid formula(tautology)                                      | 6         |
| 2.4      | Satisfiability of a set of formula                            | 6         |
| 2.4.1    | Relevance                                                     | 6         |
| 2.4.2    | Logical consequence                                           | 7         |
| 2.5      | Modelling Computations as Propositional Logic                 | 7         |
| <b>3</b> | <b>Compactness Theorem</b>                                    | <b>9</b>  |
| 3.1      | Application of the Compactness theorem for proposition logics | 12        |
| <b>4</b> | <b>Proof System</b>                                           | <b>14</b> |
| 4.1      | Hilbert Style Proof System                                    | 14        |
| 4.2      | Metatheorems                                                  | 15        |
| 4.2.1    | Monotonicity theorem                                          | 15        |
| 4.2.2    | Deduction                                                     | 16        |
| 4.3      | Resolution Proof System                                       | 17        |
| <b>5</b> | <b>Complexity</b>                                             | <b>18</b> |
| 5.1      | Turing Machine                                                | 18        |
| 5.2      | Semantics of TM                                               | 19        |
| 5.3      | Running time and Space                                        | 21        |
| 5.4      | Complexity classes                                            | 22        |
| 5.5      | Hardness                                                      | 22        |
| 5.6      | Reduction                                                     | 22        |

# 1 Syntax

**Definition 1.1.** *DSyntax decides what are allowed to be written*

## 1.1 Well-formed Formula (WFF)

A formula  $\psi$  is obtained by composing  $p \in P$  with  $c \in C$ , where  $P$  is the set of propositions and  $C$  is the set of connectors. here we use any set of complete logic, like  $C = \{\neg, \vee\}$ . This is not a precise definition as counter examples that do not form a valid formula can be formulated based on this, for example  $\neg \vee p$  means nothing. To formalize the definition we may attempt to define a language with the following grammar:

$$\Sigma = P \cup C \cup \{(\cdot, \cdot)\}$$

$$FORM : \{\psi = p \in P \mid \neg\psi \mid \psi \vee \psi \mid (\psi)\}$$

However this is a recursive definition as the definition of the a formula  $\psi$  involves itself. We need to find a declarative way to define this recursive/inductive construction.

**Definition 1.2.** *FORM is **the smallest** set of strings over  $\Sigma$  where*

1.  $P \subseteq FORM$
2.  $\forall \psi \in FORM, \neg\psi \in FORM$
3.  $\forall \psi_1, \psi_2 \in FORM, \psi_1 \vee \psi_2 \in FORM$

We can see as a construction rule, we are admitting strings into the set of formula, but we need to somewhat impose restriction on what is not acceptable.

## 1.2 Closed Set

Let's remove the attributive **the smallest** first and define another set that contains all the set that satisfy these rules listed above:

**Definition 1.3.** *Closed is a set of strings over  $\Sigma$  where*

1.  $P \subseteq Closed$
2.  $\forall \psi \in Closed, \neg\psi \in Closed$
3.  $\forall \psi_1, \psi_2 \in Closed, \psi_1 \vee \psi_2 \in Closed$

### 1.2.1 All Closed Set

And we define a set ACS (All-closed sets). We can define FORM via the following construction:

$$FORM = \bigcap ACS$$

Suppose  $BCS = \emptyset$  and from definition

$$U_0 = \bigcap BCS = \{w \in \Sigma^* \mid \forall S \in BCS, w \in S\}$$

However,  $BCS = \emptyset$ , The statement  $\forall S \in BCS, w \in S$  is vacuously true  $\forall w \in \Sigma^*$ , which means that

$$U_0 = \Sigma^* = \bigcap BCS$$

A contradiction. Therefore we have proved that ACS cannot be empty, thus  $\forall P, \exists S$  such that  $S$  is a closed set.

### 1.3 FORM is the minimum of the set of Closed

First let's prove that FORM is closed. It seems to be trivial that the intersection of closed sets is closed but formal proof is still needed. We denote a set from the intersection construction as  $T_0$ .

**Lemma 1.4.**  $T_0$  is closed

We are proving that the set  $T_0$  we get from the intersection construction still satisfy the rules that define what is closed.

*Proof.* From rule 1,

$$\begin{aligned} \forall S \in ACS, P \subseteq S \\ \rightarrow \forall p \in P, \forall S \in ACS, p \in S \\ \rightarrow \forall p \in P, p \in \bigcap ACS \\ \rightarrow P \in T_0 \end{aligned}$$

From rule 2,

$$\begin{aligned} \forall S \in ACS, T_0 \subseteq S \\ \text{let } w \in T_0, \forall S \in ACS, w \in S, \neg w \in S \\ \rightarrow \forall w \in T_0, \neg w \in \bigcap ACS \\ \rightarrow \forall w \in T_0, \neg w \in T_0 \end{aligned}$$

From rule 3,

$$\begin{aligned} \forall S \in ACS, T_0 \subseteq S \\ \text{let } w_1, w_2 \in T_0, \forall S \in ACS, w_1 \vee w_2 \in S \\ \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in \bigcap ACS \\ \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in T_0 \end{aligned}$$

□

**Lemma 1.5.**  $T_0$  is *a* smallest Closed set

*Proof.* The proof of minimality is trivial.  $T_0 = \bigcap ACS \rightarrow \forall S \in ACS, T_0 \subseteq S$

□

**Lemma 1.6.**  $T_0$  is *the* smallest Closed set (unique). In other word,

$$\forall T'_0 \in \{T \mid \forall S \in ACS, T \subseteq S\}, T'_0 = T_0$$

Now we can use the generative rule to define formula, which is called **Backus-Naur form**. Note that a context-free grammar is a type of formal language. Backus Naur form is a specification language for this type of grammar. It is used to describe language syntax.

### 1.4 Subformula

**Definition 1.7.**  $\phi_1$  is a subformula of  $\phi_2$ , denoted by

$$\phi_1 \preceq \phi_2$$

as a relation is

- reflexive:  $\forall \phi \in FORM, \phi \preceq \phi$

- *transitive*:  $\forall \phi_1, \phi_2, \phi_3 \in FORM, (\phi_1 \preceq \phi_2) \wedge (\phi_2 \preceq \phi_3) \rightarrow \phi_1 \preceq \phi_3$

*Inductive definition:*

$$\begin{aligned} \forall \phi \in FORM, \phi &\preceq (\neg \phi) \\ \forall \phi_1, \phi_2 \in FORM, \phi_1 &\preceq (\phi_1 \vee \phi_2) \\ \phi_2 &\preceq (\phi_1 \vee \phi_2) \end{aligned}$$

We might intuitively think that subformula can be defined using the notion of substring, since we have been considering formula as concatenation of proposition and connector symbols. However based on our inductive definition, we can come up with the following counter example to a definition using substring: A formula  $\phi_1$  is a subformula of  $\phi_2$  if and only if  $\phi_1$  is a well-formed formula, and is a substring to  $\phi_2$

$$p \vee q \not\preceq \neg p \vee q$$

What if we tighten the syntax by strictly enforce the use of parentheses whenever we can

## 2 Semantics

**Definition 2.1.** *Semantics describe meanings.*

### 2.1 Truth Assignment/Proposition evaluation

A truth assignment or a valuation of propositions is a function that maps from the set of propositions to true and false.

$$\begin{aligned} v : P &\rightarrow \{T, F\} \\ v &\in [P \rightarrow \{T, F\}] \end{aligned}$$

**Example 2.2.** *Consider  $v = \{p_1 \mapsto T, p_2 \mapsto F\}$ , we can derive*

$$\begin{aligned} v(p_1) &= T \\ v(p_2) &= F \\ v(\neg p_2) &= T \\ v(p_1 \vee p_2) &= T \text{ etc.} \end{aligned}$$

### 2.2 Entail/Model

We denote entailment of an evaluation function to formula as

$$v \models \psi$$

We also inductively define the following rule

$$\begin{aligned} \forall p \in P, v \models p &\iff v(p) \\ v \models \psi &\iff v \not\models \neg \psi \\ v \models \psi_1 \vee \psi_2 &\iff (v \models \psi_1) \vee (v \models \psi_2) \end{aligned}$$

#### 2.2.1 Valuation and Model

A valuation  $v$  is a model of a formula  $\phi$ , if and only if  $v(\phi) = T$ .

- $\forall \phi$  over  $P$ ,  $\exists 2^{|P|}$  valuations.
- A formula can have many or one or none models

### 2.2.2 $\models$ as a relation

**Definition 2.3.**  $\models \subseteq \{2^{|P|} \times FORM\}$  is a binary relation, between truth assignment to formula. We define as follows:

- $\forall p \in P, v \models p \iff v(p) = 1$
- $v \not\models p$  otherwise.

Consider a valuation  $v$  as a subset of  $P$  such that the propositions in  $v$  denoted by  $v(p)$  is assigned with true.  $v \in \mathcal{P}(P)$

| $\phi$ | $\psi$ | $\neg\phi$ | $(\phi \wedge \psi)$ | $(\phi \vee \psi)$ | $(\phi \rightarrow \psi)$ | $(\phi \leftrightarrow \psi)$ |
|--------|--------|------------|----------------------|--------------------|---------------------------|-------------------------------|
| 0      | 0      | 1          | 0                    | 0                  | 1                         | 1                             |
| 0      | 1      | 1          | 0                    | 1                  | 1                         | 0                             |
| 1      | 0      | 0          | 0                    | 1                  | 0                         | 0                             |
| 1      | 1      | 0          | 1                    | 1                  | 1                         | 1                             |

**Definition 2.4.**

$$p(v) = v(p)$$

A proposition can be viewed as a circuit is just a wire, thus if a proposition is assigned with true by a valuation, the valuation evaluates this proposition to be true

**Lemma 2.5.** Let  $\phi \in FORM, v \in \mathcal{P}(P), v \models \phi \iff \phi(v) = 1$

*Proof.* Base case:  $\forall \phi = p \in P, v \models \phi \iff v(p) = 1 \iff p(v) = 1$

Inductive case:

1.

□

### 2.2.3 Logical Equivalence

**Definition 2.6.** Syntax definition of logical equivalence is as follows:

$$\phi_1 \equiv \phi_2 \iff \forall v : P \mapsto \{T, F\} (v \models \phi_1), v \models \phi_2$$

**Example 2.7.**

$$p \vee (\neg p) \equiv q \vee (\neg q)$$

$$\begin{aligned} \forall v \models (p \vee (\neg p)), \\ v \models p \text{ or } v \models (\neg p) \\ v \models p \text{ or } v \not\models p \\ \forall v : P \mapsto \{T, F\}, \end{aligned}$$

*TODO*

## 2.3 Satisfiability of Formula

**Definition 2.8.** Let  $\phi \in FORM, \phi$  is satisfiable

$$\iff \exists v : P \mapsto \{T, F\}, v \models \phi$$

### 2.3.1 Valid formula(tautology)

**Definition 2.9.** *Tautology:  $\phi$  is a tautology  $\iff \forall v, v \models \phi$*

**Example 2.10.**

$$\phi = \neg(p \vee (\neg p))$$

From example 2.4, we have

$$\begin{aligned} \forall v, v \models (p \vee (\neg p)) \\ \rightarrow \forall v, v \not\models \neg(p \vee (\neg p)) \end{aligned}$$

Thus  $\phi$  is NOT satisfiable.

## 2.4 Satisfiability of a set of formula

### Set Satisfaction

**Definition 2.11.** *Let  $\Gamma \subseteq FORM$ ,*

$$v \models \Gamma \iff \forall \phi \in \Gamma, v \models \phi$$

### Set Satisfiability

**Definition 2.12.** *Let  $\Gamma \subseteq FORM$ ,  $\Gamma$  is satisfiable*

$$\iff \exists v, \forall \phi \in \Gamma, v \models \phi$$

**Example 2.13.**  $\Gamma = \emptyset$ ,  $\Gamma$  is satisfiable.

*Proof.* Consider  $v : P \mapsto \{T, F\}$ ,  $v(p) = T \forall p \in P$ ,  $\forall \phi \in \Gamma, v \models \phi$  is vacuously true.  $\square$

**Example 2.14.**  $\Gamma = FORM$ ,  $\Gamma$  is NOT satisfiable.

*Proof.* Suppose otherwise,

$$\begin{aligned} \exists v, \forall \phi \in \Gamma, v \models \phi \\ \forall \phi \in \Gamma, (\neg \phi) \in \Gamma \\ v \models (\neg \phi) \end{aligned}$$

A contradiction.  $\square$

### 2.4.1 Relevance

**Definition 2.15.** *Define set  $OCCUR(\phi) \subseteq P$ , which contains propositions appear in  $\phi$ . Inductively defined as*

$$\begin{aligned} OCCUR(\phi) = \{p\} &\iff \phi \equiv p, p \in P \\ OCCUR(\neg \phi) &= OCCUR(\phi) \\ OCCUR(\phi_1 \vee \phi_2) &= OCCUR(\phi_1) \cup OCCUR(\phi_2) \end{aligned}$$

**Lemma 2.16.** *Let  $\phi \in FORM$ , let  $v_1, v_2$  be valuations over  $P$ .*

$$\forall p \in OCCUR(\phi), v_1(p) = v_2(p) \rightarrow (v_1 \models \phi \rightarrow v_2 \models \phi)$$

$v_1, v_2$  can be different valuations over  $p' \notin OCCUR(\phi)$

*Proof.* Let  $p \in P$ ,  $\square$

### 2.4.2 Logical consequence

Semantic logical consequence:

**Definition 2.17.**

$$\Gamma \models \phi \iff \forall v \models \Gamma, v \models \phi$$

Syntactic logical consequence: (logical derivability)

**Definition 2.18.** A formula  $\phi$  is a syntactic consequence of  $\Gamma$  if there exists a formal proof (a finite sequence of formulas) ending in  $\phi$ , where each step is either a premise from  $\Gamma$ , an axiom, or follows from previous steps via an inference rule (like Modus Ponens).

**Material Implication ( $\rightarrow$ ) vs. Logical Consequence ( $\models$ )**

- Material implication is a connector within a formula,  $p \rightarrow q \equiv \neg(p \vee q)$ , and can therefore be eval to true or false in a single model.
- Logical consequence is a relation between formulas (meta-logic), it is a statement about all models.  $\{p, p \rightarrow q\} \models q$

## 2.5 Modelling Computations as Propositional Logic

Computation problems can be considered as membership check of an encoded problem instance to a language.

$$L = \{p \mid \langle M, p \rangle\}$$

Here instead of arbitrary encoding, we consider problem encoding with propositional logics only, and solve with SAT solvers by deciding whether the set of formula we form from any instance of the problem is satisfiable. That is

$$Encode : p \mapsto FORM, \forall p, \exists v \models Encode(p) \iff p \in L$$

The rules of such modelling include:

- Propositional logics only
- The final expression must one well formed fomula
- for problem of size  $n$ , the size of the formula  $\phi$  must be in  $O(poly(n))$

Note that there can be different definition on the size of the

- problem: in unary, binary etc;
- Formula: Whether by the number of nodes in the syntax tree or length of the string  
**Anything else?**

**Example 2.19.** Graph coloring problem: given a undirected graph  $G = (V, E)$ , whether there exists a  $k$  coloring of all vertices such that no neighbouring vertices have the same color.

Consider the following construction of a proposition set  $P$ :

$$P = \{p_{v,i} \mid v \in V, i \in [1, k]\}$$

The valuation of proposition  $p_{v,i}$  conceptually represents vertex  $v$  is assigned color  $i$ .

We can formulate constriants as well formed formula, and use conjunction to connect the formula to model satisfaction of multiple constraints.

**Corollary 2.20.**  $\forall \phi_1, \phi_2 \in FORM, (\phi_1 \wedge \pi_2) \in FORM$

First let us list down the constraints of this problems, including implicit ones in plain words

1. Neighbouring vertices have different colors.
2. All vertices as assigned with exactly one color.

**1.**  $\phi_{nbr}$

First lets formula the constraint between any 2 vertices,

$$\forall(\alpha, \beta) \in E, c(v_\alpha) \neq c(v_\beta)$$

And from our proposition set,  $c(v_\gamma) = i \iff \exists v, v \models p_{\gamma,i}$ . As such we have

$$\phi_{\alpha,\beta,i} = \neg(p_{\alpha,i} \wedge p_{\beta,i})$$

This models vertex  $\alpha$  and vertex  $\beta$  are not colored with color  $i$  at the same time. For every color this needs to hold, thus we have

$$\phi_{nbr(\alpha,\beta)} = \bigwedge_{i \in [1,k]} \phi_{\alpha,\beta,i}$$

For every pair of neighbouring vertices this also needs to hold. Therefore we have

$$\phi_{nbr} = \bigwedge_{(\alpha,\beta) \in E} \phi_{nbr(\alpha,\beta)}$$

**2.**  $\phi_{exactly-1}$

To model exactly 1 coloring on each vertex, we can formulate  $\phi_{\geq 1}$  and  $\phi_{\leq 1}$  first and use conjunction to connect them. Given our propositions, to make sure vertex  $v$  is at least assign a color ( $\phi_{v,\geq 1}$ ),

$$\phi_{v,\geq 1} = \bigvee_{i \in [1,k]} p_{v,i}$$

This needs to hold for every vertex, thus we have

$$\phi_{\geq 1} = \bigwedge_{v \in V} \phi_{v,\geq 1}$$

Then we need to model at most 1, (or less than 2) coloring on each vertex. Vertex  $v$  is not assignment with both color  $i$  and  $j$  can be formulated as

$$\phi_{v,i,j,\leq 1} = \neg(p_{v,i} \wedge p_{v,j})$$

This needs to hold for every distinct pair of colors, thus

$$\phi_{v,\leq 1} = \bigwedge_{i \neq j \in [1,k]} \phi_{v,i,j,\leq 1}$$

For all vertices

$$\phi_{\leq 1} = \bigwedge_{v \in V} \phi_{v,\leq 1}$$

Finally we have

$$\phi_{exactly-1} = \phi_{\leq 1} \wedge \phi_{\geq 1}$$

And the overall formula for an instance of  $G = (V, E)$  becomes

$$\phi = \phi_{nbr} \wedge \phi_{exactly-1}$$

Similar to reduction, we need also now to show that

$$\exists v : P \mapsto \{T, F\}, v \models \phi \iff \langle G = (V, E), k \rangle \in L_{color}$$

TODO

### 3 Compactness Theorem

**Definition 3.1.** *Axiom of Choice: formally*

$$\forall X, \emptyset \in X \vee \exists f, \text{dom}(f) = X \wedge \forall A \in X, f(A) \in A$$

A choice function (or selection)  $f$  defined on a collection of nonempty sets  $X$ , such that for every set  $A$  in the collection  $X$ ,  $f(A)$  is an element in  $A$ . That is, Axiom—For any set  $X$  of nonempty sets, there exists a choice function  $f$  that is defined on  $X$  and maps each set of  $X$  to an element of that set.

**Theorem 3.2.** *If  $P$  is finite, FORM over  $P$  is finite.*


**Theorem 3.3.** *If  $P$  is countable, FORM over  $P$  is countable.*

*Proof.* Given  $\forall \phi \in \text{FORM}, \phi$  is finitely long, this theorem is another way of stating: The language formed from countable alphabet  $P \cup C$  with the inductive construction rules, is countable. We can prove a stronger theorem first.  $\square$


**Theorem 3.4.** *All languages formed from a countable alphabet is countable.*

*Proof.* Consider  $L = \Sigma^*$ , we will prove that  $\Sigma$  is countable  $\rightarrow L$  is countable. We use the same diagonal proof for the countability of  $\mathbb{Q}$  to show an enumeration of all strings formable from a countable alphabet. To formalize, we use induction to prove that  $\forall i \geq 0, \Sigma^i$  is


**Rational Numbers ( $\mathbb{Q}^+$ )**



**Countable Language ( $\Sigma^{n+1}$ )**



**Countable Language ( $\Sigma^2$ )**



**Countable Language ( $\Sigma^*$ )**




Figure 1: Comparison of diagonal enumeration for numbers vs. strings.

countable

Base case:  $\Sigma^1 = \Sigma$ , countable as defined.

Inductive hypothesis:  $\Sigma^n$  is countable: we enumerate all strings in  $\Sigma^n$ , and use the same diagonal proof to show the concatenation of one symbol from the countable alphabet to

every string  $\Sigma^{n+1}$  is also countable. Thus we can give an enumeration of all  $\Sigma^i$  and use the diagonal argument again to show the existence of an enumeration of  $\Sigma^*$ .

Alternatively, Given

$$\Sigma^* = \bigcup_{i \geq 0} \Sigma^i$$

From the Axiom of Choice, which implies the union of countably many countable set is countable,  $\Sigma^*$  is countable. Since  $\forall L$  over  $\Sigma, L \subseteq \Sigma^*, \Sigma^*$  is countable, all languages over countable  $\Sigma$  is countable.  $\square$

**Definition 3.5** (Finitely Satisfiable). *A set of formulas  $\Gamma$  is said to be **finitely satisfiable** if and only if for all finite subsets  $\Gamma_0 \subseteq \Gamma$ ,  $\Gamma_0$  is satisfiable.*

**Theorem 3.6** (Compactness Theorem).

$$\Gamma \text{ is satisfiable} \iff \Gamma \text{ is finitely satisfiable}$$

*Proof.* Consider a special case where  $\Gamma$  is “ $T$ -good”, defined as:

$$\forall p \in P, (p \in \Gamma \vee \neg p \in \Gamma)$$

Claim: If  $\Gamma$  is  $T$ -good, then  $\Gamma$  is finitely satisfiable  $\implies \Gamma$  is satisfiable.

Suppose  $\Gamma$  is  $T$ -good and finitely satisfiable. We define a candidate valuation  $v_L$  such that for all  $p \in P$ :

$$v_L(p) = \begin{cases} T & \text{if } p \in \Gamma \\ F & \text{if } \neg p \in \Gamma \end{cases}$$

First, we prove that for every  $p \in P$ , **exactly one** of  $\{p, \neg p\}$  is in  $\Gamma$ . By the  $T$ -good property, at least one is in  $\Gamma$ . Suppose both  $\{p, \neg p\} \subseteq \Gamma$ . Consider the finite subset:

$$\Gamma' = \{p, \neg p\}$$

Since  $\Gamma'$  is a finite subset of a finitely satisfiable set, it must be satisfiable. However,  $\neg \exists v$  such that  $v \models \{p, \neg p\}$ , which is a contradiction. Thus,  $v_L$  is well-defined.

Now we show  $v_L \models \Gamma$ . Suppose  $\exists \phi \in \Gamma$  such that  $v_L \not\models \phi$ . Define the following finite sets based on the atoms occurring in  $\phi$ :

$$\begin{aligned} \Gamma_\phi^+ &= \{p \mid p \in \text{OCCUR}(\phi) \wedge v_L(p) = T\} \\ \Gamma_\phi^- &= \{\neg p \mid p \in \text{OCCUR}(\phi) \wedge v_L(p) = F\} \\ \Gamma_\phi &= \{\phi\} \cup \Gamma_\phi^+ \cup \Gamma_\phi^- \end{aligned}$$

Since  $\Gamma_\phi$  is a finite subset of  $\Gamma$ , it must be satisfiable by some valuation  $v$ . For this  $v$ :

- $\forall p$  such that  $p \in \Gamma_\phi^+, v(p) = T$ .
- $\forall \neg p$  such that  $\neg p \in \Gamma_\phi^-, v(\neg p) = T \implies v(p) = F$ .

By construction,  $v$  and  $v_L$  agree on all atoms in  $\text{OCCUR}(\phi)$ . By the **Relevance Lemma**,  $v(\phi) = v_L(\phi)$ . Since  $v \models \Gamma_\phi$ , we have  $v \models \phi$ . This implies  $v_L \models \phi$ , contradicting our assumption  $v_L \not\models \phi$ . Therefore,  $\forall \phi \in \Gamma, v_L \models \phi$ , which implies  $v_L \models \Gamma$ ,  $\Gamma$  is satisfiable.

Now we will generalize this to all  $\Gamma$  which may not be ‘ $T$ -good’, by showing that we can construct a ‘ $T$ -good’ finitely satisfiable set  $\Delta$ , from any finitely satisfiable  $\Gamma$ , and therefore show that  $\Gamma SAT \iff \Delta SAT$ .

We will only prove the case for countable  $P$  now. Let an enumeration of  $p \in P$  be  $\{p_1, p_2, \dots\}$ . Define finite subset  $\Gamma_i$  of  $\Gamma$  as such:

$$\begin{aligned}\Gamma_0 &= \Gamma \\ \Gamma_{i+1} &= \begin{cases} \Gamma_i \cup \{p_i\} & \text{if } \Gamma_i \cup \{p_i\} \text{ is finitely satisfiable} \\ \Gamma_i \cup \{\neg(p_i)\} & \text{Otherwise} \end{cases}\end{aligned}$$

Here we need to prove that  $\forall i$ , between  $\Gamma_i \cup \{p_i\}$  and  $\Gamma_i \cup \{\neg(p_i)\}$  at least one case is also finitely satisfiable.

**Theorem 3.7.** *Let  $\Gamma$  be finitely satisfiable set, and  $\phi$  be a formula, then  $\Gamma \cup \{\phi\}$  is finitely satisfiable or  $\Gamma \cup \{\neg(\phi)\}$  is finitely satisfiable.*

*Proof.* We prove the contrapositive of this statement: if  $\Gamma \cup \{\phi\}$  and  $\Gamma \cup \{\neg(\phi)\}$  are both not finitely satisfiable,  $\Gamma$  is not finitely satisfiable. Given both are not finitely satisfiable, there exist finite subsets  $\Gamma_1, \Gamma_2 \subseteq \Gamma$  such that  $\Gamma_1 \cup \{\phi\}$  and  $\Gamma_2 \cup \{\neg(\phi)\}$  are not satisfiable. Therefore for all valuations, it must not model either  $\Gamma_1$  or  $\{\phi\}$ , and not model either  $\Gamma_2$  or  $\{\neg(\phi)\}$ . Denote the set of valuations that models formula  $\psi$  as  $models(\psi)$ , the set of valuations that models set of formula  $\Sigma$  as  $models(\Sigma)$

$$\begin{aligned}models(\Gamma_1) \cap models(\phi) &= \emptyset \\ &\rightarrow \forall v \in models(\Gamma_1), v \not\models \phi \rightarrow v \models (\neg\phi) \\ &\rightarrow models(\Gamma_1) \subseteq models((\neg\phi)) \\ models(\Gamma_2) \cap models((\neg\phi)) &= \emptyset \\ &\rightarrow \forall v \in models(\Gamma_2), v \not\models (\neg\phi) \rightarrow v \models (\phi) \\ &\rightarrow models(\Gamma_2) \subseteq models(\phi)\end{aligned}$$

Now consider the set of valuations that models  $\Gamma_1 \cup \Gamma_2$

$$\begin{aligned}models(\Gamma_1 \cup \Gamma_2) &= models(\Gamma_1) \cap models(\Gamma_2) \\ (models(\Gamma_1) \cap models(\Gamma_2)) &\subseteq (models(\phi) \cap models((\neg\phi))) = \emptyset\end{aligned}$$

Since  $\Gamma_1 \cup \Gamma_2$  is a finite subset of  $\Gamma$  that is not satisfiable,  $\Gamma$  is not finitely satisfiable.  $\square$

Now prove that

**Lemma 3.8.**  $\forall i, \Gamma_i$  is finitely satisfiable.

*Proof.* Base case:

$$\Gamma_0 = \Gamma \text{ is finitely satisfiable by definition}$$

Inductive hypothesis: Suppose

$$\Gamma_n \text{ is finitely satisfiable}$$

By Lemma 3.7,  $\Gamma_n \cup \{p_n\}$  and  $\Gamma_n \cup \{\neg(p_n)\}$  there must be at least one that is finitely satisfiable.  $\rightarrow \Gamma_{n+1}$  is therefore finitely satisfiable.  $\square$

Next we will show that

**Lemma 3.9.**  $\Gamma_U = \bigcup_{i \geq 0} \Gamma_i$  is finitely satisfiable.

*Proof.* Consider finite set  $\Gamma' \subseteq \Gamma_U$ , since  $\Gamma'$  is finite, and  $\forall k \geq 0, \Gamma_k \subseteq \Gamma_{k+1}$  by definition,  $\exists j \geq 0, \Gamma' \subseteq \Gamma_j = \bigcup_{i \geq 0}^j \Gamma_i$ . By Lemma 3.8,  $\Gamma_j$  is satisfiable  $\rightarrow \Gamma'$  is finitely satisfiable.  $\square$

**Lemma 3.10.**  $\Gamma_U$  is satisfiable.

*Proof.* From the construction,  $\forall i \geq 0$  at least one of  $p_i$  and  $(\neg p_i)$  is in  $\bigcup_{i \geq 0} \Gamma_i$ . Since  $\bigcup_{i \geq 0} \Gamma_i$  is finitely satisfiable,  $\forall i \geq 0, p_i$  and  $(\neg p_i)$  cannot both be in  $\bigcup_{i \geq 0} \Gamma_i$ . Otherwise  $\{p_i\} \cup \{(\neg p_1)\} \subseteq \bigcup_{i \geq 0} \Gamma_i$  is not satisfiable, contradiction.  $\square$

Now we have proven that the construction proposed from a finitely satisfiable formula set  $\Gamma$  to  $\bigcup_{i \geq 0} \Gamma_i$  is also finitely satisfiable. Now to complete the proof of the compactness theorem, we need to prove that  $\exists v, v \models \Gamma$ .

*Proof.* From Lemma 3.10,  $\forall i \geq 0, (p_i \in \Gamma_U) \oplus ((\neg p_i) \in \Gamma_U), \exists v, v \models \Gamma_U$

$$v(p) : P \mapsto \{T, F\} = \begin{cases} T, & \text{if } p \in \Gamma_U \\ F, & \text{if } (\neg p) \in \Gamma_U \end{cases}$$

Define the literal set  $Literals = \{p_i, (\neg p_i) \mid i \geq 0\}$ , and function

$$\tau(p) : P \mapsto Literals = \begin{cases} p, & \text{if } p \in \Gamma_U \\ (\neg p), & \text{if } (\neg p) \in \Gamma_U \end{cases}$$

**Lemma 3.11.**

$$v \models \Gamma_U \rightarrow v \models \tau(\Gamma)$$

Let  $\phi \in \Gamma$ , Define literal set  $X = \{\tau(p) \mid p \in OCCUR(\phi)\}$  where  $X$  is a set containing single proposition literals. By definition we have

$$X \subseteq \Gamma_U \rightarrow v \models X \rightarrow \forall \psi \in X, v(\psi) = T$$

and  $OCCUR(X) = OCCUR(\phi)$  Since both  $X$  is finite,  $X \cup \{\phi\}$  is a finite subset to  $\Gamma_U$ , thus satisfiable.  $\rightarrow \exists v', v' \models X \cup \{\phi\} \rightarrow v' \models X, v' \models \phi \rightarrow \forall \psi \in X, v'(\psi) = T$ .

$$\forall p \in OCCUR(\phi), v(p) = v'(p), v' \models \phi \rightarrow v \models \phi$$

Therefore, the valuation that models enlarged construction  $\Gamma_U$  also models  $\Gamma$ , thus  $\Gamma$  is satisfiable.  $\square$

$\square$

### 3.1 Application of the Compactness theorem for proposition logics

We have already shown that we can use propositional logic to model computational problems, the compactness theorem allows us to extend problem modelling to instances of infinite size. Recall the k-colorable problem, where we used propositional logic to model any finite size instance.

**Theorem 3.12.** A graph is k-colorable iff every finite subgraph is k-colorable.

**Definition 3.13** (Infinite graph).  $G = (V, E)$  is an infinite graph if  $V$  is infinite.

**Definition 3.14** (Subgraph).  $G' = (V', E')$  is a subgraph to  $G$  if  $V' \subseteq V, E' \subseteq E$

**Definition 3.15** (Induced subgraph).  $G' = (V', E')$  is an induced subgraph to  $G$  if  $G'$  is a subgraph to  $G$ , and  $\{u, v\} \in E' \iff \{u, v\} \in E \wedge u, v \in V'$

*Proof.* Since we have formulated any finite instance of k-colorable graph as propositional logic formula, we can define infinite set

$$\Gamma_G = \{\phi_{G'}^{k\text{-colorable}} \mid G' \text{ is finite subgraph of } G\}$$

We will first prove that

**Lemma 3.16.**  $G$  is finitely  $k$ -colorable  $\rightarrow \Gamma_G$  is finitely satisfiable.

*Proof.* We will reuse the proposition definition from last chapter, where  $|V|$  and thus  $P$  is countable

$$P = \{p_{v,i} \mid v \in V, i \in [1, k]\}$$

For each instance on  $G = (V, E)$  and  $k$  colors, we formulate as such to enforce constraints on edges (No same color on adjacent vertices), and vertices (exactly 1 coloring on each vertex).

$$\phi = \phi^{nbr} \wedge \phi^{exactly-1-color}$$

Consider the infinite graph  $G = (V, E)$

$$\begin{aligned}\Gamma_G &= \Gamma_G^{nbr} \cup \Gamma_G^{exactly-1-color} \\ \Gamma_G^{exactly-1-color} &= \{\phi_v^{exactly-1-color} \mid v \in V\} \\ \phi_v^{exactly-1-color} &= \left( \bigvee_{i \in [1, k]} p_{v,i} \right) \wedge \left( \bigwedge_{i \neq j \in [1, k]} (p_{v,i} \rightarrow p_{v,j}) \right) \\ \Gamma_G^{nbr} &= \{\phi_{\{u,v\}}^{nbr} \mid \{u, v\} \in E\} \\ \phi_{\{u,v\}}^{nbr} &= \bigwedge_{i \in [1, k]} (p_{u,i} \rightarrow p_{v,i})\end{aligned}$$

Suppose the contrary, where  $G = (V, E)$  is finitely  $k$ -colorable,  $\Gamma_G$  is not finitely satisfiable,  $\exists$  finite subset  $\Gamma_0 \subseteq \Gamma_G$ ,  $\Gamma_0$  is not satisfiable. Let  $G_0 = (V_0, E_0)$  be a finite subgraph of  $G$ , such that

$$\begin{aligned}V_0 &= \{v \mid v \in V, \phi_v^{exactly-1-color} \in \Gamma_0\} \cup \{u \mid u \in V, \exists u', \phi_{\{u,u'\}}^{nbr} \in \Gamma_0\} \\ E_0 &= \{\{u, v\} \mid \{u, v\} \in E, u, v \in V_0\}\end{aligned}$$

Here we are formulating an instance of finite  $k$ -colorable induced subgraph of  $G$  from finite subset  $\Gamma_0$ . Now we need to show that  $G_0$  is not  $k$ -colorable to reach the contradiction, with another contradiction.

Suppose  $G_0$  is  $k$ -colorable, let  $C_0$  be a  $k$ -coloring of  $G_0$ , define valuation

$$v_0(p_{u,i}) = \begin{cases} T, & \text{if } C_0(u) = i \\ F, & \text{if } C_0(u) \neq i \end{cases}$$

Let

$$\Gamma'_0 = \{\phi_v^{exactly-1-color} \mid v \in V_0\} \cup \{\phi_{\{u,v\}}^{nbr} \mid \{u, v\} \in E_0\}$$

Consider

$$\Gamma_0 \subseteq \Gamma'_0$$

Since  $\Gamma'_0$  is the same construction from  $k$ -colorability to formula,  $G_0$  is  $k$ -colorable, we have

$$v_0 \models \Gamma'_0$$

By relevance lemma

$$v_0 \models \Gamma_0$$

Contradicting to our initial assumption that  $\Gamma_0$  is NOT satisfiable. □

By Compactness theorem,  $\Gamma_G$  is finitely satisfiable  $\rightarrow \Gamma_G$  is satisfiable.

**Lemma 3.17.**  $\Gamma_G$  is satisfiable  $\rightarrow G$  is  $k$ -colorable

*Proof.*  $\Gamma_G$  is satisfiable  $\rightarrow \exists v, v \models \Gamma_G$ . We can construct a coloring  $C$  from  $v$  similarly to how we constructed  $v_0$  from  $C_0$  from previous proof:

$$C(v) = i, \text{ if } v(p_{v,i}) = T$$

By construction the coloring is valid, detailed technicality is omitted. □

□

## 4 Proof System

### 4.1 Hilbert Style Proof System

Four axiom schema of the hilberts style proof system. Axiom schema means that the  $\phi, \psi, \rho$  in these schema are placeholders that can be replaced by any valid formula

1. Axiom 1: Simplification

$$\frac{}{p \rightarrow (q \rightarrow p)}$$

2. Axiom 2: Distribution

$$\frac{}{(p \rightarrow (q \rightarrow r)) \rightarrow ((p \rightarrow q) \rightarrow (p \rightarrow r))}$$

3. Axiom 3: Double Negation Elimination (DNE)

$$\frac{}{((p \rightarrow \perp) \rightarrow \perp) \rightarrow p}$$

4. Rule of Inference: Modus Ponens

$$\frac{\frac{p \rightarrow q}{q} \quad p}{q}$$

The horizontal lines are judgement lines, which means that whats below can be inferred from whats above. The first 3 rules has no assumptions or premises, meaning they are ways we can introduce arbitrary claims or assumption into our proofs.

**Definition 4.1** (Proof). *A proof of the consequence of  $\phi$  from  $\Gamma$  is a finite sequence of formula  $c_1, c_2, \dots, c_k$  where  $\forall i \leq k, c_i \in \Gamma$  or,  $c_i$  is obtained using one of the 4 axioms:*

- $\exists j, l < i, c_j \equiv c_l \rightarrow c_i$
- $\exists \phi, \psi, c_i \equiv \phi \rightarrow (\psi \rightarrow \phi)$
- $\exists \phi, \psi, \rho, c_i \equiv [\phi \rightarrow (\psi \rightarrow \rho)] \rightarrow [(\phi \rightarrow \psi) \rightarrow (\phi \rightarrow \rho)]$
- $\exists \phi, c_i \equiv ((\phi \rightarrow \perp) \rightarrow \perp) \rightarrow \phi$

**Definition 4.2** (Derivable).  *$\phi$  is derivable from  $\Gamma$  if there is a proof in HSPS of " $\phi$  is a consequence of  $\Gamma$ "*

$$\Gamma \vdash \phi$$

**Example 4.3.**

$$\Gamma = \{p, p \rightarrow q\}, \quad \phi = q$$

*Proof.* We can construct proof sequence:

$$\begin{array}{ll} c_1 = p & (p \in \Gamma) \\ c_2 = p \rightarrow q & (p \rightarrow q \in \Gamma) \\ c_3 = q & (\text{Modus Ponens on } c_1, c_2) \end{array}$$

□

Sometimes HSPS requires us to creatively introduce new hypothesis to assist in the proof.

**Example 4.4.**

$$\Gamma = \{\perp\}, \phi = p$$

*Intuitively this does not sound provable, just from  $\perp$ , but axioms that require no premises allows us to introduce helper hypothesis. In this case  $(p \rightarrow \perp)$*

*Proof.*

$$\begin{array}{ll} c_1 = \perp & (\perp \in \Gamma) \\ c_2 = \perp \rightarrow [(p \rightarrow \perp) \rightarrow \perp] & (\text{Axiom 1: Simplification}) \\ c_3 = (p \rightarrow \perp) \rightarrow \perp & (\text{Modus Ponens on } \perp, (p \rightarrow \perp)) \\ c_4 = [(p \rightarrow \perp) \rightarrow \perp] \rightarrow p & (\text{Axiom 3: Double negation}) \\ c_5 = p & (\text{Modus Ponens on } c_3 \equiv [(p \rightarrow \perp) \rightarrow \perp], c_4 \equiv ([ (p \rightarrow \perp) \rightarrow \perp ] \rightarrow p)) \end{array}$$

□

As seen from this example, HSPS can get convoluted and unintuitive for relatively straightforward consequences.

## 4.2 Metatheorems

### 4.2.1 Monotonicity theorem

**Theorem 4.5.**

$$\Gamma \vdash A, \Gamma \subseteq \Delta \implies \Delta \vdash A$$

*Proof.* Let the proof of  $A$  from  $\Gamma$  be the sequence  $\langle \phi_1, \phi_2, \dots, \phi_n \rangle, \phi_n = A$

**Base case:**  $n = 1$

- Case 1:  $\phi_1 \in \Gamma \implies \phi_1 \in \Delta \implies \Delta \vdash \phi_1$
- Case 2:  $\phi_1$  is from one of the axiom schema,  $\Delta \vdash \phi_1$

**Inductive step:**  $\forall i < n, \Gamma \vdash \phi_i, \Delta \vdash \phi_i$

- Case 1, 2:  $\phi_{i+1} \in \Gamma$  or  $\phi_{i+1}$  is from one of the axiom schema, trivial
- Case 3:  $\phi_{i+1}$  is from Modus Ponens,  $\exists \phi_j$  such that

$$\frac{\phi_j \rightarrow \phi_{i+1} \quad \phi_j}{\phi_{i+1}}$$

$\Gamma \vdash \phi_j, \Gamma \vdash \phi_j \rightarrow \phi_{i+1}$  by inductive hypothesis,  $\Delta \vdash \phi_j, \Delta \vdash \phi_j \rightarrow \phi_{i+1}$ , by MP,  $\Delta \vdash \phi_{i+1}$

□

#### 4.2.2 Deduction

**Theorem 4.6.**

$$\Gamma \cup \{A\} \vdash B \iff \Gamma \vdash A \rightarrow B$$

*Proof.* (  $\implies$  )

We will use structural induction over the length of derivation from  $\Gamma \cup \{A\}$  to  $B$ ,  $\Gamma \cup \{A\} \vdash B \implies \exists \langle \phi_1, \phi_2, \dots, \phi_n \rangle, \phi_n = B$ .

**Base case:**  $n = 1, \phi_n = B$

- Case 1:  $B$  is an axiom  $\Gamma \vdash B$ . By simplification axiom,  $\Gamma \vdash B \rightarrow (A \rightarrow B)$ , by MP,  $\Gamma \vdash A \rightarrow B$ .
- Case 2:  $B = A$ , we will use the simplification and distribution axioms:
  - $p \equiv A$
  - $q \equiv (A \rightarrow A)$
  - $r \equiv A$
  - $(A \rightarrow ((A \rightarrow A) \rightarrow A)) \rightarrow ((A \rightarrow (A \rightarrow A)) \rightarrow (A \rightarrow A))$  (Distribution)
  - $A \rightarrow ((A \rightarrow A) \rightarrow A)$  (Simplification)
  - $(A \rightarrow (A \rightarrow A)) \rightarrow (A \rightarrow A)$  (MP)
  - $A \rightarrow (A \rightarrow A)$  (Simplification)
  - $(A \rightarrow A)$  (MP)

$$\Gamma \vdash A \rightarrow A \implies \Gamma \vdash A \rightarrow B$$

**Inductive step:** Suppose  $\forall i < n, \Gamma \cup \{A\} \vdash \phi_i \implies \Gamma \vdash A \rightarrow \phi_i$

- Case 1, 2, 3:  $\phi_{i+1}$  is in  $\Gamma$  or  $\phi_{i+1}$  is from one of the axioms, trivial; If  $\phi_{i+1} = A$ , the same as Case 2 in base case
- Case 4:  $\phi_{i+1}$  is obtained from MP, where  $\exists j \leq i$ ,

$$\begin{array}{ll} \Gamma \cup \{A\} \vdash \phi_j & \implies \Gamma \vdash A \rightarrow \phi_j \\ \Gamma \cup \{A\} \vdash \phi_j \rightarrow \phi_{i+1} & \implies \Gamma \vdash A \rightarrow (\phi_j \rightarrow \phi_{i+1}) \end{array}$$

$$\frac{\phi_j \rightarrow \phi_{i+1} \quad \phi_j}{\phi_{i+1}}$$

$$\begin{array}{ll} \Gamma \vdash (A \rightarrow \phi_j) \rightarrow (A \rightarrow \phi_{i+1}) & \text{by distribution} \\ \Gamma \vdash A \rightarrow \phi_{i+1} & \text{by MP} \end{array}$$

□

**Theorem 4.7** (Soundness of HSPS). *HSPS is sound.*

$$\Gamma \vdash \phi \implies \Gamma \models \phi$$

*Proof.* Define property  $H(i) : \forall j \leq i, \Gamma \models c_j$ .

**Base case:**  $i = 1$

Suppose  $\Gamma \vdash \phi, \exists \pi = c_1, c_2, \dots, c_k, c_k = \phi$ . By definition,  $c_1 \in \Gamma$  or

- $\exists \phi, \psi, c_1 \equiv \phi \rightarrow (\psi \rightarrow \phi)$
- $\exists \phi, \psi, \rho, c_1 \equiv [\phi \rightarrow (\psi \rightarrow \rho)] \rightarrow [(\phi \rightarrow \psi) \rightarrow (\phi \rightarrow \rho)]$
- $\exists \phi, c_1 \equiv ((\phi \rightarrow \perp) \rightarrow \perp) \rightarrow \phi$

Note that Modus Ponens is not possible to yield  $c_1$  as we need at least 1 previous formula. If  $c_1 \in \Gamma, \Gamma \models c_1$ ; Else, if  $c_1$  comes from any other axiom,  $c_1$  is valid, i.e.  $\forall v, v \models c_1$ , therefore  $\Gamma \models c_1$

**Inductive step:**

Suppose  $H(i), \forall j < i, \Gamma \models c_j, \forall v \models \Gamma, v \models c_j$ .

- Case 1, 2:  $c_{i+1} \in \Gamma$ , or  $c_{i+1}$  is from an axiom,  $\Gamma \models c_{i+1}, H(i+1)$
- Case 3:  $c_{i+1}$  comes from MP:  $\exists j < i, \Gamma \vdash c_j, \Gamma \vdash c_j \rightarrow c_{i+1}$ . By inductive hypothesis,  $\forall v \models \Gamma, v \models c_j, v \models c_j \rightarrow c_{i+1}$

$$\begin{aligned} v(c_j) &= T \\ v(c_j \rightarrow c_{i+1}) &= T \equiv v(\neg c_j \vee c_{i+1}) = T \\ v(\neg c_j) &= T \text{ OR } v(c_{i+1}) = T \\ v(c_j) = T &\implies v(\neg c_j) = F \implies v(c_{i+1}) = T \\ &\implies \forall v \models \Gamma, v \models c_{i+1} \implies \Gamma \models c_{i+1} \end{aligned}$$

□

**Theorem 4.8** (Completeness of HSPS). *HSPS is complete.*

$$\Gamma \models \phi \implies \Gamma \vdash \phi$$

*Proof.*

□

### 4.3 Resolution Proof System

**Definition 4.9** (Clause). *For each proposition  $p$  define literals  $p$  and  $\neg p$ . A clause  $C$  is finite set of literals over the proposition set  $P$ .*

$$v \models C = \{\ell_1, \ell_2, \dots, \ell_k\} \iff \exists j, v \models \ell_j$$

*As such, a clause  $C$  is satisfiable if  $\exists v, v \models C$*

Note that from this definition, an empty clause (an empty set of literals) is not satisfiable, as intuitively, a valuation models a clause if there exists at least one literal in the clause such that  $v \models \ell$ .

**Definition 4.10** (Conjunctive normal form (CNF)). *A formula  $\phi$  is said to be in Conjunctive Normal Form is*

$$\phi \equiv \bigwedge_{i=1}^k C_i$$

**Definition 4.11** (Satisfiability of a set of clauses).

$$\Gamma = \{C_1, C_2, \dots\}$$

- $\Gamma$  is *satisfiable* if  $\exists v, \forall C \in \Gamma, v \models C$
- $\Gamma$  is *unsatisfiable* if  $\forall v, \exists C \in \Gamma, v \not\models C$

By this definition,

- $\Gamma = \emptyset$  is vacuously satisfiable
- $\Gamma = \{\emptyset\}$  is not satisfiable.
- Any set of clauses that contains the empty set is unsatisfiable

**Theorem 4.12.** *Each well-formed formula can be transformed to a logically equivalent CNF*

*Proof.* **Base case:**  $\phi \in P$

trivial

**Inductive Case 1:**  $\neg$

Suppose there exists a CNF  $\bigwedge_i \bigvee_j \ell_{i,j}$  logically equivalent to  $\phi$ .

$$\begin{aligned} \neg\phi &= \neg((\ell_{1,1} \vee \ell_{1,2} \vee \dots) \wedge (\ell_{2,1} \vee \ell_{2,2} \vee \dots) \wedge \dots) \\ &= \neg(\ell_{1,1} \vee \ell_{1,2} \vee \dots) \vee \neg(\ell_{2,1} \vee \ell_{2,2} \vee \dots) \vee \dots \\ &= (\neg\ell_{1,1} \wedge \neg\ell_{1,2} \wedge \dots) \vee (\neg\ell_{2,1} \wedge \neg\ell_{2,2} \wedge \dots) \vee \dots \\ &= (\neg\ell_{1,1} \vee \neg\ell_{2,1} \vee \dots) \wedge () \end{aligned}$$

**Inductive Case 2:**  $\wedge$

Suppose  $\phi \equiv \bigwedge_{i \in I} C_i$  and  $\psi \equiv \bigwedge_{j \in J} D_j$  are CNFs.

$$\begin{aligned} \phi \wedge \psi &\equiv \left( \bigwedge_{i \in I} C_i \right) \wedge \left( \bigwedge_{j \in J} D_j \right) \\ &\equiv \bigwedge_{k \in I \cup J} E_k \end{aligned}$$

where  $E_k$  is a clause from the set of clauses of  $\phi$  or  $\psi$ . Since the conjunction of two conjunctions is simply a larger conjunction of the original clauses, the result is by definition a CNF.

**Inductive Case 3:  $\vee$** 

Suppose  $\phi \equiv \bigwedge_{i \in I} C_i$  and  $\psi \equiv \bigwedge_{j \in J} D_j$  are CNFs. By applying the Distributive Law:

$$\begin{aligned}
\phi \vee \psi &\equiv \left( \bigwedge_{i \in I} C_i \right) \vee \left( \bigwedge_{j \in J} D_j \right) \\
&\equiv \bigwedge_{i \in I} \left( C_i \vee \left( \bigwedge_{j \in J} D_j \right) \right) \\
&\equiv \bigwedge_{i \in I} \left( \bigwedge_{j \in J} (C_i \vee D_j) \right) \\
&\equiv \bigwedge_{i \in I, j \in J} (C_i \vee D_j)
\end{aligned}$$

Since each  $C_i$  and  $D_j$  are disjunctions of literals, their union  $C_i \vee D_j$  is also a disjunction of literals, and thus a valid clause. The resulting expression is a conjunction of these clauses, satisfying the definition of a CNF.

**Conclusion**

By the principle of induction on the complexity of the formula, we have shown that if the sub-formulas of  $\phi$  have equivalent CNFs, then  $\neg\phi$ ,  $\phi \wedge \psi$ , and  $\phi \vee \psi$  also have equivalent CNFs. Since the set of connectives  $\{\neg, \wedge, \vee\}$  is functionally complete, every well-formed formula can be transformed into a logically equivalent CNF. We can bring all negations to literal level, and supply an additional negation to each literal to negate the whole formula.  $\square$

**Theorem 4.13** (Soundness). *The resolution proof system is sound*

$$\vdash_{RES} \Gamma \implies \Gamma \text{ is unsatisfiable}$$

*Proof.*  $\vdash_{RES} \Gamma \rightarrow$  there exists a finite sequence ends with  $C_k$ .  $\square$

**Theorem 4.14** (Completeness). *The resolution proof system is complete*

$$\Gamma \text{ is unsatisfiable} \implies \vdash_{RES} \Gamma$$

*Proof.*  $\square$

## 5 Complexity

### 5.1 Turing Machine

**Definition 5.1** (Turing Machine).

$$\mathcal{M} = \langle Q, \Sigma, \Gamma, \delta, q_0, q_{Acc}, q_{Rej}, k \rangle$$

- $Q$ : the set of states
- $\Sigma$ : finite input alphabet
- $\Gamma$ : finite worktap alphabet,  $\square \in \Gamma$
- $\delta$ : the set of transitions
- $q_0$ : starting state,  $q_0 \in Q$

- $q_{Acc}$ : accepting state,  $q_{Acc} \in Q$
- $q_{Rej}$ : rejecting state,  $q_{Rej} \in Q$  (Optional)
- $k$ : the number of worktaps (Optional)

A TM has 1 read-only input tape, may have 1 write only output tape, and  $k$  work tape. We can show that any finitely many  $k$  tapes is equivalent to 1 worktap TM. [TODO]

### Transifiton function

$$\delta = \langle (Q \times \Sigma \times \Gamma^k) \times (Q \times \{L, R, S\} \times \Gamma^k \times \{L, R, S\}^k \times \Gamma \times \{L, R, S\}) \rangle$$

A state transition

$$\tau = \langle q, \sigma, \gamma_1, \gamma_2, \dots, \gamma_k, q', d, \gamma'_1, \gamma'_2, \dots, \gamma'_k, d_1, d_2, \dots, d_k, \gamma', d_o \rangle$$

$$\begin{aligned} \tau = & \langle q, \sigma, \\ & \gamma_1, \gamma_2, \dots, \gamma_k, \\ & q', d, \\ & \gamma'_1, \gamma'_2, \dots, \gamma'_k, \\ & d_1, d_2, \dots, d_k, \\ & \gamma', d_o \rangle \end{aligned}$$

Represents a transition from state  $q$ , when the head reading input symbol  $\sigma$ , and worktap symbol  $\gamma_i$  on worktape  $i$ , transition into state  $q'$ , write worktape symbol  $\gamma'_i$ , move the input head in the direction  $d$ , and workhead  $i$  in direction  $d_i$ , writing  $\gamma'$  on the output tape and move output head in direction  $d_o$

### Non-deterministic TM

Non deterministic TM has non-deterministic transition functions

$$\delta \subseteq \langle (Q \times \Sigma \times \Gamma^k) \times (Q \times \{L, R, S\} \times \Gamma^k \times \{L, R, S\}^k \times \Gamma \times \{L, R, S\}) \rangle$$

## 5.2 Semantics of TM

**Definition 5.2** (Configuration). A configuration  $C_t$  of a Turing machine at step  $t$  captures the overall state of the TM. Now we only consider TM with no rejecting states and output tape.

$$\begin{aligned} C = & \langle q, \\ & w_{in}^1 \nabla w_{in}^2, \\ & w_{w1}^1 \nabla w_{w1}^2 \\ & w_{w2}^1 \nabla w_{w2}^2 \\ & \dots \\ & w_{wk}^1 \nabla w_{wk}^2 \rangle \end{aligned}$$

Where  $w^1 \nabla w^2 = w$ , the symbol read by the head at each configuration is the first symbol after  $\nabla$ . Note that  $\nabla$  is not a Symbol but just an indication of the head position. The initial configuration will start at the starting state  $q_0$ , input head and work heads all at the beginning, and all worktapes empty.

$$C_0 = \langle q_0, \nabla w_{in}, \underbrace{\nabla, \nabla, \dots, \nabla}_k \rangle$$

Configuration

$$C_{t+1} = \langle q, \\ w_{in}^1 \nabla w_{in}^2, \\ w_{w1}^1 \nabla w_{w1}^2 \\ w_{w2}^1 \nabla w_{w2}^2 \\ \dots \\ w_{wk}^1 \nabla w_{wk}^2 \rangle$$

is a valid successor of

$$C_t = \langle q', \\ w_{in}^{1'} \nabla w_{in}^{2'}, \\ w_{w1}^{1'} \nabla w_{w1}^{2'} \\ w_{w2}^{1'} \nabla w_{w2}^{2'} \\ \dots \\ w_{wk}^{1'} \nabla w_{wk}^{2'} \rangle$$

$$\exists \tau = \langle p, \sigma, \gamma_1, \gamma_2, \dots, \gamma_k, p', d, \gamma'_1, \gamma'_2, \dots, \gamma'_k, d_1, d_2, \dots, d_k, \gamma', d_o \rangle$$

such that:

- $\sigma$  is the first symbol after the input head:  $w_{in}^2[0]$
- $\gamma_i$  is the first symbol after work head  $i$ :  $w_{wi}^2[0]$
- - if  $d = S$ :  $w'_{in} = w_{in}$   
 $|w_{in}^1| - |w_{in}^{1'}| = 0$
  - if  $d = L$ :  $w'_{in} = w_{in}^1[: -1] \nabla w_{in}^1[-1] \cdot w_{in}^2$   
 $|w_{in}^1| - |w_{in}^{1'}| = 1$
  - if  $d = L$ :  $w'_{in} = w_{in}^1 \cdot w_{in}^2[0] \nabla w_{in}^2[1 :]$   
 $|w_{in}^1| - |w_{in}^{1'}| = -1$
- $\forall i \in [1, k]$ ,
  - if  $d_i = S$ :  $w'_{wi} = w_{wi}$   
 $|w_{wi}^1| - |w_{wi}^{1'}| = 0$
  - if  $d_i = L$ :  $w'_{wi} = w_{wi}^1[: -1] \nabla w_{wi}^1[-1] \cdot w_{wi}^2$   
 $|w_{wi}^1| - |w_{wi}^{1'}| = 1$
  - if  $d_i = L$ :  $w'_{wi} = w_{wi}^1 \cdot w_{wi}^2[0] \nabla w_{wi}^2[1 :]$   
 $|w_{wi}^1| - |w_{wi}^{1'}| = -1$

**Definition 5.3.**  $\forall w \in \Sigma^*$ ,  $\mathcal{M}$  yields some valid sequences of configurations, which is called a run  $\pi_w = C_0, C_1, \dots$  where

- $C_0$  is the initial configuration
- $\forall t \geq 1, C_t$  is a valid successor of  $C_{t-1}$

A run is accepting if

1.  $\pi$  is finite, and
2.  $\pi$  ends with a configuration  $C_n$ , where  $C_n \cdot q = q_{Acc}$

A run is rejecting if

- $\pi$  is infinite, or
- $\pi$  ends with a configuration  $C_n$ , where  $C_n.q \neq q_{Acc}$

**Definition 5.4.**  $w \in \Sigma^*$  is accepted by  $\mathcal{M}$  if  $\exists \pi_w, \pi_w$  is accepting.

Now we can bridge to the concept of languages,

**Definition 5.5.** The set of strings accepted by  $\mathcal{M}$  is the language described by  $\mathcal{M}$ , ie

$$L(\mathcal{M}) = \{w \in \Sigma^* \mid w \text{ is accepted by } \mathcal{M}\}$$

Now we will focus on decidable languages, and their recognizing non deterministic TM.

**Definition 5.6** (Terminating Turing Machine).  $\mathcal{M}$  is terminating if  $\forall w \in \Sigma^*, \forall \pi_w, \pi_w$  is finite.

### 5.3 Running time and Space

**Definition 5.7.**

$$T : \mathbb{N} \mapsto \mathbb{N}$$

The running time of  $\mathcal{M}$  is bounded by  $T$  if

$$\forall w \in \Sigma^*, \forall \pi_w, |\pi_w| \leq T(|w|)$$

**Definition 5.8** (Space usage). The space usage of at configuration  $C_i$  is

$$space(C_i) = |w_{out}| + \sum_{j=1}^k |w_{wj}|$$

**Definition 5.9.**

$$S : \mathbb{N} \mapsto \mathbb{N}$$

The space usage of  $\mathcal{M}$  is bounded by  $S$  if

$$\forall w \in \Sigma^*, \forall \pi_w, \forall C \in \pi_w, space(C) \leq S(|w|)$$

**Definition 5.10.**  $L$  is computable by  $\mathcal{M}$  if  $L = L(\mathcal{M})$

**Definition 5.11.**  $L$  is computable in time  $T$  if  $\exists \mathcal{M}$

1.  $L = L(\mathcal{M})$  and
2. the running time of  $\mathcal{M}$  is bounded by  $T$

**Definition 5.12.**  $L$  is computable in space  $S$  if  $\exists \mathcal{M}$

1.  $L = L(\mathcal{M})$  and
2. the space usage of  $\mathcal{M}$  is bounded by  $S$

**Theorem 5.13** (Church-Turing Thesis). Any model of computation is only as powerful as the Turing machine.

$$\forall L, \exists N \text{ in a new model of computation, } L(N) = L \implies \exists \mathcal{M} \in TM, L(\mathcal{M}) = L$$

## 5.4 Complexity classes

**Definition 5.14** (*NTIME*).

$$NTIME(T(\cdot)) = \{L \mid L \text{ is computable in time } T(\cdot)\}$$

**Definition 5.15** (*DTIME*).

$$DTIME(T(\cdot)) = \{L \mid \exists \text{ deterministic } \mathcal{M}, L(\mathcal{M}) = L, \mathcal{M} \text{ running time is bounded by } T(\cdot)\}$$

**Definition 5.16** (*NSPACE*).

$$NSPACE(S(\cdot)) = \{L \mid L \text{ is computable in space } S(\cdot)\}$$

**Definition 5.17** (*DSPACE*).

$$DSPACE(S(\cdot)) = \{L \mid \exists \text{ deterministic } \mathcal{M}, L(\mathcal{M}) = L, \mathcal{M} \text{ space usage is bounded by } S(\cdot)\}$$

$$\begin{aligned} P &= PTIME = \bigcup_{k \geq 0} DTIME(f_k), & f_k(n) &= n^k \\ PSPACE &= \bigcup_{k \geq 0} DSPACE(f_k), & f_k(n) &= n^k \\ NP &= NPTIME = \bigcup_{k \geq 0} NTIME(f_k), & f_k(n) &= n^k \\ NPSPACE &= \bigcup_{k \geq 0} NSPACE(f_k), & f_k(n) &= n^k \end{aligned}$$

**Theorem 5.18.**

$$PTIME \subseteq NPTIME$$

**Theorem 5.19.**

$$PSPACE \subseteq NPSPACE$$

**Theorem 5.20.**

$$NPSPACE \subseteq PSPACE$$

*Proof.*

□

**Corollary 5.21.**

$$PSPACE = NPSPACE$$

## 5.5 Hardness

**Definition 5.22** (Hardness of Turing machine).  $\mathcal{M}_1 \leq \mathcal{M}_2$  if all runs of  $\mathcal{M}_1$  are smaller than  $\mathcal{M}_2$

## 5.6 Reduction

**Theorem 5.23** (Constant speed up theorem).  $L$  is computable by  $f \implies L$  is computable by  $f'$ , where

$$f'(n) = f\left(\frac{n}{c}\right)$$

**Theorem 5.24.** For all non-deterministic turing machines  $\mathcal{M}_1$  with, there exists a deterministic turing machine  $\mathcal{M}_2$ , such that  $L(\mathcal{M}_1) = L(\mathcal{M}_2)$

**Lemma 5.25.** *For all deterministic turing machine with  $\mathcal{M}_1$  with  $k$  worktapes, there exists a deterministic turing machine  $\mathcal{M}_2$  with 1 worktape such that  $L(\mathcal{M}_1) = L(\mathcal{M}_2)$ . Same for non deterministic TM.*

Consider deterministic TM  $\mathcal{M}$  with an output tape, it induces function

$$f_{\mathcal{M}} : \Sigma^* \mapsto \Gamma^*$$

Such that

$$f_{\mathcal{M}} = \begin{cases} \text{undefined, if } \mathcal{M} \text{ does not terminate;} \\ u, \text{ if } \mathcal{M} \text{ terminates, such that in the final configuration, output tape contains } u \end{cases}$$

**Corollary 5.26.**  *$f_{\mathcal{M}}$  is total if  $\mathcal{M}$  is terminating*

**Theorem 5.27.** *A partial function  $f : \Sigma^* \mapsto \Gamma^*$  is computable is  $\exists \mathcal{M}$  that induces  $f_{\mathcal{M}}, f = f_{\mathcal{M}}$*

**Theorem 5.28.** *A total function is computable in time  $T : \mathbb{N} \mapsto \mathbb{N}$  if there exists  $\mathcal{M}$  such that  $f_{\mathcal{M}} = f$ , and  $\mathcal{M}$  is running time bounded by  $T$*

## HW1

**Definition 5.29** (Generic Closure). *Let  $U$  be a universe of elements and let  $O$  be a set of functions whose domain is  $U$  and codomain is  $\mathcal{P}(U)$ . The arity of a function  $f \in O$  is denoted by  $\text{arity}(f)$ . A function  $f$  of arity  $r$  has type*

$$f : U^r \rightarrow \mathcal{P}(U).$$

*Let  $I \subseteq U$  be an initial set. A set  $S \subseteq U$  is said to be  $(I, O)$ -closed if:*

1.  $I \subseteq S$ .
2. *For every  $f \in O$  with  $\text{arity}(f) = r$  and for all  $e_1, \dots, e_r \in U$ , if  $e_1, \dots, e_r \in S$ , then  $f(e_1, \dots, e_r) \subseteq S$ .*

**Lemma 5.30.** *Let  $U$  be a universe,  $O$  a set of operators, and  $I \subseteq U$  an initial set. There exists a smallest  $(I, O)$ -closed set.*

Define  $\mathcal{P}(U)$  as a collection of all ordering of all elements in the power set  $\mathcal{P}(U)$

## Q1

Claim: The smallest  $(I, O)$ -closed set can be constructed by taking the intersection of all  $(I, O)$ -closed set, ie.

$$AIOCS = \{S \subseteq U \mid S \text{ is } (I, O)\text{-closed}\}$$

$$T_0 = \bigcap AIOCS$$

### Existence

First we prove the existence of such a set.

*Proof.* Suppose otherwise,

$$T_0 = \bigcap AIOCS = \emptyset$$

$$T_0 = \{x \in \mathcal{P}(U) \mid \forall S \in AIOCS, x \in S\}$$

$$T_0 = \emptyset \rightarrow x \in S \text{ is vacuously true } \forall x \in \mathcal{P}(U) \rightarrow T_0 = \mathcal{P}(U)$$

contradiction. □

### Closed

Now we prove that  $T_0$  is  $(I, O)$ -closed.

*Proof.* Property 1:

$$\begin{aligned} \forall S \in AIOCS, I &\subseteq S \\ &\rightarrow \forall i \in I, \forall S \in AIOCS, i \in S \\ &\rightarrow \forall i \in I, i \in \bigcap AIOCS \\ &\rightarrow \forall i \in I, i \in T_0 \end{aligned}$$

Property 2:

$$\begin{aligned} \forall S \in AIOCS, T_0 &\subseteq S \\ \text{Let } e_1, e_2, \dots, e_r &\in T_0, \forall S \in AIOCS, e_1, e_2, \dots, e_r \in S \\ &\rightarrow \exists f \in O, f(e_1, e_2, \dots, e_r) \in S \\ &\rightarrow \forall e_1, e_2, \dots, e_r \in T_0, \exists f \in O, f(e_1, e_2, \dots, e_r) \in \bigcap AIOCS \\ &\rightarrow f(e_1, e_2, \dots, e_r) \in T_0 \end{aligned}$$

□

### Uniqueness

Claim:

$$\forall T'_0 \in \{T \in \mathcal{P}(U) \mid \forall S \in AIOCS, T \subseteq S\}, T'_0 = T_0$$

*Proof.* Suppose otherwise,  $\exists T'_0 \in AIOCS, T'_0 \neq T_0$ .

$$\begin{aligned} T_0, T'_0 \in AIOCS &\rightarrow \forall S \in AIOCS, T_0 \subseteq S, T'_0 \subseteq S \\ &\rightarrow T_0 \subseteq T'_0, T'_0 \subseteq T_0 \\ &\rightarrow T_0 = T'_0 \end{aligned}$$

□

## Q2

For each of the following strings, prove or disprove that it is a well-formed formula:

1.  $w_1 = \epsilon$  (the empty string).
2.  $w_2 = (\neg p)$  where  $p \in P$ .
3.  $w_3 = (p\neg)$  where  $p \in P$ .

### 2.1

$$w_1 = \epsilon \notin FORM$$

Claim:  $\forall w \in FORM, |w| > 0$  I'd like to use definition 2.19 and lemma 2.21 which I proved later without using the conclusion from this question.

$$FORM = INDFORM = \bigcup_{i \geq 0} FORM_i$$

*Proof.* Base case:

$$\begin{aligned} FORM_0 &= P \\ \forall \phi \in FORM_0, \phi &= p \in P \\ &\rightarrow \forall \phi \in FORM_i, |\phi| > 0 \end{aligned}$$

Inductive Hypothesis: Suppose

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| > 0$$

$$\begin{aligned} FORM_{i+1} &= FORM_i \cup \{(\neg\phi) \mid \phi \in FORM_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in FORM_i\} \\ \forall \psi \in FORM_{i+1}, \exists \phi \in FORM_i, |\psi| &= |\phi| \text{ or } |\psi| = |\phi| + 3 \text{ or} \\ \exists \phi_1, \phi_2 \in FORM_i, |\psi| &= |\phi_1| + |\phi_2| + 3 \end{aligned}$$

□

### 2.2

$$p \in P \rightarrow p \in FORM \rightarrow (\neg p) \in FORM$$

### 2.3

$(p \neg) \notin FORM$

*Proof.* Suppose otherwise,  $(p \neg \epsilon) \in FORM$

$$\begin{aligned} \forall \phi \in FORM, (\neg \phi) \in FORM \\ \rightarrow \epsilon \in FORM \end{aligned}$$

Contradiction from 2.1 □

### Q3

Fix  $U = \mathbb{R}$ . For each of the following sets  $S$ , define a finite initial set  $I \subseteq \mathbb{R}$  and a set of operators  $O$  with  $|O| < 10$  such that the smallest  $(I, O)$ -closed set is  $S$ .

1.  $S = \{0, 1, 2, 3, \dots\}$ .
2.  $S = \{1, 3, 5, 7, \dots\}$ .
3.  $S = \mathbb{Q}$ .
4.  $S = \{\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \frac{5}{3}, \frac{7}{3}, \dots\}$ .

**Answer:**

1.  $\mathbb{N} : I = \{0\}, O = \{f(x) = x + 1\}$
2.  $Odd : I = \{1\}, O = \{f(x) = x + 2\}$
3.  $\mathbb{Q} : I = \{0\}, O = \{f(x) = -x, f(x) = x + 1, f(x, y) = \begin{cases} \frac{x}{y} & \text{if } y \neq 0 \\ \emptyset & \text{if } y = 0 \end{cases}\}$
4.  $\{\frac{x}{3} \mid x \in \mathbb{N}\} \setminus \mathbb{N} : I = \{\frac{1}{3}, \frac{2}{3}\}, O = \{f(x) = x + 1\}$

## Finite Directed Graphs

**Definition 5.31** (Graph Reachability). *Let  $G = (V, E)$  be a directed graph. A path from  $s$  to  $t$  is a sequence of vertices  $\pi = u_1, \dots, u_k$  such that:*

1.  $u_1 = s$ ,
2.  $(u_i, u_{i+1}) \in E$  for all  $1 \leq i < k$ ,
3.  $u_k = t$ .

*The length of  $\pi$  is  $k - 1$ . We say that  $t$  is reachable from  $s$  if  $s = t$  or such a path exists. The set of ancestors of  $t$  is defined as*

$$Reach_t = \{v \in V \mid t \text{ is reachable from } v\}.$$

### Q4

Let  $G = (V, E)$  be a directed graph and let  $t \in V$ . Fix the universe  $U = V$ .

1. Define an initial set  $I$  (with  $|I| \leq 2$ ) and a unary operator  $f$  such that the smallest  $(I, \{f\})$ -closed set is the set of vertices reachable from  $t$ .
2. Let  $d \in \mathbb{N} \setminus \{0\}$  and  $r < d$ . Define  $I$  and a unary operator  $f$  such that the smallest  $(I, \{f\})$ -closed set is the set of vertices reachable from  $s$  via paths of length  $\ell$  with  $\ell \equiv r \pmod{d}$ .

#### 4.1

$$I = \{t\}, O = \{f(v) = succ(v)\}$$

#### 4.2

$$I = \{s\} \cup \{v \in V \mid v \in succ^r(s)\}, O = \{f(v) = succ^d(v)\}$$

#### Q5

Let  $G = (V, E)$  be a directed graph and fix  $U = V \times V$ . Define an initial set  $I$  with  $|I| \leq |V|$  and a binary operator  $f$  such that the smallest  $(I, \{f\})$ -closed set is

$$AllReachPairs = \{(u, v) \mid v \text{ is reachable from } u\}.$$

#### 5.1

$$I = \{(v, v) \mid v \in V\}, O = \{f((v_1, v_2), (v_3, v_4)) \mapsto \begin{cases} \{(v_1, v_4)\} & \text{if } v_2 = v_3 \\ \emptyset & \text{if } v_2 \neq v_3 \end{cases}\}$$

### Inductive Definitions

**Definition 5.32** (Inductive Well-Formed Formulae). *Let  $P$  be a set of propositions and  $C = \{\neg, \vee\}$ . Define sets  $FORM_i$  as follows:*

$$FORM_0 = P,$$

$$FORM_{i+1} = FORM_i \cup \{(\neg w) \mid w \in FORM_i\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in FORM_i\}.$$

Define

$$INDFORM = \bigcup_{i \geq 0} FORM_i.$$

#### Q6

1. Show that  $FORM_i \subseteq FORM_{i+1}$  for all  $i$ .
2. Prove or disprove that  $FORM_i = FORM_{i+1}$  for some  $i$ .
3. Prove or disprove that  $FORM_i$  is  $(P, C)$ -closed for all  $i$ .
4. Prove or disprove that  $FORM_i$  is  $(P, C)$ -closed for some  $i$ .

#### 6.1

Base case:  $FORM_0 \subseteq FORM_1$  as

$$FORM_1 = FORM_0 \cup \{(\neg w) \mid w \in FORM_0\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in FORM_0\}$$

Wait this is just trivial

#### 6.2

Claim:

$$\exists i \geq 0, FORM_i = FORM_{i+1}$$

It is obvious that  $\forall i \geq 0, FORM_i$  is finite. Suppose

### 6.3

Claim:

$$\forall i \geq 0, FORM_i \text{ is (P,C)-closed}$$

*Proof.* Suppose otherwise, consider  $FORM_0$ ,

$$FORM_0 = P$$

$$\forall p \in P, (\neg p) \in FORM_0$$

Contradiction □

### 6.4

Claim:

$$\exists i \geq 0, FORM_i \text{ is (P,C)-closed}$$

*Proof.*

**Lemma 5.33.**

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

*The longest formula in  $FORM_i$  has size  $2^{i+3} - 3$ . Parentheses are counted in string length*

*Proof.* Base case:

$$FORM_0 = P$$

$$\forall \phi \in FORM_0, |\phi| = 1$$

Inductive hypothesis:

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

Consider all new strings formed from the longest string  $\phi \in FORM_i$ ,

$$|\phi| = 2^{i+3} - 3$$

$$|(\neg \phi)| = 2^{i+3}$$

$$|(\phi \vee \phi)| = 2 \times (2^{i+3} - 3) + 3 = 2^{(i+1)+3} - 3$$

□

Suppose  $\exists i \geq 0, FORM_i$  is (P,C)-closed, and from lemma 2.20 the longest formula in  $FORM_i$  has size  $|\phi_{max}| = 2^{i+3} - 3$ .  $\rightarrow \forall \phi \in FORM_i, (\neg \phi) \in FORM_i \rightarrow \phi'_{max} = (\neg \phi_{max}) \in FORM_i, |\phi'_{max}| > 2^{i+3} - 3$ , a contradiction.

□

**Lemma 5.34.**  $FORM = INDFORM$ .

## Q7

Prove the lemma by showing:

1.  $INDFORM$  is  $(P, C)$ -closed.
2. For every  $(P, C)$ -closed set  $S$ ,  $INDFORM \subseteq S$ .

### 7.1

Claim:  $INDFORM$  is (P,C)-closed

*Proof.* **Claim:**  $P \subseteq \text{INDFORM}$

First condition of (P,C)-closed set:

$$\text{FORM}_0 \subseteq \text{INDFORM} \rightarrow P \subseteq \text{INDFORM}$$

**Claim:**  $\forall w \in \text{INDFORM}, (\neg w) \in \text{INDFORM}$

$$\begin{aligned} \text{INDFORM} &= \bigcup_{i \geq 0} \text{FORM}_i \\ &\rightarrow \forall i \geq 0, \text{FORM}_i, \text{FORM}_{i+1} \in \text{INDFORM} \\ &\quad \forall w \in \text{INDFORM}, \exists j \geq 0, w \in \text{FORM}_j \\ &\rightarrow (\neg w) \in \text{FORM}_{j+1} \rightarrow (\neg w) \in \text{INDFORM} \end{aligned}$$

**Claim:**  $\forall w_1, w_2 \in \text{INDFORM}, (w_1 \vee w_2) \in \text{INDFORM}$

Case 1:  $\exists i \geq 0, w_1, w_2 \in \text{FORM}_i$

$$w_1, w_2 \in \text{FORM}_i \rightarrow (w_1 \vee w_2) \in \text{FORM}_{i+1} \rightarrow (w_1 \vee w_2) \in \text{INDFORM}$$

Case 2:  $\nexists i \geq 0, w_1, w_2 \in \text{FORM}_i$ . WLOG, suppose  $w_1 \in \text{FORM}_i, w_2 \in \text{FORM}_j, i < j$

$$\forall w \in \text{FORM}_n, w \in \text{FORM}_{n+1} \text{ By induction } w_1 \in \text{FORM}_i \rightarrow w_1, w_2 \in \text{FORM}_j$$

Back to case 1. □

## 7.2

$$\forall (\text{P,C})\text{-closed set } S, \text{INDFORM} \subseteq S$$

To prove this, we only need to prove that  $\text{INDFORM} \subseteq S_{\min} = \bigcap_S$  is (P,C)-closed  $S$ .

*Proof.* Base case:

$$\bigcup_{i=0}^0 \text{FORM}_i = \text{FORM}_0 = P \subseteq S_{\min}$$

Inductive hypothesis: Suppose

$$\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min}$$

$$\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min} \rightarrow \text{FORM}_n \in S_{\min}$$

$$S_{\min} \text{ is (P,C)-closed} \rightarrow \forall i \in [1, n], \forall \phi \in \text{FORM}_n, (\neg \phi) \in S_{\min},$$

$$\forall i \in [0, n], \forall \phi_1, \phi_2 \in \text{FORM}_i, (\phi_1 \vee \phi_2) \in S_{\min}$$

$$\rightarrow \bigcup_{i=0}^n \text{FORM}_i \cup \{(\neg \phi) \mid \forall \phi \in \bigcup_{i=0}^n \text{FORM}_i\} \cup \{(\phi_1 \vee \phi_2) \mid \forall \phi_1, \phi_2 \in \bigcup_{i=0}^n \text{FORM}_i\} \subseteq S_{\min}$$

$$\rightarrow \bigcup_{i=0}^{n+1} \text{FORM}_i \subseteq S_{\min}$$

□

**Definition 5.35** (Alternative Inductive Definition). Define  $AltFORM_0 = P$  and for  $i \geq 0$ :

$$\begin{aligned} AltFORM_{i+1} = & \{(\neg w) \mid w \in AltFORM_i\} \\ & \cup \{(w_1 \vee w_2) \mid w_1 \in AltFORM_i, \exists j \leq i, w_2 \in AltFORM_j\} \\ & \cup \{(w_1 \vee w_2) \mid w_2 \in AltFORM_i, \exists j \leq i, w_1 \in AltFORM_j\}. \end{aligned}$$

Define

$$AltINDFORM = \bigcup_{i \geq 0} AltFORM_i.$$

### Question 8

1. Show that  $FORM_i = \bigcup_{j=0}^i AltFORM_j$  for all  $i$ .
2. Show that  $INDFORM = AltINDFORM$ .

### 8.1

**Lemma 5.36.**

*Proof.* Base case:

$$FORM_0 = \bigcup_{j=0}^0 AltFORM_j$$

$$FORM_0 = P = AltFORM_0$$

Inductive hypothesis: Suppose

$$\forall j \in [0, i], FORM_i = \bigcup_{k=0}^i AltFORM_k$$

$$\begin{aligned} FORM_{i+1} &= \bigcup_{k=0}^i AltFORM_k \cup \{(\neg \phi) \mid \phi \in \bigcup_{k=0}^i AltFORM_k\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i AltFORM_k\} \\ &\supseteq \bigcup_{k=0}^i AltFORM_k \cup \{(\neg \phi) \mid \phi \in AltFORM_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i AltFORM_k\} \end{aligned}$$

$$\bigcup_{k=0}^{i+1} AltFORM_k \subseteq FORM_{i+1}$$

Now we prove the other way:

$$\forall \phi \in FORM_{i+1}, \phi \in \bigcup_{k=0}^i AltFORM_k \text{ or } \phi = (\neg \psi), \psi \in \bigcup_{k=0}^i AltFORM_k \text{ or } \phi = (\psi_1 \vee \psi_2), \psi_1, \psi_2 \in \bigcup_{k=0}^i AltFORM_k$$

Case 1:

$$\phi \in \bigcup_{k=0}^i AltFORM_k \rightarrow \phi \in \bigcup_{k=0}^{i+1} AltFORM_k$$

Case 2:

$$\begin{aligned}
\psi &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j \leq i, \psi \in \text{AltFORM}_j \\
\phi &= (\neg\psi), \psi \in \text{AltFORM}_j, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

Case 3:

$$\begin{aligned}
\psi_1, \psi_2 &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j, l \leq i, \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l \\
\phi &= (\psi_1 \vee \psi_2), \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k \\
\text{FORM}_{i+1} &\subseteq \bigcup_{k=0}^{i+1} \text{AltFORM}_k \rightarrow \text{FORM}_{i+1} = \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

□

## Formation Sequences

**Definition 5.37** (Formation Sequence). *A  $(P, C)$ -formation sequence is a nonempty finite sequence  $\rho = w_1, \dots, w_k$  such that for each  $i$ :*

- $w_i \in P$ , or
- $w_i = (\neg w_j)$  for some  $j < i$ , or
- $w_i = (w_j \vee w_k)$  for some  $j, k < i$ .

*The sequence belongs to  $w$  if  $w_k = w$ .*

### Q9

Give two distinct formation sequences that belong to

$$w = ((\neg p_1) \vee ((\neg p_3) \vee (\neg(p_2 \vee (\neg p_1)))))$$

Two distinct formation sequences  $\rho_1, \rho_2$  are:

$\rho_1$ :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= (\neg w_1) \\
w_3 &= p_2 \\
w_4 &= (w_3 \vee w_2) \\
w_5 &= (\neg w_4) \\
w_6 &= (p_3) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_7 \vee w_5) \\
w_9 &= (w_2 \vee w_8)
\end{aligned}$$

$\rho_2$ :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= p_2 \\
w_3 &= p_3 \\
w_4 &= (\neg w_3) \\
w_5 &= (\neg w_1) \\
w_6 &= (w_2 \vee w_5) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_4 \vee w_7) \\
w_9 &= (w_5 \vee w_8)
\end{aligned}$$

These are distinct because the components at index  $i = 2$  differ  $((\neg p_1)$  vs  $p_2$ ).

**Lemma 5.38.**  $INDFORM = \{w \in \Sigma^* \mid w \text{ is formation-sequence-well-formed}\}$ .

## Q10

Prove the lemma by showing:

1. Every formation-sequence-well-formed string is in  $INDFORM$ .
2. Every  $w \in INDFORM$  has a formation sequence.

Before attempting Q10 from the above definition of formation-sequence-well-formed, I will prove the following lemma using the definition below first:

**Definition 5.39.** Define  $ExFORM_0 = P$

$$\begin{aligned}
ExFORM_{i+1} &= (\{(\neg w) \mid w \in ExFORM_i\} \\
&\cup \{(w_1 \vee w_2) \mid w_1 \in ExFORM_i, \exists j \leq i, w_2 \in ExFORM_j\} \\
&\cup \{(w_1 \vee w_2) \mid w_2 \in ExFORM_i, \exists j \leq i, w_1 \in ExFORM_j\} \\
&)\setminus \bigcup_{j=0}^i ExFORM_j
\end{aligned}$$

**Lemma 5.40.**

$$\bigcup_{j=0}^i ExFORM_j = FORM_i$$

*Proof.* Base case:

$$ExFORM_0 = P = FORM_0$$

Inductive Hypothesis: Suppose  $\forall n \geq 0$ ,

$$\bigcup_{i=0}^n ExFORM_i = FORM_n$$

From definition 2.19,

$$\begin{aligned} FORM_{n+1} &= \bigcup_{i=0}^n ExFORM_i \cup \{(\neg w) \mid w \in \bigcup_{i=0}^n ExFORM_i\} \\ &\cup \{(w_1 \vee w_2) \mid w_1 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_2 \in \bigcup_{i=0}^n ExFORM_j\} \\ &\cup \{(w_1 \vee w_2) \mid w_2 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_1 \in \bigcup_{i=0}^n ExFORM_j\} \end{aligned}$$

**Definition 5.41.**

$$\begin{aligned} FromFORM : n &\mapsto \{(\neg w) \mid w \in \bigcup_{i=0}^n ExFORM_i\} \\ &\cup \{(w_1 \vee w_2) \mid w_1 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_2 \in \bigcup_{i=0}^n ExFORM_j\} \\ &\cup \{(w_1 \vee w_2) \mid w_2 \in \bigcup_{i=0}^n ExFORM_i, \exists j \leq i, w_1 \in \bigcup_{i=0}^n ExFORM_j\} \end{aligned}$$

**Lemma 5.42.**

$$FORM_{n+1} = \bigcup_{i=0}^n ExFORM_i \cup FromFORM(n)$$

**Lemma 5.43.**

$$\begin{aligned} ExFORM_{n+1} &= (\bigcup_{i=0}^n ExFORM_i \cup FromFORM(n)) \setminus \bigcup_{i=0}^n ExFORM_i \\ ExFORM_{n+1} &= FromFORM(n) \setminus \bigcup_{i=0}^n ExFORM_i \end{aligned}$$

From Lemma 5.42 and Lemma 5.43, we have

$$\begin{aligned} \bigcup_{i=0}^{n+1} ExFORM_i &= \bigcup_{i=0}^n ExFORM_i \cup ExFORM_{n+1} \\ &= \bigcup_{i=0}^n ExFORM_i \cup (FromFORM(n) \setminus \bigcup_{i=0}^n ExFORM_i) \\ &= \bigcup_{i=0}^n ExFORM_i \cup FromFORM(n) \\ &= FORM_{n+1} \end{aligned}$$

□

**Lemma 5.44.**

$$\bigcup_{i \geq 0} ExFORM_i = \bigcup_{i \geq 0} FORM_i$$

*Proof.*  $\bigcup_{i \geq 0} ExFORM_i \subseteq \bigcup_{i \geq 0} FORM_i$  $\forall n \geq 0$ , From Lemma 2.28

$$\phi \in \bigcup_{i=0}^n ExFORM_i \rightarrow \phi \in FORM_n \rightarrow \phi \in \bigcup_{i=0}^n FORM_i$$

$$\bigcup_{i \geq 0} ExFORM_i \supseteq \bigcup_{i \geq 0} FORM_i$$

Corollary from Lemma 2.28

□

**Corollary 5.45.**

$$\bigcup_{i \geq 0} ExFORM_i = INDFORM$$

 $\forall \phi \in ExFORM_i \rightarrow \phi$  is well-formed formula

## 10.1, 2

Claim:

$$FORMSeq = \{w \in \Sigma^* \mid w \text{ is formation-sequence-well-formed}\} = INDFORM$$

*Proof.* Let  $\rho_i$  denotes the set of  $\rho$  that contains a sequence of  $i$  strings  $w_1, w_2, \dots, w_i$   
Define

$$\begin{aligned} str : \rho &\mapsto w, str(\rho = w_1, w_2, \dots, w_k) = w_k \\ str : \rho_i &\mapsto \{w\}, str(\rho = w_1, w_2, \dots, w_k) = \{w \mid w = str(\rho), \rho \in \rho_i\} \end{aligned}$$

Claim:

$$\bigcup_{k \geq 1} str(\rho_k) = \bigcup_{i \geq 0} ExFORM_i$$

*Proof.*  $\bigcup_{k \geq 1} \rho_k \subseteq \bigcup_{i \geq 0} ExFORM_i$  $\forall w \in FORMSeq, \exists \rho = w_1, w_2, \dots, w_k, w = w_k$ 

Base case:

$$\bigcup_{k=1}^1 str(\rho_1) = \{w \mid w \in P\} = \bigcup_{i=0}^0 ExFORM_i$$

Inductive hypothesis: Suppose

$$\bigcup_{j=1}^{k+1} str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$$

$$\bigcup_{j=1}^{k+2} str(\rho_j) = \bigcup_{j=1}^{k+1} str(\rho_j) \cup \{w \mid \exists \rho \in \rho_{k+2}, w = str(\rho)\}$$

By definition,  $w_{k+2} = str(\rho)$  for some  $\rho \in \rho_{k+2}$ , one of the 3 below holds:

1.  $w_{k+2} \in P \rightarrow w_{k+2} \in \bigcup_{i \geq 0} ExFORM_i$
2.  $w_{k+2} = (\neg w_n), n < k+2 \rightarrow w_n \in \bigcup_{j=1}^n str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$
3.  $w_{k+2} = (w_m \vee w_n), m, n < k+2 \rightarrow w_m, w_n \in \bigcup_{j=1}^{\max(m,n)} str(\rho_j) \subseteq \bigcup_{i=0}^k ExFORM_i$

$$\bigcup_{k \geq 1} \rho_k \supseteq \bigcup_{i \geq 0} ExFORM_i$$

Base case is the same as above.

Inductive hypothesis: (this will be used in Q11)

**Lemma 5.46.**

$$\bigcup_{i=0}^k ExFORM_i \subseteq \bigcup_{j=1}^{2(k+1)} str(\rho_j)$$

$\forall w \in ExFORM_{k+1}$ , one of the 3 below holds: 1.

$$w \in P \rightarrow w \in ExFORM_0 \rightarrow w \in \bigcup_{i=0}^k ExFORM_i \subseteq \bigcup_{j=1}^{k+1} str(\rho_j)$$

2.

$$\begin{aligned} w = (\neg w_n), w_n \in ExFORM_k &\rightarrow \exists \rho = w_1, w_2, \dots, w_k \in \bigcup_{j=1}^{k+1} \rho_j, w_n = str(\rho) \\ &\rightarrow \exists \rho' \in \rho_{k+2}, \rho' = w_1, w_2, \dots, w_k, (\neg w_n) \\ &\rightarrow w \in str(\rho_{k+2}) \rightarrow w \in \bigcup_{j=1}^{k+2} str(\rho_j) \\ &\rightarrow w \in \bigcup_{j=1}^{2(k+1)} str(\rho_j) \end{aligned}$$

3. (Concat finite sequences that yield formula composing disjunctions gives finite sequence)

$$\begin{aligned} w = (w_m \vee w_n), WLOG, w_m \in ExFORM_k, \exists q \leq k, w_n \in ExFORM_q \\ &\rightarrow \exists \rho^m \in \rho_{k+1} = w_1^m, w_2^m, \dots, w_k^m, \\ &\quad \rho^n \in \rho_{q+1} = w_1^n, w_2^n, \dots, w_q^n \in \bigcup_{j=1}^{k+1} \rho_j, w_m = str(\rho^m), w_n = str(\rho^n) \\ &\rightarrow \exists \rho' = w_1^m, w_2^m, \dots, w_k^m, w_1^n, w_2^n, \dots, w_q^n, (w_k^m \vee w_q^n) \in \rho_{k+q+2} \\ &\rightarrow w \in \bigcup_{j=1}^{k+q+1} str(\rho_j) \\ &\rightarrow w \in \bigcup_{j=1}^{2(k+1)} str(\rho_j) \end{aligned}$$

Therefore,

$$\bigcup_{k \geq 1} str(\rho_k) = \bigcup_{i \geq 0} ExFORM_i$$

□

By Corollary 5.45,

$$\bigcup_{k \geq 1} str(\rho_k) = INDFORM$$

□

## Q11

**Lemma 5.47.** *Every well-formed formula  $\phi$  is of one of the following forms:*

1.  $\phi \in P$ ,
2.  $\phi = (\neg\psi)$ ,
3.  $\phi = (\psi_1 \vee \psi_2)$ .

*Proof.* From Corollary 5.45,

$$\forall \phi \in INDFORM, \exists k \geq 0, \phi \in ExFORM_k$$

Case  $x = 0$ :

$$x = 0 \rightarrow \phi \in ExFORM_0 = P$$

Case  $x > 0$ :

$$\begin{aligned} & \phi \in ExFORM_x \text{ and } \forall y \in [0, x), \phi \notin ExFORM_y \\ & \forall \phi \in ExFORM_x, \phi \in P \text{ or} \\ & \phi \in (\{(\neg\psi) \mid \psi \in ExFORM_{x-1}\} \\ & \quad \cup \{(\psi_1 \vee \psi_2) \mid \psi_1 \in ExFORM_{x-1}, \exists j \leq x-1, \psi_2 \in ExFORM_j\}) \end{aligned}$$

Since we define  $ExFORM_i$  to be pair wise disjoint, we guarantee that

$$\phi \in ExFORM_k \iff k \text{ is the minimum depth needed to form } \phi \text{ in the syntax tree}$$

Intuitively,  $x$  is the minimum depth required to formulate  $\phi$ , thus with depth  $d \in [0, x)$ ,  $\phi$  cannot be formed

Suppose  $\phi \in P$ ,

$$\phi \in ExFORM_0, \phi \in ExFORM_x, x > 0$$

Contradiction. Therefore

$$\phi \in (\{(\neg\psi) \mid \psi \in ExFORM_{x-1}\} \cup \{(\psi_1 \vee \psi_2) \mid \psi_1 \in ExFORM_{x-1}, \exists j \leq x-1, \psi_2 \in ExFORM_j\})$$

Now we will prove  $\forall \phi \in INDFORM$

$$(\phi \in \{(\neg\psi) \mid \psi \in ExFORM_{x-1}\}) \oplus (\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in ExFORM_{x-1}, \exists j \leq x-1, \psi_2 \in ExFORM_j\})$$

Suppose we enforce the use of parentheses for all subformula, that is except  $\phi \in P$ , all formula starts with '(' . If for some  $i$ ,  $\phi \in \{(\neg\psi) \mid \psi \in ExFORM_i\}$ , ' $\neg$ ' must be at index 1. Likewise, if for some  $i$ ,  $\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in ExFORM_i, \exists j \leq i, \psi_2 \in ExFORM_j\}$ , ' $\vee$ ' must not be at index 1. Thus  $\nexists \phi \in INDFORM, \phi \in \{(\neg\psi) \mid \psi \in ExFORM_{x-1}\}$  and  $\phi \in \{(\psi_1 \vee \psi_2) \mid \psi_1 \in ExFORM_i, \exists j \leq i, \psi_2 \in ExFORM_j\}$  □

## HW2

**Lemma 5.48** (Characterization of Well-Formed Formulae). *Let  $P$  be a set of propositions and  $C = \{\neg, \vee\}$ . Every well-formed formula  $\phi$  over  $(P, C)$  is of exactly one of the following forms:*

1.  $\phi \in P$ ,
2.  $\phi = (\neg\psi)$  for some well-formed formula  $\psi$ ,
3.  $\phi = (\psi_1 \vee \psi_2)$  for some well-formed formulae  $\psi_1, \psi_2$ .

**Structural Induction Principle.** A property  $P(\phi)$  holds for all well-formed formulae if:

- $P(p)$  holds for all  $p \in P$ ,
- $P(\psi)$  implies  $P(\neg\psi)$ ,
- $P(\psi_1)$  and  $P(\psi_2)$  imply  $P(\psi_1 \vee \psi_2)$ .

## Q1

### 1.1

**Theorem 5.49.** *Let  $\alpha$  be a well-formed formula. Let  $c(\alpha)$  denote the number of occurrences of binary connective symbols ( $\wedge, \vee, \rightarrow, \leftrightarrow$ ) in  $\alpha$ , and let  $s(\alpha)$  denote the number of occurrences of sentence symbols in  $\alpha$ . Then*

$$s(\alpha) = c(\alpha) + 1.$$

*Proof. Base Case:*

Let  $\alpha = p$  for some proposition  $p \in \mathcal{P}$ .

- $s(p) = 1$  and  $c(p) = 0$ .
- Since  $1 = 0 + 1$ ,  $P(p)$  holds for all atomic propositions.

**Inductive Case 1 (Negation):**

Suppose  $P(\psi)$  holds, i.e.,  $s(\psi) = c(\psi) + 1$ . Let  $\alpha = (\neg\psi)$ .

- $s(\neg\psi) = s(\psi)$  no new sentence symbols are added.
- $c(\neg\psi) = c(\psi) + 1$  (not binary connective).
- Thus,  $s(\alpha) = s(\psi) = c(\psi) + 1 = c(\alpha) + 1$ .  $P(\alpha)$  holds.

**Inductive Case 2 (Binary Connectives):**

Suppose  $P(\psi_1)$  and  $P(\psi_2)$  hold. Let  $\alpha = (\psi_1 c \psi_2)$  where  $c \in \{\vee, \rightarrow\}$ .

- $s(\alpha) = s(\psi_1) + s(\psi_2)$  (sum of occurrences in sub-formulae).
- $c(\alpha) = c(\psi_1) + c(\psi_2) + 1$  (sum of binary connectives plus the new  $\star$ ).
- By the inductive hypothesis:  $s(\alpha) = (c(\psi_1) + 1) + (c(\psi_2) + 1)$ .
- Rearranging gives  $s(\alpha) = (c(\psi_1) + c(\psi_2) + 1) + 1$ .
- Therefore,  $s(\alpha) = c(\alpha) + 1$ .  $P(\alpha)$  holds.

□

## 1.2

We define the property  $P(\alpha)$  as follows: the length of  $\alpha$ , denoted  $L(\alpha)$ , is of the form  $4k + 1$  and the number of sentence symbols,  $s(\alpha)$ , is  $k + 1$  for some  $k \in \mathbb{N}$

*Proof. Base Case:*

Let  $\alpha = p$  for some proposition  $p \in \mathcal{P}$

- The length  $L(p) = 1 = 4 \cdot 0 + 1$
- The number of sentence symbols  $s(p) = 1$
- Thus,  $P(p)$  holds.

**Inductive Case:**

Suppose  $P(\psi_1)$  and  $P(\psi_2)$  hold. Then  $L(\psi_1) = 4k_1 + 1$  and  $s(\psi_1) = k_1 + 1$ . Also  $L(\psi_2) = 4k_2 + 1$  and  $s(\psi_2) = k_2 + 1$ . Let  $\alpha = (\psi_1 c \psi_2)$  where  $c$  is a binary connective

**Length of  $\alpha$ :**

- The string  $\alpha$  consists of: one left parenthesis,  $\psi_1$ , one binary connective,  $\psi_2$ , and one right parenthesis
- $L(\alpha) = 1 + L(\psi_1) + 1 + L(\psi_2) + 1$ .
- $L(\alpha) = 3 + (4k_1 + 1) + (4k_2 + 1) = 4k_1 + 4k_2 + 5$ .
- $L(\alpha) = 4(k_1 + k_2 + 1) + 1$ .
- Let  $k = k_1 + k_2 + 1$ , we see  $L(\alpha) = 4k + 1$ , odd

**Number of sentence symbols:**

- $s(\alpha) = s(\psi_1) + s(\psi_2) = (k_1 + 1) + (k_2 + 1) = k_1 + k_2 + 2$ .
- Using  $k = k_1 + k_2 + 1$  from above,  $s(\alpha) = (k_1 + k_2 + 1) + 1 = k + 1$
- $k + 1 = \frac{4k+4}{4} > \frac{4k+1}{4}$

□

## Q2

Recall the following derived connectives:

- $\phi_1 \wedge \phi_2 \equiv \neg((\neg\phi_1) \vee (\neg\phi_2))$ ,
- $\perp \equiv \neg(p \vee \neg p)$  for some  $p \in P$ ,
- $\phi_1 \rightarrow \phi_2 \equiv (\neg\phi_1) \vee \phi_2$ .

Using structural induction, show that for every formula  $\phi$  over  $(P, \{\neg, \vee\})$  with  $P \neq \emptyset$ , there exists a formula  $\psi$  using only connectives  $\{\perp, \rightarrow\}$  such that  $\phi$  and  $\psi$  are logically equivalent.

*Proof.* We will show that every formula formed over  $\{\neg, \vee\}$  can be formed using  $\{\perp, \rightarrow\}$ , and denote the logically equivalent formula using the later connector set from  $\phi$  as  $\psi$ . Define property  $P(\phi)$  as follows: there exists a logically equivalent formula over  $P$  using the connector set  $\{\perp, \rightarrow\}$ .

**Base Case:**

- $\forall \phi = p \in P, \psi = p$

**Inductive Case 1:**

Suppose  $P(\phi)$  for some formula,  $\exists \psi \equiv \phi$

- $(\neg \phi) = (\neg \phi) \vee ((\neg \phi) \vee (\neg \phi)) = \phi \rightarrow \perp = \psi \rightarrow \perp$

**Inductive Case 2:**

Suppose  $P(\phi_1), P(\phi_2)$  for some formula,  $\exists \psi_1 \equiv \phi_1, \psi_2 \equiv \phi_2$

- $(\phi_1 \vee \phi_2) = ((\neg(\neg \phi_1)) \vee \phi_2) = ((\neg \phi_1) \rightarrow \phi_2)$
- From inductive case 1,  $((\neg \phi_1) \rightarrow \phi_2) = ((\phi_1 \rightarrow \perp) \rightarrow \phi_2) = ((\psi_1 \rightarrow \perp) \rightarrow \psi_2)$

□

## Q3

**Definition 5.50** (Conjunctive Normal Form). *A formula  $\phi$  over  $(P, \{\neg, \vee, \wedge\})$  is in conjunctive normal form (CNF) if*

$$\phi = C_1 \wedge C_2 \wedge \cdots \wedge C_m \quad (m \geq 1),$$

where each clause  $C_i$  has the form

$$C_i = \ell_{i,1} \vee \ell_{i,2} \vee \cdots \vee \ell_{i,k_i},$$

and each literal  $\ell_{i,j}$  is either  $p$  or  $\neg p$  for some  $p \in P$ .

Using structural induction, show that for every well-formed formula  $\phi$ , there exists a logically equivalent formula  $\psi$  in CNF.

*Proof.* Define property  $P(\phi)$  to be there exists a CNF  $\psi$  that is logically equivalent to  $\phi$

**Base Case:**

$\phi = p$ . An atomic proposition  $p$  is a clause containing a single literal. A single clause is, by definition, a CNF. Thus,  $P(p)$  holds.

**Inductive Case 1  $\neg$ :**

Assume  $P(\phi)$  holds. If  $\phi$  has a CNF, we can first convert  $\neg \phi$  into NNF [https://en.wikipedia.org/wiki/Negation\\_normal\\_form](https://en.wikipedia.org/wiki/Negation_normal_form)

- $\neg(A \wedge B) \equiv \neg A \vee \neg B$
- $\neg(A \vee B) \equiv \neg A \wedge \neg B$
- $\neg \neg A \equiv A$

**Inductive Case 2  $\vee$ :**

Assume  $P(\phi_1)$  and  $P(\phi_2)$  hold. Let  $\psi_1 = \bigwedge_i C_i$  and  $\psi_2 = \bigwedge_j D_j$  be CNFs.

- $\phi_1 \vee \phi_2 \equiv (\bigwedge_i C_i) \vee (\bigwedge_j D_j)$ .
- By the Distributive Law  $(A \vee (B \wedge C) \equiv (A \vee B) \wedge (A \vee C))$ , we distribute the  $\vee$  over the  $\wedge$ :
- $\phi_1 \vee \phi_2 \equiv \bigwedge_i \bigwedge_j (C_i \vee D_j)$ .
- Since the disjunction of two clauses  $C_i \vee D_j$  is also a clause, the final expression is a CNF.

□

**Q4**

**Definition 5.51** (Occurs). Define  $OCCUR : \text{FORM} \rightarrow \mathcal{P}(P)$  inductively by:

- If  $\phi = p \in P$ , then  $OCCUR(\phi) = \{p\}$ ,
- If  $\phi = \neg\psi$ , then  $OCCUR(\phi) = OCCUR(\psi)$ ,
- If  $\phi = \psi_1 \oplus \psi_2$  with  $\oplus \in \{\wedge, \vee\}$ , then  $OCCUR(\phi) = OCCUR(\psi_1) \cup OCCUR(\psi_2)$ .

**Lemma 5.52** (Relevance Lemma). Let  $\phi$  be a propositional formula and let  $v_1, v_2 : P \rightarrow \{\text{true}, \text{false}\}$  be valuations such that  $v_1(p) = v_2(p)$  for all  $p \in OCCUR(\phi)$ . Then  $v_1 \models \phi$  if and only if  $v_2 \models \phi$ .

Prove this lemma using structural induction.

*Proof.* We define the property  $P(\phi)$  as: for any  $v_1, v_2$ , if  $v_1(p) = v_2(p) \forall p \in OCCUR(\phi)$ ,  $v_1 \models \phi \iff v_2 \models \phi$ .

**Base Case:**

$\phi = p$  for some  $p \in P$

- $OCCUR(p) = \{p\}$ ,  $v_1(p) = v_2(p)$
- $v_1 \models p \iff v_1(p) = \text{true}$  and  $v_2 \models p \iff v_2(p) = \text{true}$
- $v_1(p) = v_2(p)$ ,  $v_1 \models p \iff v_2 \models p$

**Inductive Step 1 (negation):**

Assume  $P(\psi)$  holds. Let  $\phi = \neg\psi$ .

- $OCCUR(\phi) = OCCUR(\psi)$ ,  $v_1 \models \psi \iff v_2 \models \psi$
- $v_1 \models \neg\psi \iff v_1 \not\models \psi$  and  $v_2 \models \neg\psi \iff v_2 \not\models \psi$
- Since the truth values of  $\psi$  are the same, so do their negation.

### Inductive Step 2 (Binary connective):

Assume  $P(\psi_1)$  and  $P(\psi_2)$  hold.

- Let  $\phi = \psi_1 \oplus \psi_2$  for  $\oplus \in \{\wedge, \vee\}$
- $OCCUR(\phi) = OCCUR(\psi_1) \cup OCCUR(\psi_2)$ , if  $v_1, v_2$  agree on  $OCCUR(\phi)$ , they agree on both  $OCCUR(\psi_1)$  and  $OCCUR(\psi_2)$
- $v_1 \models \psi_1 \iff v_2 \models \psi_1$  and  $v_1 \models \psi_2 \iff v_2 \models \psi_2$
- the satisfaction of  $\psi_1 \oplus \psi_2$  depends only on the truth values of  $\psi_1$  and  $\psi_2$ , and those values are the same for  $v_1$  and  $v_2$ , thus  $v_1 \models \phi \iff v_2 \models \phi$

□

## Q5

**Lemma 5.53** (Generalized Relevance Lemma). *Let  $\phi$  be a propositional formula and let  $v_1, v_2$  be valuations such that  $v_1(p) = v_2(p)$  for all  $p$  in some set  $S \supseteq OCCUR(\phi)$ . Then  $v_1 \models \phi$  if and only if  $v_2 \models \phi$ .*

Prove this lemma.

*Proof.* Let  $\phi$  be a formula and  $S$  be a set such that  $S \supseteq OCCUR(\phi)$ . Suppose  $v_1(p) = v_2(p) \forall p \in S$ . Since  $S$  contains all elements of  $OCCUR(\phi)$ , the condition  $v_1(p) = v_2(p) \forall p \in S \implies v_1(p) = v_2(p) \forall p \in OCCUR(\phi)$ . By the Relevance lemma, if  $v_1$  and  $v_2$  agree on  $\forall p \in OCCUR(\phi)$ , then  $v_1 \models \phi \iff v_2 \models \phi$ . □

## Q6

**Lemma 5.54** (Relevance Lemma for Logical Equivalence). *Let  $\phi_1, \phi_2$  be propositional formulae. The following are equivalent:*

- For all valuations  $v_1, v_2$  agreeing on  $OCCUR(\phi_1) \cup OCCUR(\phi_2)$ , we have  $v_1 \models \phi_1 \iff v_2 \models \phi_2$ .
- $\phi_1$  and  $\phi_2$  are logically equivalent.

Prove this lemma.

*Proof.* We prove the equivalence of the two statements by showing both implications.

( $\Leftarrow$ ) Suppose  $\phi_1$  and  $\phi_2$  are logically equivalent.  $\forall v, v \models \phi_1 \iff v \models \phi_2$ . Consider  $v_1, v_2$  that agree on  $OCCUR(\phi_1) \cup OCCUR(\phi_2)$ , we have:

- $v_1 \models \phi_1 \iff v_1 \models \phi_2$
- $v_2 \models \phi_1 \iff v_2 \models \phi_2$

Trivial

( $\Rightarrow$ ) Suppose  $\forall v_1, v_2$  agreeing on  $S = OCCUR(\phi_1) \cup OCCUR(\phi_2)$ , we have  $v_1 \models \phi_1 \iff v_2 \models \phi_2$ . Let  $v$  be any valuation. We will show that  $v \models \phi_1 \iff v \models \phi_2$ . Define  $v_1 = v$  and  $v_2 = v$ . These valuations trivially agree on  $S$ .

- $v_1 \models \phi_1 \iff v_2 \models \phi_2$
- $v \models \phi_1 \iff v \models \phi_2$
- $\phi_1$  and  $\phi_2$  are logically equivalent

□

## Q7

Let  $G = (V, E)$  be an undirected graph and  $k \in \mathbb{N}$ . Construct a propositional formula  $\phi_G$  such that:

1.  $|\phi|$  is polynomial in  $k + |V| + |E|$ ,
2.  $\phi$  is satisfiable if and only if  $G$  has a vertex cover of size at most  $k$ .

Formally prove correctness of the encoding.

*Proof.* Define the proposition set  $P$ :

$$\begin{aligned} P &= P_{in} \cup P_{out} \cup P_{count} \\ P_{in} &= \{p_{v,in} \mid v \in V\} \\ P_{out} &= \{p_{v,out} \mid v \in V\} \\ P_{count} &= \{c_{i,j} \mid i, j \in [0, |V|]\} \\ |P| &\in O(|V|^2) \end{aligned}$$

There are constraint we need to consider:

$$\forall v \in V, v \in V' \text{ or } v \in V \setminus V'$$

$$\phi_{in} = \bigwedge_{v \in V} (p_{v,in} \vee p_{v,out})$$

$$|\phi_{in}| \in O(|V|)$$

$$\forall v \in V, v \notin V' \text{ or } v \in V \setminus V'$$

$$\phi_{out} = \bigwedge_{v \in V} \neg(p_{v,in} \wedge p_{v,out})$$

$$|\phi_{out}| \in O(|V|)$$

**All edges covered**

$$\phi_e = \bigwedge_{\{u,v\} \in E} (p_{u,in} \vee p_{v,in})$$

$$|\phi_e| \in O(|V|^2)$$

$$|V'| \leq k$$

We use the following facts to define inductive rules:

- The number of true proposition is non decreasing as  $i$  increases in the sequence  $p_1, \dots, p_i \rightarrow$  if the  $c_{i,j}$  is true,  $\forall \ell \geq i, c_{\ell,j}$  is true.
- if the  $c_{i,j}$  is false,  $\forall \ell \geq j, c_{i,\ell}$  is false

Inductively define  $c_{i,j}$ :

$$[c_{i-1,j} \vee (c_{i-1,j-1} \wedge p_{i,in})] \leftrightarrow c_{i,j}$$

We use bidirectional implication to prevent vacuous truthity of  $c_{i,j}$  when conditions do not hold. And the last constraint can be represented by  $\neg(c_{|V|,k+1})$

$$|\{c_{i,j} \mid i \in [1, |V|], j \in [1, k+1]\}| \in O(|V| \cdot k)$$

Putting everything together we can encode  $k$ -vertex cover using

$$\phi = \phi_{in} \wedge \phi_{out} \wedge \phi_e \wedge (\neg c_{|V|,k+1})$$

□

### Correctness of encoding

To prove the correctness of the encoding, we will show:

**Theorem 5.55.**  $\phi$  is satisfiable  $\rightarrow \exists k$ -vertex cover on  $G$

*Proof.*  $\phi$  is satisfiable  $\rightarrow \exists v, v \models \phi, \rightarrow$

$$\begin{aligned} v &\models \phi_{in} \\ v &\models \phi_{out} \\ v &\models \phi_e \\ v &\models (\neg c_{|V|,k+1}) \end{aligned}$$

As per construction, this guarantees  $\forall v \in V$ ,  $v$  is either in the vertex cover or exclusively;  $\forall e = \{u, v\} \in E$ , either  $u$  or  $v$  inclusively are in the vertex cover, and the size of the cover is at most  $k$   $\square$

**Theorem 5.56.**  $\exists k$ -vertex cover on  $G \rightarrow \phi$  is satisfiable

*Proof.* Suppose  $G = (V, E)$  has a vertex cover  $V' \subseteq V$  such that  $|V'| \leq k$ . We construct a valuation  $v$  as follows:

1. For each  $u \in V$ , set  $v(p_{u,in}) = \text{true}$  if  $u \in V'$ , and  $v(p_{u,in}) = \text{false}$  otherwise
2. Set  $v(p_{u,out}) = \neg v(p_{u,in}) \implies \phi_{in} \wedge \phi_{out}$  is satisfiable
3. Since  $V'$  is a vertex cover, for every edge  $\{u, v\} \in E$ , at least one of  $u$  or  $v$  must be in  $V'$ . Thus,  $v(p_{u,in}) \vee v(p_{v,in})$  is true for all edges, satisfying  $\phi_e$ .
4. Define  $v(c_{i,j})$  to be true if and only if  $|\{u_1, \dots, u_i\} \cap V'| \geq j$ .
  - the recurrence  $[c_{i-1,j} \vee (c_{i-1,j-1} \wedge p_{i,in})] \leftrightarrow c_{i,j}$  is satisfied because the count of vertices in the cover up to  $i$  increases only if the  $i$ -th vertex is in the cover.
  - Since  $|V'| \leq k$ , the total count of vertices in  $V'$  after considering all  $|V|$  vertices cannot be  $k+1$ . Thus,  $v(c_{|V|,k+1}) = \text{false}$ , which satisfies  $\neg c_{|V|,k+1}$ .

Since all components are satisfied,  $\phi$  is satisfiable.  $\square$

## Q8

Given a partial  $N^2 \times N^2$  Sudoku grid  $S$ , construct a propositional formula  $\phi_S$  of size polynomial in  $N^2$  such that  $\phi_S$  is satisfiable if and only if  $S$  has a completion to a valid Sudoku grid. Prove correctness.

*Proof.* Define proposition set  $P = \{p_{i,j,k} \mid i, j, k \in [1, N]\}$ , representing cell  $(i, j)$  is assigned the number  $k$ . We need to satisfy the following constraints:

**Cell: At most 1 value assigned**

$$\phi_{cell, \leq 1} = \bigwedge_{i=1}^N \bigwedge_{j=1}^N \bigwedge_{1 \leq k \leq m \leq N} (\neg p_{i,j,k} \vee \neg p_{i,j,m})$$

**Cell: At least 1 value assigned**

$$\phi_{cell, \geq 1} = \bigwedge_{i=1}^N \bigwedge_{j=1}^N \bigvee_{k=1}^N p_{i,j,k}$$

**Row: At most 1 value assigned**

$$\phi_{row, \leq 1} = \bigwedge_{i=1}^N \bigwedge_{k=1}^N \bigwedge_{1 \leq j < m \leq N} (\neg p_{i,j,k} \vee \neg p_{i,m,k})$$

**Column: At most 1 value assigned**

$$\phi_{col, \leq 1} = \bigwedge_{j=1}^N \bigwedge_{k=1}^N \bigwedge_{1 \leq i < m \leq N} (\neg p_{i,j,k} \vee \neg p_{m,j,k})$$

**Block: At most 1 value assigned**

Let  $n = \sqrt{N}$ . For each block starting at row  $r \in \{0, \dots, n-1\}$  and column  $c \in \{0, \dots, n-1\}$ :

$$\phi_{block, \leq 1} = \bigwedge_{r=0}^{n-1} \bigwedge_{c=0}^{n-1} \bigwedge_{k=1}^N \bigwedge_{(i,j) \neq (i',j')} (\neg p_{rn+i, cn+j, k} \vee \neg p_{rn+i', cn+j', k})$$

where  $1 \leq i, j, i', j' \leq n$ .

**Initial Setup**

Let  $S$  be the set of triples  $(i, j, k)$  such that cell  $(i, j)$  is pre-assigned value  $k$ :

$$\phi_{clues} = \bigwedge_{(i,j,k) \in S} p_{i,j,k}$$

The complete formula is:

$$\phi_{Sudoku} = \phi_{cell, \geq 1} \wedge \phi_{cell, \leq 1} \wedge \phi_{row, \leq 1} \wedge \phi_{col, \leq 1} \wedge \phi_{block, \leq 1} \wedge \phi_{clues}$$

□

## Correctness of Encoding

**Theorem 5.57.** *The formula  $\phi_{Sudoku}$  is satisfiable  $\iff$  the Sudoku puzzle has a valid solution*

*Proof.* ( $\Rightarrow$ ) Suppose  $\phi_{Sudoku}$  is satisfiable under valuation  $v$ . We construct a Sudoku assignment by setting cell  $(i, j) = k$  if  $v(p_{i,j,k}) = \text{true}$ .

- By  $\phi_{cell, \geq 1}$  and  $\phi_{cell, \leq 1}$ , every cell is assigned exactly one value.
- By  $\phi_{row, \leq 1}$ , no value  $k$  is repeated in a row. Since there are  $N$  cells in a row and  $N$  possible values, each value must appear exactly once.
- The same logic applies to  $\phi_{col, \leq 1}$  and  $\phi_{block, \leq 1}$  for columns and  $n \times n$  subgrids.
- $\phi_{clues}$  ensures the solution respects the starting digits.

( $\Leftarrow$ ) Suppose a valid Sudoku solution exists. Define a valuation  $v$  such that  $v(p_{i,j,k}) = \text{true}$  if cell  $(i, j)$  contains  $k$ , and false otherwise.

- Because the solution is valid, each cell has one number (satisfying cell constraints).
- Because rows, columns, and blocks contain no duplicates, the "at most one" ( $\leq 1$ ) constraints for each are satisfied.
- Because the solution contains the initial clues,  $\phi_{clues}$  is satisfied.

Therefore,  $v \models \phi_{Sudoku}$ .

□

## Background: Compactness

**Definition 5.58** (Satisfiable). A set  $\Gamma$  of propositional formulae is satisfiable if there exists a valuation  $v$  such that  $v \models \Gamma$ .

**Definition 5.59** (Finitely Satisfiable). A set  $\Gamma$  is finitely satisfiable if every finite  $\Gamma' \subseteq \Gamma$  is satisfiable.

**Theorem 5.60** (Compactness Theorem). A set  $\Gamma$  of propositional formulae is satisfiable if and only if it is finitely satisfiable.

### Q11

**Lemma 5.61.** If  $\Gamma$  is satisfiable, then  $\Gamma$  is finitely satisfiable.

*Proof.*  $\Gamma$  is satisfiable  $\implies \exists v, \forall \phi \in \Gamma, v \models \phi$   
 $\forall$  finite subset  $\Gamma' \subseteq \Gamma, \forall \psi \in \Gamma', \psi \in \Gamma \implies \forall \Gamma' \subseteq \Gamma, \forall \psi \in \Gamma', v \models \psi$  □

### Q12

**Definition 5.62** ( $P$ -complete).  $\Gamma$  is  $P$ -complete if for every  $p \in P$ , either  $p \in \Gamma$  or  $\neg p \in \Gamma$ .

**Definition 5.63** ( $P$ -consistent).  $\Gamma$  is  $P$ -consistent if for no  $p \in P$  do both  $p$  and  $\neg p$  belong to  $\Gamma$ .

**Lemma 5.64.** If  $\Gamma$  is finitely satisfiable and  $P$ -complete, then  $\Gamma$  is  $P$ -consistent.

1. Prove the lemma.

*Proof.* Suppose otherwise,  $\Gamma$  is  $P$ -complete, finitely satisfiable and NOT  $P$ -consistent,  $\exists$  finite set  $\Gamma' \subseteq \Gamma, \exists p \in P, p \in \Gamma', (\neg p) \in \Gamma' \implies \Gamma'$  is not satisfiable, a contradiction. □

### 12.2

- $P = \{p, \}$
- $\Gamma = \emptyset$

### 12.3

- $P = \{p\}$
- $\Gamma = \{(\neg(p \vee (\neg p)))\}$

### 12.4

I cannot

### Q13

Let  $\Gamma$  be finitely satisfiable and let  $\phi$  be a formula.

1. Prove that at least one of  $\Gamma \cup \{\phi\}$  or  $\Gamma \cup \{\neg\phi\}$  is finitely satisfiable.
2. Give examples as requested.

**Lemma 5.65.** at least one of  $\Gamma \cup \{\phi\}$  or  $\Gamma \cup \{\neg\phi\}$  is finitely satisfiable.

*Proof.* We will prove the contrapositive of this statement: if neither  $\Gamma \cup \{\phi\}$  nor  $\Gamma \cup \{(\neg\phi)\}$  is satisfiable,  $\Gamma$  is not finitely satisfiable. Consider finite subset  $\Gamma_1, \Gamma_2 \subseteq \Gamma$ , and  $\phi \in FORM$ ,  $\Gamma_1 \cup \{\phi\}$  and  $\Gamma_2 \cup \{(\neg\phi)\}$  are both not satisfiable,  $\forall v, v \not\models (\Gamma_1 \cup \{\phi\}) \implies v \not\models \Gamma_1$  or  $v \not\models \phi$ , and  $\forall v, v \not\models (\Gamma_2 \cup \{(\neg\phi)\}) \implies v \not\models \Gamma_2$  or  $v \not\models (\neg\phi)$

Rewrite with respect to the set of all valuations, define

- $V_1 = \{v : P \mapsto \{T, F\} \mid v \models \Gamma_1\}$
- $V_2 = \{v : P \mapsto \{T, F\} \mid v \models \Gamma_2\}$
- $V_\phi = \{v : P \mapsto \{T, F\} \mid v \models \{\phi\}\}$
- $V_{(\neg\phi)} = \{v : P \mapsto \{T, F\} \mid v \not\models \{\phi\}\}$

$$V_1 \cap V_\phi = \emptyset \implies \forall v \in V_1, v \not\models \phi \implies v \models (\neg\phi) \implies V_1 \subseteq V_{(\neg\phi)}$$

$$V_2 \cap V_{(\neg\phi)} = \emptyset \implies \forall v \in V_2, v \not\models (\neg\phi) \implies v \models \phi \implies V_2 \subseteq V_\phi$$

Now consider also a finite subset to  $\Gamma, \Gamma_1 \cup \Gamma_2$ , define

- $V_U = \{v : P \mapsto \{T, F\} \mid v \models (\Gamma_1 \cup \Gamma_2)\}$

$$V_U = (V_1 \cap V_2) \subseteq (V_\phi \cup V_{(\neg\phi)}) = \emptyset \implies \nexists v, v \models (\Gamma_1 \cup \Gamma_2) \implies \Gamma \text{ is not finitely satisfiable. } \square$$

## 13.2

- $\Gamma = \{p\}, \phi = q$

## 13.3

- $\Gamma = \{p, (\neg q)\}, \phi = q$

## Q14, 15

Let  $P = \{p_1, p_2, \dots\}$  and define  $\Gamma_0, \Gamma_1, \dots$

**Lemma 5.66.**  $\Gamma_U = \bigcup_{i \geq 0} \Gamma_i$  is finitely satisfiable.

**Lemma 5.67.**

$$\forall i \geq 0, \Gamma_i \text{ is finitely satisfiable}$$

*Proof.* Base case:  $i = 0, \Gamma_i = \Gamma_0 = \Gamma$  is finitely satisfiable by definition.

Inductive Hypothesis:  $\Gamma_n$  is finitely satisfiable. By Lemma 5.65,  $\Gamma_n \cup \{p_n\}$  is finitely satisfiable or  $\Gamma_n \cup \{(\neg p_n)\}$  is finitely satisfiable.  $\square$

*Proof.* Consider finite set  $\Gamma' \subseteq \Gamma_U$ , since  $\Gamma'$  is finite, and  $\forall k \geq 0, \Gamma_k \subseteq \Gamma_{k+1}$  by definition,  $\exists j \geq 0, \Gamma' \subseteq \Gamma_j = \bigcup_{i \geq 0}^j \Gamma_i$ . By Lemma 5.61,  $\Gamma_j$  is satisfiable  $\rightarrow \Gamma'$  is finitely satisfiable.  $\square$

*Proof.* From the construction,  $\forall i \geq 0$  at least one of  $p_i$  and  $(\neg p_i)$  is in  $\bigcup_{i \geq 0} \Gamma_i$ . Since  $\bigcup_{i \geq 0} \Gamma_i$  is finitely satisfiable,  $\forall i \geq 0, p_i$  and  $(\neg p_i)$  cannot both be in  $\bigcup_{i \geq 0} \Gamma_i$ . Otherwise  $\{p_i\} \cup \{(\neg p_1)\} \subseteq \bigcup_{i \geq 0} \Gamma_i$  is not satisfiable, contradiction.  $\square$

Now we have proven that the construction proposed from a finitely satisfiable formula set  $\Gamma$  to  $\bigcup_{i \geq 0} \Gamma_i$  is also finitely satisfiable. Now to complete the proof of the compactness theorem, we need to prove that  $\exists v, v \models \Gamma$ .

*Proof.* From Lemma 5.66,  $\forall i \geq 0, (p_i \in \Gamma_U) \oplus ((\neg p_i) \in \Gamma_U), \exists v, v \models \Gamma_U$

$$v(p) : P \mapsto \{T, F\} = \begin{cases} T, & \text{if } p \in \Gamma_U \\ F, & \text{if } (\neg p) \in \Gamma_U \end{cases}$$

Define the literal set  $Literals = \{p_i, (\neg p_i) \mid i \geq 0\}$ , and function

$$\tau(p) : P \mapsto Literals = \begin{cases} p, & \text{if } p \in \Gamma_U \\ (\neg p), & \text{if } (\neg p) \in \Gamma_U \end{cases}$$

**Lemma 5.68.**

$$v \models \Gamma_U \rightarrow v \models \tau(\Gamma)$$

Let  $\phi \in \Gamma$ , Define literal set  $X = \{\tau(p) \mid p \in OCCUR(\phi)\}$  where  $X$  is a set containing single proposition literals. By definition we have

$$X \subseteq \Gamma_U \rightarrow v \models X \rightarrow \forall \psi \in X, v(\psi) = T$$

and  $OCCUR(X) = OCCUR(\phi)$  Since both  $X$  is finite,  $X \cup \{\phi\}$  is a finite subset to  $\Gamma_U$ , thus satisfiable.  $\rightarrow \exists v', v' \models X \cup \{\phi\} \rightarrow v' \models X, v' \models \phi \rightarrow \forall \psi \in X, v'(\psi) = T$ .

$$\forall p \in OCCUR(\phi), v(p) = v'(p), v' \models \phi \rightarrow v \models \phi$$

□

## Q16

Let  $\top$  and  $\perp$  be the 0-place connectives (truth and falsity). For a wff  $\phi$  and sentence symbol  $A$ , let  $\phi_{\top}^A$  and  $\phi_{\perp}^A$  denote the substitution of  $A$  with  $\top$  and  $\perp$  respectively. Define the formula  $\phi_A^* = (\phi_{\top}^A \vee \phi_{\perp}^A)$ .

*Proof.* **1.**  $\phi \models \phi_A^*$ .

Let  $v$  be a valuation such that  $v \models \phi$ .

- If  $v(A) = T \implies v(\phi) = v(\phi_{\top}^A)$ .  $v \models \phi \implies v \models \phi_{\top}^A$ .
- If  $v(A) = F \implies v(\phi) = v(\phi_{\perp}^A)$ .  $v \models \phi \implies v \models \phi_{\perp}^A$ .
- either way,  $v \models (\phi_{\top}^A \vee \phi_{\perp}^A)$  Thus,  $\phi \models \phi_A^*$ .

**2. If  $\phi \models \psi$  and  $A$  does not occur in  $\psi$ , then  $\phi_A^* \models \psi$ .**

Let  $v \models \phi_A^* \implies v \models \phi_{\top}^A$  or  $v \models \phi_{\perp}^A$ .

- Case 1:  $v \models \phi_{\top}^A$ . Let  $v'$  be valuation  $v[A \mapsto \text{true}] \implies v' \models \phi$ .  $\phi \models \psi \implies v' \models \psi$ .  $A \notin OCCUR(\psi) \implies \psi$  is independent of  $A \implies v \models \psi$ .
- Case 2: similar
- Thus,  $\phi_A^* \models \psi$

**3.  $\phi$  is satisfiable iff  $\phi_A^*$  is satisfiable.**

- $(\Rightarrow)$   $\phi$  is satisfiable  $\Rightarrow \exists v \models \phi$ . By part 1,  $\phi \models \phi_A^* \Rightarrow v \models \phi_A^*$
- $(\Leftarrow)$   $\phi_A^*$  is satisfiable  $\Rightarrow \exists v, v \models \phi_{\top}^A$  or  $v \models \phi_{\perp}^A$ .  $v \models \phi_{\top}^A \Rightarrow v[A \mapsto \text{true}] \models \phi$ .  
 $v \models \phi_{\perp}^A \Rightarrow v[A \mapsto \text{false}] \models \phi$

□

**Theorem 5.69** (Interpolation Theorem). *If  $\alpha \models \beta$ , then there exists a formula  $\gamma$  such that  $\alpha \models \gamma \models \beta$ , and  $OCCUR(\gamma) \subseteq OCCUR(\alpha) \cap OCCUR(\beta)$ .*

*Proof.* Let  $S = OCCUR(\alpha) \setminus OCCUR(\beta)$ ,

- Let  $\alpha_0 = \alpha$ .
- $\forall i \in [1, n]$ , let  $\alpha_i = (\alpha_{i-1})_{A_i}^*$ .
- By part 1,  $\alpha_{i-1} \models \alpha_i$ , by transitivity  $\alpha \models \alpha_n$ .
- By part 2,  $\alpha \models \beta$  and  $A_1 \notin OCCUR(\beta) \Rightarrow \alpha_1 \models \beta$ . Repeat  $\forall A_i \in S, \alpha_n \models \beta$ .
- $\gamma = \alpha_n$  contains no propositions from  $S$ , meaning  $OCCUR(\gamma) \subseteq OCCUR(\alpha) \cap OCCUR(\beta)$ .

□

## HW3

### Hilbert-style Proof System

The Hilbert-style (or Frege) proof system for propositional logic comprises of 3 axiom schema and the modus ponens rule of inference

- **Axiom Schema 1:**  $\varphi \rightarrow (\psi \rightarrow \varphi)$
- **Axiom Schema 2:**  $(\varphi \rightarrow (\psi \rightarrow \rho)) \rightarrow ((\varphi \rightarrow \psi) \rightarrow (\varphi \rightarrow \rho))$
- **Axiom Schema 3:**  $((\varphi \rightarrow \perp) \rightarrow \perp) \rightarrow \varphi$
- **Modus Ponens:** From  $\varphi$  and  $\varphi \rightarrow \psi$ , infer  $\psi$

### Question 1

Let  $\Gamma, \Gamma'$  be sets of propositional logic formulae and let  $\varphi$  be a formula. Show that if  $\Gamma \vdash \varphi$  and if  $\Gamma \subseteq \Gamma'$ , then  $\Gamma' \vdash \varphi$

### Theorem 1 (Deduction Theorem)

Let  $\Gamma$  be a set of propositional logic formulae and  $\varphi, \psi$  be propositional logic formulae.  $\Gamma \vdash (\varphi \rightarrow \psi)$  iff  $\Gamma \cup \{\varphi\} \vdash \psi$

### Question 2

Prove Theorem 1. Do not use the completeness theorem (Theorem 3) to establish this! We will instead use Theorem 1 to establish Theorem 3 in later questions. *Hint: One direction is easier. For the other direction, consider a proof  $\pi = \psi_1, \psi_2, \dots, \psi_n$  of  $\psi$  from  $\Gamma \cup \{\varphi\}$ . Use induction on the index  $i$  to show that the formula  $(\varphi \rightarrow \psi_i)$  can be proved from  $\Gamma$*

### Question 3

Prove the following (in the Hilbert proof system). Here,  $\varphi, \psi$ , and  $\rho$  are propositional logic formulae, and  $\Gamma$  is a set of propositional logic formulae

- (a)  $\vdash \varphi \rightarrow \varphi$
- (b)  $\vdash (\psi \rightarrow \varphi) \rightarrow \varphi$ , where  $\psi = \varphi \rightarrow \perp$
- (c)  $\{\perp\} \vdash \varphi$
- (d)  $\vdash \psi \rightarrow (\varphi \rightarrow \rho)$  where  $\psi = \varphi \rightarrow \perp$

### Theorem 2 (Soundness of Hilbert-style Proof System)

Let  $\Gamma$  be a set of propositional logic formulae and  $\varphi$  be a propositional logic formula. If  $\Gamma \vdash \varphi$ , then  $\Gamma \models \varphi$

### Question 4

Prove Theorem 2. *Hint: Given a proof, inductively establish that every formula in the proof is a logical consequence of the assumptions.*

### Theorem 3 (Completeness of Hilbert-style Proof System)

Let  $\Gamma$  be a set of propositional logic formulae and  $\varphi$  be a propositional logic formula. If  $\Gamma \models \varphi$ , then  $\Gamma \vdash \varphi$

### Theorem 4 (Weak Completeness of Hilbert-style Proof System)

Let  $\varphi$  be a propositional logic formula. If  $\models \varphi$ , then  $\vdash \varphi$ .

### Question 5

Prove that two notions of a set being inconsistent are identical. That is, prove that the following two statements are equivalent

1. There is a formula  $\varphi$  such that  $\Gamma \vdash \varphi$  and  $\Gamma \vdash (\varphi \rightarrow \perp)$
2. For every formula  $\rho$ ,  $\Gamma \vdash \rho$

### Lemma 1

Let  $\Gamma$  be a set of formulae and  $\varphi$  be a formula. If  $\Gamma \cup \{\varphi \rightarrow \perp\}$  is inconsistent, then  $\Gamma \vdash \varphi$

### Lemma 2

Let  $\Gamma$  be a set of formulae and  $\varphi$  be a formula. If  $\Gamma \cup \{\varphi\}$  is inconsistent, then  $\Gamma \vdash (\varphi \rightarrow \perp)$

### Lemma 3

Let  $\Gamma$  be a set of formulae. If  $\Gamma$  is inconsistent, then there is a finite subset  $\Gamma' \subseteq_{fin} \Gamma$  of  $\Gamma$  such that  $\Gamma'$  is also inconsistent

### Question 6

- (a) Prove Lemma 1
- (b) Prove Lemma 2
- (c) Let  $\Gamma$  be a set of formulae and  $\varphi$  be a formula. Show that, if  $\Gamma$  is consistent, then at least one of  $\Gamma \cup \{\varphi\}$  or  $\Gamma \cup \{\varphi \rightarrow \perp\}$  is consistent
- (d) Let  $\Gamma, \Gamma'$  be sets of formulae such that  $\Gamma \subseteq \Gamma'$ . If  $\Gamma'$  is consistent, then so is  $\Gamma$
- (e) Prove Lemma 3

### Question 7

- (a) Prove, using induction, that  $\Delta_i$  is consistent, for every  $i$
- (b) Prove that  $\Delta$  is consistent
- (c) Prove that  $\Delta$  is negation-complete, i.e., for every formula  $\psi$ , we have  $(\psi \rightarrow \perp) \in \Delta$  iff  $(\psi \notin \Delta)$

- (d) Use the fact that  $\Delta$  is negation-complete to show that for every formula  $\psi$ ,  $\psi \in \Delta$  iff  $\Delta \vdash \psi$

### Question 8

- (a) Show that for any formula  $\psi$ ,  $v \models \psi$  iff  $\psi \in \Delta$
- (b) Show that  $v \models (\varphi \rightarrow \perp)$  and thus  $\not\models \varphi$

### Question 9

Prove Theorem 3 using Theorem 4, the Compactness theorem and the Deduction theorem

### Theorem 5 (Soundness of Resolution Proof System)

Let  $\Gamma$  be a set of clauses (over propositions  $P$ ). If there is a resolution proof of  $\Gamma$ , then  $\Gamma$  is unsatisfiable.

### Question 10

Prove Theorem 5, *Hint: Consider a resolution proof  $\pi = C_1, C_2, \dots, C_n$  of  $\Gamma$ . Use induction to establish that for every  $1 \leq i \leq n$  if a valuation  $v$  is such that  $v \models \Gamma$ , then  $v \models C_i$ .*

### Theorem 6 (Completeness of Resolution Proof System)

Let  $\Gamma$  be a set of clauses (over propositions  $P$ ). If  $\Gamma$  is unsatisfiable, then there is a resolution proof of  $\Gamma$

### Question 11

Let  $\Gamma$  be an unsatisfiable set of clauses Show that, if  $\Gamma$  is unsatisfiable, then there is a finite set  $\Gamma_0 \subseteq_{fin} \Gamma$  such that  $\Gamma_0$  is unsatisfiable.

### Question 12

Let  $\Gamma_1, \Gamma_2$  be sets of clauses such that  $\Gamma_1 \subseteq \Gamma_2$ . Show that, if there is a resolution proof of  $\Gamma_1$ , then there is also a resolution proof of  $\Gamma_2$

### Question 13

Inductively show that  $\Delta_i$  is unsatisfiable, for every  $0 \leq i \leq k$ . Also show that  $\Delta_i \neq \emptyset$  for every  $0 \leq i \leq k$

### Question 14

Inductively show that  $\Delta_i$  does not contain the literals  $\{p_1, \neg p_1, \dots, p_i, \neg p_i\}$  for every  $1 \leq i \leq k$

### Question 15

Finish the proof of Theorem 6 by explicitly constructing a resolution proof for  $\Gamma$  and establishing its correctness

### Question 16

Assume that the Hilbert-style Proof System is sound and complete for propositional logic. Use these facts to establish that the Compactness Theorem holds true for propositional logic

### Question 17

Assume that the Resolution Proof System is sound and complete for propositional logic. Use these facts to establish that the Compactness Theorem holds true for propositional logic