

Week 2-3: Propositional Logic

Wang Xiyu

1 Syntax

Definition 1.1. *Syntax decides what are allowed to be written*

1.1 Well-formed Formula (WFF)

A formula ψ is obtained by composing $p \in P$ with $c \in C$, where P is the set of propositions and C is the set of connectors. here we use any set of complete logic, like $C = \{\neg, \vee\}$. This is not a precise definition as counter examples that do not form a valid formula can be formulated based on this, for example $\neg \vee p$ means nothing. To formalize the definition we may attempt to define a language with the following grammar:

$$\Sigma = P \cup C \cup \{(\,,\,)\}$$

$$FORM : \{\psi = p \in P | \neg\psi | \psi \vee \psi | (\psi)\}$$

However this is a recursive definition as the definition of the a formula ψ involves itself. We need to find a declarative way to define this recursive/inductive construction.

Definition 1.2. *FORM is **the smallest** set of strings over Σ where*

1. $P \subseteq FORM$
2. $\forall \psi \in FORM, \neg\psi \in FORM$
3. $\forall \psi_1, \psi_2 \in FORM, \psi_1 \vee \psi_2 \in FORM$

We can see as a construction rule, we are admitting strings into the set of formula, but we need to somewhat impose restriction on what is not acceptable.

1.2 Closed Set

Let's remove the attributive **the smallest** first and define another set that contains all the set that satisfy these rules listed above:

Definition 1.3. *Closed is a set of strings over Σ where*

1. $P \subseteq Closed$
2. $\forall \psi \in Closed, \neg\psi \in Closed$
3. $\forall \psi_1, \psi_2 \in Closed, \psi_1 \vee \psi_2 \in Closed$

1.2.1 All Closed Set

And we define a set ACS (All-closed sets). We can define FORM via the following constriction:

$$FORM = \bigcap ACS$$

Suppose $BCS = \emptyset$ and from definition

$$U_0 = \bigcap BCS = \{w \in \Sigma^* | \forall S \in BCS, w \in S\}$$

However, $BCS = \emptyset$, The statement $\forall S \in BCS, w \in S$ is vacuously true $\forall w \in \Sigma^*$, which means that

$$U_0 = \Sigma^* = \bigcap BCS$$

A contradiction. Therefore we have proved that ACS cannot be empty, thus $\forall P, \exists S$ such that S is a closed set.

1.3 FORM is the minimum of the set of Closed

First let's prove that FORM is closed. It seems to be trivial that the intersection of closed sets is closed but formal proof is still needed. We denote a set from the intersection construction as T_0 .

Lemma 1.4. T_0 is closed

We are proving that the set T_0 we get from the intersection construction still satisfy the rules that define what is closed.

Proof. From rule 1,

$$\begin{aligned} & \forall S \in ACS, P \subseteq S \\ & \rightarrow \forall p \in P, \forall S \in ACS, p \in S \\ & \rightarrow \forall p \in P, p \in \bigcap ACS \\ & \rightarrow P \in T_0 \end{aligned}$$

From rule 2,

$$\begin{aligned} & \forall S \in ACS, T_0 \subseteq S \\ & \text{let } w \in T_0, \forall S \in ACS, w \in S, \neg w \in S \\ & \rightarrow \forall w \in T_0, \neg w \in \bigcap ACS \\ & \rightarrow \forall w \in T_0, \neg w \in T_0 \end{aligned}$$

From rule 3,

$$\begin{aligned} & \forall S \in ACS, T_0 \subseteq S \\ & \text{let } w_1, w_2 \in T_0, \forall S \in ACS, w_1 \vee w_2 \in S \\ & \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in \bigcap ACS \\ & \rightarrow \forall w_1, w_2 \in T_0, w_1 \vee w_2 \in T_0 \end{aligned}$$

□

Lemma 1.5. T_0 is a smallest Closed set

Proof. The proof of minimality is trivial. $T_0 = \bigcap ACS \rightarrow \forall S \in ACS, T_0 \subseteq S$

□

Lemma 1.6. T_0 is *the* smallest Closed set (unique). In other word,

$$\forall T'_0 \in \{T | \forall S \in ACS, T \subseteq S\}, T'_0 = T_0$$

Proof. Suppose otherwise, $\exists T'_0 \in ACS \neq T_0$.

$$\begin{aligned} & T_0, T'_0 \in ACS \rightarrow \forall S \in ACS, T_0 \subseteq S, T'_0 \subseteq S \\ & \rightarrow T_0 \subseteq T'_0 \wedge T'_0 \subseteq T_0 \\ & \rightarrow T_0 = T'_0 \end{aligned}$$

A contradiction.

□

Now we can use the generative rule to define formula, which is called **Backus-Naur form**. Note that a context-free grammar is a type of formal language. Backus Naur form is a specification language for this type of grammar. It is used to describe language syntax.

1.4 Subformula

Definition 1.7. ϕ_1 is a subformula of ϕ_2 , denoted by

$$\phi_1 \preceq \phi_2$$

as a relation is

- *reflexive:* $\forall \phi \in FORM, \phi \preceq \phi$
- *transitive:* $\forall \phi_1, \phi_2, \phi_3 \in FORM, (\phi_1 \preceq \phi_2) \wedge (\phi_2 \preceq \phi_3) \rightarrow \phi_1 \preceq \phi_3$

Inductive definition:

$$\begin{aligned} \forall \phi \in FORM, \phi \preceq (\neg \phi) \\ \forall \phi_1, \phi_2 \in FORM, \phi_1 \preceq (\phi_1 \vee \phi_2) \\ \phi_2 \preceq (\phi_1 \vee \phi_2) \end{aligned}$$

We might intuitively think that subformula can be defined using the notion of substring, since we have been considering formula as concatenation of proposition and connector symbols. However based on our inductive definition, we can come up with the following counter example to a definition using substring: A formula ϕ_1 is a subformula of ϕ_2 if and only if ϕ_1 is a well-formed formula, and is a substring to ϕ_2

$$p \vee q \not\preceq \neg p \vee q$$

What if we tighten the syntax by strictly enforce the use of parentheses whenever we can

2 Semantics

Definition 2.1. *Semantics describe meanings.*

2.1 Truth Assignment/Proposition evaluation

A truth assignment or a valuation of propositions is a function that maps from the set of propositions to true and false.

$$\begin{aligned} v : P \rightarrow \{T, F\} \\ v \in [P \rightarrow \{T, F\}] \end{aligned}$$

Example 2.2. Consider $v = \{p_1 \mapsto T, p_2 \mapsto F\}$, we can derive

$$\begin{aligned} v(p_1) &= T \\ v(p_2) &= F \\ v(\neg p_2) &= T \\ v(p_1 \vee p_2) &= T \text{ etc.} \end{aligned}$$

2.2 Entail/Model

We denote entailment of an evaluation function to formula as

$$v \models \psi$$

We also inductively define the following rule

$$\begin{aligned} \forall p \in P, v \models p &\iff v(p) \\ v \models \psi &\iff v \not\models \neg \psi \\ v \models \psi_1 \vee \psi_2 &\iff (v \models \psi_1) \vee (v \models \psi_2) \end{aligned}$$

2.3 Logical Equivalence

Definition 2.3. *Syntax definition of logical equivalence is as follows:*

$$\phi_1 \equiv \phi_2 \iff \forall v : P \mapsto \{T, F\} (v \models \phi_1), v \models \phi_2$$

Example 2.4.

$$p \vee (\neg p) \equiv q \vee (\neg q)$$

$$\begin{aligned} \forall v \models (p \vee (\neg p)), \\ v \models p \text{ or } v \models (\neg p) \\ v \models p \text{ or } v \not\models p \\ \forall v : P \mapsto \{T, F\}, \end{aligned}$$

TODO

2.4 Satisfiability of Formula

Definition 2.5. *Let $\phi \in FORM$, ϕ is satisfiable*

$$\iff \exists v : P \mapsto \{T, F\}, v \models \phi$$

Definition 2.6. *Tautology: ϕ is a tautology $\iff \forall v, v \models \phi$*

Example 2.7.

$$\phi = \neg(p \vee (\neg p))$$

From example 2.4, we have

$$\begin{aligned} \forall v, v \models (p \vee (\neg p)) \\ \rightarrow \forall v, v \not\models \neg(p \vee (\neg p)) \end{aligned}$$

Thus ϕ is NOT satisfiable.

2.5 Satisfiability of a set of formula

Set Satisfication

Definition 2.8. *Let $\Gamma \subseteq FORM$,*

$$v \models \Gamma \iff \forall \phi \in \Gamma, v \models \phi$$

Set Satisfiability

Definition 2.9. *Let $\Gamma \subseteq FORM$, Γ is satisfiable*

$$\iff \exists v, \forall \phi \in \Gamma, v \models \phi$$

Example 2.10. $\Gamma = \emptyset$, Γ is satisfiable.

Proof. Consider $v : P \mapsto \{T, F\}$, $v(p) = T \forall p \in P$, $\forall \phi \in \Gamma$, $v \models \phi$ is vacuously true. □

Example 2.11. $\Gamma = FORM$, Γ is NOT satisfiable.

Proof. Suppose otherwise,

$$\begin{aligned} \exists v, \forall \phi \in \Gamma, v \models \phi \\ \forall \phi \in \Gamma, (\neg \phi) \in \Gamma \\ v \models (\neg \phi) \end{aligned}$$

A contradiction. □

2.5.1 Relevance

Definition 2.12. Define set $OCCUR(\phi) \subseteq P$, which contains propositions appear in ϕ . Inductively defined as

$$\begin{aligned} OCCUR(\phi) &= \{p\} \iff \phi \equiv p, p \in P \\ OCCUR(\neg\phi) &= OCCUR(\phi) \\ OCCUR(\phi_1 \vee \phi_2) &= OCCUR(\phi_1) \cup OCCUR(\phi_2) \end{aligned}$$

Lemma 2.13. Let $\phi \in FORM$, let v_1, v_2 be valuations over P .

$$\forall p \in OCCUR(\phi), v_1(p) = v_2(p) \rightarrow (v_1 \models \phi \rightarrow v_2 \models \phi)$$

v_1, v_2 can be different valuations over $p' \notin OCCUR(\phi)$

2.6 Modelling Computations as Propositional Logic

Computation problems can be considered as membership check of an encoded problem instance to a language.

$$L = \{p | \langle M, p \rangle\}$$

Here instead of arbitrary encoding, we consider problem encoding with propositional logics only, and solve with SAT solvers by deciding whether the set of formula we form from any instance of the problem is satisfiable. That is

$$Encode : p \mapsto FORM, \forall p, \exists v \models Encode(p) \iff p \in L$$

The rules of such modelling include:

- Propositional logics only
- The final expression must one well formed fomula
- for problem of size n , the size of the formula ϕ must be in $O(poly(n))$

Note that there can be different definition on the size of the

- problem: in unary, binary etc;
- Formula: Whether by the number of nodes in the syntax tree or length of the string **Anything else?**

Example 2.14. *Graph coloring problem: given a undirected graph $G = (V, E)$, whether there exists a k coloring of all vertices such that no neighbouring vertices have the same color. Consider the following construction of a proposition set P :*

$$P = \{p_{v,i} | v \in V, i \in [1, k]\}$$

The valuation of proposition $p_{v,i}$ conceptually represents vertex v is assigned color i .

We can formulate constriants as well formed formula, and use conjunction to connect the formula to model satisfaction of multiple constraints.

Corollary 2.15. $\forall \phi_1, \phi_2 \in FORM, (\phi_1 \wedge \phi_2) \in FORM$

First let us list down the constraints of this problems, including implicit ones in plain words

1. *Neighbouring vertices have different colors.*
2. *All vertices as assigned with exactly one color.*

1. ϕ_{nbr}

First lets formula the constraint between any 2 vertices,

$$\forall(\alpha, \beta) \in E, c(v_\alpha) \neq c(v_\beta)$$

And from our proposition set, $c(v_\gamma) = i \iff \exists v, v \models p_{\gamma,i}$. As such we have

$$\phi_{\alpha,\beta,i} = \neg(p_{\alpha,i} \wedge p_{\beta,i})$$

This models vertex α and vertex β are not colored with color i at the same time. For every color this needs to hold, thus we have

$$\phi_{nbr(\alpha,\beta)} = \bigwedge_{i \in [1,k]} \phi_{\alpha,\beta,i}$$

For every pair of neighbouring vertices this also needs to hold. Therefore we have

$$\phi_{nbr} = \bigwedge_{(\alpha,\beta) \in E} \phi_{nbr(\alpha,\beta)}$$

2. $\phi_{exactly-1}$

To model exactly 1 coloring on each vertex, we can formulate $\phi_{\geq 1}$ and $\phi_{\leq 1}$ first and use conjunction to connect them. Given our propositions, to make sure vertex v is at least assign a color ($\phi_{v,\geq 1}$),

$$\phi_{v,\geq 1} = \bigvee_{i \in [1,k]} p_{v,i}$$

This needs to hold for every vertex, thus we have

$$\phi_{\geq 1} = \bigwedge_{v \in V} \phi_{v,\geq 1}$$

Then we need to model at most 1, (or less than 2) coloring on each vertex. Vertex v is not assignment with both color i and j can be formulated as

$$\phi_{v,i,j,\leq 1} = \neg(p_{v,i} \wedge p_{v,j})$$

This needs to hold for every distinct pair of colors, thus

$$\phi_{v,\leq 1} = \bigwedge_{i \neq j \in [1,k]} \phi_{v,i,j,\leq 1}$$

For all vertices

$$\phi_{\leq} = \bigwedge_{v \in V} \phi_{v,\leq}$$

Finally we have

$$\phi_{exactly-1} = \phi_{\leq 1} \wedge \phi_{\geq 1}$$

And the overall formula for an instance of $G = (V, E)$ becomes

$$\phi = \phi_{nbr} \wedge \phi_{exactly-1}$$

Similar to reduction, we need also now to show that

$$\exists v : P \mapsto \{T, F\}, v \models \phi \iff \langle G = (V, E), k \rangle \in L_{color}$$

TODO

Tutorial

Definition 2.16 (Generic Closure). Let U be a universe of elements and let O be a set of functions whose domain is U and codomain is $\mathcal{P}(U)$. The arity of a function $f \in O$ is denoted by $\text{arity}(f)$. A function f of arity r has type

$$f : U^r \rightarrow \mathcal{P}(U).$$

Let $I \subseteq U$ be an initial set. A set $S \subseteq U$ is said to be (I, O) -closed if:

1. $I \subseteq S$.
2. For every $f \in O$ with $\text{arity}(f) = r$ and for all $e_1, \dots, e_r \in U$, if $e_1, \dots, e_r \in S$, then $f(e_1, \dots, e_r) \subseteq S$.

Lemma 2.17. Let U be a universe, O a set of operators, and $I \subseteq U$ an initial set. There exists a smallest (I, O) -closed set.

Define $\mathcal{P}(U)$ as a collection of all ordering of all elements in the power set $\mathcal{P}(U)$

Q1

Claim: The smallest (I, O) -closed set can be constructed by taking the intersection of all (I, O) -closed set, ie.

$$AIOCS = \{S \subseteq U \mid S \text{ is } (I, O)\text{-closed}\}$$

$$T_0 = \bigcap AIOCS$$

Existence

First we prove the existence of such a set.

Proof. Suppose otherwise,

$$T_0 = \bigcap AIOCS = \emptyset$$

$$T_0 = \{x \in \mathcal{P}(U) \mid \forall S \in AIOCS, x \in S\}$$

$$T_0 = \emptyset \rightarrow x \in S \text{ is vacuously true } \forall x \in \mathcal{P}(U) \rightarrow T_0 = \mathcal{P}(U)$$

contradiction. □

Closed

Now we prove that T_0 is (I, O) -closed.

Proof. Property 1:

$$\begin{aligned} & \forall S \in AIOCS, I \subseteq S \\ & \rightarrow \forall i \in I, \forall S \in AIOCS, i \in S \\ & \rightarrow \forall i \in I, i \in \bigcap AIOCS \\ & \rightarrow \forall i \in I, i \in T_0 \end{aligned}$$

Property 2:

$$\begin{aligned} & \forall S \in AIOCS, T_0 \subseteq S \\ & \text{Let } e_1, e_2, \dots, e_r \in T_0, \forall S \in AIOCS, e_1, e_2, \dots, e_r \in S \\ & \rightarrow \exists f \in O, f(e_1, e_2, \dots, e_r) \in S \\ & \rightarrow \forall e_1, e_2, \dots, e_r \in T_0, \exists f \in O, f(e_1, e_2, \dots, e_r) \in \bigcap AIOCS \\ & \rightarrow f(e_1, e_2, \dots, e_r) \in T_0 \end{aligned}$$

□

Uniqueness

Claim:

$$\forall T'_0 \in \{T \in \mathcal{P}(U) \mid \forall S \in AIOCS, T \subseteq S\}, T'_0 = T_0$$

Proof. Suppose otherwise, $\exists T'_0 \in AIOCS, T'_0 \neq T_0$.

$$\begin{aligned} T_0, T'_0 \in AIOCS &\rightarrow \forall S \in AIOCS, T_0 \subseteq S, T'_0 \subseteq S \\ &\rightarrow T_0 \subseteq T'_0, T'_0 \subseteq T_0 \\ &\rightarrow T_0 = T'_0 \end{aligned}$$

□

Q2

For each of the following strings, prove or disprove that it is a well-formed formula:

1. $w_1 = \varepsilon$ (the empty string).
2. $w_2 = (\neg p)$ where $p \in P$.
3. $w_3 = (p\neg)$ where $p \in P$.

2.1

$$w_1 = \varepsilon \notin FORM$$

Claim: $\forall w \in FORM, |w| > 0$ I'd like to use definition 2.19 and lemma 2.21 which I proved later without using the conclusion from this question.

$$FORM = INDFORM = \bigcup_{i \geq 0} FORM_i$$

Proof. Base case:

$$\begin{aligned} FORM_0 &= P \\ \forall \phi \in FORM_0, \phi &= p \in P \\ &\rightarrow \forall \phi \in FORM_i, |\phi| > 0 \end{aligned}$$

Inductive Hypothesis: Suppose

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| > 0$$

$$\begin{aligned} FORM_{i+1} &= FORM_i \cup \{(\neg\phi) \mid \phi \in FORM_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in FORM_i\} \\ &\quad \forall \psi \in FORM_{i+1}, \exists \phi \in FORM_i, |\psi| = |\phi| \text{ or } |\psi| = |\phi| + 3 \text{ or} \\ &\quad \exists \phi_1, \phi_2 \in FORM_i, |\psi| = |\phi_1| + |\phi_2| + 3 \end{aligned}$$

□

2.2

$$p \in P \rightarrow p \in FORM \rightarrow (\neg p) \in FORM$$

2.3

$(p \neg) \notin FORM$

Proof. Suppose otherwise, $(p \neg \epsilon) \in FORM$

$$\begin{aligned} \forall \phi \in FORM, (\neg \phi) \in FORM \\ \rightarrow \epsilon \in FORM \end{aligned}$$

Contradiction from 2.1 □

Q3

Fix $U = \mathbb{R}$. For each of the following sets S , define a finite initial set $I \subseteq \mathbb{R}$ and a set of operators O with $|O| < 10$ such that the smallest (I, O) -closed set is S .

1. $S = \{0, 1, 2, 3, \dots\}$.
2. $S = \{1, 3, 5, 7, \dots\}$.
3. $S = \mathbb{Q}$.
4. $S = \{\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, \frac{5}{3}, \frac{7}{3}, \dots\}$.

Answer:

1. $\mathbb{N} : I = \{0\}, O = \{f(x) = x + 1\}$
2. $Odd : I = \{1\}, O = \{f(x) = x + 2\}$
3. $\mathbb{Q} : I = \{0\}, O = \{f(x) = -x, f(x) = x + 1, f(x, y) = \begin{cases} \frac{x}{y} & \text{if } y \neq 0 \\ \emptyset & \text{if } y = 0 \end{cases}\}$
4. $\{\frac{x}{3} | x \in \mathbb{N}\} \setminus \mathbb{N} : I = \{\frac{1}{3}, \frac{2}{3}\}, O = \{f(x) = x + 1\}$

Finite Directed Graphs

Definition 2.18 (Graph Reachability). *Let $G = (V, E)$ be a directed graph. A path from s to t is a sequence of vertices $\pi = u_1, \dots, u_k$ such that:*

1. $u_1 = s$,
2. $(u_i, u_{i+1}) \in E$ for all $1 \leq i < k$,
3. $u_k = t$.

The length of π is $k - 1$. We say that t is reachable from s if $s = t$ or such a path exists. The set of ancestors of t is defined as

$$Reach_t = \{v \in V \mid t \text{ is reachable from } v\}.$$

Q4

Let $G = (V, E)$ be a directed graph and let $t \in V$. Fix the universe $U = V$.

1. Define an initial set I (with $|I| \leq 2$) and a unary operator f such that the smallest $(I, \{f\})$ -closed set is the set of vertices reachable from t .
2. Let $d \in \mathbb{N} \setminus \{0\}$ and $r < d$. Define I and a unary operator f such that the smallest $(I, \{f\})$ -closed set is the set of vertices reachable from s via paths of length ℓ with $\ell \equiv r \pmod{d}$.

4.1

$$I = \{t\}, O = \{f(v) = \text{succ}(v)\}$$

4.2

$$I = \{s\} \cup \{v \in V \mid v \in \text{succ}^r(s)\}, O = \{f(v) = \text{succ}^d(v)\}$$

Q5

Let $G = (V, E)$ be a directed graph and fix $U = V \times V$. Define an initial set I with $|I| \leq |V|$ and a binary operator f such that the smallest $(I, \{f\})$ -closed set is

$$\text{AllReachPairs} = \{(u, v) \mid v \text{ is reachable from } u\}.$$

5.1

$$I = \{(v, v) \mid v \in V\}, O = \{f((v_1, v_2), (v_3, v_4)) \mapsto \begin{cases} \{(v_1, v_4)\} & \text{if } v_2 = v_3 \\ \emptyset & \text{if } v_2 \neq v_3 \end{cases}\}$$

Inductive Definitions

Definition 2.19 (Inductive Well-Formed Formulae). *Let P be a set of propositions and $C = \{\neg, \vee\}$. Define sets FORM_i as follows:*

$$\begin{aligned} \text{FORM}_0 &= P, \\ \text{FORM}_{i+1} &= \text{FORM}_i \cup \{(\neg w) \mid w \in \text{FORM}_i\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in \text{FORM}_i\}. \end{aligned}$$

Define

$$\text{INDFORM} = \bigcup_{i \geq 0} \text{FORM}_i.$$

Q6

1. Show that $\text{FORM}_i \subseteq \text{FORM}_{i+1}$ for all i .
2. Prove or disprove that $\text{FORM}_i = \text{FORM}_{i+1}$ for some i .
3. Prove or disprove that FORM_i is (P, C) -closed for all i .
4. Prove or disprove that FORM_i is (P, C) -closed for some i .

6.1

Base case: $\text{FORM}_0 \subseteq \text{FORM}_1$ as

$$\text{FORM}_1 = \text{FORM}_0 \cup \{(\neg w) \mid w \in \text{FORM}_0\} \cup \{(w_1 \vee w_2) \mid w_1, w_2 \in \text{FORM}_0\}$$

Wait this is just trivial

6.2

Claim:

$$\exists i \geq 0, \text{FORM}_i = \text{FORM}_{i+1}$$

It is obvious that $\forall i \geq 0, \text{FORM}_i$ is finite. Suppose

6.3

Claim:

$$\forall i \geq 0, FORM_i \text{ is (P,C)-closed}$$

Proof. Suppose otherwise, consider $FORM_0$,

$$\begin{aligned} FORM_0 &= P \\ \forall p \in P, (\neg p) &\in FORM_0 \end{aligned}$$

Contradiction □

6.4

Claim:

$$\exists i \geq 0, FORM_i \text{ is (P,C)-closed}$$

Proof.

Lemma 2.20.

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

The longest formula in $FORM_i$ has size $2^{i+3} - 3$. Parentheses are counted in string length

Proof. Base case:

$$\begin{aligned} FORM_0 &= P \\ \forall \phi \in FORM_0, |\phi| &= 1 \end{aligned}$$

Inductive hypothesis:

$$\forall i \geq 0, \forall \phi \in FORM_i, |\phi| \leq 2^{i+3} - 3$$

Consider all new strings formed from the longest string $\phi \in FORM_i$,

$$\begin{aligned} |\phi| &= 2^{i+3} - 3 \\ |(\neg\phi)| &= 2^{i+3} \\ |(\phi \vee \phi)| &= 2 \times (2^{i+3} - 3) + 3 = 2^{(i+1)+3} - 3 \end{aligned}$$

□

Suppose $\exists i \geq 0, FORM_i$ is (P,C)-closed, and from lemma 2.20 the longest formula in $FORM_i$ has size $|\phi_{max}| = 2^{i+3} - 3$. $\rightarrow \forall \phi \in FORM_i, (\neg\phi) \in FORM_i \rightarrow \phi'_{max} = (\neg\phi_{max}) \in FORM_i, |\phi'_{max}| > 2^{i+3} - 3$, a contradiction.

□

Lemma 2.21. $FORM = INDFORM$.

Q7

Prove the lemma by showing:

1. $INDFORM$ is (P, C) -closed.
2. For every (P, C) -closed set S , $INDFORM \subseteq S$.

7.1

Claim: $INDFORM$ is (P,C)-closed

Proof. **Claim:** $P \subseteq \text{INDFORM}$

First condition of (P,C)-closed set:

$$\text{FORM}_0 \subseteq \text{INDFORM} \rightarrow P \subseteq \text{INDFORM}$$

Claim: $\forall w \in \text{INDFORM}, (\neg w) \in \text{INDFORM}$

$$\text{INDFORM} = \bigcup_{i \geq 0} \text{FORM}_i \rightarrow \forall i \geq 0, \text{FORM}_i, \text{FORM}_{i+1} \in \text{INDFORM}$$

$$\forall w \in \text{INDFORM}, \exists j \geq 0, w \in \text{FORM}_j \rightarrow (\neg w) \in \text{FORM}_{j+1} \rightarrow (\neg w) \in \text{INDFORM}$$

Claim: $\forall w_1, w_2 \in \text{INDFORM}, (w_1 \vee w_2) \in \text{INDFORM}$

Case 1: $\exists i \geq 0, w_1, w_2 \in \text{FORM}_i$

$$w_1, w_2 \in \text{FORM}_i \rightarrow (w_1 \vee w_2) \in \text{FORM}_{i+1} \rightarrow (w_1 \vee w_2) \in \text{INDFORM}$$

Case 2: $\nexists i \geq 0, w_1, w_2 \in \text{FORM}_i$. WLOG, suppose $w_1 \in \text{FORM}_i, w_2 \in \text{FORM}_j, i < j$

$$\forall w \in \text{FORM}_n, w \in \text{FORM}_{n+1} \text{ By induction } w_1 \in \text{FORM}_i \rightarrow w_1, w_2 \in \text{FORM}_j$$

Back to case 1. □

7.2

$$\forall (\text{P,C})\text{-closed set } S, \text{INDFORM} \subseteq S$$

To prove this, we only need to prove that $\text{INDFORM} \subseteq S_{\min} = \bigcap_{S \text{ is } (\text{P,C})\text{-closed}} S$.

Proof. Base case:

$$\bigcup_{i=0}^0 \text{FORM}_i = \text{FORM}_0 = P \subseteq S_{\min}$$

Inductive hypothesis: Suppose

$$\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min}$$

$$\bigcup_{i=0}^n \text{FORM}_i \subseteq S_{\min} \rightarrow \text{FORM}_n \in S_{\min}$$

$$S_{\min} \text{ is } (\text{P,C})\text{-closed} \rightarrow \forall i \in [1, n], \forall \phi \in \text{FORM}_n, (\neg \phi) \in S_{\min},$$

$$\forall i \in [0, n], \forall \phi_1, \phi_2 \in \text{FORM}_i, (\phi_1 \vee \phi_2) \in S_{\min}$$

$$\rightarrow \bigcup_{i=0}^n \text{FORM}_i \cup \{(\neg \phi) | \forall \phi \in \bigcup_{i=0}^n \text{FORM}_i\} \cup \{(\phi_1 \vee \phi_2) | \forall \phi_1, \phi_2 \in \bigcup_{i=0}^n \text{FORM}_i\} \subseteq S_{\min}$$

$$\bigcup_{i=0}^{n+1} \text{FORM}_i \subseteq S_{\min}$$

□

Definition 2.22 (Alternative Inductive Definition). Define $\text{AltFORM}_0 = P$ and for $i \geq 0$:

$$\begin{aligned}\text{AltFORM}_{i+1} = & \{(\neg w) \mid w \in \text{AltFORM}_i\} \\ & \cup \{(w_1 \vee w_2) \mid w_1 \in \text{AltFORM}_i, \exists j \leq i, w_2 \in \text{AltFORM}_j\} \\ & \cup \{(w_1 \vee w_2) \mid w_2 \in \text{AltFORM}_i, \exists j \leq i, w_1 \in \text{AltFORM}_j\}.\end{aligned}$$

Define

$$\text{AltINDFORM} = \bigcup_{i \geq 0} \text{AltFORM}_i.$$

Question 8

1. Show that $\text{FORM}_i = \bigcup_{j=0}^i \text{AltFORM}_j$ for all i .
2. Show that $\text{INDFORM} = \text{AltINDFORM}$.

8.1

Lemma 2.23.

Proof. Base case:

$$\text{FORM}_0 = \bigcup_{j=0}^0 \text{AltFORM}_j$$

$$\text{FORM}_0 = P = \text{AltFORM}_0$$

Inductive hypothesis: Suppose

$$\forall j \in [0, i], \text{FORM}_i = \bigcup_{k=0}^i \text{AltFORM}_k$$

$$\begin{aligned}\text{FORM}_{i+1} &= \bigcup_{k=0}^i \text{AltFORM}_k \cup \{(\neg \phi) \mid \phi \in \bigcup_{k=0}^i \text{AltFORM}_k\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i \text{AltFORM}_k\} \\ &\supseteq \bigcup_{k=0}^i \text{AltFORM}_k \cup \{(\neg \phi) \mid \phi \in \text{AltFORM}_i\} \cup \{(\phi_1 \vee \phi_2) \mid \phi_1, \phi_2 \in \bigcup_{k=0}^i \text{AltFORM}_k\}\end{aligned}$$

$$\bigcup_{k=0}^{i+1} \text{AltFORM}_k \subseteq \text{FORM}_{i+1}$$

Now we prove the other way:

$$\forall \phi \in \text{FORM}_{i+1}, \phi \in \bigcup_{k=0}^i \text{AltFORM}_k \text{ or } \phi = (\neg \psi), \psi \in \bigcup_{k=0}^i \text{AltFORM}_k \text{ or } \phi = (\psi_1 \vee \psi_2), \psi_1, \psi_2 \in \bigcup_{k=0}^i \text{AltFORM}_k$$

Case 1:

$$\phi \in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k$$

Case 2:

$$\begin{aligned}
\psi &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j \leq i, \psi \in \text{AltFORM}_j \\
\phi &= (\neg\psi), \psi \in \text{AltFORM}_j, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

Case 3:

$$\begin{aligned}
\psi_1, \psi_2 &\in \bigcup_{k=0}^i \text{AltFORM}_k \rightarrow \exists j, l \leq i, \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l \\
\phi &= (\psi_1 \vee \psi_2), \psi_1 \in \text{AltFORM}_j, \psi_2 \in \text{AltFORM}_l, \phi \in \text{AltFORM}_{j+1} \\
&\rightarrow \phi \in \bigcup_{k=0}^{i+1} \text{AltFORM}_k \\
\text{FORM}_{i+1} &\subseteq \bigcup_{k=0}^{i+1} \text{AltFORM}_k \rightarrow \text{FORM}_{i+1} = \bigcup_{k=0}^{i+1} \text{AltFORM}_k
\end{aligned}$$

□

Formation Sequences

Definition 2.24 (Formation Sequence). *A (P, C) -formation sequence is a nonempty finite sequence $\rho = w_1, \dots, w_k$ such that for each i :*

- $w_i \in P$, or
- $w_i = (\neg w_j)$ for some $j < i$, or
- $w_i = (w_j \vee w_k)$ for some $j, k < i$.

The sequence belongs to w if $w_k = w$.

Q9

Give two distinct formation sequences that belong to

$$w = ((\neg p_1) \vee ((\neg p_3) \vee (\neg(p_2 \vee (\neg p_1)))))$$

Two distinct formation sequences ρ_1, ρ_2 are:

ρ_1 :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= (\neg w_1) \\
w_3 &= p_2 \\
w_4 &= (w_3 \vee w_2) \\
w_5 &= (\neg w_4) \\
w_6 &= (p_3) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_7 \vee w_5) \\
w_9 &= (w_2 \vee w_8)
\end{aligned}$$

ρ_2 :

$$\begin{aligned}
w_1 &= p_1 \\
w_2 &= p_2 \\
w_3 &= p_3 \\
w_4 &= (\neg w_3) \\
w_5 &= (\neg w_1) \\
w_6 &= (w_2 \vee w_5) \\
w_7 &= (\neg w_6) \\
w_8 &= (w_4 \vee w_7) \\
w_9 &= (w_5 \vee w_8)
\end{aligned}$$

These are distinct because the components at index $i = 2$ differ $((\neg p_1)$ vs p_2).

Lemma 2.25. $INDFORM = \{w \in \Sigma^* \mid w \text{ is formation-sequence-well-formed}\}$.

Question 10

Prove the lemma by showing:

1. Every formation-sequence-well-formed string is in $INDFORM$.
2. Every $w \in INDFORM$ has a formation sequence.

Lemma 2.26. *Every well-formed formula φ is of one of the following forms:*

1. $\varphi \in P$,
2. $\varphi = (\neg \psi)$,
3. $\varphi = (\psi_1 \vee \psi_2)$.

Question 11

Prove the lemma above using the previous results.