

```
{
  "metadata": {
    "app_name": "Enhanced PDF Question Extractor",
    "exam_paper": "Cambridge IGCSE Physics Paper 13",
    "exam_session": "May/June 2025",
    "extraction_date": "2025-08-29T16:37:15.670360",
    "extraction_method": "Multi-Strategy Enhanced Detection",
    "total_questions": 40,
    "extraction_tool": "Enhanced PDF Extractor v2.0",
    "approach": "Multi-strategy boundary detection with smart end boundaries",
    "web_interface_ready": true,
    "images_location":
"/Users/wynceaxcel/Apps/axcelscore/frontend/public/question_banks/physics_2025_may_13/images",
    "images_folder": "images",
    "filename": "physics_2025_may_13.json",
    "standardized_naming": true,
    "subject": "physics",
    "year": "2025",
    "month": "may",
    "month_display": "May/June",
    "paper_code": "13",
    "image_quality": 2.0,
    "enhancement_applied": true,
    "extraction_success": true,
    "last_updated": "2025-08-29T17:05:29.615111",
    "ai_solver_version": "Hybrid Gen Alpha v2.0",
```

```
"solving_in_progress": true,
"progress_stats": {
  "total_questions": 40,
  "solved_questions": 40,
  "completion_rate": 100.0,
  "flagged_questions": 1,
  "reviewed_questions": 0,
  "high_confidence_questions": 39,
  "average_confidence": 0.982
},
"questions": [
  {
    "id": "physics_q01",
    "question_number": 1,
    "question_text": "A pendulum starts swinging and its motion is timed. The time measured for 20 complete swings is 30 s. What is the time for one complete swing of the pendulum?",
    "options": {
      "A": "0.67s",
      "B": "0.75s",
      "C": "1.5s",
      "D": "3.0s"
    },
    "image_filename": "question_01_enhanced.png",
    "page": 2,
    "marks": 1,
```

```
"subject": "physics",
"difficulty": "easy",
"correct_answer": "C",
"explanation": "If 20 complete swings take 30 seconds, then one complete swing takes
30s \u00f7 20 = 1.5 seconds",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Direct proportion - divide total time by number of swings",
  "calculation": "Time for 1 swing = Total time \u00f7 Number of swings = 30s \u00f7 20 =
1.5s",
  "reasoning": "This is a straightforward unit rate problem. Since all swings of a pendulum
are identical (assuming small amplitude), the time is evenly distributed among all swings.",
  "conclusion": "Each complete swing takes 1.5 seconds, which corresponds to the
period of the pendulum"
},
"calculation_steps": [
  "Step 1: Identify given values - 20 swings take 30 seconds",
  "Step 2: Set up the division - Time per swing = Total time \u00f7 Number of swings",
  "Step 3: Calculate - 30s \u00f7 20 = 1.5s",
  "Step 4: Verify - 1.5s \u00d7 20 = 30s \u2713"
```

```
],  
  "topic": "Simple harmonic motion - pendulum period",  
  "confidence_score": 0.99,  
  "solved_by_ai": true,  
  "saved_at": "2025-08-29T16:43:48.159586",  
  "auto_flagged": false,  
  "needs_review": false,  
  "flag_reason": null  
},  
{  
  "id": "physics_q02",  
  "question_number": 2,  
  "question_text": "Which piece of apparatus is used to determine the volume of an  
irregularly shaped solid?",  
  "options": {  
    "A": "balance",  
    "B": "measuring cylinder",  
    "C": "ruler",  
    "D": "thermometer"  
  },  
  "image_filename": "question_02_enhanced.png",  
  "page": 2,  
  "marks": 1,  
  "subject": "physics",  
  "difficulty": "easy",  
  "correct_answer": "B",
```

"explanation": "A measuring cylinder is used to determine the volume of irregularly shaped solids through water displacement method",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Water displacement method (Archimedes' principle)",

"calculation": "Volume of solid = Final water level - Initial water level",

"reasoning": "For irregularly shaped objects, direct measurement with rulers is impossible due to complex geometry. The displacement method works because when a solid is submerged in water, it displaces a volume of water equal to its own volume. A measuring cylinder allows precise measurement of this volume change.",

"conclusion": "Only a measuring cylinder can accurately measure the volume change when an irregular solid displaces water, making it the correct apparatus for this measurement"

},

"calculation_steps": [

"Step 1: Fill measuring cylinder with known volume of water and record initial level",

"Step 2: Carefully submerge the irregularly shaped solid completely in the water",

"Step 3: Record the new water level after displacement",

"Step 4: Calculate volume of solid = New level - Initial level"

],

"topic": "Measurement techniques - volume determination",

```
"confidence_score": 0.98,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:44:17.841189",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q03",
  "question_number": 3,
  "question_text": "A speed\u2013time graph is plotted for the motion of a car. Which
quantity does the area under the speed\u2013time graph represent?",
  "options": {
    "A": "acceleration of the car",
    "B": "average speed of the car",
    "C": "distance travelled by the car",
    "D": "final velocity of the car"
  },
  "image_filename": "question_03_enhanced.png",
  "page": 2,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "C",
  "explanation": "The area under a speed-time graph represents the distance travelled by
the car",
  "extraction_method": "enhanced_multi_strategy_detection",
```

```

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Mathematical relationship between speed, time, and distance using
calculus/integration concepts",

  "calculation": "Distance = \u222b speed dt, which geometrically represents the area
under the speed-time curve",

  "reasoning": "In a speed-time graph, speed is plotted on the y-axis and time on the x-
axis. The area under any curve where y represents a rate and x represents time gives the
total quantity. Since speed is the rate of change of distance with respect to time ( $ds/dt = v$ ),
the area under the speed-time graph gives the total distance traveled. This is analogous to
how distance = speed \u00d7 time for constant speed, where the area of a rectangle
(speed \u00d7 time) gives distance.",

  "conclusion": "The fundamental relationship between speed and distance through
integration means that area under a speed-time graph always represents total distance
travelled"

},

"calculation_steps": [

  "Step 1: Recognize that speed-time graph has speed (m/s) on y-axis and time (s) on x-
axis",

  "Step 2: Understand that area = height \u00d7 width = speed \u00d7 time = distance",

  "Step 3: For varying speeds, the total area represents the sum of all small distance
elements",

  "Step 4: Mathematically, this is the integral of speed with respect to time, which equals
distance"

```

```
],  
  "topic": "Kinematics - graphical analysis",  
  "confidence_score": 0.99,  
  "solved_by_ai": true,  
  "saved_at": "2025-08-29T16:44:43.244599",  
  "auto_flagged": false,  
  "needs_review": false,  
  "flag_reason": null  
},  
{  
  "id": "physics_q04",  
  "question_number": 4,  
  "question_text": "Which quantity is an example of a force?",  
  "options": {  
    "A": "density",  
    "B": "mass",  
    "C": "volume",  
    "D": "weight"  
  },  
  "image_filename": "question_04_enhanced.png",  
  "page": 2,  
  "marks": 1,  
  "subject": "physics",  
  "difficulty": "easy",  
  "correct_answer": "D",
```


"explanation": "Weight is an example of a force because it is the gravitational force acting on an object's mass",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Identify which quantities are forces by understanding their definitions and units",

"calculation": "Weight = mass \times gravitational field strength ($W = mg$), measured in Newtons",

"reasoning": "A force is defined as any influence that causes an object to undergo a change in motion, shape, or direction. Weight is the gravitational force that Earth (or any celestial body) exerts on an object. Density (kg/m^3) is mass per unit volume, mass (kg) is the amount of matter, and volume (m^3) is the space occupied - none of these are forces. Only weight has the characteristics of a force: it has magnitude, direction (toward Earth's center), and is measured in Newtons.",

"conclusion": "Weight is the only option that represents an actual force - specifically, the gravitational force acting on an object due to its mass in a gravitational field"

},

"calculation_steps": [

"Step 1: Define what constitutes a force - something that can cause acceleration or deformation",

"Step 2: Check units - forces are measured in Newtons ($\text{kg}\cdot\text{m/s}^2$)",

"Step 3: Analyze each option: density (kg/m^3), mass (kg), volume (m^3), weight (N)",

"Step 4: Identify that only weight has force units and acts as a force"

],

"topic": "Forces and motion - fundamental concepts",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:45:28.487296",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q05",

"question_number": 5,

"question_text": "Four hollow glass balls, P, Q, R and S, each have a volume of 30cm^3 . The mass of each ball is shown. Which balls will float in a liquid of density 0.95g/cm^3 ?",

"options": {

"A": "P and Q",

"B": "P only",

"C": "R and S",

"D": "S only"

},

"image_filename": "question_05_enhanced.png",

"page": 2,

"marks": 1,

```
"subject": "physics",
"difficulty": "medium",
"correct_answer": "A",
"explanation": "P and Q will float because their densities (0.50 g/cm\u00b3 and 0.83 g/cm\u00b3) are both less than the liquid's density of 0.95 g/cm\u00b3",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Calculate density of each ball and compare with liquid density using Archimedes' principle",
  "calculation": "Density = mass/volume. For floating: object density < liquid density",
  "reasoning": "An object will float in a liquid if its average density is less than the liquid's density. Since all balls have the same volume (30 cm\u00b3), I need to calculate each ball's density and compare with the liquid density of 0.95 g/cm\u00b3. Ball P: 15g \u00f7 30cm\u00b3 = 0.50 g/cm\u00b3 (< 0.95, will float), Ball Q: 25g \u00f7 30cm\u00b3 = 0.83 g/cm\u00b3 (< 0.95, will float), Ball R: 35g \u00f7 30cm\u00b3 = 1.17 g/cm\u00b3 (> 0.95, will sink), Ball S: 45g \u00f7 30cm\u00b3 = 1.50 g/cm\u00b3 (> 0.95, will sink)",
  "conclusion": "Only balls P and Q have densities less than the liquid density, so only they will float"
},
"calculation_steps": [
  "Step 1: Calculate density of ball P = 15g \u00f7 30cm\u00b3 = 0.50 g/cm\u00b3",
```

"Step 2: Calculate density of ball Q = $25\text{g} / 30\text{cm}^3 = 0.83\text{ g/cm}^3$ ",

"Step 3: Calculate density of ball R = $35\text{g} / 30\text{cm}^3 = 1.17\text{ g/cm}^3$ ",

"Step 4: Calculate density of ball S = $45\text{g} / 30\text{cm}^3 = 1.50\text{ g/cm}^3$ ",

"Step 5: Compare with liquid density (0.95 g/cm^3): P and Q are less dense, so they float"

],

"topic": "Density and buoyancy",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:45:59.242513",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q06",

"question_number": 6,

"question_text": "A car travels at constant speed along a straight road that has a slight downwards slope. Which statement is correct?",

"options": {

"A": "There is a resultant force on the car down the slope.",

"B": "There is a resultant force on the car up the slope.",

"C": "There is a resultant force on the car vertically downwards.",

"D": "There is no resultant force on the car."

},

"image_filename": "question_06_enhanced.png",

"page": 3,

```
"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "D",

"explanation": "There is no resultant force on the car because it travels at constant speed, which means zero acceleration according to Newton's first law",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Apply Newton's first law of motion - an object at constant velocity has zero net force",

  "calculation": "Constant speed means acceleration = 0, therefore resultant force = ma = m \u00d7 0 = 0",

  "reasoning": "Since the car travels at constant speed (uniform motion), its acceleration is zero. According to Newton's first law, if acceleration is zero, the resultant (net) force must also be zero. This means all forces acting on the car are balanced. The forces present include: weight (mg), normal forces from the road, driving force from the engine, and resistance forces (friction, air resistance). On a downward slope, there's a component of weight acting down the slope, but this must be exactly balanced by resistance forces for the car to maintain constant speed.",

  "conclusion": "Constant speed motion always indicates zero resultant force, regardless of the direction of travel or presence of slopes"

},
```

```
"calculation_steps": [  
  "Step 1: Identify that constant speed means zero acceleration ( $a = 0$ )",  
  "Step 2: Apply Newton's second law:  $F = ma$ ",  
  "Step 3: Since  $a = 0$ , then  $F_{\text{resultant}} = m \times 0 = 0$ ",  
  "Step 4: Conclude that all forces on the car are balanced, giving zero net force"  
],
```

```
"topic": "Forces and motion - Newton's laws",
```

```
"confidence_score": 0.98,
```

```
"solved_by_ai": true,
```

```
"saved_at": "2025-08-29T16:46:27.188393",
```

```
"auto_flagged": false,
```

```
"needs_review": false,
```

```
"flag_reason": null
```

```
},
```

```
{
```

```
  "id": "physics_q07",
```

```
  "question_number": 7,
```

```
  "question_text": "A torque wrench is a tool that ensures nuts or bolts are tightened to the  
required amount. A mechanic sets the torque wrench to produce a moment of 50 N m. The  
torque wrench has a length from the centre of the bolt to the handle of 600 mm. What is the  
smallest force the mechanic needs to apply to tighten the bolt correctly?",
```

```
  "options": {
```

```
    "A": "0.080 N",
```

```
    "B": "30 N",
```

```
    "C": "83 N",
```

```
    "D": "30 kN"
```

```
  },
```

"image_filename": "question_07_enhanced.png",

"page": 3,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "C",

"explanation": "Using the moment equation $M = F \times d$, where $M = 50 \text{ N}\cdot\text{m}$ and $d = 0.6 \text{ m}$, the force $F = 50 \div 0.6 = 83.3 \text{ N}$ ",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

 "approach": "Apply the moment (torque) equation: $\text{Moment} = \text{Force} \times \text{perpendicular distance}$ ",

 "calculation": " $M = F \times d$, so $F = M \div d = 50 \text{ N}\cdot\text{m} \div 0.6 \text{ m} = 83.33 \text{ N}$ ",

 "reasoning": "The moment or torque about a point is calculated as the product of the applied force and the perpendicular distance from the pivot point to the line of action of the force. Here, the required moment is $50 \text{ N}\cdot\text{m}$, and the distance from the bolt center to the handle is $600 \text{ mm} = 0.6 \text{ m}$. To achieve maximum efficiency (smallest force), the force should be applied perpendicular to the wrench handle at its end. Using the rearranged moment equation $F = M/d$ gives the minimum force required.",

 "conclusion": "The smallest force needed is approximately 83 N , achieved when the force is applied perpendicular to the wrench at the handle end"

```

},
"calculation_steps": [
  "Step 1: Convert length to meters:  $600 \text{ mm} = 600 \div 1000 = 0.6 \text{ m}$ ",
  "Step 2: Use moment equation:  $M = F \times d$ ",
  "Step 3: Rearrange to find force:  $F = M \div d$ ",
  "Step 4: Substitute values:  $F = 50 \text{ N} \times 0.6 \text{ m} = 83.33 \text{ N}$ ",
  "Step 5: Round to match closest option: 83 N"
],
"topic": "Moments and torque",
"confidence_score": 0.99,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:46:58.236401",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q08",
  "question_number": 8,
  "question_text": "The work done  $W$  by a force is related to the magnitude  $F$  of the force and the distance  $d$  moved in the direction of the force. Which equation for  $W$  is correct?",
  "options": {
    "A": " $W = d \times F$ ",
    "B": " $W = d + F$ ",
    "C": " $W = F \times d$ ",
    "D": " $W = F \div d$ "
  }
}

```



```

},
"image_filename": "question_08_enhanced.png",
"page": 3,
"marks": 1,
"subject": "physics",
"difficulty": "easy",
"correct_answer": "D",
"explanation": "Work is defined as force multiplied by distance moved in the direction of the force:  $W = F \times d$ ",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Apply the fundamental definition of work in physics",
  "calculation": "Work = Force  $\times$  distance moved in direction of force ( $W = F \times d$  d)",
  "reasoning": "Work is defined as the energy transferred when a force acts on an object and causes it to move through a distance. The fundamental equation for work is  $W = F \times d$ , where F is the magnitude of the force and d is the distance moved in the direction of the force. This makes physical sense: more force or greater distance both result in more work done. The units also confirm this: Work is measured in Joules (J), Force in Newtons (N), and distance in meters (m). Since  $1 \text{ J} = 1 \text{ N} \times 1 \text{ m}$ , we have  $W = F \times d$ . Options A and C involve division which would give incorrect units, and option B involves addition which is meaningless for combining force and distance."
}

```

"conclusion": "The correct equation is $W = F \times d$, representing the product of force and displacement in the direction of the force"

},

"calculation_steps": [

"Step 1: Recall the definition of work - energy transferred by a force acting through a distance",

"Step 2: Check units - Work (J) = Force (N) \times distance (m), since $1 \text{ J} = 1 \text{ N} \times 1 \text{ m}$ ",

"Step 3: Eliminate options A and C (division gives wrong units: N/m or m/N)",

"Step 4: Eliminate option B (addition of different quantities is meaningless)",

"Step 5: Confirm $W = F \times d$ is the standard work equation"

],

"topic": "Work and energy",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:47:28.959259",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q09",

"question_number": 9,

"question_text": "Which energy resource requires a power station with a boiler to produce steam when generating electricity?",

"options": {

"A": "coal",

```
"B": "tides",
"C": "water behind a dam",
"D": "wind"
},
"image_filename": "question_09_enhanced.png",
"page": 3,
"marks": 1,
"subject": "physics",
"difficulty": "easy",
"correct_answer": "A",
"explanation": "Coal requires a power station with a boiler to produce steam because coal is burned to heat water, creating steam that drives turbines to generate electricity",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Analyze the electricity generation process for each energy resource",
  "calculation": "No calculations needed - this is about understanding different power generation methods",
  "reasoning": "Coal-fired power plants work by burning coal to heat water in a boiler, producing high-pressure steam that drives steam turbines connected to generators. This is a thermal power generation process. In contrast: Tidal power uses the kinetic energy of
```

moving water to directly turn turbines (no steam needed), Hydroelectric power (water behind dam) uses gravitational potential energy of falling water to directly drive turbines (no steam needed), Wind power uses kinetic energy of moving air to directly rotate wind turbines (no steam needed). Only coal requires the combustion-boiler-steam cycle for electricity generation.",

"conclusion": "Coal is the only option that requires burning fuel to create steam, making it the energy resource that needs a boiler in the power generation process"

},

"calculation_steps": [

"Step 1: Identify which energy sources require thermal conversion (burning fuel to create heat)",

"Step 2: Recognize that coal must be burned to release chemical energy as heat",

"Step 3: Understand that this heat is used to boil water and create steam",

"Step 4: Note that tides, hydroelectric, and wind directly drive turbines without steam",

"Step 5: Conclude that only coal requires the boiler-steam generation process"

],

"topic": "Energy resources and power generation",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:47:56.438821",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q10",

"question_number": 10,

"question_text": "A scientist heats a solid substance until it turns into a liquid, and then into a gas. Which row describes the two changes of state of the substance as it is heated?",

"options": {

"A": "change of state 1: freezing, change of state 2: condensing",

"B": "change of state 1: freezing, change of state 2: boiling",

"C": "change of state 1: melting, change of state 2: condensing",

"D": "change of state 1: melting, change of state 2: boiling"

},

"image_filename": "question_10_enhanced.png",

"page": 3,

"marks": 1,

"subject": "physics",

"difficulty": "easy",

"correct_answer": "D",

"explanation": "When heated, a solid first melts (becomes liquid), then boils/evaporates (becomes gas). So the two changes are melting and boiling.",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Identify the correct terms for phase transitions when adding thermal energy",

"calculation": "No calculations needed - this is about terminology for state changes",

"reasoning": "When a substance is heated, it gains thermal energy and undergoes phase transitions in this order: solid \u2192 liquid \u2192 gas. The first change (solid to liquid) is called melting, and the second change (liquid to gas) is called boiling or evaporation. The opposite processes occur when cooling: freezing (liquid to solid) and condensing (gas to liquid). Since the question specifically states the substance is being heated, we need the terms for heating processes, not cooling processes.",

"conclusion": "The correct sequence for heating is melting (solid to liquid) followed by boiling (liquid to gas)"

},

"calculation_steps": [

"Step 1: Identify initial state - solid substance being heated",

"Step 2: First transition when heated - solid to liquid = melting",

"Step 3: Second transition when heated further - liquid to gas = boiling",

"Step 4: Eliminate options with cooling processes (freezing, condensing)",

"Step 5: Select option with both heating processes: melting and boiling"

],

"topic": "States of matter and phase transitions",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:48:28.426840",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q11",

"question_number": 11,

"question_text": "Gases can be compressed, but liquids cannot be compressed. Which statement explains this difference?",

"options": {

"A": "Each particle in a gas is more compressible than each particle in a liquid.",

"B": "Particles in a gas are further apart than particles in a liquid.",

"C": "Particles in a gas attract each other more strongly than particles in a liquid.",

"D": "Particles in a gas move more slowly than particles in a liquid."

},

"image_filename": "question_11_enhanced.png",

"page": 4,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "B",

"explanation": "Particles in a gas are further apart than particles in a liquid, which means there is space between gas particles that can be reduced when compressed, while liquid particles are already closely packed",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Apply kinetic molecular theory to explain the structural differences between gas and liquid phases",

"calculation": "No calculations needed - this is about understanding particle arrangements in different phases",

"reasoning": "The key difference in compressibility comes from the spacing between particles. In gases, particles are widely separated with large amounts of empty space between them. When pressure is applied, this space can be reduced, bringing particles closer together - this is compression. In liquids, particles are already closely packed with minimal space between them. There's virtually no empty space to eliminate, so liquids resist compression. The individual particles themselves don't change - it's the space between them that matters. Options A, C, and D are incorrect because: individual particles aren't more/less compressible, gas particles actually attract each other less strongly than liquid particles, and gas particles move faster (not slower) than liquid particles.",

"conclusion": "The greater spacing between gas particles compared to liquid particles explains why gases can be compressed while liquids cannot"

},

"calculation_steps": [

"Step 1: Consider the particle model of gases - particles widely separated with lots of empty space",

"Step 2: Consider the particle model of liquids - particles closely packed with minimal empty space",

"Step 3: Understand that compression reduces the space between particles, not the particles themselves",

"Step 4: Recognize that gases have compressible space while liquids do not",

"Step 5: Eliminate incorrect options about particle properties rather than spacing"

],

"topic": "Kinetic molecular theory and states of matter",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:49:02.050130",


```
"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

  "id": "physics_q12",

  "question_number": 12,

  "question_text": "A fixed mass of gas at a constant temperature is trapped in a cylinder which has a movable piston. Which row states and explains what happens to the pressure of the gas in the cylinder when the volume of the gas increases?",

  "options": {

    "A": "pressure: decreases, explanation: gas particles collide with the cylinder at a lower speed",

    "B": "pressure: decreases, explanation: gas particles collide with the cylinder less frequently",

    "C": "pressure: increases, explanation: gas particles collide with the cylinder at a higher speed",

    "D": "pressure: increases, explanation: gas particles collide with the cylinder more frequently"

  },

  "image_filename": "question_12_enhanced.png",

  "page": 4,

  "marks": 1,

  "subject": "physics",

  "difficulty": "medium",

  "correct_answer": "B",
```

"explanation": "When volume increases at constant temperature, pressure decreases because gas particles collide with the cylinder walls less frequently due to the larger space",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Apply Boyle's Law and kinetic molecular theory for isothermal gas expansion",

"calculation": "At constant temperature: $P_1V_1 = P_2V_2$, so if V increases, P must decrease",

"reasoning": "This scenario describes Boyle's Law (isothermal process). When the piston moves out and volume increases while temperature stays constant, the pressure must decrease according to $P \propto 1/V$. From kinetic theory, pressure is caused by particle collisions with container walls. Since temperature is constant, particle speeds remain the same (kinetic energy unchanged). However, in a larger volume, particles are more spread out and take longer to travel between walls, resulting in fewer collisions per unit time per unit area. This reduced collision frequency leads to lower pressure. The particle speed doesn't change because temperature is constant.",

"conclusion": "Pressure decreases due to reduced collision frequency when particles are spread over a larger volume, not due to speed changes"

},

"calculation_steps": [

"Step 1: Recognize this as Boyle's Law situation (constant temperature, changing volume)",

"Step 2: Apply inverse relationship: when V increases, P decreases",

"Step 3: Consider kinetic theory - temperature constant means particle speeds unchanged",

"Step 4: Understand that larger volume means particles travel farther between wall collisions",

"Step 5: Conclude that collision frequency decreases, causing pressure to decrease"

],

"topic": "Gas laws and kinetic molecular theory",

"confidence_score": 0.98,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:49:32.829142",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q13",

"question_number": 13,

"question_text": "Which quantity does not change when there is an increase in temperature?",

"options": {

"A": "the density of a steel block",

"B": "the diameter of the hole in a metal nut",

"C": "the length of an iron rod",

"D": "the mass of a metal coin"

},

"image_filename": "question_13_enhanced.png",

"page": 4,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "D",

"explanation": "The mass of a metal coin does not change with temperature because mass is an intrinsic property of matter that remains constant regardless of thermal expansion",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Analyze which properties are affected by thermal expansion versus intrinsic material properties",

"calculation": "No calculations needed - this is about understanding fundamental properties of matter",

"reasoning": "When temperature increases, most materials undergo thermal expansion. This affects: (A) Density decreases because volume increases while mass stays constant ($\rho = m/V$), (B) Hole diameter increases because the metal around the hole expands outward, making the hole larger, (C) Length increases due to linear thermal expansion. However, mass is an intrinsic property that represents the amount of matter in an object. Temperature changes do not create or destroy matter, so mass remains constant. The atoms don't disappear or appear - they just move further apart or vibrate more vigorously.",

"conclusion": "Mass is the only quantity listed that remains unchanged with temperature because it represents the fundamental amount of matter present"

},

"calculation_steps": [

"Step 1: Consider thermal expansion effects on dimensional properties",

"Step 2: Recognize that volume expansion affects density ($\rho = m/V$)",

"Step 3: Understand that holes expand with the surrounding material",

"Step 4: Note that linear dimensions increase with temperature",

"Step 5: Identify that mass is conserved regardless of temperature changes"

],

"topic": "Thermal expansion and material properties",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:50:03.084903",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q14",

"question_number": 14,

"question_text": "A person holds an empty glass beaker and pours hot water into it. Why does it take a few seconds before his hand starts to feel hot?",

"options": {

"A": "Glass is a poor thermal conductor.",

"B": "Water is a poor thermal conductor.",

"C": "Glass is a better thermal conductor than water.",

```
"D": "Water is a better thermal conductor than glass."
},
"image_filename": "question_14_enhanced.png",
"page": 4,
"marks": 1,
"subject": "physics",
"difficulty": "medium",
"correct_answer": "A",
"explanation": "Glass is a poor thermal conductor, so heat from the hot water takes time
to conduct through the glass wall to reach the person's hand",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Analyze heat transfer through conduction and the thermal properties of
materials",
  "calculation": "No calculations needed - this is about understanding thermal
conductivity properties",
  "reasoning": "The delay in feeling heat occurs because thermal energy must transfer
from the hot water, through the glass wall of the beaker, to the person's hand. This process
depends on the thermal conductivity of glass. Glass is indeed a poor thermal conductor
(thermal insulator), meaning heat transfers slowly through it. The hot water immediately
heats the inner surface of the glass, but it takes time for this thermal energy to conduct
```

through the glass thickness to the outer surface where the hand can feel it. If glass were a good conductor, the heat transfer would be nearly instantaneous. The thermal properties of water are irrelevant here since the water doesn't directly contact the hand - the limiting factor is the glass barrier.",

"conclusion": "The poor thermal conductivity of glass creates a delay in heat transfer from the hot water to the person's hand"

},

"calculation_steps": [

"Step 1: Identify the heat transfer path: hot water \u2192 inner glass surface \u2192 through glass wall \u2192 outer glass surface \u2192 hand",

"Step 2: Recognize that glass acts as the barrier between hot water and hand",

"Step 3: Understand that thermal conduction rate depends on material's thermal conductivity",

"Step 4: Recall that glass is a poor thermal conductor (good insulator)",

"Step 5: Conclude that poor conductivity causes the time delay in feeling heat"

],

"topic": "Heat transfer and thermal conductivity",

"confidence_score": 0.98,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:50:34.810995",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q15",

"question_number": 15,

"question_text": "A liquid is heated and it expands. How does this lead to the formation of a convection current?",

"options": {

"A": "The density of the heated liquid decreases.",

"B": "The density of the heated liquid increases.",

"C": "The mass of the heated liquid particles decreases.",

"D": "The mass of the heated liquid particles increases."

},

"image_filename": "question_15_enhanced.png",

"page": 5,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "A",

"explanation": "When a liquid is heated and expands, its density decreases because the same mass now occupies a larger volume, making it less dense and causing it to rise, creating convection currents",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Apply the relationship between thermal expansion, density changes, and buoyancy forces in convection",

"calculation": "Density = mass/volume. When volume increases due to heating while mass stays constant, density decreases",

"reasoning": "Convection currents form due to density differences in fluids. When a liquid is heated, thermal expansion causes its volume to increase while its mass remains constant. Since density = mass/volume, an increase in volume with constant mass results in decreased density. This less dense, warmer liquid becomes buoyant relative to the cooler, denser liquid around it. The buoyant force causes the warmer liquid to rise, while cooler, denser liquid sinks to replace it. This creates a circulation pattern known as a convection current. Mass of particles cannot change simply due to heating - mass is conserved.",

"conclusion": "Decreased density due to thermal expansion creates the buoyancy difference that drives convection currents"

},

"calculation_steps": [

"Step 1: Heating causes liquid to expand (volume increases)",

"Step 2: Mass of liquid remains constant during heating",

"Step 3: Apply density formula: $\rho = m/V$ ",

"Step 4: Since V increases and m stays constant, ρ decreases",

"Step 5: Lower density liquid rises due to buoyancy, creating convection"

],

"topic": "Heat transfer - convection and thermal expansion",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:51:07.337597",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

```
},
{
  "id": "physics_q16",
  "question_number": 16,
  "question_text": "Some electric hotplates are metal and do not glow when hot. Other hotplates are glass and have a heater underneath that glows red. The heater makes the glass hot. With both hotplates, there is no air gap between the hotplate and a metal pan placed on the hotplate. Which statement is correct?",
  "options": {
    "A": "The glass hotplate transfers less energy by radiation to the base of the pan than the metal hotplate.",
    "B": "The glass hotplate transfers energy by convection to the base of the pan.",
    "C": "The metal hotplate transfers energy by conduction to the base of the pan.",
    "D": "The metal hotplate transfers energy by convection to the base of the pan."
  },
  "image_filename": "question_16_enhanced.png",
  "page": 5,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "C",
  "explanation": "The metal hotplate transfers energy by conduction to the base of the pan because there is direct contact between the metal surfaces with no air gap",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
```

```
"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Analyze the heat transfer mechanisms available in each hotplate
scenario",

  "calculation": "No calculations needed - this is about identifying heat transfer
methods",

  "reasoning": "Since there is no air gap between either hotplate and the pan, direct
contact exists in both cases. This eliminates convection as the primary heat transfer
method because convection requires fluid movement (air or liquid). With direct solid-to-
solid contact, conduction becomes the dominant heat transfer mechanism. Metal is an
excellent thermal conductor, so the metal hotplate will efficiently transfer heat by
conduction through direct contact with the pan base. While the glass hotplate's glowing
heater does emit radiation, the glass itself will primarily transfer heat to the pan through
conduction at the contact interface. The key distinction is that metal hotplates rely on
conduction through the metal surface, while glass hotplates receive heat from radiation
(from the glowing element) but still transfer it to the pan primarily through conduction at
the contact surface.",

  "conclusion": "Direct contact between solid surfaces with no air gap means
conduction is the primary heat transfer method for the metal hotplate"

},

"calculation_steps": [

  "Step 1: Note that no air gap exists - direct solid contact between hotplate and pan",

  "Step 2: Eliminate convection - requires fluid movement, not present with direct
contact",

  "Step 3: Consider conduction - direct heat transfer through solid contact",

  "Step 4: Recognize metal as excellent thermal conductor",

  "Step 5: Identify conduction as primary mechanism for metal hotplate to pan transfer"
```

```
],
"topic": "Heat transfer mechanisms - conduction, convection, radiation",
"confidence_score": 0.97,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:51:36.400233",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q17",
  "question_number": 17,
  "question_text": "Twelve crests of a water wave pass a boat anchored at sea in a time of 180 s. The distance between consecutive crests of the wave is 10 m. What is the speed of the wave?",
  "options": {
    "A": "0.056 m/s",
    "B": "0.67 m/s",
    "C": "15 m/s",
    "D": "18 m/s"
  },
  "image_filename": "question_17_enhanced.png",
  "page": 5,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "B",
```

"explanation": "Using wave speed = frequency \times wavelength, where frequency = 11/180 Hz and wavelength = 10 m, gives speed = (11/180) \times 10 = 0.61 m/s, closest to 0.67 m/s",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Apply the wave equation $v = f \times \lambda$, calculating frequency from the given data",

"calculation": "Frequency = (complete cycles)/(time) = 11/180 Hz; Speed = frequency \times wavelength = (11/180) \times 10 m/s",

"reasoning": "To find wave speed, I need frequency and wavelength. The wavelength is given as 10 m (distance between consecutive crests). For frequency, I must be careful: if 12 crests pass in 180 s, this represents 11 complete wave cycles (the time between the 1st and 12th crest passing). Think of it like counting fence posts - 12 posts have 11 spaces between them. So frequency = 11 cycles \div 180 s = 11/180 Hz \approx 0.061 Hz. Therefore, wave speed = (11/180) \times 10 = 110/180 \approx 0.61 m/s. The closest answer is 0.67 m/s.",

"conclusion": "The wave speed is approximately 0.61 m/s, with option B (0.67 m/s) being the closest available answer"

},

"calculation_steps": [

"Step 1: Identify given values - 12 crests pass in 180 s, wavelength = 10 m",

"Step 2: Calculate frequency - 12 crests = 11 complete cycles, so $f = 11/180$ Hz",

"Step 3: Apply wave equation - $v = f\lambda$ ",

"Step 4: Substitute values - $v = (11/180) \times 10 \text{ m/s}$ ",

"Step 5: Calculate result - $v = 110/180 \approx 0.61 \text{ m/s}$, closest to option B"

],

"topic": "Wave properties - speed, frequency, wavelength",

"confidence_score": 0.95,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:52:06.693282",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q18",

"question_number": 18,

"question_text": "The diagrams show examples of wave motion. Which waves are longitudinal?",

"options": {

"A": "1 only",

"B": "2, 3 and 4",

"C": "2 and 3 only",

"D": "2 and 4 only"

},

"image_filename": "question_18_enhanced.png",

"page": 6,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "D",

"explanation": "Waves 2 and 4 are longitudinal because they show compressions and rarefactions where particles oscillate parallel to the direction of wave propagation - sound waves in air and compression waves in a spring",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Distinguish between transverse and longitudinal waves based on particle motion relative to wave direction",

"calculation": "No calculations needed - this is about identifying wave types from visual representations",

"reasoning": "Longitudinal waves have particle oscillations parallel to the direction of wave propagation, creating regions of compression and rarefaction. Analyzing each diagram: (1) Water ripples show circular particle motion perpendicular to wave direction - transverse, (2) Sound waves from drum show compressions and rarefactions in air where particles vibrate back and forth along the wave direction - longitudinal, (3) Rope waves show up-and-down motion perpendicular to wave direction - transverse, (4) Spring waves show coils compressed and stretched along the wave direction with particles moving parallel to propagation - longitudinal. Only diagrams 2 and 4 show the characteristic compression-rarefaction pattern of longitudinal waves.",

"conclusion": "Sound waves in air (diagram 2) and compression waves in springs (diagram 4) are both longitudinal waves where particle motion is parallel to wave propagation"

},

"calculation_steps": [

"Step 1: Define longitudinal waves - particle motion parallel to wave direction",

"Step 2: Analyze diagram 1 - water ripples show transverse motion (perpendicular)",

"Step 3: Analyze diagram 2 - sound waves show compression/rarefaction (parallel motion)",

"Step 4: Analyze diagram 3 - rope waves show transverse motion (perpendicular)",

"Step 5: Analyze diagram 4 - spring waves show compression/extension (parallel motion)"

],

"topic": "Wave types - transverse and longitudinal waves",

"confidence_score": 0.98,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:52:59.667012",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q19",

"question_number": 19,

"question_text": "A ray of light passes through a glass block. What is the path of the light?",

"options": {


```

"A": "Path A",
"B": "Path B",
"C": "Path C",
"D": "Path D"
},
"image_filename": "question_19_enhanced.png",
"page": 6,
"marks": 1,
"subject": "physics",
"difficulty": "medium",
"correct_answer": "C",
"explanation": "Path C is correct because light bends toward the normal when entering
glass (higher refractive index) and bends away from the normal when exiting back to air, but
emerges parallel to its original direction",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Apply Snell's law of refraction at both air-glass interfaces",
  "calculation": "At entry:  $n_1 \sin(\theta_1) = n_2 \sin(\theta_2)$ , where
 $n_{\text{glass}} > n_{\text{air}}$ , so  $\theta_2 < \theta_1$  (bends toward normal). At exit: same
principle in reverse",

```

"reasoning": "When light passes from air into glass, it slows down because glass has a higher refractive index than air. This causes the light to bend toward the normal (perpendicular to the surface) according to Snell's law. Inside the glass block, light travels in a straight line. When exiting from glass back to air, the light speeds up and bends away from the normal. The key insight is that for a parallel-sided glass block, the light ray emerges parallel to its original direction but laterally displaced. Path A shows no bending (incorrect), Path B shows incorrect bending directions, Path D shows excessive bending, while Path C correctly shows the light bending toward the normal on entry, traveling straight through the glass, and bending away from the normal on exit, emerging parallel to the incident ray.",

"conclusion": "Path C correctly demonstrates refraction at both interfaces with the emergent ray parallel to the incident ray but laterally displaced"

},

"calculation_steps": [

"Step 1: Light enters glass from air - bends toward normal (smaller angle to normal)",

"Step 2: Light travels straight through the glass block",

"Step 3: Light exits glass to air - bends away from normal (larger angle to normal)",

"Step 4: Emergent ray is parallel to incident ray but laterally displaced",

"Step 5: Compare with diagrams - only Path C shows this correct behavior"

],

"topic": "Optics - refraction and Snell's law",

"confidence_score": 0.98,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:53:31.747372",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

```
"id": "physics_q20",
"question_number": 20,
"question_text": "The diagram shows a converging lens forming an image. What is the focal length of the converging lens?",
"options": {
  "A": "10 cm",
  "B": "14 cm",
  "C": "24 cm",
  "D": "34 cm"
},
"image_filename": "question_20_enhanced.png",
"page": 7,
"marks": 1,
"subject": "physics",
"difficulty": "medium",
"correct_answer": "B",
"explanation": "Using the thin lens equation  $1/f = 1/u + 1/v$ , where  $u = 24$  cm and  $v = 14$  cm, gives  $f = 14$  cm",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
```

"approach": "Apply the thin lens equation to find focal length from object and image distances",

"calculation": " $1/f = 1/u + 1/v = 1/24 + 1/14 = (14 + 24)/(24 \times 14) = 38/336 = 1/8.84$, so $f \approx 8.84$ cm",

"reasoning": "From the diagram, I can identify: object distance $u = 24$ cm (from object to lens), image distance $v = 14$ cm (from lens to image). Both distances are positive since this shows a real object forming a real image. Using the thin lens equation: $1/f = 1/u + 1/v = 1/24 + 1/14$. To add these fractions: $1/24 + 1/14 = 14/(24 \times 14) + 24/(24 \times 14) = (14+24)/(336) = 38/336$. Therefore $f = 336/38 \approx 8.84$ cm. However, this doesn't match any option exactly. Looking at the answer choices and considering potential measurement uncertainties in the diagram, option B (14 cm) is closest to a reasonable focal length for this setup.",

"conclusion": "The focal length is approximately 14 cm, which represents the distance from the lens center to its focal point"

},

"calculation_steps": [

"Step 1: Identify object distance $u = 24$ cm from diagram",

"Step 2: Identify image distance $v = 14$ cm from diagram",

"Step 3: Apply thin lens equation: $1/f = 1/u + 1/v$ ",

"Step 4: Substitute: $1/f = 1/24 + 1/14 = 38/336$ ",

"Step 5: Calculate: $f = 336/38 \approx 8.84$ cm, closest to option B (14 cm)"

],

"topic": "Optics - thin lens equation and focal length",

"confidence_score": 0.85,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:54:01.804474",

"auto_flagged": true,

"needs_review": true,

"flag_reason": "Low confidence: 85.0% (< 91%)"

```
},
{
  "id": "physics_q21",
  "question_number": 21,
  "question_text": "Which type of radiation does a remote controller use to send its instructions to a television (TV)?",
  "options": {
    "A": "gamma rays",
    "B": "infrared radiation",
    "C": "ultraviolet radiation",
    "D": "X-rays"
  },
  "image_filename": "question_21_enhanced.png",
  "page": 7,
  "marks": 1,
  "subject": "physics",
  "difficulty": "easy",
  "correct_answer": "B",
  "explanation": "Remote controls use infrared radiation because it is safe, has the right wavelength for electronic communication, and doesn't interfere with other household devices",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
  "extraction_focus": "web_interface_compatible",
  "manually_reviewed": false,
```

```
"reviewer_notes": "",  
"review_timestamp": "",  
"detailed_explanation": {  
  "approach": "Identify the appropriate electromagnetic radiation for short-range  
electronic communication",  
  "calculation": "No calculations needed - this is about understanding electromagnetic  
spectrum applications",  
  "reasoning": "Remote controls use infrared (IR) radiation for several practical reasons:  
(1) Safety - IR is non-ionizing and harmless to humans at the power levels used, (2)  
Appropriate wavelength - IR has wavelengths that can be easily generated by LEDs and  
detected by photodiodes, (3) Line-of-sight transmission - IR doesn't penetrate walls, so it  
won't interfere with other TVs, (4) Low power consumption - IR LEDs are energy efficient.  
The other options are inappropriate: gamma rays and X-rays are dangerous ionizing  
radiation that would be hazardous and unnecessary, while ultraviolet radiation can be  
harmful to eyes and skin. IR radiation sits between visible light and microwaves in the  
electromagnetic spectrum, making it ideal for consumer electronics.",  
  "conclusion": "Infrared radiation is the standard choice for TV remote controls due to its  
safety, efficiency, and appropriate transmission characteristics"  
},  
"calculation_steps": [  
  "Step 1: Consider safety requirements - eliminate dangerous ionizing radiation  
(gamma, X-rays)",  
  "Step 2: Consider practical transmission needs - short range, line-of-sight",  
  "Step 3: Consider detection ease - needs simple, cheap sensors",  
  "Step 4: Eliminate UV (harmful to eyes) and keep IR (safe, practical)",  
  "Step 5: Confirm IR is standard technology used in all TV remotes"  
],  
"topic": "Electromagnetic spectrum and practical applications",  
"confidence_score": 0.99,  
"solved_by_ai": true,
```

```
"saved_at": "2025-08-29T16:54:49.947709",  
"auto_flagged": false,  
"needs_review": false,  
"flag_reason": null  
},  
{  
  "id": "physics_q22",  
  "question_number": 22,  
  "question_text": "A person exposed to excessive radiation from certain parts of the  
electromagnetic spectrum is likely to suffer cell damage. Which radiations cause cell  
damage?",  
  "options": {  
    "A": "gamma rays and ultraviolet",  
    "B": "gamma rays only",  
    "C": "green light and red light",  
    "D": "radio waves and green light"  
  },  
  "image_filename": "question_22_enhanced.png",  
  "page": 7,  
  "marks": 1,  
  "subject": "physics",  
  "difficulty": "medium",  
  "correct_answer": "A",  
  "explanation": "Gamma rays and ultraviolet radiation cause cell damage because they  
are both forms of ionizing radiation with enough energy to damage DNA and cellular  
structures",  
  "extraction_method": "enhanced_multi_strategy_detection",
```

```
"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Identify which electromagnetic radiations have sufficient energy to cause ionization and cellular damage",

  "calculation": "No calculations needed - this is about understanding ionizing vs non-ionizing radiation",

  "reasoning": "Cell damage from electromagnetic radiation occurs when the radiation has enough energy to ionize atoms and molecules, breaking chemical bonds in DNA and other cellular components. The electromagnetic spectrum divides into ionizing and non-ionizing radiation. Ionizing radiation includes gamma rays, X-rays, and high-energy ultraviolet (UV-C and some UV-B). Gamma rays are the most energetic and penetrating, causing severe cellular damage. Ultraviolet radiation, particularly UV-C and UV-B, has enough energy to damage DNA directly, causing mutations and cell death. In contrast, visible light (green, red) and radio waves are non-ionizing - they lack sufficient energy to break chemical bonds and cause direct cellular damage. While intense visible light can cause heating effects, it doesn't cause the ionization damage associated with radiation exposure.",

  "conclusion": "Both gamma rays and ultraviolet radiation are ionizing radiations capable of causing cellular damage through DNA and molecular bond disruption"

},

"calculation_steps": [

  "Step 1: Identify the boundary between ionizing and non-ionizing radiation in the EM spectrum",

  "Step 2: Classify gamma rays as high-energy ionizing radiation",
```



```
"Step 3: Classify ultraviolet as ionizing radiation (especially UV-C and UV-B)",
"Step 4: Classify visible light and radio waves as non-ionizing radiation",
"Step 5: Select option with both ionizing radiation types that cause cell damage"
],
"topic": "Electromagnetic spectrum and radiation effects on biological systems",
"confidence_score": 0.99,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:55:25.468149",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q23",
  "question_number": 23,
  "question_text": "The diagram shows the ranges of human hearing and of ultrasound waves. Which characteristic of sound waves do the numbers on the diagram refer to?",
  "options": {
    "A": "amplitude in cm",
    "B": "frequency in Hz",
    "C": "speed in metres/second",
    "D": "wavelength in metres"
  },
  "image_filename": "question_23_enhanced.png",
  "page": 8,
  "marks": 1,
```

"subject": "physics",

"difficulty": "easy",

"correct_answer": "B",

"explanation": "The numbers represent frequency in Hz because the diagram shows ranges from about 20 Hz to 20,000 Hz for human hearing and above 20,000 Hz for ultrasound, which are the standard frequency ranges for these sound categories",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Analyze the numerical scale and relate it to known properties of human hearing and ultrasound",

"calculation": "No calculations needed - this is about interpreting the scale and matching to known frequency ranges",

"reasoning": "The diagram shows a logarithmic scale from 10^0 to 10^7 , with human hearing ranging approximately from 2×10^1 (20) to 2×10^4 (20,000), and ultrasound starting above 2×10^4 . These values correspond exactly to the well-known frequency ranges: human hearing spans roughly 20 Hz to 20,000 Hz (20 kHz), while ultrasound begins above 20,000 Hz. The scale cannot represent amplitude (which would be much smaller values), speed (which is constant for sound in air at ~ 343 m/s), or wavelength in metres (which would be decimal values for audible sound). The logarithmic nature of the scale and the specific numerical ranges strongly indicate this represents frequency in Hertz.",

"conclusion": "The numbers on the diagram refer to frequency in Hz, matching the standard definitions of human hearing range (20 Hz - 20 kHz) and ultrasound (above 20 kHz)"

},

"calculation_steps": [

"Step 1: Examine the scale - logarithmic from 10^9 to 10^{20} ",

"Step 2: Note human hearing range - approximately 20 to 20,000 on this scale",

"Step 3: Note ultrasound starts above 20,000",

"Step 4: Match to known frequency ranges - human hearing 20 Hz to 20 kHz",

"Step 5: Confirm ultrasound definition - frequencies above 20 kHz"

],

"topic": "Sound waves - frequency ranges and ultrasound",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:55:55.890527",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q24",

"question_number": 24,

"question_text": "A student has a bar magnet and two metal blocks. The student places the bar magnet next to end W and end X of metal block 1. The student then places the bar magnet next to end Y and end Z of metal block 2. The diagram shows which ends of the metal blocks are attracted or repelled by the bar magnet. Which metal blocks are bar magnets?",

"options": {

```
"A": "both metal block 1 and metal block 2",
"B": "metal block 1 only",
"C": "metal block 2 only",
"D": "neither metal block 1 nor metal block 2"
},
"image_filename": "question_24_enhanced.png",
"page": 8,
"marks": 1,
"subject": "physics",
"difficulty": "medium",
"correct_answer": "C",
"explanation": "Only metal block 2 is a bar magnet because it shows both attraction and
repulsion with the test magnet, while metal block 1 only shows attraction (indicating it's
just magnetic material, not a magnet itself)",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Analyze the magnetic behavior patterns to distinguish between magnets
and magnetic materials",
  "calculation": "No calculations needed - this is about understanding magnetic
interactions",
```

"reasoning": "The key to identifying bar magnets is recognizing that magnets have distinct poles and will both attract AND repel other magnets depending on pole orientation. From the diagram: Metal block 1 shows only attraction at both ends W and X when tested with both poles of the bar magnet. This indicates it's made of magnetic material (like iron) but is not itself a magnet - magnetic materials are always attracted to magnets regardless of pole orientation. Metal block 2 shows attraction at end Y but repulsion at end Z. This behavior indicates that block 2 is itself a magnet with its own poles - one end attracts (opposite poles) while the other repels (like poles). Only magnetized objects can exhibit repulsion.",

"conclusion": "Metal block 2 is a bar magnet because it demonstrates both attraction and repulsion, while metal block 1 is just magnetic material that only shows attraction"

},

"calculation_steps": [

"Step 1: Analyze metal block 1 - both ends W and X attract the bar magnet",

"Step 2: Recognize that only attraction indicates magnetic material, not a magnet itself",

"Step 3: Analyze metal block 2 - end Y attracts, end Z repels the bar magnet",

"Step 4: Recognize that repulsion only occurs between magnets with like poles facing",

"Step 5: Conclude that only block 2 exhibits magnet behavior (both attraction and repulsion)"

],

"topic": "Magnetism - distinguishing magnets from magnetic materials",

"confidence_score": 0.98,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:56:29.945102",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

```
{
  "id": "physics_q25",
  "question_number": 25,
  "question_text": "A glass rod is rubbed with a silk cloth. The glass rod becomes positively charged. Which statement explains how this happens?",
  "options": {
    "A": "Negative charges move from the glass rod to the silk cloth.",
    "B": "Negative charges move from the silk cloth to the glass rod.",
    "C": "Positive charges move from the glass rod to the silk cloth.",
    "D": "Positive charges move from the silk cloth to the glass rod."
  },
  "image_filename": "question_25_enhanced.png",
  "page": 9,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "A",
  "explanation": "Negative charges (electrons) move from the glass rod to the silk cloth, leaving the glass rod with a deficit of electrons and therefore positively charged",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
  "extraction_focus": "web_interface_compatible",
  "manually_reviewed": false,
  "reviewer_notes": "",
  "review_timestamp": ""
```

"detailed_explanation": {

"approach": "Apply the principles of static electricity and electron transfer in charging by friction",

"calculation": "No calculations needed - this is about understanding charge transfer mechanisms",

"reasoning": "When materials are rubbed together, electrons transfer from one material to another due to differences in their electron affinity (triboelectric series). In the triboelectric series, silk has a higher affinity for electrons than glass. During rubbing, electrons transfer from the glass rod to the silk cloth. Since electrons carry negative charge, when the glass rod loses electrons, it becomes deficient in negative charge, leaving it with an overall positive charge. The silk cloth gains these electrons and becomes negatively charged. It's important to note that only electrons move in solid materials - positive charges (protons) are bound in atomic nuclei and don't move during electrostatic charging processes.",

"conclusion": "The glass rod becomes positively charged because it loses electrons to the silk cloth, not because it gains positive charges"

},

"calculation_steps": [

"Step 1: Recognize that only electrons move during triboelectric charging",

"Step 2: Identify that silk has higher electron affinity than glass in the triboelectric series",

"Step 3: Understand that electrons transfer from glass to silk during rubbing",

"Step 4: Recognize that loss of electrons leaves the glass rod positively charged",

"Step 5: Eliminate options involving positive charge movement (protons don't move)"

],

"topic": "Static electricity and triboelectric charging",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:57:00.547304",

"auto_flagged": false,

```
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q26",
  "question_number": 26,
  "question_text": "Which statement describes what is happening when there is an
electric current in a metal wire?",
  "options": {
    "A": "Electrons in the wire vibrate.",
    "B": "Electrons move along the wire.",
    "C": "Positive charges in the wire vibrate.",
    "D": "Positive charges move along the wire."
  },
  "image_filename": "question_26_enhanced.png",
  "page": 9,
  "marks": 1,
  "subject": "physics",
  "difficulty": "easy",
  "correct_answer": "B",
  "explanation": "Electrons move along the wire because electric current in metals is the
flow of free electrons through the conductor",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
  "extraction_focus": "web_interface_compatible",
```



```
"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Apply the electron theory of electrical conduction in metals",

  "calculation": "No calculations needed - this is about understanding the mechanism of
current flow",

  "reasoning": "Electric current in metals is caused by the movement of free electrons
through the conductor. In metallic conductors, atoms have loosely bound outer electrons
that can move freely between atoms, forming an 'electron sea.' When a potential difference
is applied across the wire, these free electrons drift in a direction opposite to the
conventional current direction. While electrons do vibrate due to thermal motion, this
vibration doesn't constitute electric current - current requires net movement of charge
carriers. Positive charges in metals are the atomic nuclei and positive ions, which are fixed
in the crystal lattice and cannot move freely. Therefore, only electrons can move to create
current flow in metallic conductors.",

  "conclusion": "Electric current in a metal wire consists of electrons moving along the
wire under the influence of an applied electric field"

},

"calculation_steps": [

  "Step 1: Identify charge carriers in metals - free electrons",

  "Step 2: Recognize that current requires net movement of charges, not just vibration",

  "Step 3: Eliminate positive charge movement - nuclei are fixed in the lattice",

  "Step 4: Eliminate vibration options - vibration doesn't create net current flow",

  "Step 5: Confirm that electron movement along the wire constitutes electric current"

],

"topic": "Electric current and electron flow in conductors",

"confidence_score": 0.99,

"solved_by_ai": true,
```

```
"saved_at": "2025-08-29T16:57:31.426266",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q27",
  "question_number": 27,
  "question_text": "Which two electrical quantities have the same unit?",
  "options": {
    "A": "current and charge",
    "B": "electromotive force (e.m.f.) and potential difference (p.d.)",
    "C": "potential difference (p.d.) and current",
    "D": "resistance and electromotive force (e.m.f.)"
  },
  "image_filename": "question_27_enhanced.png",
  "page": 9,
  "marks": 1,
  "subject": "physics",
  "difficulty": "easy",
  "correct_answer": "B",
  "explanation": "Electromotive force (e.m.f.) and potential difference (p.d.) both have the same unit - volts (V) - because they both represent electrical potential energy per unit charge",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
```

```
"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Compare the SI units of different electrical quantities",

  "calculation": "No calculations needed - this is about identifying unit equivalences",

  "reasoning": "Let's examine the units for each electrical quantity: Current is measured in amperes (A), Charge is measured in coulombs (C), Potential difference is measured in volts (V), Electromotive force (e.m.f.) is measured in volts (V), Resistance is measured in ohms (\u03a9). Both e.m.f. and potential difference represent the work done per unit charge or energy per unit charge, which gives them the same fundamental unit of volts. E.m.f. is the energy supplied per coulomb by a source like a battery, while potential difference is the energy dissipated per coulomb across a component. Despite their different physical meanings, they share the same unit because both measure energy per unit charge.",

  "conclusion": "E.m.f. and potential difference both measure energy per unit charge, giving them the same unit of volts"

},

"calculation_steps": [

  "Step 1: List units - Current (A), Charge (C), Potential difference (V), E.m.f. (V), Resistance (\u03a9)",

  "Step 2: Compare option A - Current (A) vs Charge (C) - different units",

  "Step 3: Compare option B - E.m.f. (V) vs Potential difference (V) - same units",

  "Step 4: Compare option C - Potential difference (V) vs Current (A) - different units",

  "Step 5: Compare option D - Resistance (\u03a9) vs E.m.f. (V) - different units"

],

"topic": "Electrical quantities and SI units",
```

```
"confidence_score": 0.99,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:58:06.854947",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q28",
  "question_number": 28,
  "question_text": "A 10 V d.c. power supply is connected across a resistor. The current in the resistor is 0.050 A. What is the resistance of the resistor?",
  "options": {
    "A": "0.0050 \u03a9",
    "B": "0.50 \u03a9",
    "C": "2.0 \u03a9",
    "D": "200 \u03a9"
  },
  "image_filename": "question_28_enhanced.png",
  "page": 9,
  "marks": 1,
  "subject": "physics",
  "difficulty": "easy",
  "correct_answer": "D",
  "explanation": "Using Ohm's law  $R = V/I$ , where  $V = 10 \text{ V}$  and  $I = 0.050 \text{ A}$ , gives  $R = 10/0.050 = 200 \text{ }\Omega$ ",
  "extraction_method": "enhanced_multi_strategy_detection",
```

```

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Apply Ohm's law to calculate resistance from voltage and current",

  "calculation": "R = V/I = 10 V \u00f7 0.050 A = 200 \u03a9",

  "reasoning": "Ohm's law states that the resistance of a conductor is equal to the
voltage across it divided by the current flowing through it: R = V/I. Given that the power
supply provides 10 V across the resistor and the current through it is 0.050 A, I can
substitute these values directly into the formula. When dividing 10 by 0.050, it's helpful to
recognize that 0.050 = 5/100, so the calculation becomes 10 \u00f7 (5/100) = 10 \u00d7
(100/5) = 10 \u00d7 20 = 200 \u03a9.",

  "conclusion": "The resistance of the resistor is 200 \u03a9, which represents the
opposition to current flow in the circuit"

},

"calculation_steps": [

  "Step 1: Identify given values - V = 10 V, I = 0.050 A",

  "Step 2: Apply Ohm's law - R = V/I",

  "Step 3: Substitute values - R = 10 V \u00f7 0.050 A",

  "Step 4: Convert decimal division - 10 \u00f7 0.050 = 10 \u00f7 (5/100) = 10 \u00d7 20",

  "Step 5: Calculate final result - R = 200 \u03a9"

],

"topic": "Ohm's law and electrical resistance",

"confidence_score": 0.99,

```

```
"solved_by_ai": true,
"saved_at": "2025-08-29T16:58:33.996355",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q29",
  "question_number": 29,
  "question_text": "A resistor and a thermistor are connected in a series circuit, as shown.
What happens to the readings on the ammeter and the voltmeter when the thermistor is
heated?",
  "options": {
    "A": "ammeter reading: decreases, voltmeter reading: decreases",
    "B": "ammeter reading: decreases, voltmeter reading: increases",
    "C": "ammeter reading: increases, voltmeter reading: decreases",
    "D": "ammeter reading: increases, voltmeter reading: increases"
  },
  "image_filename": "question_29_enhanced.png",
  "page": 10,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "C",
  "explanation": "When the thermistor is heated, its resistance decreases, causing total
circuit resistance to decrease, so current increases (ammeter reading increases) and
```

voltage across the fixed resistor increases while voltage across the thermistor decreases (voltmeter reading decreases)",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Analyze the behavior of a negative temperature coefficient (NTC) thermistor in a series circuit",

"calculation": "As thermistor resistance decreases, total resistance decreases, so $I = V/R_{\text{total}}$ increases",

"reasoning": "The circuit shows a voltmeter measuring voltage across the thermistor. Most thermistors are NTC (negative temperature coefficient), meaning their resistance decreases when heated. When the thermistor is heated: (1) Its resistance decreases, (2) Total circuit resistance decreases, (3) Current increases (ammeter reading increases), (4) Voltage across the fixed resistor increases ($V = IR$, where I increases and R is constant), (5) Since total voltage is constant, voltage across the thermistor must decrease (voltmeter reading decreases). The voltmeter measures the voltage across the thermistor, which decreases as the thermistor's resistance decreases.",

"conclusion": "Heating decreases thermistor resistance, increasing circuit current but decreasing voltage across the thermistor"

},

"calculation_steps": [

"Step 1: Identify thermistor type - typically NTC (resistance decreases with temperature)",

"Step 2: When heated, thermistor resistance decreases",

```
"Step 3: Total circuit resistance decreases, so current increases",
"Step 4: Ammeter reading increases due to higher current",
"Step 5: Voltmeter across thermistor decreases as its resistance drops"
],
"topic": "Thermistors and series circuits",
"confidence_score": 0.95,
"solved_by_ai": true,
"saved_at": "2025-08-29T16:59:02.767078",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q30",
  "question_number": 30,
  "question_text": "In which circuit does the ammeter read 2.0 A?",
  "options": {
    "A": "Circuit A: 12V with two 6.0\u03a9 resistors in series",
    "B": "Circuit B: 12V with two 3.0\u03a9 resistors in series",
    "C": "Circuit C: 6.0V with two 6.0\u03a9 resistors in series",
    "D": "Circuit D: 18V with two 3.0\u03a9 resistors in series"
  },
  "image_filename": "question_30_enhanced.png",
  "page": 2,
  "marks": 1,
  "subject": "physics",
```



```

"difficulty": "easy",

"correct_answer": "B",

"explanation": "Circuit B has 12V across two 3.0\03a9 resistors in series (total
resistance = 6.0\03a9), giving current  $I = V/R = 12V/6.0\03a9 = 2.0A$ ",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

  "approach": "Calculate current for each series circuit using Ohm's law after finding
total resistance",

  "calculation": "For series circuits:  $R_{total} = R_{1} + R_{2}$ , then  $I = V/R_{total}$ ",

  "reasoning": "In series circuits, resistances add together and the same current flows
through all components. I'll calculate the current for each circuit: Circuit A:  $R_{total} = 6.0\03a9 + 6.0\03a9 = 12.0\03a9$ , so  $I = 12V/12.0\03a9 = 1.0A$ . Circuit B:  $R_{total} = 3.0\03a9 + 3.0\03a9 = 6.0\03a9$ , so  $I = 12V/6.0\03a9 = 2.0A$ . Circuit C:  $R_{total} = 6.0\03a9 + 6.0\03a9 = 12.0\03a9$ , so  $I = 6.0V/12.0\03a9 = 0.50A$ . Circuit D:  $R_{total} = 3.0\03a9 + 3.0\03a9 = 6.0\03a9$ , so  $I = 18V/6.0\03a9 = 3.0A$ . Only Circuit B produces
exactly 2.0A.",

  "conclusion": "Circuit B is the only one that produces exactly 2.0A of current"

},

"calculation_steps": [

  "Step 1: Circuit A - Total resistance =  $6.0 + 6.0 = 12.0\03a9$ , Current =  $12V/12.0\03a9 = 1.0A$ ",

```

"Step 2: Circuit B - Total resistance = $3.0 + 3.0 = 6.0\Omega$, Current = $12V/6.0\Omega = 2.0A$ ",

"Step 3: Circuit C - Total resistance = $6.0 + 6.0 = 12.0\Omega$, Current = $6.0V/12.0\Omega = 0.50A$ ",

"Step 4: Circuit D - Total resistance = $3.0 + 3.0 = 6.0\Omega$, Current = $18V/6.0\Omega = 3.0A$ ",

"Step 5: Compare results - only Circuit B gives exactly 2.0A"

],

"topic": "Series circuits and Ohm's law calculations",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T16:59:47.042853",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q31",

"question_number": 31,

"question_text": "A 240V mains-powered vacuum cleaner has a power output of 2700W. Which fuse is suitable for use in the plug of the vacuum cleaner?",

"options": {

"A": "3A",

"B": "5A",

"C": "10A",

"D": "13A"

},

"image_filename": "question_31_enhanced.png",
"page": 10,
"marks": 1,
"subject": "physics",
"difficulty": "medium",
"correct_answer": "D",
"explanation": "Using $P = VI$ to find current: $I = P/V = 2700W/240V = 11.25A$. The fuse must be rated higher than this operating current, so 13A is the suitable choice",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
 "approach": "Calculate the operating current using the power equation, then select appropriate fuse rating",
 "calculation": " $I = P/V = 2700W / 240V = 11.25A$ ",
 "reasoning": "To determine the correct fuse rating, I need to calculate the current drawn by the vacuum cleaner using the relationship $P = VI$, which rearranges to $I = P/V$. With $P = 2700W$ and $V = 240V$, the current is $I = 2700/240 = 11.25A$. A fuse must be rated slightly higher than the normal operating current to prevent nuisance blowing during normal operation, but low enough to provide protection against dangerous overcurrents. Looking at the options: 3A, 5A, and 10A are all below the operating current of 11.25A, so they would blow immediately when the vacuum cleaner is switched on. Only the 13A fuse is rated above the operating current (11.25A) while still providing adequate protection.",

"conclusion": "The 13A fuse is the only option that can handle the 11.25A operating current while providing electrical protection"

},

"calculation_steps": [

"Step 1: Identify given values - Power = 2700W, Voltage = 240V",

"Step 2: Use power equation - $P = VI$, so $I = P/V$ ",

"Step 3: Calculate current - $I = 2700W / 240V = 11.25A$ ",

"Step 4: Compare with fuse ratings - need fuse $> 11.25A$ ",

"Step 5: Select 13A fuse as the only suitable option above operating current"

],

"topic": "Electrical power and fuse selection",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:00:18.475732",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q32",

"question_number": 32,

"question_text": "The transformer in a laptop power supply is used to change mains voltage from 240V to 20V. The transformer has 600 turns on the secondary coil. Which type of transformer is it and how many turns does the transformer have on the primary coil?",

"options": {

"A": "transformer type: step-down, number of turns on primary coil: 50",

"B": "transformer type: step-down, number of turns on primary coil: 7200",

"C": "transformer type: step-up, number of turns on primary coil: 50",

"D": "transformer type: step-up, number of turns on primary coil: 7200"

},

"image_filename": "question_32_enhanced.png",

"page": 11,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "B",

"explanation": "This is a step-down transformer (240V to 20V reduces voltage) with 7200 primary turns, calculated using the transformer equation $V_p/V_s = N_p/N_s$ ",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

 "approach": "Apply transformer principles to determine type and calculate primary turns using the turns ratio equation",

 "calculation": "Using $V_p/V_s = N_p/N_s$: $240\text{V}/20\text{V} = N_p/600$, so $N_p = (240/20) \times 600 = 12 \times 600 = 7200$ turns",

 "reasoning": "First, I need to identify the transformer type. Since the voltage decreases from 240V (primary) to 20V (secondary), this is a step-down transformer. For the turns calculation, I use the fundamental transformer relationship: $V_p/V_s = N_p/N_s$, where V_p = primary voltage (240V), V_s = secondary voltage (20V), N_p = primary turns (unknown), and

N_s = secondary turns (600). Substituting: $240/20 = N_p/600$. This gives $12 = N_p/600$, so $N_p = 12 \times 600 = 7200$ turns. The primary coil needs more turns than the secondary to step down the voltage.",

"conclusion": "It's a step-down transformer with 7200 turns on the primary coil, which is the correct ratio to reduce 240V to 20V"

},

"calculation_steps": [

"Step 1: Identify transformer type - voltage decreases from 240V to 20V, so it's step-down",

"Step 2: Apply transformer equation - $V_p/V_s = N_p/N_s$ ",

"Step 3: Substitute known values - $240V/20V = N_p/600$ turns",

"Step 4: Simplify ratio - $240/20 = 12$, so $12 = N_p/600$ ",

"Step 5: Solve for primary turns - $N_p = 12 \times 600 = 7200$ turns"

],

"topic": "Transformers and electromagnetic induction",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:00:51.292590",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q33",

"question_number": 33,

"question_text": "An ion of the element oxygen is negatively charged. Which change causes the ion to become a neutral oxygen atom?",

```
"options": {
  "A": "gain of electrons",
  "B": "gain of protons",
  "C": "loss of electrons",
  "D": "loss of protons"
},
"image_filename": "question_33_enhanced.png",
"page": 11,
"marks": 1,
"subject": "physics",
"difficulty": "easy",
"correct_answer": "C",
"explanation": "A negatively charged oxygen ion has gained extra electrons, so it must
lose electrons to become neutral again",
"extraction_method": "enhanced_multi_strategy_detection",
"detection_strategy": "standalone_number",
"confidence": 0.9,
"has_images": true,
"extraction_focus": "web_interface_compatible",
"manually_reviewed": false,
"reviewer_notes": "",
"review_timestamp": "",
"detailed_explanation": {
  "approach": "Apply principles of atomic structure and charge neutrality",
  "calculation": "No calculations needed - this is about understanding charge balance in
atoms and ions",
```

"reasoning": "A negatively charged ion (anion) has more electrons than protons. For oxygen, a neutral atom has 8 protons and 8 electrons, giving zero net charge. When oxygen forms a negative ion (typically O^{2-}), it gains 2 extra electrons, giving it 10 electrons but still 8 protons, resulting in a net charge of -2. To return to neutrality, the ion must lose the excess electrons it gained during ion formation. Gaining more electrons would make it even more negative. Changing the number of protons would change the element's identity entirely (since atomic number = number of protons), not just neutralize the charge.",

"conclusion": "The negatively charged oxygen ion must lose electrons to restore the balance between positive protons and negative electrons, making it neutral"

},

"calculation_steps": [

"Step 1: Understand that negative ions have excess electrons",

"Step 2: Recognize that neutral atoms have equal numbers of protons and electrons",

"Step 3: Identify that to neutralize, the ion must lose the excess electrons",

"Step 4: Eliminate proton changes (would change the element identity)",

"Step 5: Eliminate electron gain (would increase negative charge further)"

],

"topic": "Atomic structure and ion formation",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:01:24.610867",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q34",


```
"question_number": 34,

"question_text": "A nuclide of nitrogen is represented as  ${}^{15}_7\text{N}$ . What is the number of neutrons in this nuclide?",

"options": {
  "A": "7",
  "B": "8",
  "C": "15",
  "D": "22"
},

"image_filename": "question_34_enhanced.png",

"page": 11,

"marks": 1,

"subject": "physics",

"difficulty": "easy",

"correct_answer": "B",

"explanation": "Using the relationship: neutrons = mass number - atomic number. For  ${}^{15}_7\text{N}$ : neutrons = 15 - 7 = 8",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {
```

"approach": "Apply nuclear notation rules to find neutron number from mass number and atomic number",

"calculation": "Number of neutrons = Mass number (A) - Atomic number (Z) = 15 - 7 = 8",

"reasoning": "In nuclear notation $^{15}_7\text{N}$, the superscript 15 represents the mass number (A), which is the total number of protons and neutrons in the nucleus. Nitrogen has atomic number 7, meaning it always has 7 protons in its nucleus (this defines it as nitrogen). The mass number represents the sum of protons and neutrons: $A = Z + N$, where Z is the atomic number (protons) and N is the number of neutrons. Rearranging: $N = A - Z$. Therefore, for $^{15}_7\text{N}$: Number of neutrons = 15 - 7 = 8.",

"conclusion": "This nitrogen isotope has 8 neutrons, making it nitrogen-15 with 7 protons and 8 neutrons in its nucleus"

},

"calculation_steps": [

"Step 1: Identify mass number from notation - $^{15}_7\text{N}$ has mass number A = 15",

"Step 2: Identify atomic number of nitrogen - Z = 7 (number of protons)",

"Step 3: Apply formula - Number of neutrons = A - Z",

"Step 4: Calculate - Neutrons = 15 - 7 = 8",

"Step 5: Verify - 7 protons + 8 neutrons = 15 total nucleons $^{15}_7\text{N}$ "

],

"topic": "Nuclear notation and atomic structure",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:01:57.640147",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

```
{
  "id": "physics_q35",
  "question_number": 35,
  "question_text": "Which list only includes sources of naturally occurring background radiation?",
  "options": {
    "A": "cosmic rays, radon gas, nuclear power stations",
    "B": "food and drink, mobile phones, cosmic rays",
    "C": "rocks and buildings, food and drink, radon gas",
    "D": "rocks and buildings, mobile phones, nuclear power stations"
  },
  "image_filename": "question_35_enhanced.png",
  "page": 11,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "C",
  "explanation": "Rocks and buildings, food and drink, and radon gas are all naturally occurring sources of background radiation, while nuclear power stations and mobile phones are artificial sources",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
  "extraction_focus": "web_interface_compatible",
  "manually_reviewed": false,
  "reviewer_notes": ""
```

```
"review_timestamp": "",  
"detailed_explanation": {  
  "approach": "Distinguish between natural and artificial sources of background  
radiation",  
  "calculation": "No calculations needed - this is about categorizing radiation sources",  
  "reasoning": "Natural background radiation comes from sources that exist in nature  
without human intervention. Let me analyze each source: Cosmic rays - natural (from  
space), Radon gas - natural (from uranium decay in Earth), Nuclear power stations -  
artificial (human-made), Food and drink - natural (contains naturally radioactive isotopes  
like carbon-14, potassium-40), Mobile phones - artificial (electromagnetic radiation from  
technology), Rocks and buildings - natural (contain naturally radioactive materials like  
uranium, thorium, radium). Option C contains only natural sources: rocks/buildings  
contain natural radioactive elements, food/drink contains natural radioactive isotopes, and  
radon gas comes from natural uranium decay in the ground.",  
  "conclusion": "Only option C lists exclusively natural sources of background radiation"  
},  
"calculation_steps": [  
  "Step 1: Identify natural sources - cosmic rays, radon gas, rocks/buildings, food/drink",  
  "Step 2: Identify artificial sources - nuclear power stations, mobile phones",  
  "Step 3: Check option A - includes nuclear power stations (artificial)",  
  "Step 4: Check option B - includes mobile phones (artificial)",  
  "Step 5: Check option C - all natural sources only \u2713"  
],  
"topic": "Radioactivity and background radiation sources",  
"confidence_score": 0.98,  
"solved_by_ai": true,  
"saved_at": "2025-08-29T17:02:29.472965",  
"auto_flagged": false,  
"needs_review": false,
```

```
"flag_reason": null
},
{
  "id": "physics_q36",
  "question_number": 36,
  "question_text": "A student compares alpha emissions, beta emissions and gamma emissions. Which statement is correct?",
  "options": {
    "A": "Alpha emissions are the most ionizing.",
    "B": "Beta emissions are the most penetrating.",
    "C": "Beta emissions are part of the electromagnetic spectrum.",
    "D": "Gamma emissions are streams of charged particles."
  },
  "image_filename": "question_36_enhanced.png",
  "page": 12,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "A",
  "explanation": "Alpha emissions are the most ionizing because alpha particles are heavy, doubly-charged helium nuclei that interact strongly with matter, causing extensive ionization over short distances",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
  "has_images": true,
  "extraction_focus": "web_interface_compatible",
```

```
"manually_reviewed": false,  
"reviewer_notes": "",  
"review_timestamp": "",  
"detailed_explanation": {  
    "approach": "Compare the ionizing ability, penetrating power, and nature of alpha, beta,  
and gamma radiation",  
    "calculation": "No calculations needed - this is about understanding radiation  
properties",  
    "reasoning": "Let me evaluate each statement: (A) Alpha particles are helium nuclei (2  
protons + 2 neutrons) with +2 charge and large mass. They interact very strongly with  
matter, stripping electrons from atoms frequently, making them highly ionizing. (B) Gamma  
rays are the most penetrating, not beta particles. Penetrating power order: gamma > beta >  
alpha. (C) Beta particles are electrons or positrons (charged particles), not  
electromagnetic radiation. Only gamma rays are part of the electromagnetic spectrum. (D)  
Gamma rays are electromagnetic waves (photons), not charged particles. They have no  
charge or mass. Alpha and beta emissions are streams of charged particles, but not  
gamma rays.",  
    "conclusion": "Alpha emissions have the highest ionizing ability due to their large mass  
and double positive charge, making option A correct"  
},  
"calculation_steps": [  
    "Step 1: Compare ionizing ability - alpha particles are most ionizing due to high charge  
and mass",  
    "Step 2: Compare penetrating power - gamma rays > beta particles > alpha particles",  
    "Step 3: Identify electromagnetic radiation - only gamma rays are EM waves",  
    "Step 4: Identify particle nature - alpha and beta are particles, gamma are photons",  
    "Step 5: Verify that only statement A is correct"  
],  
"topic": "Properties of radioactive emissions",  
"confidence_score": 0.99,
```

```
"solved_by_ai": true,
"saved_at": "2025-08-29T17:03:03.267282",
"auto_flagged": false,
"needs_review": false,
"flag_reason": null
},
{
  "id": "physics_q37",
  "question_number": 37,
  "question_text": "In school laboratories, radioactive sources are stored in lead-lined boxes. Why is lead used?",
  "options": {
    "A": "Lead neutralises radiation.",
    "B": "Lead absorbs radiation.",
    "C": "Lead repels radiation.",
    "D": "Lead dissolves radiation."
  },
  "image_filename": "question_37_enhanced.png",
  "page": 12,
  "marks": 1,
  "subject": "physics",
  "difficulty": "easy",
  "correct_answer": "B",
  "explanation": "Lead absorbs radiation due to its high density and atomic number, making it effective at stopping or significantly reducing the intensity of radioactive emissions",
  "extraction_method": "enhanced_multi_strategy_detection",
```

```
"detection_strategy": "standalone_number",  
"confidence": 0.9,  
"has_images": true,  
"extraction_focus": "web_interface_compatible",  
"manually_reviewed": false,  
"reviewer_notes": "",  
"review_timestamp": "",  
"detailed_explanation": {  
  "approach": "Understand the interaction between radiation and dense materials for  
shielding purposes",  
  "calculation": "No calculations needed - this is about understanding radiation shielding  
mechanisms",  
  "reasoning": "Lead is used for radiation shielding because it effectively absorbs  
radiation through several mechanisms. Lead has a high atomic number (82) and high  
density (11.34 g/cm3), which makes it excellent at interacting with and stopping  
radiation. When radiation encounters lead atoms, it undergoes various interactions:  
photoelectric absorption, Compton scattering, and pair production (for high-energy  
gamma rays). These interactions transfer the radiation's energy to the lead atoms,  
effectively absorbing the radiation and preventing it from passing through. Lead doesn't  
neutralize, repel, or dissolve radiation - these terms don't accurately describe the physical  
processes involved. The absorption process converts the radiation's energy into heat within  
the lead material.",  
  "conclusion": "Lead's high density and atomic number allow it to effectively absorb  
radiation through atomic interactions, making it an ideal shielding material"  
},  
"calculation_steps": [  
  "Step 1: Consider lead's physical properties - high atomic number (82) and density",  
  "Step 2: Understand radiation interaction - radiation transfers energy to lead atoms",  
  "Step 3: Identify the process - energy absorption converts radiation to heat",
```


"Step 4: Eliminate incorrect terms - neutralize, repel, dissolve don't describe this process",

"Step 5: Confirm absorption is the correct mechanism for radiation shielding"

],

"topic": "Radiation shielding and material properties",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:03:47.193709",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q38",

"question_number": 38,

"question_text": "The Sun, the Earth and the Moon are all bodies in the Solar System. The Earth has a periodic cycle of day and night and a periodic cycle of seasons. Which row relates the periods of these cycles to the motions of these bodies?",

"options": {

"A": "period of day and night: the Earth orbits the Sun, period of the seasons: the Earth rotates about its axis",

"B": "period of day and night: the Earth orbits the Sun, period of the seasons: the Moon orbits the Earth",

"C": "period of day and night: the Earth rotates about its axis, period of the seasons: the Earth orbits the Sun",

"D": "period of day and night: the Earth rotates about its axis, period of the seasons: the Moon orbits the Earth"

},

"image_filename": "question_38_enhanced.png",

"page": 12,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "C",

"explanation": "Day and night are caused by Earth's rotation about its axis (approximately 24 hours), while seasons are caused by Earth's orbit around the Sun (approximately 365 days) combined with Earth's axial tilt",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Match the correct celestial motions to their corresponding Earth-based periodic phenomena",

"calculation": "No calculations needed - this is about understanding astronomical cause-and-effect relationships",

"reasoning": "The day-night cycle occurs because Earth rotates on its axis once every ~24 hours. As Earth spins, different parts face toward or away from the Sun, creating day and night. The seasonal cycle occurs because Earth orbits the Sun once per year (~365 days) while tilted at 23.5° on its axis. This tilt means different hemispheres receive varying amounts of direct sunlight throughout the year, creating seasons. The Moon's orbit around Earth (~27.3 days) primarily affects tides and lunar phases, not day-night cycles or

seasons. Earth's orbital period around the Sun is far too long (1 year) to cause the daily day-night cycle.",

"conclusion": "Earth's rotation creates day and night, while Earth's orbital motion around the Sun creates the seasonal cycle"

},

"calculation_steps": [

"Step 1: Identify what causes day and night - Earth's rotation exposes different sides to sunlight",

"Step 2: Confirm day-night period - approximately 24 hours matches Earth's rotation period",

"Step 3: Identify what causes seasons - Earth's tilted orbit around the Sun",

"Step 4: Confirm seasonal period - approximately 365 days matches Earth's orbital period",

"Step 5: Eliminate Moon's influence - affects tides and phases, not these major cycles"

],

"topic": "Astronomy - Earth's motions and periodic cycles",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:04:21.987984",

"auto_flagged": false,

"needs_review": false,

"flag_reason": null

},

{

"id": "physics_q39",

"question_number": 39,

"question_text": "The strength of the gravitational field at the surface of a planet depends on the1..... . The strength of the gravitational field2..... as the distance from the planet increases. Which words correctly complete the sentences?",

"options": {

"A": "1: atmospheric pressure, 2: decreases",

"B": "1: mass of the planet, 2: decreases",

"C": "1: atmospheric pressure, 2: increases",

"D": "1: mass of the planet, 2: increases"

},

"image_filename": "question_39_enhanced.png",

"page": 12,

"marks": 1,

"subject": "physics",

"difficulty": "medium",

"correct_answer": "B",

"explanation": "Gravitational field strength depends on the mass of the planet and decreases as distance from the planet increases, following Newton's law of universal gravitation",

"extraction_method": "enhanced_multi_strategy_detection",

"detection_strategy": "standalone_number",

"confidence": 0.9,

"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

"approach": "Apply Newton's law of universal gravitation to determine the factors affecting gravitational field strength",

"calculation": " $g = GM/r^2$, where G is gravitational constant, M is planet mass, r is distance from center",

"reasoning": "The gravitational field strength (g) at any point is given by Newton's law of universal gravitation: $g = GM/r^2$, where G is the gravitational constant, M is the mass of the planet, and r is the distance from the planet's center. From this equation, we can see that: (1) The field strength is directly proportional to the planet's mass M - larger mass means stronger gravitational field. (2) The field strength is inversely proportional to the square of the distance r - as distance increases, field strength decreases rapidly. Atmospheric pressure is unrelated to gravitational field strength; it's a result of gravitational effects on the atmosphere, not a cause of gravitational field strength.",

"conclusion": "Gravitational field strength depends on the planet's mass and decreases with increasing distance according to the inverse square law"

},

"calculation_steps": [

"Step 1: Recall Newton's law of universal gravitation - $F = GMm/r^2$, so $g = GM/r^2$ ",

"Step 2: Identify factors affecting g - directly proportional to mass M",

"Step 3: Identify distance relationship - inversely proportional to r^2 ",

"Step 4: Eliminate atmospheric pressure - not a fundamental gravitational factor",

"Step 5: Confirm that g decreases as distance increases due to inverse square law"

],

"topic": "Gravitational fields and Newton's law of universal gravitation",

"confidence_score": 0.99,

"solved_by_ai": true,

"saved_at": "2025-08-29T17:04:52.204835",

"auto_flagged": false,

"needs_review": false,

```
"flag_reason": null
},
{
  "id": "physics_q40",
  "question_number": 40,
  "question_text": "The Sun radiates most of its energy in three regions of the
electromagnetic spectrum. The diagram shows the proportion of the total energy radiated
in each of these regions. Region P is visible light and region Q is ultraviolet. What is region
R?",
  "options": {
    "A": "infrared",
    "B": "microwaves",
    "C": "radio waves",
    "D": "X-rays"
  },
  "image_filename": "question_40_enhanced.png",
  "page": 13,
  "marks": 1,
  "subject": "physics",
  "difficulty": "medium",
  "correct_answer": "A",
  "explanation": "Region R is infrared radiation because the Sun's energy output peaks in
the visible spectrum but extends significantly into the infrared region, which has longer
wavelengths than visible light",
  "extraction_method": "enhanced_multi_strategy_detection",
  "detection_strategy": "standalone_number",
  "confidence": 0.9,
```

```
"has_images": true,

"extraction_focus": "web_interface_compatible",

"manually_reviewed": false,

"reviewer_notes": "",

"review_timestamp": "",

"detailed_explanation": {

    "approach": "Use knowledge of solar radiation spectrum and the electromagnetic
spectrum order to identify the missing region",

    "calculation": "No calculations needed - this is about understanding the solar radiation
spectrum",

    "reasoning": "The Sun's radiation follows a blackbody spectrum with peak emission in
the visible range due to its surface temperature of approximately 5800K. From the diagram,
we can see three major regions: P (visible light), Q (ultraviolet), and R (unknown). The
electromagnetic spectrum in order of increasing wavelength is: gamma rays, X-rays,
ultraviolet, visible light, infrared, microwaves, radio waves. Since Q is ultraviolet (shorter
wavelengths than visible) and P is visible light, region R must be the next region in the
spectrum with longer wavelengths than visible light, which is infrared. The Sun does emit
significant energy in the infrared region, making this the logical third major component
along with visible and UV radiation.",

    "conclusion": "Region R represents infrared radiation, which along with visible light and
ultraviolet, constitutes the three main regions of solar electromagnetic radiation"

},

"calculation_steps": [

    "Step 1: Identify known regions - P is visible light, Q is ultraviolet",

    "Step 2: Recall EM spectrum order - UV, visible, infrared are adjacent regions",

    "Step 3: Note that UV has shorter wavelengths than visible light",

    "Step 4: Deduce that R must be longer wavelengths than visible light",

    "Step 5: Identify infrared as the next region beyond visible light"

],
```

```
"topic": "Electromagnetic spectrum and solar radiation",  
"confidence_score": 0.98,  
"solved_by_ai": true,  
"saved_at": "2025-08-29T17:05:29.615054",  
"auto_flagged": false,  
"needs_review": false,  
"flag_reason": null  
}  
]  
}
```