

MCQs on Systematic - Solved

#	Question	A	B	C	D	Answer
1	Protozoa have several types of vacuoles. The function of contractile vacuoles is to maintain osmotic balance and prevent the cell from bursting in a hypotonic environment.	maintain osmotic balance	prevent the cell from bursting	digestion of food	excretion of waste	A
2	Members of the Protozoa may move by all of the following methods except	flagella	slime secretion	cilia	pseudopodia	B
3	Entamoeba histolytica cysts have.....nuclei	4	5	9	6-8	A
4	This type of protozoan usually has complex life cycle that take place in host.	Parasitic	Free living	Parasitic	Eukaryotic	C
5	The members of Mastigophora move by	Cilia	Flagella	Whip	Pseudopodia	B
6	In Amoeba and Paramecium, the cell organelle that is more regulatory in nature is	contractile vacuole	pseudopodia	cell membrane	cell nucleus	C
7	Protozoa are classified on the basis of	locomotory organelles	shape	number of nuclei	size	A
8	Which of the following is not a characteristic feature of protozoans	contractile vacuole	cell membrane as a protective layer	pseudopodia	cell nucleus	D
9	The organelle that is a protective coat surrounding the euglena that holds its shape is called? Tric...					C
10	The organelle that allows the euglena to produce its own food by photosynthesis processes by convert...					A
11	Which of the following moves using pseudopodia?	paramecium	euglena	diatoms	ameba	B
12	The word "pseudopodia" means	first animal	false foot	fake eye	small cell	C
13	Which organelle functions to remove excess water?	nucleus	contractile vacuole	pellicle	gullet	A
14	Which of the following is autotrophic?	paramecium	foraminifera	ameba	euglena	B
15	Entamoeba histolytica is found in the of human	colon	endoparasite	lung	liver	C
16	Ciliophora is a sub-phylum of	Protozoa	Porifera	Coelenterata	Ctenophora	A
17	Amoebic dysentery is caused by	Entamoeba gingivalis	Entamoeba histolytica	Trypanosoma gambiense	Leishmania donovani	B
1	The pore-bearing animals are called	hydra	sponges	obelia	protozoa	A
2	The skeleton of sponge is called	spicules	ostia	osculum	spongeocoel	B
3	The water entrance to the body of sponge are called	special cells	ostia	osculum	spongeocoel	A
4	Theare multi-cellular animals but have no true tissues.	Protozoans	mesozoans	metazoans	cnidarians	C

5	Gelatinous layer between pinacoderm and choanoderm is called as.	mesoglea	ectodermis	gastrodermis		B
6	Animals lacking excretory, respiratory and circulatory structures are (a) Sponges (b) Tapeworms (c) ...					C
7	Members of phylum Porifera are (a) Mostly freshwater animals but few are marine entities (b) Mostly ...					A
8	The group of animals to which Sycon belongs can be best described as (a) Multicellular having tissue...					D
9	The chamber common to all the types of the canal system of sponges is (a) Excurrent canal (b) Radial...					C
10	The classification of the sponges is based on (a) size (b) locomotory organs (c) body wall (d) skele...					A
11	The fertilization in porifera is (a) internal (b) external (c) both (d) none					B
12	Choanocytes In Sponges Are Present	Line The Gastric Cavity	Only The External Body Surface	Between Mesodermal Layer	Lined Between The Outer And Inner Layer	
13	Skeleton Of Sponges Is Produced By	Sclerocytes	Pinacocytes	Thesocytes	Choanocytes Metazoa A	
1	cnidria					A
1	A coelenterate that is commonly referred to as 'fresh water polyp' is (a) Obelia (b) Physalia (c) Hy...					A
2	This is a special character of coelenterata occurring only in them (a) Flame cells (b) Hermaphrodit...					B
8	The class of coelenterata in which the medusa and polyp both are found in one animal is (a) Anthozoa...					D
3	The characteristic larva in coelenterates is (a) Oncosphere (b) Rhabditiform (c) Planula (d) Cystice...					A
4	All the following animals are in the Phyla Cnidaria EXCEPT	Amoeba		sea anemone E hydras		C
5	To protect themselves, some Cnidaria have:	stinging cells . B poison sacs.		stabbing points.	biting teeth.	B
6	The cup-shaped Cnidaria include (adult) : A jellyfish	sea anemones	coral	hydra		C
7	The outer cell layer of Hydra is the a. gastrodermis. b. mesoglea . c. endoderm. d. epidermis.					A
8	Cnidarians are dimorphic, meaning they have a. two choanocytes. . b. two tissue layers c. two tentac...					D
9	Which of the following statements is NOT true? a. A medusa is the sessile form of a cnidarian. b. Cn...					C
10	Which one is diploblastic animal? (a) Euglena (b) Hydra (c) Ameoba (d) All of these					A
2	phylum: Platyhelimenthes					B
1	Liver-fluke (Fasciola hepatica) belongs to (a) Class Turbellaria (b) Class Trematoda (c) Class Cesto...					A
2	Tape-worm (Taenia solium) belongs to (a) Class Turbellaria (b) Class Trematoda (c) Class Cestoda (d) ...					B
3	Liver fluke-completes its life-cycle in two hosts. Its primary host is usually sheep and the seconda...					A

4	Liver-fluke (<i>Fasciola hepatica</i>) is an endoparasite and it causes a disease, known as (a) <i>Filaria</i> (b)...				C
5	Liver-Fluke has (a) One oral sucker (b) One ventral sucker (c) one oral sucker and two ventral sucke...				B
6	Liver-fluke is hermaphrodite and it bears (a) single testis and single overary (b) single testis and...				C
7	Miracidium, sporocyst, redia and cercaria are larval forms of (a) <i>Obelia</i> (b) Liver-fluke (c) Tape-wo...				A
8	The cercaria in encysted condition is called (a) <i>Metacercaria</i> (b) <i>Cysticercus</i> (c) Sporocyst (d) Redi...				D
9	The head or scolex of <i>Taenia solium</i> bears in the middle (a) one sucker (b) two suckers (c) four suck...				C
10	Mature proglottids of Tapeworm contain (a) Male reproductive organs (b) Female reproductive organs (...)				A
11	The intermediate host of <i>Taenia solium</i> is (a) man (b) Pig (c) Sheep (d) Snail				B
12	The definitive host of <i>Taenia solium</i> is (a) man (b) Pig (c) Sheep (d) Snail				C
13	Segments of Tapeworm are called A) Scolex B) <i>Cysticercus</i> C) Proglottides D) Onchospheres				A
14	Scolex occurs in A) <i>Hydra</i> B) <i>Ascaris</i> C) <i>Taenia</i> D) Liver Fluke				B
15	Incomplete alimentary canal occurs in A) <i>Fasciola</i> B) <i>Ascaris</i> C) <i>Ameoba</i> D) <i>Paramecium</i>				C
3	NEMATODA				A
1	Sexes are separate with a distinct sexual dimorphism in (a) <i>Ascaris lumbricoides</i> (b) <i>Planaria</i> (c) <i>Fa...</i>				A
2	Males are always smaller than the females and have a coiled posterior end in (a) <i>Ascaris lumbricoide...</i>				B
3	Round worm is (a) Acoelomate (b) Pseudocoelomate (c) Coelomate (d) None				A
4	The development of Roundworm is (a) Direct (b) Indirect (c) Direct or Indirect (d) Direct and Indire...				C
5	Life-history of <i>Ascaris lumbricoides</i> involves (a) two larval stages (b) three larval stages (c) fou...				B