

# Algolab 2015

## Radiation Therapy Solution

Antonis Thomas

16 December 2015

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1, \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1, \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling

LP

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling
- Implementation Details

LP

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,  $p(t_j) < 0$  for  $j = 1, \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling
- Implementation Details

LP

number types, etc

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling
- Implementation Details
- Generating the polynomials

LP

number types, etc



# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling
- Implementation Details
- Generating the polynomials

LP

number types, etc

nested for loops

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1 \dots, t$

## Example: Polynomial of degree 2

$$1c_0 + x_1c_1 + x_2c_2 + x_3c_3 + x_1^2c_4 + x_1x_2c_5 + x_1x_3c_6 + x_2^2c_7 + x_2x_3c_8 + x_3^2c_9$$

## Solution

- Modeling
- Implementation Details
- Generating the polynomials
- Find the best degree

LP

number types, etc

nested for loops

# Radiation Therapy

- healthy cells  $h_i$  and tumor cells  $t_j$  in  $\mathbb{R}^3$   
( $1 \leq h + t \leq 50$ )
- **Polynomial**  $p$ : of minimal degree  $d$ , s.t.  
 $p(h_i) > 0$  for  $i = 1, \dots, h$ ,       $p(t_j) < 0$  for  $j = 1, \dots, t$

## Example: Polynomial of degree 2

$$1C_0 + X_1C_1 + X_2C_2 + X_3C_3 + X_1^2C_4 + X_1X_2C_5 + X_1X_3C_6 + X_2^2C_7 + X_2X_3C_8 + X_3^2C_9$$

## Solution

- Modeling LP
- Implementation Details number types, etc
- Generating the polynomials nested for loops
- Find the best degree binary search on  $d$

# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

- Option 1 (additional variable  $0 \leq \epsilon \leq 1$ )

$$p(h_i) \geq \epsilon \text{ for } i = 1, \dots, h, \quad p(t_j) \leq 0 \text{ for } j = 1 \dots, t$$

Cells can be separated if there exists a *positive*  $\epsilon$

# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

- Option 1 (additional variable  $0 \leq \epsilon \leq 1$ )

$$p(h_i) \geq \epsilon \text{ for } i = 1, \dots, h, \quad p(t_j) \leq 0 \text{ for } j = 1 \dots, t$$

Cells can be separated if there exists a *positive*  $\epsilon$

- Option 2 (scaling)

$$p(h_i) \geq 1 \text{ for } i = 1, \dots, h, \quad p(t_j) \leq -1 \text{ for } j = 1 \dots, t$$

# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

- Option 1 (additional variable  $0 \leq \epsilon \leq 1$ )  
 $p(h_i) \geq \epsilon$  for  $i = 1, \dots, h$ ,  $p(t_j) \leq 0$  for  $j = 1 \dots, t$   
Cells can be separated if there exists a *positive*  $\epsilon$
- Option 2 (scaling)  
 $p(h_i) \geq 1$  for  $i = 1, \dots, h$ ,  $p(t_j) \leq -1$  for  $j = 1 \dots, t$

## Implementation Details

- **Number types**  
Observe  $|x_1| < 2^{10}$ . Therefore  $x_1^{30} \approx 2^{300}$ .

# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

- Option 1 (additional variable  $0 \leq \epsilon \leq 1$ )  
 $p(h_i) \geq \epsilon$  for  $i = 1, \dots, h$ ,  $p(t_j) \leq 0$  for  $j = 1 \dots, t$   
Cells can be separated if there exists a *positive*  $\epsilon$
- Option 2 (scaling)  
 $p(h_i) \geq 1$  for  $i = 1, \dots, h$ ,  $p(t_j) \leq -1$  for  $j = 1 \dots, t$

## Implementation Details

- **Number types**

Observe  $|x_1| < 2^{10}$ . Therefore  $x_1^{30} \approx 2^{300}$ .

**IT and ET:** any exact type, like

`CGAL::Gmpz` or `CGAL::Gmpzf` or `CGAL::MP_Float`



# Modeling

**Problem:** we can not use  $>$  or  $<$  for the constraints:

- Option 1 (additional variable  $0 \leq \epsilon \leq 1$ )

$$p(h_i) \geq \epsilon \text{ for } i = 1, \dots, h, \quad p(t_j) \leq 0 \text{ for } j = 1 \dots, t$$

Cells can be separated if there exists a *positive*  $\epsilon$

- Option 2 (scaling)

$$p(h_i) \geq 1 \text{ for } i = 1, \dots, h, \quad p(t_j) \leq -1 \text{ for } j = 1 \dots, t$$

## Implementation Details

- **Number types**

Observe  $|x_1| < 2^{10}$ . Therefore  $x_1^{30} \approx 2^{300}$ .

**IT and ET:** any exact type, like

`CGAL::Gmpz` or `CGAL::Gmpzf` or `CGAL::MP_Float`

- **Cycling**

Use `CGAL::QP_BLAND`

# Generating the polynomials

```
• for (int i = 0; i <= degree; ++i)
    for (int j = 0; j <= degree-i; ++j)
        for (int k = 0; k <= degree-i-j; ++k) {
            /* term x_1^i * x_2^j * x_3^k */
            pw(p.x1, i) * pw(p.x2, j) * pw(p.x3, k);
        }
```

For efficiency, you can also precompute the values of  
 $\text{pw}(p.x[1-3], i)$  for  $i = 0, \dots, 30$

# Finding the minimum degree $d$

- To find the minimum degree  $d$ :

# Finding the minimum degree $d$

- To find the minimum degree  $d$ :
  - ▶ Fast exponentiation to find upper bound  $u$ ;

# Finding the minimum degree $d$

- To find the minimum degree  $d$ :
  - ▶ Fast exponentiation to find upper bound  $u$ ;
  - ▶ Binary search in the interval  $[\frac{u}{2}, u]$ .