

Tutorial 1

Vectors and matrices

Homework

1. Vectors, Questions 1–11: Provide detailed solutions to any five subparts of your choice. The five subparts may come from different questions; you do not need to complete all subparts of a selected question.
2. Vectors, Question 12: Provide detailed solutions to Parts 2, 3, and 4.
3. Matrices, Questions 2–5: Provide detailed solutions to any ten subparts of your choice from Questions 2–5. The ten subparts may be selected independently across these questions.
4. Matrices, Questions 8–11: Provide detailed solutions to any ten subparts of your choice from Questions 8–11. The ten subparts may be selected independently across these questions.
5. Matrices, Question 12: Select any three subparts from Parts 1–11 and provide detailed solutions. For each selected subpart, solve both the 3×3 case and the 4×4 case.

1 Vectors

Question 1. Given points A, B, C , find the missing coordinate of B such that

a $\overrightarrow{AB}$ and $\overrightarrow{AC}$ are orthogonal

b $\overrightarrow{AB}$ and $\overrightarrow{AC}$ are parallel

1. $A = (3, 5)$, $B = (2, y)$, $C = (2, 8)$
2. $A = (-2, 5)$, $B = (1, y)$, $C = (4, -3)$
3. $A = (1, 5)$, $B = (-1, y)$, $C = (2, -3)$
4. $A = (2, 1)$, $B = (x, -2)$, $C = (1, 3)$

Question 2. Consider the triangle with vertices A, B , and C , $\triangle ABC$. Define the vectors

$$\mathbf{a} = \overrightarrow{BC}, \quad \mathbf{b} = \overrightarrow{AC}, \quad \mathbf{c} = \overrightarrow{AB}.$$

Let M, N , and P denote the midpoints of the sides BC, AC , and AB , respectively.

Determine the expressions for the vectors $\overrightarrow{AM}$, $\overrightarrow{BN}$, and $\overrightarrow{CP}$ in terms of $\mathbf{a}$, $\mathbf{b}$, and $\mathbf{c}$. Subsequently, rewrite these expressions using only the vectors $\mathbf{a}$ and $\mathbf{b}$.

Question 3. Given points A, B , find x such that the norm of the vector $\overrightarrow{AB}$, $\|\overrightarrow{AB}\|$ equals the specified value d .

1. $A = (2, -3)$, $B = (x, 0)$, $d = 5$
2. $A = (1, 4)$, $B = (x, 1)$, $d = 4$
3. $A = (1, 5)$, $B = (1, x)$, $d = 6$

Question 4. Prove that the triangle $\triangle ABC$ is isosceles.

1. $A = (-3, -2)$, $B = (1, 4)$, $C = (-5, 0)$.
2. $A = (7, -3, 6)$, $B = (11, -5, 3)$, $C = (10, -7, 8)$.

Question 5. Given vectors $\mathbf{a} = (3, -2)$ and $\mathbf{b} = (-1, 5)$, find vector $\mathbf{c}$ such that:

- $\mathbf{a} \cdot \mathbf{c} = 17$.
- $\mathbf{b} \cdot \mathbf{c} = 3$.

Question 6. Find all vectors $\mathbf{a}$ that are orthogonal to the given vector $\mathbf{b}$ and have norm d .

1. $\mathbf{b} = (3, 4)$, $d = 15$
2. $\mathbf{b} = (-3, 2)$, $d = 10$
3. $\mathbf{b} = (1, 4)$, $d = 8$
4. $\mathbf{b} = (-1, 2)$, $d = 7$
5. $\mathbf{b} = (2, -5)$, $d = 21$

Question 7. Find the vector $\mathbf{c}$.

1. From the equation:

$$2\mathbf{c} + 3\mathbf{a} = \mathbf{b}, \quad \text{where } \mathbf{a} = (-1, 2), \quad \mathbf{b} = (0, -2).$$

2. From the equation:

$$3\mathbf{c} - \mathbf{a} = 2\mathbf{b}, \quad \text{where } \mathbf{a} = (3, -1), \quad \mathbf{b} = (0, 2).$$

3. From the equation:

$$2\mathbf{c} - 2\mathbf{a} = \mathbf{b}, \quad \text{where } \mathbf{a} = (3, -1), \quad \mathbf{b} = (0, 2).$$

4. From the equation:

$$3\mathbf{c} + 5\mathbf{a} = 4\mathbf{b}, \quad \text{where } \mathbf{a} = (2, 4), \quad \mathbf{b} = (1, -2).$$

Question 8. Calculate the side lengths of the triangle $\triangle ABC$. Additionally, find the coordinates of a fourth point D such that the quadrilateral $ABCD$ forms a parallelogram.

1. $A = (-4, -2), B = (-1, 4), C = (2, 2)$
2. $A = (5, 1), B = (4, 2), C = (-1, 4)$
3. $A = (3, 2), B = (7, 4), C = (5, 6)$
4. $A = (-1, 2), B = (-3, 4), C = (-3, -3)$

Question 9. Calculate the lengths of the sides and diagonals of the specified quadrilateral $ABCD$, classify its type.

1. $A = (8, -4), B = (5, -6), C = (1, -4), D = (4, 2)$
2. $A = (-2, -3), B = (-5, -7), C = (-1, -10), D = (2, -6)$
3. $A = (7, 5), B = (-6, 2), C = (3, -1), D = (6, 1)$

Question 10. Find the interior angles of the triangle $\triangle ABC$, where

$$A = (5\sqrt{3}, 5), B = (-\sqrt{3}, 1), C = (0, 0)$$

Question 11. Identify which of the following vector pairs are parallel.

1. $(-2, 3, 1), (6, -4, -3)$
2. $(10, -8, 9, 0, 24), \left(\frac{5}{6}, -\frac{2}{3}, \frac{3}{4}, 0, 2\right)$
3. $(8, -6, -9, -8, -2), \left(-\frac{32}{3}, 8, 12, \frac{32}{3}, \frac{8}{3}\right)$
4. $(0, 0, 2, 3, 5), \left(0, 1, \frac{4}{3}, 2, \frac{10}{3}\right)$

Question 12. Compute $\text{proj}_{\mathbf{a}} \mathbf{b}$, $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$, $\text{proj}_{\mathbf{b}} \mathbf{a}$ and $\mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a}$.

1. $\mathbf{a} = (2, 1, 5), \mathbf{b} = (1, 4, -5)$
2. $\mathbf{a} = (6, 2, 4), \mathbf{b} = (1, -5, 4)$

3. $\mathbf{a} = (1, 0, 1, -2)$, $\mathbf{b} = (5, -1, 0, 2)$

4. $\mathbf{a} = (-2, 3, -4, 1)$, $\mathbf{b} = (5, 1, -8, 2)$

Question 13. For any $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ with $\mathbf{a} \neq \mathbf{0}$, show that $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$ is orthogonal to $\mathbf{a}$.

Question 14. Take any $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$, where both vectors are nonzero.

1. Suppose $\mathbf{a}$ and $\mathbf{b}$ are orthogonal. Find $\text{proj}_{\mathbf{a}} \mathbf{b}$.
2. Suppose $\mathbf{a}$ and $\mathbf{b}$ are parallel. Find $\text{proj}_{\mathbf{a}} \mathbf{b}$.

2 Matrices

Question 1. Let

$$A = \begin{pmatrix} -4 & 2 & 3 \\ 0 & 5 & -1 \\ 6 & 1 & 2 \end{pmatrix} \quad B = \begin{pmatrix} 6 & -1 & 0 \\ 2 & 2 & -4 \\ 3 & -1 & 1 \end{pmatrix} \quad C = \begin{pmatrix} 5 & -1 \\ -3 & 4 \end{pmatrix}$$

$$D = \begin{pmatrix} -7 & 1 & -4 \\ 3 & -2 & 8 \end{pmatrix} \quad E = \begin{pmatrix} 3 & -3 & 5 \\ 1 & 0 & -2 \\ 6 & 7 & -2 \end{pmatrix} \quad F = \begin{pmatrix} 8 & -1 \\ 2 & 0 \\ 5 & -3 \end{pmatrix}$$

Compute each of the following, if it is defined.

- | | | |
|------------------|----------------------|-----------------------------------|
| 1. $A + B$ | 2. $2A - 3E - B$ | 3. $2C^\top - 3F$ |
| 4. $C + D$ | 5. $2D - 3F$ | 6. $5(F^\top - D^\top)$ |
| 7. $4A$ | 8. $A^\top + E^\top$ | 9. $((B - A)^\top + E^\top)^\top$ |
| 10. $2A - 3B$ | 11. $(A + E)^\top$ | 12. $A - B + E$ |
| 13. $C + 3F - E$ | 14. $4D + 3F^\top$ | |

For Questions 2-5, let

$$A = \begin{pmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 4 & -1 \\ 0 & 2 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{pmatrix}$$

$$D = \begin{pmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{pmatrix}, \quad E = \begin{pmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{pmatrix}$$

Question 2. Compute each of the following, if it is defined.

- | | |
|---------------|------------------------|
| 1. $D + E$ | 2. $-3(D + 2E)$ |
| 3. $D - E$ | 4. $A - A$ |
| 5. $5A$ | 6. $\text{tr}(D)$ |
| 7. $-7D$ | 8. $\text{tr}(D - 3E)$ |
| 9. $2B - C$ | 10. $4\text{tr}(7B)$ |
| 11. $4E - 2D$ | 12. $\text{tr}(A)$ |

Question 3. Compute each of the following, if it is defined.

- | | |
|---------------------------------------|-------------------------------|
| 1. $2A^\top + C$ | 2. $2E^\top - 3D^\top$ |
| 3. $D^\top - E^\top$ | 4. $(2E^\top - 3D^\top)^\top$ |
| 5. $(D - E)^\top$ | 6. $(CD)E$ |
| 7. $B^\top + 5C^\top$ | 8. $C(AB)$ |
| 9. $\frac{1}{2}C^\top - \frac{1}{4}A$ | 10. $\text{tr}(DE^\top)$ |
| 11. $B - B^\top$ | 12. $\text{tr}(BC)$ |

Question 4. Compute each of the following, if it is defined.

- | | |
|--|-----------------------------------|
| 1. AB | 2. BA |
| 3. $(3E)D$ | 4. $(AB)C$ |
| 5. $A(BC)$ | 6. $CC^\top$ |
| 7. $(DA)^\top$ | 8. $(C^\top B)A^\top$ |
| 9. $\text{tr}(DD^\top)$ | 10. $\text{tr}(4E^\top - D)$ |
| 11. $\text{tr}(C^\top A^\top + 2E^\top)$ | 12. $\text{tr}((EC^\top)^\top A)$ |

Question 5. Compute each of the following, if it is defined.

- | | |
|---------------------------|---------------------------------|
| 1. $(2D^\top - E)A$ | 2. $(BA^\top - 2C)^\top$ |
| 3. $(4B)C + 2B$ | 4. $B^\top(CC^\top - A^\top A)$ |
| 5. $(-AC)^\top + 5D^\top$ | 6. $D^\top E^\top - (ED)^\top$ |

Question 6. Given matrices

$$A \in \mathcal{M}_{4 \times 5}, \quad B \in \mathcal{M}_{4 \times 5}, \quad C \in \mathcal{M}_{5 \times 2}, \quad D \in \mathcal{M}_{4 \times 2}, \quad E \in \mathcal{M}_{5 \times 4},$$

for each of the cases below, determine whether the given computation is defined. If it is, specify the size of the resulting matrix.

- | | | |
|--------------|------------------|--------------------|
| 1. BA | 2. $A - 3E^\top$ | 3. $BC - 3D$ |
| 4. $AB^\top$ | 5. $E(5B + A)$ | 6. $D^\top(BE)$ |
| 7. $AC + D$ | 8. $CD^\top$ | 9. $B^\top D + ED$ |
| 10. $E(AC)$ | 11. DC | 12. $BA^\top + D$ |

Question 7. Determine which of the following matrices are square, diagonal, upper or lower triangular, symmetric, or skew-symmetric. Compute the transpose of each matrix.

$$A = \begin{pmatrix} -1 & 4 \\ 0 & 1 \\ 6 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}, \quad C = \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} -1 \\ 4 \\ 2 \end{pmatrix}$$

$$E = \begin{pmatrix} 0 & 0 & 6 \\ 0 & -6 & 0 \\ -6 & 0 & 0 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

$$G = \begin{pmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

$$H = \begin{pmatrix} 0 & -1 & 6 & 2 \\ 1 & 0 & -7 & 1 \\ -6 & 7 & 0 & -4 \\ -2 & -1 & 4 & 0 \end{pmatrix}$$

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$K = \begin{pmatrix} 1 & 2 & 3 & 4 \\ -2 & 1 & 5 & 6 \\ -3 & -5 & 1 & 7 \\ -4 & -6 & -7 & 1 \end{pmatrix}$$

$$L = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

$$N = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$Q = \begin{pmatrix} -2 & 0 & 0 \\ 4 & 0 & 0 \\ -1 & 2 & 3 \end{pmatrix}$$

$$R = \begin{pmatrix} 6 & 2 \\ 3 & -2 \\ -1 & 0 \end{pmatrix}$$

For Questions 8-11, let

$$A = \begin{pmatrix} -2 & 3 \\ 6 & 5 \\ 1 & -4 \end{pmatrix}$$

$$B = \begin{pmatrix} -5 & 3 & 6 \\ 3 & 8 & 0 \\ -2 & 0 & 4 \end{pmatrix}$$

$$C = \begin{pmatrix} 11 & -2 \\ -4 & -2 \\ 3 & -1 \end{pmatrix}$$

$$D = \begin{pmatrix} -1 & 4 & 3 & 7 \\ 2 & 1 & 7 & 5 \\ 0 & 5 & 5 & -2 \end{pmatrix}$$

$$E = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

$$F = \begin{pmatrix} 9 & -3 \\ 5 & -4 \\ 2 & 0 \\ 8 & -3 \end{pmatrix}$$

$$G = \begin{pmatrix} 5 & 1 & 0 \\ 0 & -2 & -1 \\ 1 & 0 & 3 \end{pmatrix}$$

$$H = \begin{pmatrix} 6 & 3 & 1 \\ 1 & -15 & -5 \\ -2 & -1 & 10 \end{pmatrix}$$

$$J = \begin{pmatrix} 8 \\ -1 \\ 4 \end{pmatrix}$$

$$K = \begin{pmatrix} 2 & 1 & -5 \\ 0 & 2 & 7 \end{pmatrix}$$

$$L = \begin{pmatrix} 10 & 9 \\ 8 & 7 \end{pmatrix}$$

$$M = \begin{pmatrix} 7 & -1 \\ 11 & 3 \end{pmatrix}$$

$$N = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

$$P = \begin{pmatrix} 3 & -1 \\ 4 & 7 \end{pmatrix}$$

$$Q = \begin{pmatrix} 1 & 4 & -1 & 6 \\ 8 & 7 & -3 & 3 \end{pmatrix}$$

$$R = (-3, \ 6, \ -2)$$

$$S = (6, \ -4, \ 3, \ 2)$$

$$T = (4, \ -1, \ 7)$$

Question 8. Compute each of the following, if it is defined.

- | | |
|--------------|-------------|
| 1. AB | 2. BA |
| 3. JM | 4. DF |
| 5. RJ | 6. JR |
| 7. RT | 8. SF |
| 9. KN | 10. F^2 |
| 11. B^2 | 12. E^3 |
| 13. $(TJ)^3$ | 14. $D(FK)$ |
| 15. $(CL)G$ | |

Question 9. Determine which of the following pairs of matrices commute.

- | | |
|----------------|----------------|
| 1. L and M | 2. F and Q |
| 3. G and H | 4. R and J |
| 5. A and K | 6. N and P |

Question 10. Find the specified row or column of the matrix product for the given matrices.

- The second row of the product BG

2. The third column of the product DE
3. The first column of the product SE
4. The third row of the product FQ

Question 11. Verify whether the following computations can be performed. If they are possible, identify which of the specified equalities hold. Provide justification by naming the relation and referencing the mathematical properties that confirm or disprove the validity of the equality.

1. $(RG)H = R(GH)$
2. $LP = PL$
3. $E(FK) = (EF)K$
4. $K(A + C) = KA + KC$
5. $(QF)^\top = F^\top Q^\top$
6. $L(ML) = L^2M$
7. $GC + HC = (G + H)C$
8. $R(J + T^\top) = RJ + RT^\top$
9. $(AK)^\top = A^\top K^\top$
10. $(Q + F^\top)E^\top = QE^\top + (EF)^\top$

Question 12. Given matrices

$$A = \begin{pmatrix} 3 & -2 & 7 \\ 6 & 5 & 4 \\ 0 & 4 & 9 \end{pmatrix}, \quad B = \begin{pmatrix} 6 & -2 & 4 \\ 0 & 1 & 3 \\ 7 & 7 & 5 \end{pmatrix};$$

or

$$A = \begin{pmatrix} 1 & 5 & -2 & 1 \\ 0 & 1 & -2 & 3 \\ 1 & 1 & 2 & 1 \\ 0 & 0 & 2 & -4 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 & 5 & -1 \\ 2 & 1 & 2 & 1 \\ 4 & -2 & -1 & 0 \\ -1 & 2 & 1 & 3 \end{pmatrix}$$

Compute the specified row or column of the matrix product of the given matrices, both for the 3×3 and 4×4 cases:

1. The first row of the product AB
2. The third row of the product AB
3. The second column of the product AB
4. The first column of the product BA

5. The second column of the product BA
6. The third row of the product AA
7. The third column of the product AA
8. The first column of the product AA
9. The second column of the product BB
10. The second row of the product BB
11. The third column of the product BAA

Question 13.

1. Find a non-diagonal matrix A such that $A^2 = I_2$.
2. Find a non-diagonal matrix A such that $A^2 = I_3$ (modify the result from part (a)).
3. Find a non-identity matrix A such that $A^3 = I_3$.