

# Tutorial 1

## Vectors and matrices

### Homework

1. Vectors, Questions 1–11: Provide detailed solutions to any five subparts of your choice. The five subparts may come from different questions; you do not need to complete all subparts of a selected question.
2. Vectors, Question 12: Provide detailed solutions to Parts 2, 3, and 4.
3. Matrices, Questions 2–5: Provide detailed solutions to any ten subparts of your choice from Questions 2–5. The ten subparts may be selected independently across these questions.
4. Matrices, Questions 8–11: Provide detailed solutions to any ten subparts of your choice from Questions 8–11. The ten subparts may be selected independently across these questions.
5. Matrices, Question 12: Select any three subparts from Parts 1–11 and provide detailed solutions. For each selected subpart, solve both the  $3 \times 3$  case and the  $4 \times 4$  case.

## 1 Vectors

**Question 1.** Given points  $A, B, C$ , find the missing coordinate of  $B$  such that

a  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  are orthogonal

b  $\overrightarrow{AB}$  and  $\overrightarrow{AC}$  are parallel

1.  $A = (3, 5)$ ,  $B = (2, y)$ ,  $C = (2, 8)$
2.  $A = (-2, 5)$ ,  $B = (1, y)$ ,  $C = (4, -3)$
3.  $A = (1, 5)$ ,  $B = (-1, y)$ ,  $C = (2, -3)$
4.  $A = (2, 1)$ ,  $B = (x, -2)$ ,  $C = (1, 3)$

*Solution.*

1. We have

$$\mathbf{a} := \overrightarrow{AB} = (-1, y - 5), \quad \mathbf{b} := \overrightarrow{AC} = (-1, 3)$$

- a. If  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal, then

$$\mathbf{a} \cdot \mathbf{b} = 1 + 3y - 15 = 0 \implies y = \frac{14}{3}$$

- b. If  $\mathbf{a}$  and  $\mathbf{b}$  are parallel, then  $\exists \alpha \in \mathbb{R}$ ,  $\alpha \neq 0$  s.t.

$$\mathbf{a} = \alpha \mathbf{b} \implies (-1, y - 5) = \alpha (-1, 3) \implies (-1, y - 5) = (-\alpha, 3\alpha),$$

which gives

$$\alpha = 1, \quad y - 5 = 3\alpha \implies y = 8.$$

2. We have

$$\mathbf{a} := \overrightarrow{AB} = (3, y-5), \quad \mathbf{b} := \overrightarrow{AC} = (6, -8)$$

a. If  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal, then

$$\mathbf{a} \cdot \mathbf{b} = 18 - 8y + 40 = 0 \implies y = \frac{40+18}{8} = \frac{29}{4}.$$

b. If  $\mathbf{a}$  and  $\mathbf{b}$  are parallel, then  $\exists \alpha \in \mathbb{R}, \alpha \neq 0$  s.t.

$$\mathbf{a} = \alpha \mathbf{b} \implies (3, y-5) = \alpha (6, -8) \implies (3, y-5) = (6\alpha, -8\alpha),$$

which gives

$$\alpha = \frac{1}{2}, \quad y-5 = -8\alpha \implies y = -4 + 5 = 1.$$

3. We have

$$\mathbf{a} := \overrightarrow{AB} = (-2, y-5), \quad \mathbf{b} := \overrightarrow{AC} = (1, -8)$$

a. If  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal, then

$$\mathbf{a} \cdot \mathbf{b} = -2 - 8y + 40 = 0 \implies y = \frac{38}{8} = \frac{19}{4}.$$

b. If  $\mathbf{a}$  and  $\mathbf{b}$  are parallel, then  $\exists \alpha \in \mathbb{R}, \alpha \neq 0$  s.t.

$$\mathbf{a} = \alpha \mathbf{b} \implies (-2, y-5) = \alpha (1, -8) \implies (-2, y-5) = (\alpha, -8\alpha),$$

which gives

$$\alpha = -2, \quad y-5 = -8\alpha \implies y = 16 + 5 = 21.$$

4. We have

$$\mathbf{a} := \overrightarrow{AB} = (x-2, -3), \quad \mathbf{b} := \overrightarrow{AC} = (-1, 2)$$

a. If  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal, then

$$\mathbf{a} \cdot \mathbf{b} = 2 - x - 6 = 0 \implies x = -4.$$

b. If  $\mathbf{a}$  and  $\mathbf{b}$  are parallel, then  $\exists \alpha \in \mathbb{R}, \alpha \neq 0$  s.t.

$$\mathbf{a} = \alpha \mathbf{b} \implies (x-2, -3) = \alpha (-1, 2) \implies (x-2, -3) = (-\alpha, 2\alpha),$$

which gives

$$\alpha = -\frac{3}{2}, \quad x-2 = -\alpha \implies x = \frac{7}{2}.$$

**Question 2.** Consider the triangle with vertices  $A, B$ , and  $C$ ,  $\triangle ABC$ . Define the vectors


$$\mathbf{a} = \overrightarrow{BC}, \quad \mathbf{b} = \overrightarrow{AC}, \quad \mathbf{c} = \overrightarrow{AB}.$$

Let  $M, N$ , and  $P$  denote the midpoints of the sides  $BC, AC$ , and  $AB$ , respectively.

Determine the expressions for the vectors  $\overrightarrow{AM}$ ,  $\overrightarrow{BN}$ , and  $\overrightarrow{CP}$  in terms of  $\mathbf{a}$ ,  $\mathbf{b}$ , and  $\mathbf{c}$ . Subsequently, rewrite these expressions using only the vectors  $\mathbf{a}$  and  $\mathbf{b}$ .

*Solution.*

$$\overrightarrow{AM} = \overrightarrow{AC} + \overrightarrow{CM} = \mathbf{b} - \frac{1}{2}\mathbf{a}, \quad \overrightarrow{BN} = \overrightarrow{BC} + \overrightarrow{CN} = \mathbf{a} - \frac{1}{2}\mathbf{b}, \quad \overrightarrow{CP} = \overrightarrow{CB} + \overrightarrow{BP} = -\mathbf{a} - \frac{1}{2}\mathbf{c}$$



Since

$$\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC},$$

we have

$$\mathbf{c} + \mathbf{a} = \mathbf{b} \implies \mathbf{c} = \mathbf{b} - \mathbf{a},$$

which gives

$$\overrightarrow{CP} = -\mathbf{a} - \frac{1}{2}\mathbf{c} = -\mathbf{a} - \frac{1}{2}\mathbf{b} + \frac{1}{2}\mathbf{a} = -\frac{1}{2}\mathbf{a} - \frac{1}{2}\mathbf{b}$$

**Question 3.** Given points  $A, B$ , find  $x$  such that the norm of the vector  $\overrightarrow{AB}$ ,  $\|\overrightarrow{AB}\|$  equals the specified value  $d$ .

1.  $A = (2, -3)$ ,  $B = (x, 0)$ ,  $d = 5$
2.  $A = (1, 4)$ ,  $B = (x, 1)$ ,  $d = 4$
3.  $A = (1, 5)$ ,  $B = (1, x)$ ,  $d = 6$

*Solution.*

1. By definition

$$\|\overrightarrow{AB}\| = \|(x - 2, 3)\| = \sqrt{(x - 2)^2 + 9}.$$

We have

$$(x - 2)^2 + 9 = d^2 = 25 \implies x^2 + 4 - 4x - 16 = 0 \implies x^2 - 4x - 12 = 0 \implies x = 6, -2$$

2. By definition

$$\|\overrightarrow{AB}\| = \|(x - 1, -3)\| = \sqrt{(x - 1)^2 + 9}.$$

We have

$$(x - 1)^2 + 9 = d^2 = 16 \implies x^2 - 2x - 6 = 0 \implies x = 1 - \sqrt{7}, 1 + \sqrt{7}$$

3. By definition

$$\|\overrightarrow{AB}\| = \|(0, x - 5)\| = \sqrt{(x - 5)^2}.$$

We have

$$(x - 5)^2 = d^2 = 6^2 \implies x = 6 + 5, -6 + 5 \implies x = 11, -1$$

**Question 4.** Prove that the triangle  $\triangle ABC$  is isosceles.

1.  $A = (-3, -2), B = (1, 4), C = (-5, 0)$ .
2.  $A = (7, -3, 6), B = (11, -5, 3), C = (10, -7, 8)$ .

*Solution.*

1. Let

$$\mathbf{a} := \overrightarrow{AB} = (4 \ 6), \quad \mathbf{b} := \overrightarrow{BC} = (-6 \ -4), \quad \mathbf{c} := \overrightarrow{AC} = (-2 \ 2).$$

And let  $\theta_1, \theta_2, \theta_3$  denote the angles  $\angle BAC, \angle ABC, \angle BCA$  respectively. Then

$$\begin{aligned} \cos \theta_1 &= \frac{\mathbf{a} \cdot \mathbf{c}}{\|\mathbf{a}\| \|\mathbf{c}\|} = \frac{-8 + 12}{\sqrt{16 + 36} \sqrt{4 + 4}} = \frac{4}{\sqrt{52} \sqrt{8}} = \frac{1}{\sqrt{26}} \\ \cos \theta_2 &= \frac{(-\mathbf{a}) \cdot \mathbf{b}}{\|-\mathbf{a}\| \|\mathbf{b}\|} = \frac{24 + 24}{\sqrt{52} \sqrt{52}} = \frac{48}{52} = \frac{12}{13} \\ \cos \theta_3 &= \frac{(-\mathbf{b}) \cdot (-\mathbf{c})}{\|-\mathbf{b}\| \|\mathbf{c}\|} = \frac{12 - 8}{\sqrt{52} \sqrt{8}} = \frac{1}{\sqrt{26}} \end{aligned}$$

Since  $\cos \theta_1 = \cos \theta_3$  and both angles lie in the range  $(0, \pi)$ , they must be equal.

2. Let

$$\mathbf{a} := \overrightarrow{AB} = (4 \ -2 \ -3), \quad \mathbf{b} := \overrightarrow{BC} = (-1 \ -2 \ 5), \quad \mathbf{c} := \overrightarrow{AC} = (3 \ -4 \ 2).$$

And let  $\theta_1, \theta_2, \theta_3$  denote the angles  $\angle BAC, \angle ABC, \angle BCA$  respectively. Then

$$\begin{aligned} \cos \theta_1 &= \frac{\mathbf{a} \cdot \mathbf{c}}{\|\mathbf{a}\| \|\mathbf{c}\|} = \frac{12 + 8 - 6}{\sqrt{16 + 4 + 9} \sqrt{9 + 16 + 4}} = \frac{14}{\sqrt{29} \sqrt{29}} = \frac{14}{29} \\ \cos \theta_2 &= \frac{(-\mathbf{a}) \cdot \mathbf{b}}{\|-\mathbf{a}\| \|\mathbf{b}\|} = \frac{4 - 4 + 15}{\sqrt{16 + 4 + 9} \sqrt{1 + 4 + 25}} = \frac{15}{\sqrt{29} \sqrt{30}} = \frac{\sqrt{870}}{58} \\ \cos \theta_3 &= \frac{(-\mathbf{b}) \cdot (-\mathbf{c})}{\|-\mathbf{b}\| \|\mathbf{c}\|} = \frac{-3 + 8 + 10}{\sqrt{30} \sqrt{29}} = \frac{15}{\sqrt{29} \sqrt{30}} = \frac{\sqrt{870}}{58} \end{aligned}$$

Since  $\cos \theta_2 = \cos \theta_3$  and both angles lie in the range  $(0, \pi)$ , they must be equal.

**Question 5.** Given vectors  $\mathbf{a} = (3, -2)$  and  $\mathbf{b} = (-1, 5)$ , find vector  $\mathbf{c}$  such that:

- $\mathbf{a} \cdot \mathbf{c} = 17$ .
- $\mathbf{b} \cdot \mathbf{c} = 3$ .

*Solution.* Let  $\mathbf{c} = (c_1, c_2)$ . Then by the definition of dot product, we have

$$\mathbf{a} \cdot \mathbf{c} = 3c_1 - 2c_2 = 17, \tag{1}$$

$$\mathbf{b} \cdot \mathbf{c} = -c_1 + 5c_2 = 3. \tag{2}$$

Next, multiply Equation (2) by 3 and add it to Equation (1) to eliminate  $c_1$

$$(15 - 2)c_2 = 26 \implies c_2 = \frac{26}{13} = 2.$$

Substituting  $c_2 = 2$  into Equation (2), we find:

$$c_1 = 5c_2 - 3 = 10 - 3 = 7.$$

We have  $\mathbf{c} = (7, 2)$

**Question 6.** Find all vectors  $\mathbf{a}$  that are orthogonal to the given vector  $\mathbf{b}$  and have norm  $d$ .

1.  $\mathbf{b} = (3, 4), d = 15$
2.  $\mathbf{b} = (-3, 2), d = 10$
3.  $\mathbf{b} = (1, 4), d = 8$
4.  $\mathbf{b} = (-1, 2), d = 7$
5.  $\mathbf{b} = (2, -5), d = 21$

*Solution.* Let  $\mathbf{a} = (a_1, a_2)$

1. By the given conditions, we have

$$\mathbf{a} \cdot \mathbf{b} = 3a_1 + 4a_2 = 0, \quad \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = 15.$$

Then we have the following two equations:

$$a_1 = -\frac{4}{3}a_2 \tag{3}$$

$$a_1^2 + a_2^2 = 15^2 \tag{4}$$

Substitute  $a_1$  using Equation (3) in Equation (4) gives

$$a_2^2 \left( \frac{16}{9} + 1 \right) = 225 \implies a_2^2 = 225 \times \frac{9}{25} = 81 \implies a_2 = \pm 9$$

Together with Equation (3) we have

$$a_1 = -\frac{4}{3} \times 9 = -12, \text{ or } a_1 = \frac{4}{3} \times 9 = 12.$$

Thus

$$\mathbf{a} = (-12, 9) \text{ or } (12, -9).$$

2. By the given conditions, we have

$$\mathbf{a} \cdot \mathbf{b} = -3a_1 + 2a_2 = 0, \quad \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = 10.$$

Then we have the following two equations:

$$a_1 = \frac{2}{3}a_2 \tag{5}$$

$$a_1^2 + a_2^2 = 100 \tag{6}$$

Substitute  $a_1$  using Equation (5) in Equation (6) gives

$$a_2^2 \left( \frac{4}{9} + 1 \right) = 100 \implies a_2^2 = 100 \times \frac{9}{13} = \frac{900}{13} \implies a_2 = \pm \frac{30\sqrt{13}}{13}$$

Together with Equation (5) we have

$$a_1 = \frac{2}{3} \times \frac{30\sqrt{13}}{13} = \frac{20\sqrt{13}}{13}, \text{ or } a_1 = -\frac{2}{3} \times \frac{30\sqrt{13}}{13} = -\frac{20\sqrt{13}}{13}.$$

Thus

$$\mathbf{a} = \left( \frac{20\sqrt{13}}{13}, \frac{30\sqrt{13}}{13} \right) \text{ or } \left( -\frac{20\sqrt{13}}{13}, -\frac{30\sqrt{13}}{13} \right).$$

3. By the given conditions, we have

$$\mathbf{a} \cdot \mathbf{b} = a_1 + 4a_2 = 0, \quad \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = 8.$$

Then we have the following two equations:

$$a_1 = -4a_2 \tag{7}$$

$$a_1^2 + a_2^2 = 64 \tag{8}$$

Substitute  $a_1$  using Equation (7) in Equation (8) gives

$$a_2^2(16 + 1) = 64 \implies a_2^2 = \frac{64}{17} \implies a_2 = \pm 8\frac{\sqrt{17}}{17}$$

Together with Equation (7) we have

$$a_1 = (-4) \times 8\frac{\sqrt{17}}{17} = -32\frac{\sqrt{17}}{17}, \text{ or } a_1 = 4 \times 8\frac{\sqrt{17}}{17} = 32\frac{\sqrt{17}}{17}.$$

Thus

$$\mathbf{a} = \left( 32\frac{\sqrt{17}}{17}, -8\frac{\sqrt{17}}{17} \right) \text{ or } \left( -32\frac{\sqrt{17}}{17}, 8\frac{\sqrt{17}}{17} \right).$$

4. By the given conditions, we have

$$\mathbf{a} \cdot \mathbf{b} = -a_1 + 2a_2 = 0, \quad \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = 7.$$

Then we have the following two equations:

$$a_1 = 2a_2 \tag{9}$$

$$a_1^2 + a_2^2 = 49 \tag{10}$$

Substitute  $a_1$  using Equation (9) in Equation (10) gives

$$a_2^2(4 + 1) = 49 \implies a_2^2 = \frac{49}{5} \implies a_2 = \pm 7\frac{\sqrt{5}}{5}$$

Together with Equation (9) we have

$$a_1 = 2 \times 7\frac{\sqrt{5}}{5} = 14\frac{\sqrt{5}}{5}, \text{ or } a_1 = -2 \times 7\frac{\sqrt{5}}{5} = -14\frac{\sqrt{5}}{5}.$$

Thus

$$\mathbf{a} = \left( 14\frac{\sqrt{5}}{5}, 7\frac{\sqrt{5}}{5} \right) \text{ or } \left( -14\frac{\sqrt{5}}{5}, -7\frac{\sqrt{5}}{5} \right).$$

5. By the given conditions, we have

$$\mathbf{a} \cdot \mathbf{b} = 2a_1 - 5a_2 = 0, \quad \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2} = 21.$$

Then we have the following two equations:

$$a_1 = \frac{5}{2}a_2 \quad (11)$$

$$a_1^2 + a_2^2 = 441 \quad (12)$$

Substitute  $a_1$  using Equation (11) in Equation (12) gives

$$a_2^2 \left( \frac{25}{4} + 1 \right) = 441 \implies a_2^2 = \frac{441 \times 4}{29} \implies a_2 = \pm 42 \frac{\sqrt{29}}{29}$$

Together with Equation (11) we have

$$a_1 = \frac{5}{2} \times 42 \frac{\sqrt{29}}{29} = 105 \frac{\sqrt{29}}{29}, \text{ or } a_1 = -\frac{5}{2} \times 42 \frac{\sqrt{29}}{29} = -105 \frac{\sqrt{29}}{29}.$$

Thus

$$\mathbf{a} = \left( 105 \frac{\sqrt{29}}{29}, 42 \frac{\sqrt{29}}{29} \right) \text{ or } \left( -105 \frac{\sqrt{29}}{29}, -42 \frac{\sqrt{29}}{29} \right).$$

**Question 7.** Find the vector  $\mathbf{c}$ .

1. From the equation:

$$2\mathbf{c} + 3\mathbf{a} = \mathbf{b}, \quad \text{where } \mathbf{a} = (-1, 2), \quad \mathbf{b} = (0, -2).$$

2. From the equation:

$$3\mathbf{c} - \mathbf{a} = 2\mathbf{b}, \quad \text{where } \mathbf{a} = (3, -1), \quad \mathbf{b} = (0, 2).$$

3. From the equation:

$$2\mathbf{c} - 2\mathbf{a} = \mathbf{b}, \quad \text{where } \mathbf{a} = (3, -1), \quad \mathbf{b} = (0, 2).$$

4. From the equation:

$$3\mathbf{c} + 5\mathbf{a} = 4\mathbf{b}, \quad \text{where } \mathbf{a} = (2, 4), \quad \mathbf{b} = (1, -2).$$

*Solution.* Let  $\mathbf{c} = (c_1, c_2)$ .

1. By the given conditions, we have

$$\begin{aligned} 2(c_1, c_2) + 3(-1, 2) &= (0, -2) \\ (2c_1 - 3, 2c_2 + 6) &= (0, -2), \end{aligned}$$

which gives

$$2c_1 - 3 = 0, \quad 2c_2 + 6 = -2.$$

And

$$c_1 = \frac{3}{2}, \quad c_2 = -4, \quad \mathbf{c} = \left( \frac{3}{2}, -4 \right)$$

2. By the given conditions, we have

$$\begin{aligned} 3(c_1, c_2) - (3, -1) &= 2(0, 2) \\ (3c_1 - 3, 3c_2 + 1) &= (0, 4), \end{aligned}$$

which gives

$$3c_1 - 3 = 0, \quad 3c_2 + 1 = 4.$$

And

$$c_1 = 1, \quad c_2 = 1, \quad \mathbf{c} = (1, 1)$$

3. By the given conditions, we have

$$\begin{aligned} 2(c_1, c_2) - 2(3, -1) &= (0, 2) \\ (2c_1 - 6, 2c_2 + 2) &= (0, 2), \end{aligned}$$

which gives

$$2c_1 - 6 = 0, \quad 2c_2 + 2 = 2.$$

And

$$c_1 = 3, \quad c_2 = 0, \quad \mathbf{c} = (3, 0)$$

4. By the given conditions, we have

$$\begin{aligned} 3(c_1, c_2) + 5(2, 4) &= 4(1, -2) \\ (3c_1 + 10, 3c_2 + 20) &= (4, -8), \end{aligned}$$

which gives

$$3c_1 + 10 = 4, \quad 3c_2 + 20 = -8.$$

And

$$c_1 = -2, \quad c_2 = -\frac{28}{3}, \quad \mathbf{c} = (-2, -\frac{28}{3})$$

**Question 8.** Calculate the side lengths of the triangle  $\triangle ABC$ . Additionally, find the coordinates of a fourth point  $D$  such that the quadrilateral  $ABCD$  forms a parallelogram.

1.  $A = (-4, -2), B = (-1, 4), C = (2, 2)$
2.  $A = (5, 1), B = (4, 2), C = (-1, 4)$
3.  $A = (3, 2), B = (7, 4), C = (5, 6)$
4.  $A = (-1, 2), B = (-3, 4), C = (-3, -3)$

*Solution.* Let  $D = (d_1, d_2)$ . Then  $\overrightarrow{DC} = \overrightarrow{AB}$ .

1. The side lengths of the triangle are given by

$$\begin{aligned} \|\overrightarrow{AB}\| &= \|(-1 + 4, 4 + 2)\| = \|(3, 6)\| = \sqrt{9 + 36} = \sqrt{45} \\ \|\overrightarrow{BC}\| &= \|(2 + 1, 2 - 4)\| = \|(3, -2)\| = \sqrt{9 + 4} = \sqrt{13} \\ \|\overrightarrow{AC}\| &= \|(2 + 4, 2 + 2)\| = \|(6, 4)\| = \sqrt{36 + 16} = \sqrt{52} \end{aligned}$$

$\overrightarrow{DC} = \overrightarrow{AB}$  gives

$$(2 - d_1, 2 - d_2) = (3, 6) \implies d_1 = 2 - 3 = -1, \quad d_2 = 2 - 6 = -4.$$

We have  $D = (-1, -4)$ .

2. The side lengths of the triangle are given by

$$\begin{aligned}\|\overrightarrow{AB}\| &= \|(4-5, 2-1)\| = \|(-1, 1)\| = \sqrt{1+1} = \sqrt{2} \\ \|\overrightarrow{BC}\| &= \|(-1-4, 4-2)\| = \|(-5, 2)\| = \sqrt{25+4} = \sqrt{29} \\ \|\overrightarrow{AC}\| &= \|(-1-5, 4-1)\| = \|(-6, 3)\| = \sqrt{36+9} = \sqrt{45}\end{aligned}$$

$\overrightarrow{DC} = \overrightarrow{AB}$  gives

$$(-1-d_1, 4-d_2) = (-1, 1) \implies d_1 = 0, d_2 = 4-1 = 3.$$

We have  $D = (0, 3)$ .

3. The side lengths of the triangle are given by

$$\begin{aligned}\|\overrightarrow{AB}\| &= \|(7-3, 4-2)\| = \|(4, 2)\| = \sqrt{16+4} = \sqrt{20} \\ \|\overrightarrow{BC}\| &= \|(5-7, 6-4)\| = \|(-2, 2)\| = \sqrt{4+4} = \sqrt{8} \\ \|\overrightarrow{AC}\| &= \|(5-3, 6-2)\| = \|(2, 4)\| = \sqrt{4+16} = \sqrt{20}\end{aligned}$$

$\overrightarrow{DC} = \overrightarrow{AB}$  gives

$$(5-d_1, 6-d_2) = (4, 2) \implies d_1 = 1, d_2 = 6-2 = 4.$$

We have  $D = (1, 4)$ .

4. The side lengths of the triangle are given by

$$\begin{aligned}\|\overrightarrow{AB}\| &= \|(-3+1, 4-2)\| = \|(-2, 2)\| = \sqrt{4+4} = \sqrt{8} \\ \|\overrightarrow{BC}\| &= \|(-3+3, -3-4)\| = \|(0, -7)\| = \sqrt{0+49} = 7 \\ \|\overrightarrow{AC}\| &= \|(-3+1, -3-2)\| = \|(-2, -5)\| = \sqrt{4+25} = \sqrt{29}\end{aligned}$$

$\overrightarrow{DC} = \overrightarrow{AB}$  gives

$$(-3-d_1, -3-d_2) = (-2, 2) \implies d_1 = -1, d_2 = -5.$$

We have  $D = (-1, -5)$ .

**Question 9.** Calculate the lengths of the sides and diagonals of the specified quadrilateral  $ABCD$ , classify its type.

1.  $A = (8, -4)$ ,  $B = (5, -6)$ ,  $C = (1, -4)$ ,  $D = (4, 2)$
2.  $A = (-2, -3)$ ,  $B = (-5, -7)$ ,  $C = (-1, -10)$ ,  $D = (2, -6)$
3.  $A = (7, 5)$ ,  $B = (-6, 2)$ ,  $C = (3, -1)$ ,  $D = (6, 1)$

*Solution.*

1. The length of the four sides are given by

$$\begin{aligned}\|AB\| &= \|(-3, -2)\| = \sqrt{9+4} = \sqrt{13} \\ \|BC\| &= \|(-4, 2)\| = \sqrt{16+4} = \sqrt{20} \\ \|CD\| &= \|(3, 6)\| = \sqrt{9+36} = \sqrt{45} \\ \|DA\| &= \|(4, -6)\| = \sqrt{16+36} = \sqrt{52}\end{aligned}$$

The length of the diagonals are given by

$$\|AC\| = \|(-7, 0)\| = 7, \quad \|BD\| = \|(-1, 8)\| = \sqrt{1+64} = \sqrt{65}$$

This quadrilateral is an irregular quadrilateral.

2. The length of the four sides are given by

$$\begin{aligned}\|AB\| &= \|(-3, -4)\| = \sqrt{9+16} = 5 \\ \|BC\| &= \|(4, -3)\| = \sqrt{16+9} = 5 \\ \|CD\| &= \|(3, 4)\| = \sqrt{9+16} = 5 \\ \|DA\| &= \|(-4, 3)\| = \sqrt{16+9} = 5\end{aligned}$$

The length of the diagonals are given by

$$\|AC\| = \|(-1, -7)\| = \sqrt{50}, \quad \|BD\| = \|(7, 1)\| = \sqrt{50}$$

Furthermore, we have

$$\begin{aligned}\overrightarrow{BA} \cdot \overrightarrow{BC} &= (3, 4) \cdot (4, -3) = 0 \\ \overrightarrow{AB} \cdot \overrightarrow{AD} &= (-3, -4) \cdot (4, -3) = 0 \\ \overrightarrow{DA} \cdot \overrightarrow{DC} &= (-4, 3) \cdot (-3, -4) = 0 \\ \overrightarrow{CD} \cdot \overrightarrow{CB} &= (3, 4) \cdot (-4, 3) = 0\end{aligned}$$

showing that the four angles are right angles. Thus this quadrilateral is a square.

3. The length of the four sides are given by

$$\begin{aligned}\|AB\| &= \|(-13, -3)\| = \sqrt{169+9} = \sqrt{178} \\ \|BC\| &= \|(9, -3)\| = \sqrt{81+9} = \sqrt{90} \\ \|CD\| &= \|(3, 2)\| = \sqrt{9+4} = \sqrt{13} \\ \|DA\| &= \|(1, 4)\| = \sqrt{1+16} = \sqrt{17}\end{aligned}$$

The length of the diagonals are given by

$$\|AC\| = \|(-4, -6)\| = \sqrt{16+36} = \sqrt{52}, \quad \|BD\| = \|(12, -1)\| = \sqrt{144+1} = \sqrt{145}$$

This quadrilateral is an irregular quadrilateral.

**Question 10.** Find the interior angles of the triangle  $\triangle ABC$ , where

$$A = (5\sqrt{3}, 5), \quad B = (-\sqrt{3}, 1), \quad C = (0, 0)$$

*Solution.* Let

$$\mathbf{a} := \overrightarrow{AB} = (-6\sqrt{3}, -4), \quad \mathbf{b} := \overrightarrow{BC} = (\sqrt{3}, -1), \quad \mathbf{c} := \overrightarrow{AC} = (-5\sqrt{3}, -5).$$

And let  $\theta_1, \theta_2, \theta_3$  denote the angles  $\angle BAC, \angle ABC, \angle BCA$  respectively. Then

$$\begin{aligned}\cos \theta_1 &= \frac{\mathbf{a} \cdot \mathbf{c}}{\|\mathbf{a}\| \|\mathbf{c}\|} = \frac{90+20}{\sqrt{108+16}\sqrt{75+25}} = \frac{110}{\sqrt{124} \times 10} = \frac{11}{2\sqrt{31}} \implies \theta_1 \approx 8.95^\circ \\ \cos \theta_2 &= \frac{(-\mathbf{a}) \cdot \mathbf{b}}{\|-\mathbf{a}\| \|\mathbf{b}\|} = \frac{18-4}{\sqrt{124}\sqrt{3+1}} = \frac{7}{2\sqrt{31}} \implies \theta_2 \approx 51.05^\circ \\ \cos \theta_3 &= \frac{(-\mathbf{b}) \cdot (-\mathbf{c})}{\|-\mathbf{b}\| \|\mathbf{c}\|} = \frac{-15+5}{2 \times 10} = -\frac{1}{2} \implies \theta_3 \approx 120^\circ\end{aligned}$$

**Question 11.** Identify which of the following vector pairs are parallel.

1.  $(-2, 3, 1), (6, -4, -3)$
2.  $(10, -8, 9, 0, 24), \left(\frac{5}{6}, -\frac{2}{3}, \frac{3}{4}, 0, 2\right)$
3.  $(8, -6, -9, -8, -2), \left(-\frac{32}{3}, 8, 12, \frac{32}{3}, \frac{8}{3}\right)$
4.  $(0, 0, 2, 3, 5), \left(0, 1, \frac{4}{3}, 2, \frac{10}{3}\right)$

*Solution.*

1. Suppose the two vectors were parallel, then there exists  $\alpha \in \mathbb{R}$  such that

$$(-2, 3, 1) = \alpha(6, -4, -3).$$

In particular, we have

$$-2 = 6\alpha \implies \alpha = -\frac{1}{3}, \quad 3 = -4\alpha \implies \alpha = -\frac{3}{4},$$

a contradiction. Therefore, the two vectors are not parallel.

2. Those two vectors are parallel because

$$(10, -8, 9, 0, 24) = 12 \left(\frac{5}{6}, -\frac{2}{3}, \frac{3}{4}, 0, 2\right)$$

3. Those two vectors are parallel because

$$(8, -6, -9, -8, -2) = -\frac{3}{4} \left(-\frac{32}{3}, 8, 12, \frac{32}{3}, \frac{8}{3}\right)$$

4. Suppose the two vectors were parallel, then there exists  $\alpha \in \mathbb{R}, \alpha \neq 0$ , such that

$$(0, 0, 2, 3, 5) = \alpha \left(0, 1, \frac{4}{3}, 2, \frac{10}{3}\right).$$

In particular,

$$0 = \alpha \times 1,$$

which implies  $\alpha = 0$ , contradicting  $\alpha \neq 0$ .

**Question 12.** Compute  $\text{proj}_{\mathbf{a}} \mathbf{b}$ ,  $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$ ,  $\text{proj}_{\mathbf{b}} \mathbf{a}$  and  $\mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a}$ .

1.  $\mathbf{a} = (2, 1, 5), \mathbf{b} = (1, 4, -5)$
2.  $\mathbf{a} = (6, 2, 4), \mathbf{b} = (1, -5, 4)$
3.  $\mathbf{a} = (1, 0, 1, -2), \mathbf{b} = (5, -1, 0, 2)$
4.  $\mathbf{a} = (-2, 3, -4, 1), \mathbf{b} = (5, 1, -8, 2)$

*Solution.*

1. We have

$$\begin{aligned}\text{proj}_{\mathbf{a}} \mathbf{b} &= \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\|^2} \mathbf{a} = \frac{2+4-25}{4+1+25} \mathbf{a} = \frac{-19}{30} (2, 1, 5) = \left(-\frac{19}{15}, -\frac{19}{30}, -\frac{19}{6}\right), \\ \mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b} &= \left(1 + \frac{19}{15}, 4 + \frac{19}{30}, -5 + \frac{19}{6}\right) = \left(\frac{34}{15}, \frac{139}{30}, -\frac{11}{6}\right) \\ \text{proj}_{\mathbf{b}} \mathbf{a} &= \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{b}\|^2} \mathbf{b} = \frac{-19}{1+16+25} (1, 4, -5) = -\frac{19}{42} (1, 4, -5) = \left(-\frac{19}{42}, -\frac{38}{21}, \frac{95}{42}\right) \\ \mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(2 + \frac{19}{42}, 1 + \frac{38}{21}, 5 - \frac{95}{42}\right) = \left(\frac{103}{42}, \frac{59}{21}, \frac{115}{42}\right).\end{aligned}$$

2. Answers:

$$\begin{aligned}\text{proj}_{\mathbf{a}} \mathbf{b} &= \left(\frac{9}{7}, \frac{3}{7}, \frac{6}{7}\right), & \mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b} &= \left(-\frac{2}{7}, -\frac{38}{7}, \frac{22}{7}\right), \\ \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(\frac{2}{7}, -\frac{10}{7}, \frac{8}{7}\right), & \mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(\frac{40}{7}, \frac{24}{7}, \frac{20}{7}\right)\end{aligned}$$

3. Answers:

$$\begin{aligned}\text{proj}_{\mathbf{a}} \mathbf{b} &= \left(\frac{1}{6}, 0, \frac{1}{6}, -\frac{1}{3}\right), & \mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b} &= \left(\frac{29}{6}, -1, -\frac{1}{6}, \frac{7}{3}\right), \\ \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(\frac{1}{6}, -\frac{1}{30}, 0, \frac{1}{15}\right), & \mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(\frac{5}{6}, \frac{1}{30}, 1, -\frac{31}{15}\right)\end{aligned}$$

4. Answers:

$$\begin{aligned}\text{proj}_{\mathbf{a}} \mathbf{b} &= \left(-\frac{9}{5}, \frac{27}{10}, -\frac{18}{5}, \frac{9}{10}\right), & \mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b} &= \left(\frac{34}{5}, -\frac{17}{10}, -\frac{22}{5}, \frac{11}{10}\right), \\ \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(\frac{135}{94}, \frac{27}{94}, -\frac{108}{47}, \frac{27}{47}\right), & \mathbf{a} - \text{proj}_{\mathbf{b}} \mathbf{a} &= \left(-\frac{323}{94}, \frac{255}{94}, -\frac{80}{47}, \frac{20}{47}\right)\end{aligned}$$

**Question 13.** For any  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$  with  $\mathbf{a} \neq \mathbf{0}$ , show that  $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$  is orthogonal to  $\mathbf{a}$ .

*Solution.* Take any  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$  with  $\mathbf{a} \neq \mathbf{0}$ . Then

$$\mathbf{a} \cdot (\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}) = \mathbf{a} \cdot \mathbf{b} - \mathbf{a} \cdot \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\|^2} \mathbf{a} = \mathbf{a} \cdot \mathbf{b} - \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\|^2} (\mathbf{a} \cdot \mathbf{a}) = \mathbf{a} \cdot \mathbf{b} - \mathbf{a} \cdot \mathbf{b} = 0.$$

Therefore,  $\mathbf{b} - \text{proj}_{\mathbf{a}} \mathbf{b}$  is orthogonal to  $\mathbf{a}$ .

**Question 14.** Take any  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^n$ , where both vectors are nonzero.

1. Suppose  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal. Find  $\text{proj}_{\mathbf{a}} \mathbf{b}$ .

2. Suppose  $\mathbf{a}$  and  $\mathbf{b}$  are parallel. Find  $\text{proj}_{\mathbf{a}} \mathbf{b}$ .

*Solution.*

1. If  $\mathbf{a}$  and  $\mathbf{b}$  are orthogonal, by definition, we have  $\mathbf{a} \cdot \mathbf{b} = 0$ . Then

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\|^2} \mathbf{a} = \mathbf{0}.$$

2. If  $\mathbf{a}$  and  $\mathbf{b}$  are parallel, by definition, there exists  $\alpha \in \mathbb{R}$ ,  $\alpha \neq 0$  such that  $\mathbf{b} = \alpha \mathbf{a}$ . Then

$$\text{proj}_{\mathbf{a}} \mathbf{b} = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\|^2} \mathbf{a} = \frac{\alpha \mathbf{a} \cdot \mathbf{a}}{\|\mathbf{a}\|^2} \mathbf{a} = \alpha \mathbf{a} = \mathbf{b}.$$

## 2 Matrices

**Question 1.** Let

$$A = \begin{pmatrix} -4 & 2 & 3 \\ 0 & 5 & -1 \\ 6 & 1 & 2 \end{pmatrix} \quad B = \begin{pmatrix} 6 & -1 & 0 \\ 2 & 2 & -4 \\ 3 & -1 & 1 \end{pmatrix} \quad C = \begin{pmatrix} 5 & -1 \\ -3 & 4 \end{pmatrix}$$

$$D = \begin{pmatrix} -7 & 1 & -4 \\ 3 & -2 & 8 \end{pmatrix} \quad E = \begin{pmatrix} 3 & -3 & 5 \\ 1 & 0 & -2 \\ 6 & 7 & -2 \end{pmatrix} \quad F = \begin{pmatrix} 8 & -1 \\ 2 & 0 \\ 5 & -3 \end{pmatrix}$$

Compute each of the following, if it is defined.

- |                  |                      |                                   |
|------------------|----------------------|-----------------------------------|
| 1. $A + B$       | 2. $2A - 3E - B$     | 3. $2C^\top - 3F$                 |
| 4. $C + D$       | 5. $2D - 3F$         | 6. $5(F^\top - D^\top)$           |
| 7. $4A$          | 8. $A^\top + E^\top$ | 9. $((B - A)^\top + E^\top)^\top$ |
| 10. $2A - 3B$    | 11. $(A + E)^\top$   | 12. $A - B + E$                   |
| 13. $C + 3F - E$ | 14. $4D + 3F^\top$   |                                   |

*Solution.*

$$1. A + B = \begin{pmatrix} 2 & 1 & 3 \\ 2 & 7 & -5 \\ 9 & 0 & 3 \end{pmatrix} \quad 2. 2A - 3E - B = \begin{pmatrix} -23 & 14 & -9 \\ -5 & 8 & 8 \\ -9 & -18 & 9 \end{pmatrix}$$

3.  $2C^\top - 3F$  is not defined

4.  $C + D$  is not defined

5.  $2D - 3F$  is not defined

6.  $5(F^\top - D^\top)$  is not defined

$$7. 4A = \begin{pmatrix} -16 & 8 & 12 \\ 0 & 20 & -4 \\ 24 & 4 & 8 \end{pmatrix} \quad 8. A^\top + E^\top = \begin{pmatrix} -1 & 1 & 12 \\ -1 & 5 & 8 \\ 8 & -3 & 0 \end{pmatrix}$$

$$9. ((B - A)^\top + E^\top)^\top = \begin{pmatrix} 13 & -6 & 2 \\ 3 & -3 & -5 \\ 3 & 5 & -3 \end{pmatrix} \quad 10. 2A - 3B = \begin{pmatrix} -26 & 7 & 6 \\ -6 & 4 & 10 \\ 3 & 5 & 1 \end{pmatrix}$$

$$11. (A + E)^\top = \begin{pmatrix} -1 & 1 & 12 \\ -1 & 5 & 8 \\ 8 & -3 & 0 \end{pmatrix} \quad 12. A - B + E = \begin{pmatrix} -7 & 0 & 8 \\ -1 & 3 & 1 \\ 9 & 9 & -1 \end{pmatrix}$$

13.  $C + 3F - E$  is not defined

$$14. 4D + 3F^\top = \begin{pmatrix} -4 & 10 & -1 \\ 9 & -8 & 23 \end{pmatrix}$$

**For Questions 2-5, let**

$$A = \begin{pmatrix} 3 & 0 \\ -1 & 2 \\ 1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 4 & -1 \\ 0 & 2 \end{pmatrix}, \quad C = \begin{pmatrix} 1 & 4 & 2 \\ 3 & 1 & 5 \end{pmatrix}$$

$$D = \begin{pmatrix} 1 & 5 & 2 \\ -1 & 0 & 1 \\ 3 & 2 & 4 \end{pmatrix}, \quad E = \begin{pmatrix} 6 & 1 & 3 \\ -1 & 1 & 2 \\ 4 & 1 & 3 \end{pmatrix}$$

**Question 2.** Compute each of the following, if it is defined.

- |               |                        |
|---------------|------------------------|
| 1. $D + E$    | 2. $-3(D + 2E)$        |
| 3. $D - E$    | 4. $A - A$             |
| 5. $5A$       | 6. $\text{tr}(D)$      |
| 7. $-7D$      | 8. $\text{tr}(D - 3E)$ |
| 9. $2B - C$   | 10. $4\text{tr}(7B)$   |
| 11. $4E - 2D$ | 12. $\text{tr}(A)$     |

*Solution.*

|                                                                                 |                                                                                                    |
|---------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|
| 1. $D + E = \begin{pmatrix} 7 & 6 & 5 \\ -2 & 1 & 3 \\ 7 & 3 & 7 \end{pmatrix}$ | 2. $-3(D + 2E) = \begin{pmatrix} -39 & -21 & -24 \\ 9 & -6 & -15 \\ -33 & -12 & -30 \end{pmatrix}$ |
|---------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------|

|                                                                                     |                                                                    |
|-------------------------------------------------------------------------------------|--------------------------------------------------------------------|
| 3. $D - E = \begin{pmatrix} -5 & 4 & -1 \\ 0 & -1 & -1 \\ -1 & 1 & 1 \end{pmatrix}$ | 4. $A - A = \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{pmatrix}$ |
|-------------------------------------------------------------------------------------|--------------------------------------------------------------------|

|                                                                    |                       |
|--------------------------------------------------------------------|-----------------------|
| 5. $5A = \begin{pmatrix} 15 & 0 \\ -5 & 10 \\ 5 & 5 \end{pmatrix}$ | 6. $\text{tr}(D) = 5$ |
|--------------------------------------------------------------------|-----------------------|

|                                                                                          |                              |
|------------------------------------------------------------------------------------------|------------------------------|
| 7. $-7D = \begin{pmatrix} -7 & -35 & -14 \\ 7 & 0 & -7 \\ -21 & -14 & -28 \end{pmatrix}$ | 8. $\text{tr}(D - 3E) = -25$ |
|------------------------------------------------------------------------------------------|------------------------------|

|                            |                            |
|----------------------------|----------------------------|
| 9. $2B - C$ is not defined | 10. $4\text{tr}(7B) = 168$ |
|----------------------------|----------------------------|

|                                                                                       |                                                                     |
|---------------------------------------------------------------------------------------|---------------------------------------------------------------------|
| 11. $4E - 2D = \begin{pmatrix} 22 & -6 & 8 \\ -2 & 4 & 6 \\ 10 & 0 & 4 \end{pmatrix}$ | 12. $\text{tr}(A)$ is not defined, since $A$ is not a square matrix |
|---------------------------------------------------------------------------------------|---------------------------------------------------------------------|

**Question 3.** Compute each of the following, if it is defined.

- |                                       |                               |
|---------------------------------------|-------------------------------|
| 1. $2A^\top + C$                      | 2. $2E^\top - 3D^\top$        |
| 3. $D^\top - E^\top$                  | 4. $(2E^\top - 3D^\top)^\top$ |
| 5. $(D - E)^\top$                     | 6. $(CD)E$                    |
| 7. $B^\top + 5C^\top$                 | 8. $C(AB)$                    |
| 9. $\frac{1}{2}C^\top - \frac{1}{4}A$ | 10. $\text{tr}(DE^\top)$      |
| 11. $B - B^\top$                      | 12. $\text{tr}(BC)$           |

*Solution.*

$$1. 2A^\top + C = \begin{pmatrix} 7 & 2 & 4 \\ 3 & 5 & 7 \end{pmatrix}$$

$$2. 2E^\top - 3D^\top = \begin{pmatrix} 9 & 1 & -1 \\ -13 & 2 & -4 \\ 0 & 1 & -6 \end{pmatrix}$$

$$3. D^\top - E^\top = \begin{pmatrix} -5 & 0 & -1 \\ 4 & -1 & 1 \\ -1 & -1 & 1 \end{pmatrix}$$

$$4. (2E^\top - 3D^\top)^\top = \begin{pmatrix} 9 & -13 & 0 \\ 1 & 2 & 1 \\ -1 & -4 & -6 \end{pmatrix}$$

$$5. (D - E)^\top = \begin{pmatrix} -5 & 0 & -1 \\ 4 & -1 & 1 \\ -1 & -1 & 1 \end{pmatrix}$$

$$6. (CD)E = \begin{pmatrix} 65 & 26 & 69 \\ 185 & 69 & 182 \end{pmatrix}$$

$$7. B^\top + 5C^\top \text{ is not defined}$$

$$8. C(AB) = \begin{pmatrix} 4 & 19 \\ 52 & 1 \end{pmatrix}$$

$$9. \frac{1}{2}C^\top - \frac{1}{4}A = \begin{pmatrix} -0.25 & 1.5 \\ 2.25 & 0.0 \\ 0.75 & 2.25 \end{pmatrix}$$

$$10. \text{tr}(DE^\top) = 46$$

$$11. B - B^\top = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

$$12. \text{tr}(BC) \text{ is not defined, } BC \text{ is not a square matrix}$$

**Question 4.** Compute each of the following, if it is defined.

$$1. AB$$

$$2. BA$$

$$3. (3E)D$$

$$4. (AB)C$$

$$5. A(BC)$$

$$6. CC^\top$$

$$7. (DA)^\top$$

$$8. (C^\top B)A^\top$$

$$9. \text{tr}(DD^\top)$$

$$10. \text{tr}(4E^\top - D)$$

$$11. \text{tr}(C^\top A^\top + 2E^\top)$$

$$12. \text{tr}((EC^\top)^\top A)$$

*Solution.*

$$1. AB = \begin{pmatrix} 12 & -3 \\ -4 & 5 \\ 4 & 1 \end{pmatrix}$$

2.  $BA$  is not defined

$$3. (3E)D = \begin{pmatrix} 42 & 108 & 75 \\ 12 & -3 & 21 \\ 36 & 78 & 63 \end{pmatrix}$$

$$4. (AB)C = \begin{pmatrix} 3 & 45 & 9 \\ 11 & -11 & 17 \\ 7 & 17 & 13 \end{pmatrix}$$

$$5. A(BC) = \begin{pmatrix} 3 & 45 & 9 \\ 11 & -11 & 17 \\ 7 & 17 & 13 \end{pmatrix}$$

$$6. CC^\top = \begin{pmatrix} 21 & 17 \\ 17 & 35 \end{pmatrix}$$

$$7. (DA)^\top = \begin{pmatrix} 0 & -2 & 11 \\ 12 & 1 & 8 \end{pmatrix}$$

$$8. (C^\top B)A^\top = \begin{pmatrix} 12 & 6 & 9 \\ 48 & -20 & 14 \\ 24 & 8 & 16 \end{pmatrix}$$

$$9. \text{tr}(DD^\top) = 61$$

$$10. \text{tr}(4E^\top - D) = 35$$

$$11. \text{tr}(C^\top A^\top + 2E^\top) = 28$$

$$12. \text{tr}((EC^\top)^\top A) = 99$$

**Question 5.** Compute each of the following, if it is defined.

$$1. (2D^\top - E)A$$

$$2. (BA^\top - 2C)^\top$$

$$3. (4B)C + 2B$$

$$4. B^\top(CC^\top - A^\top A)$$

$$5. (-AC)^\top + 5D^\top$$

$$6. D^\top E^\top - (ED)^\top$$

*Solution.*

$$1. (2D^\top - E)A = \begin{pmatrix} -6 & -3 \\ 36 & 0 \\ 4 & 7 \end{pmatrix}$$

$$2. (BA^\top - 2C)^\top = \begin{pmatrix} 10 & -6 \\ -14 & 2 \\ -1 & -8 \end{pmatrix}$$

3.  $(4B)C + 2B$  is not defined

$$4. B^\top(CC^\top - A^\top A) = \begin{pmatrix} 40 & 72 \\ 26 & 42 \end{pmatrix}.$$

$$5. (-AC)^\top + 5D^\top = \begin{pmatrix} 2 & -10 & 11 \\ 13 & 2 & 5 \\ 4 & -3 & 13 \end{pmatrix}$$

$$6. D^\top E^\top - (ED)^\top = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

**Question 6.** Given matrices

$$A \in \mathcal{M}_{4 \times 5}, \quad B \in \mathcal{M}_{4 \times 5}, \quad C \in \mathcal{M}_{5 \times 2}, \quad D \in \mathcal{M}_{4 \times 2}, \quad E \in \mathcal{M}_{5 \times 4},$$

for each of the cases below, determine whether the given computation is defined. If it is, specify the size of the resulting matrix.

$$1. BA$$

$$2. A - 3E^\top$$

$$3. BC - 3D$$

$$4. AB^\top$$

$$5. E(5B + A)$$

$$6. D^\top(BE)$$

$$7. AC + D$$

$$8. CD^\top$$

$$9. B^\top D + ED$$

$$10. E(AC)$$

$$11. DC$$

$$12. BA^\top + D$$

*Solution.*

- |                                           |                                               |                                                 |
|-------------------------------------------|-----------------------------------------------|-------------------------------------------------|
| 1. $BA$ is not defined                    | 2. $A - 3E^\top \in \mathcal{M}_{4 \times 5}$ | 3. $BC - 3D \in \mathcal{M}_{4 \times 2}$       |
| 4. $AB^\top \in \mathcal{M}_{4 \times 4}$ | 5. $E(5B + A) \in \mathcal{M}_{5 \times 5}$   | 6. $D^\top(BE) \in \mathcal{M}_{2 \times 4}$    |
| 7. $AC + D \in \mathcal{M}_{4 \times 2}$  | 8. $CD^\top \in \mathcal{M}_{5 \times 4}$     | 9. $B^\top D + ED \in \mathcal{M}_{5 \times 2}$ |
| 10. $E(AC) \in \mathcal{M}_{5 \times 2}$  | 11. $DC$ is not defined                       | 12. $BA^\top + D$ is not defined                |

**Question 7.** Determine which of the following matrices are square, diagonal, upper or lower triangular, symmetric, or skew-symmetric. Compute the transpose of each matrix.

$$A = \begin{pmatrix} -1 & 4 \\ 0 & 1 \\ 6 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}, \quad C = \begin{pmatrix} -1 & 1 \\ -1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} -1 \\ 4 \\ 2 \end{pmatrix}$$

$$E = \begin{pmatrix} 0 & 0 & 6 \\ 0 & -6 & 0 \\ -6 & 0 & 0 \end{pmatrix}$$

$$F = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix}$$

$$G = \begin{pmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{pmatrix}$$

$$H = \begin{pmatrix} 0 & -1 & 6 & 2 \\ 1 & 0 & -7 & 1 \\ -6 & 7 & 0 & -4 \\ -2 & -1 & 4 & 0 \end{pmatrix}$$

$$J = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$K = \begin{pmatrix} 1 & 2 & 3 & 4 \\ -2 & 1 & 5 & 6 \\ -3 & -5 & 1 & 7 \\ -4 & -6 & -7 & 1 \end{pmatrix}$$

$$L = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

$$M = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

$$N = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

$$P = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

$$Q = \begin{pmatrix} -2 & 0 & 0 \\ 4 & 0 & 0 \\ -1 & 2 & 3 \end{pmatrix}$$

$$R = \begin{pmatrix} 6 & 2 \\ 3 & -2 \\ -1 & 0 \end{pmatrix}$$

*Solution.*

$$A^\top = \begin{pmatrix} -1 & 0 & 6 \\ 4 & 1 & 0 \end{pmatrix}, \quad B^\top = \begin{pmatrix} 2 & 0 \\ 0 & -1 \end{pmatrix}, \quad C^\top = \begin{pmatrix} -1 & -1 \\ 1 & 1 \end{pmatrix}, \quad D^\top = (-1, 4, 2)$$

$$\begin{aligned}
E^\top &= \begin{pmatrix} 0 & 0 & -6 \\ 0 & -6 & 0 \\ 6 & 0 & 0 \end{pmatrix} & F^\top &= \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \end{pmatrix} & G^\top &= \begin{pmatrix} 6 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 6 \end{pmatrix} \\
H^\top &= \begin{pmatrix} 0 & 1 & -6 & -2 \\ -1 & 0 & 7 & -1 \\ 6 & -7 & 0 & 4 \\ 2 & 1 & -4 & 0 \end{pmatrix} & J^\top &= \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix} & K^\top &= \begin{pmatrix} 1 & -2 & -3 & -4 \\ 2 & 1 & -5 & -6 \\ 3 & 5 & 1 & -7 \\ 4 & 6 & 7 & 1 \end{pmatrix} \\
L^\top &= \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{pmatrix} & M^\top &= \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} & N^\top &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
P^\top &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} & Q^\top &= \begin{pmatrix} -2 & 4 & -1 \\ 0 & 0 & 2 \\ 0 & 0 & 3 \end{pmatrix} & R^\top &= \begin{pmatrix} 6 & 3 & -1 \\ 2 & -2 & 0 \end{pmatrix}
\end{aligned}$$

- Square matrices:  $B, C, E, F, G, H, J, K, L, M, N, P, Q$
- Diagonal matrices:  $B, G, N$
- Upper triangular matrices:  $B, G, L, N$
- Lower triangular matrices:  $B, G, M, N, Q$
- Symmetric matrices:  $B, F, G, J, N, P$
- Skew-symmetric matrices:  $H$

For Questions 8-11, let

$$A = \begin{pmatrix} -2 & 3 \\ 6 & 5 \\ 1 & -4 \end{pmatrix}$$

$$B = \begin{pmatrix} -5 & 3 & 6 \\ 3 & 8 & 0 \\ -2 & 0 & 4 \end{pmatrix}$$

$$C = \begin{pmatrix} 11 & -2 \\ -4 & -2 \\ 3 & -1 \end{pmatrix}$$

$$D = \begin{pmatrix} -1 & 4 & 3 & 7 \\ 2 & 1 & 7 & 5 \\ 0 & 5 & 5 & -2 \end{pmatrix}$$

$$E = \begin{pmatrix} 1 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 \end{pmatrix}$$

$$F = \begin{pmatrix} 9 & -3 \\ 5 & -4 \\ 2 & 0 \\ 8 & -3 \end{pmatrix}$$

$$G = \begin{pmatrix} 5 & 1 & 0 \\ 0 & -2 & -1 \\ 1 & 0 & 3 \end{pmatrix}$$

$$H = \begin{pmatrix} 6 & 3 & 1 \\ 1 & -15 & -5 \\ -2 & -1 & 10 \end{pmatrix}$$

$$J = \begin{pmatrix} 8 \\ -1 \\ 4 \end{pmatrix}$$

$$K = \begin{pmatrix} 2 & 1 & -5 \\ 0 & 2 & 7 \end{pmatrix}$$

$$L = \begin{pmatrix} 10 & 9 \\ 8 & 7 \end{pmatrix}$$

$$M = \begin{pmatrix} 7 & -1 \\ 11 & 3 \end{pmatrix}$$

$$N = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

$$P = \begin{pmatrix} 3 & -1 \\ 4 & 7 \end{pmatrix}$$

$$Q = \begin{pmatrix} 1 & 4 & -1 & 6 \\ 8 & 7 & -3 & 3 \end{pmatrix}$$

$$R = (-3, 6, -2)$$

$$S = (6, -4, 3, 2)$$

$$T = (4, -1, 7)$$

**Question 8.** Compute each of the following, if it is defined.

1.  $AB$

2.  $BA$

3.  $JM$

4.  $DF$

5.  $RJ$

6.  $JR$

7.  $RT$

8.  $SF$

9.  $KN$

10.  $F^2$

11.  $B^2$

12.  $E^3$

13.  $(TJ)^3$

14.  $D(FK)$

15.  $(CL)G$

*Solution.*

1.  $AB$  is not defined

2.  $BA = \begin{pmatrix} 34 & -24 \\ 42 & 49 \\ 8 & -22 \end{pmatrix}$

3.  $JM$  is not defined

4.  $DF = \begin{pmatrix} 73 & -34 \\ 77 & -25 \\ 19 & -14 \end{pmatrix}$

5.  $RJ = (-38)$

6.  $JR = \begin{pmatrix} -24 & 48 & -16 \\ 3 & -6 & 2 \\ -12 & 24 & -8 \end{pmatrix}$

7.  $RT$  is not defined

8.  $SF = (56, -8)$

9.  $KN$  is not defined

10.  $F^2$  is not defined

11.  $B^2 = \begin{pmatrix} 22 & 9 & -6 \\ 9 & 73 & 18 \\ 2 & -6 & 4 \end{pmatrix}$

12.  $E^3 = \begin{pmatrix} 5 & 3 & 2 & 5 \\ 4 & 1 & 3 & 1 \\ 1 & 1 & 0 & 2 \\ 4 & 1 & 3 & 1 \end{pmatrix}$

13.  $(TJ)^3 = (226981)$

14.  $D(FK) = \begin{pmatrix} 146 & 5 & -603 \\ 154 & 27 & -560 \\ 38 & -9 & -193 \end{pmatrix}$

15.  $(CL)G$  is not defined

**Question 9.** Determine which of the following pairs of matrices commute.

1.  $L$  and  $M$

2.  $F$  and  $Q$

3.  $G$  and  $H$

4.  $R$  and  $J$

5.  $A$  and  $K$

6.  $N$  and  $P$

*Solution.*

1.  $L$  and  $M$  do not commute

$$LM = \begin{pmatrix} 169 & 17 \\ 133 & 13 \end{pmatrix}, \quad ML = \begin{pmatrix} 62 & 56 \\ 134 & 120 \end{pmatrix}$$

2.  $F$  and  $Q$  are not square matrices, so they do not commute

3.  $H$  and  $G$  commute

$$GH = HG = \begin{pmatrix} 31 & 0 & 0 \\ 0 & 31 & 0 \\ 0 & 0 & 31 \end{pmatrix}$$

4.  $R$  and  $J$  are not square matrices, so they do not commute

5.  $A$  and  $K$  are not square matrices, so they do not commute

6.  $N$  and  $P$  commute because

$$NP = PN = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

**Question 10.** Find the specified row or column of the matrix product for the given matrices.

1. The second row of the product  $BG$

2. The third column of the product  $DE$

3. The first column of the product  $SE$
4. The third row of the product  $FQ$

*Solution.*

1. The second row of  $BG$  is given by

$$(3, \ 8, \ 0) G = (15, \ -13, \ -8)$$

2. The third column of  $DE$  is given by

$$D \begin{pmatrix} 0 \\ 1 \\ 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 11 \\ 6 \\ 3 \end{pmatrix}$$

3. The first column of  $SE$  is given by

$$S \begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix} = (4)$$

4. The third row of  $FQ$  is given by

$$(2, \ 0) Q = (2, \ 8, \ -2, \ 12)$$

**Question 11.** Verify whether the following computations can be performed. If they are possible, identify which of the specified equalities hold. Provide justification by naming the relation and referencing the mathematical properties that confirm or disprove the validity of the equality.

1.  $(RG)H = R(GH)$
2.  $LP = PL$
3.  $E(FK) = (EF)K$
4.  $K(A + C) = KA + KC$
5.  $(QF)^\top = F^\top Q^\top$
6.  $L(ML) = L^2M$
7.  $GC + HC = (G + H)C$
8.  $R(J + T^\top) = RJ + RT^\top$
9.  $(AK)^\top = A^\top K^\top$
10.  $(Q + F^\top)E^\top = QE^\top + (EF)^\top$

*Solution.*

1.  $(RG)H$  and  $R(GH)$  are both well-defined. Furthermore, it follows from the associative law of matrix multiplication that  $(RG)H = R(GH)$ . We have

$$(RG)H = R(GH) = \begin{pmatrix} -93 & 186 & -62 \end{pmatrix}$$

2.  $LP$  and  $PL$  are both well-defined. But they are not equal

$$LP = \begin{pmatrix} 66 & 53 \\ 52 & 41 \end{pmatrix}, \quad PL = \begin{pmatrix} 22 & 20 \\ 96 & 85 \end{pmatrix}$$

3.  $E(FK)$  and  $(EF)K$  are both well-defined. Furthermore, it follows from the associative law of matrix multiplication that  $E(FK) = (EF)K$ . We have

$$E(FK) = (EF)K = \begin{pmatrix} 44 & 2 & -180 \\ 22 & 5 & -76 \\ 16 & 2 & -61 \\ 22 & 5 & -76 \end{pmatrix}$$

4.  $K(A + C)$  and  $KA + KC$  are both well-defined. Furthermore, it follows from the distributive law of matrix multiplication over addition that  $K(A + C) = KA + KC$ . We have

$$K(A + C) = KA + KC = \begin{pmatrix} 0 & 30 \\ 32 & -29 \end{pmatrix}$$

5.  $(QF)^\top$  and  $F^\top Q^\top$  are both well-defined. Furthermore, it follows from the definition of matrix transpose and matrix multiplication that  $(QF)^\top = F^\top Q^\top$ . We have

$$(QF)^\top = F^\top Q^\top = \begin{pmatrix} 75 & 125 \\ -37 & -61 \end{pmatrix}$$

6.  $L(ML)$  and  $L^2M$  are both well-defined. However, they are not equal.

$$L(ML) = \begin{pmatrix} 1826 & 1640 \\ 1434 & 1288 \end{pmatrix}, \quad L^2M = \begin{pmatrix} 2887 & 287 \\ 2283 & 227 \end{pmatrix}.$$

7.  $GC + HC$  and  $(G + H)C$  are both well-defined. Furthermore, it follows from the distributive law of matrix multiplication over addition that  $GC + HC = (G + H)C$ . We have

$$GC + HC = (G + H)C = \begin{pmatrix} 108 & -31 \\ 61 & 38 \\ 32 & -9 \end{pmatrix}$$

8.  $R(J + T^\top)$  and  $RJ + RT^\top$  are both well-defined. Furthermore, it follows from the distributive law of matrix multiplication over addition that  $R(J + T^\top) = RJ + RT^\top$ . We have

$$R(J + T^\top) = RJ + RT^\top = \begin{pmatrix} -70 \end{pmatrix}.$$

9.  $(AK)^\top$  and  $A^\top K^\top$  are both well-defined. However, the equality does not hold. We have

$$(AK)^\top = \begin{pmatrix} -4 & 12 & 2 \\ 4 & 16 & -7 \\ 31 & 5 & -33 \end{pmatrix}, \quad A^\top K^\top = \begin{pmatrix} -3 & 19 \\ 31 & -18 \end{pmatrix}.$$

10. Both sides of the equality are well-defined. First, it follows from the distributive law of matrix multiplication over addition that  $(Q + F^\top)E^\top = QE^\top + F^\top E^\top$ . Then it follows from the definition of matrix transpose and matrix multiplication that  $F^\top E^\top = (EF)^\top$ .

We have

$$(Q + F^\top)E^\top = QE^\top + (EF)^\top = \begin{pmatrix} 33 & 11 & 14 & 11 \\ 8 & 2 & 0 & 2 \end{pmatrix}$$

**Question 12.** Given matrices

$$A = \begin{pmatrix} 3 & -2 & 7 \\ 6 & 5 & 4 \\ 0 & 4 & 9 \end{pmatrix}, \quad B = \begin{pmatrix} 6 & -2 & 4 \\ 0 & 1 & 3 \\ 7 & 7 & 5 \end{pmatrix};$$

or

$$A = \begin{pmatrix} 1 & 5 & -2 & 1 \\ 0 & 1 & -2 & 3 \\ 1 & 1 & 2 & 1 \\ 0 & 0 & 2 & -4 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 1 & 5 & -1 \\ 2 & 1 & 2 & 1 \\ 4 & -2 & -1 & 0 \\ -1 & 2 & 1 & 3 \end{pmatrix}$$

Compute the specified row or column of the matrix product of the given matrices, both for the  $3 \times 3$  and  $4 \times 4$  cases:

1. The first row of the product  $AB$
2. The third row of the product  $AB$
3. The second column of the product  $AB$
4. The first column of the product  $BA$
5. The second column of the product  $BA$
6. The third row of the product  $AA$
7. The third column of the product  $AA$
8. The first column of the product  $AA$
9. The second column of the product  $BB$
10. The second row of the product  $BB$
11. The third column of the product  $BAA$

*Solution.*

1. The first row of the product  $AB$  is given by

$$(3, -2, 7)B = 3(6, -2, 4) - 2(0, 1, 3) + 7(7, 7, 5) = (67 \ 41 \ 41)$$

or

$$(1 \ 5 \ -2 \ 1)B = (1 \ 12 \ 18 \ 7)$$

2. The third row of the product  $AB$  is given by

$$(0, 4, 9)B = (63 \ 67 \ 57)$$

or

$$(1 \ 1 \ 2 \ 1)B = (9 \ 0 \ 6 \ 3)$$

3. The second column of the product  $AB$  is given by

$$A \begin{pmatrix} -2 \\ 1 \\ 7 \end{pmatrix} = \begin{pmatrix} 41 \\ 21 \\ 67 \end{pmatrix}$$

or

$$A \begin{pmatrix} 1 \\ 1 \\ -2 \\ 2 \end{pmatrix} = \begin{pmatrix} 12 \\ 11 \\ 0 \\ -12 \end{pmatrix}$$

4. The first column of the product  $BA$  is given by

$$B \begin{pmatrix} 3 \\ 6 \\ 0 \end{pmatrix} = \begin{pmatrix} 6 \\ 6 \\ 63 \end{pmatrix}$$

or

$$B \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 5 \\ 4 \\ 3 \\ 0 \end{pmatrix}$$

5. The second column of the product  $BA$  is given by

$$\begin{pmatrix} -6 \\ 17 \\ 41 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} 6 \\ 13 \\ 17 \\ -2 \end{pmatrix}$$

6. The third row of the product  $AA$  is given by

$$(24 \ 56 \ 97), \quad \text{or} \quad (3 \ 8 \ 2 \ 2)$$

7. The third column of the product  $AA$  is given by

$$\begin{pmatrix} 76 \\ 98 \\ 97 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} -14 \\ 0 \\ 2 \\ -4 \end{pmatrix}$$

8. The first column of the product  $AA$  is given by

$$\begin{pmatrix} -3 \\ 48 \\ 24 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} -1 \\ -2 \\ 3 \\ 2 \end{pmatrix}$$

9. The second column of the product  $BB$  is given by

$$\begin{pmatrix} 14 \\ 22 \\ 28 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} -11 \\ 1 \\ 4 \\ 5 \end{pmatrix}$$

10. The second row of the product  $BB$  is given by

$$(21 \ 22 \ 18), \quad \text{or} \quad (9 \ 1 \ 11 \ 2)$$

11. The third column of the product  $BAA$  is given by

$$\begin{pmatrix} 648 \\ 389 \\ 1703 \end{pmatrix}, \quad \text{or} \quad \begin{pmatrix} 14 \\ -28 \\ -58 \\ 4 \end{pmatrix}$$

### Question 13.

1. Find a non-diagonal matrix  $A$  such that  $A^2 = I_2$ .
2. Find a non-diagonal matrix  $A$  such that  $A^2 = I_3$  (modify the result from part (a)).
3. Find a non-identity matrix  $A$  such that  $A^3 = I_3$ .

*Solution.*

1. Let  $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ , then we have

$$A^2 = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} = \begin{pmatrix} a_{11}^2 + a_{12}a_{21} & a_{11}a_{12} + a_{12}a_{22} \\ a_{21}a_{11} + a_{22}a_{21} & a_{21}a_{12} + a_{22}^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Consequently,

$$a_{11}^2 + a_{12}a_{21} = 1, \quad a_{11}a_{12} + a_{12}a_{22} = 0, \quad a_{21}a_{11} + a_{22}a_{21} = 0, \quad a_{21}a_{12} + a_{22}^2 = 1.$$

Since  $A$  is not diagonal,  $a_{12} \neq 0$  or  $a_{21} \neq 0$ . Then

$$a_{11} = -a_{22}, \quad a_{11}^2 = 1 - a_{12}a_{21}.$$

If we let

$$a_{12} = a_{21} = 1, \quad a_{11} = a_{22} = 0,$$

the matrix

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

is one matrix that satisfies  $A^2 = I_2$ . Similarly,

$$a_{12} = a_{21} = \frac{1}{2}, \quad a_{11} = \frac{\sqrt{3}}{2}, \quad a_{22} = -\frac{\sqrt{3}}{2}$$

gives

$$A = \frac{1}{2} \begin{pmatrix} \sqrt{3} & 1 \\ 1 & -\sqrt{3} \end{pmatrix}$$

2.

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

3.

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$