

Logic (PHIL 2080, COMP 2620, COMP 6262)
Chapter: Sequents, Semantics, and Propositional Natural
Deduction — Conjunction, Implication, Theorems

Yoshihiro Maruyama



Australian
National
University

Introduction

Recap: Sequents

- We want to know when one logical formula follows logically from another.
- Suppose we know that “ p is true’ (technically: it is *interpreted* as true), and we know that $p \rightarrow q$ holds as well. Then we can logically conclude that q also holds!
- We can express this with *sequents*: $p, p \rightarrow q \models q$
- These conclusions can be arbitrarily complicated, however! I.e., it might not be obvious that the conclusion follows from the premises.
- Two ways to show validity of sequents: semantics (validity by meaning) and syntactic proof system (validity by following deductive rules; natural deduction introduced soon).

Sequents

Introduction

In general, a sequent is of the following form with X a set of formulae and A a single formula:

$$X \models A$$

- Read it: A follows from X ; or X entails A .
- For example, “ q follows from p and $p \rightarrow q$ ”
- We write down:
but that just abbreviates:

$$\begin{array}{ccc} p, p \rightarrow q & \models & q \\ \underbrace{\{p, p \rightarrow q\}}_X & \models & \underbrace{q}_A \end{array}$$

- Also $X, Y \models A$ abbreviates $X \cup Y \models A$,

Semantically Valid Sequents

Definition:

$X \models A$ means the sequent is *valid*. This is the case if and only if:

- *A is true for every interpretation for which all the formulae in X are true. Or, equivalently:*
- *There is no interpretation that makes X true, but not A.*

How to check/test/prove $X \models A$? Create the truth tables!

- Create a table t_X for all formulae in X (all need to be true)
- Create another table t_A for A and check validity criterion.

Checking Validity, Example 2

$$\text{Show } \overbrace{(p \vee q) \rightarrow r, p}^X \models \overbrace{(p \rightarrow r) \wedge (q \rightarrow r)}^A$$

Table t_X for premises:

p	q	r	$p \vee q$	$(p \vee q) \rightarrow r$	X
0	0	0	0	1	0
0	0	1	0	1	0
0	1	0	1	0	0
0	1	1	1	1	0
1	0	0	1	0	0
1	0	1	1	1	1
1	1	0	1	0	0
1	1	1	1	1	1

Table t_A for conclusion:

p	q	r	$p \rightarrow r$	$q \rightarrow r$	A
0	0	0	1	1	1
0	0	1	1	1	1
0	1	0	1	0	0
0	1	1	1	1	1
1	0	0	0	1	0
1	0	1	1	1	1
1	1	0	0	0	0
1	1	1	1	1	1

Recall the definition: The sequent is valid if all interpretations that make X true also make A true!

Checking Validity, Example 2

$$\text{Show } \overbrace{(p \vee q) \rightarrow r, p}^X \models \overbrace{(p \rightarrow r) \wedge (q \rightarrow r)}^A$$

Table t_X for premises:

p	q	r	$p \vee q$	$(p \vee q) \rightarrow r$	X
0	0	0	0	1	0
0	0	1	0	1	0
0	1	0	1	0	0
0	1	1	1	1	0
1	0	0	1	0	0
1	0	1	1	1	1
1	1	0	1	0	0
1	1	1	1	1	1

Table t_A for conclusion:

p	q	r	$p \rightarrow r$	$q \rightarrow r$	A
0	0	0	1	1	1
0	0	1	1	1	1
0	1	0	1	0	0
0	1	1	1	1	1
1	0	0	0	1	0
1	0	1	1	1	1
1	1	0	0	0	0
1	1	1	1	1	1

Only two interpretations exist that make all $x \in X$ true:

$$1 \quad I_1(p) = I_1(r) = 1, I_1(q) = 0$$

$$2 \quad I_2(p) = I_2(q) = I_2(r) = 1$$

Both of them make A true! Thus, $X \models A$.

Natural Deduction

Natural Deduction and Derivations

- Natural deduction is pure syntax manipulation and acts as *proof system* with a formal notion of proof as a mathematical entity (cf. informal proof in ordinary math).
- Natural Deduction exploits *derivations* (or formal proofs).
- A derivation is a finite sequence of formulae, which are derived from each other based on elementary formula manipulations (“1-step inference rules”)
- Syntax (proof system) vs. semantics is arguably the most important distinction in formal logic.

Syntax of Sequents

- From now on, we write $X \vdash A$ rather than $X \models A$.
- $X \vdash A$ means A syntactically follows from X , i.e., you can formally prove the conclusion A using X as assumptions (within the system of natural deduction).
- $X \models A$ means A semantically follows from X .

Conjunction

The 1-Step Rules: And-Elimination

What are the 1-step rules for dealing with conjunction?

Elimination rule:

$$\frac{A \wedge B}{A} \wedge E$$

$$\frac{A \wedge B}{B} \wedge E$$

Which reads: If we derived $A \wedge B$, we can derive both A and B .

The 1-Step Rules: And-Introduction

What are the 1-step rules for dealing with conjunction?

Introduction rule:

$$\frac{A \quad B}{A \wedge B} \wedge I$$

Which reads: If we derived A and we derived B , we can derive $A \wedge B$.

Proof Syntax / Notation: Overview

- How to write down proofs?
- There are many different notations that describe the same thing
- We introduce two:
 - Tree-like representation of the applied rules
(just since it's another standard)
 - list-like representation (only use that one!)

Proof Syntax / Notation: Tree- and List-like Representations

- Assume we want to prove $p \wedge q \vdash q \wedge p$
- In the tree-like format:

$$\frac{\frac{p \wedge q}{q} \wedge E \quad \frac{p \wedge q}{p} \wedge E}{q \wedge p} \wedge I$$

- Leaves are assumptions, root is conclusion
- Advantages: Makes the proof structure obvious
- In exercises, etc: *Do not use it, unless we ask you to!*

Proof Syntax / Notation: Tree- and List-like Representations

- Assume we want to prove $p \wedge q \vdash q \wedge p$
- In the list format:

α_1	(1)	$p \wedge q$	A	\equiv	$p \wedge q \vdash p \wedge q$
α_1	(2)	q	1 $\wedge E$	\equiv	$p \wedge q \vdash q$
α_1	(3)	p	1 $\wedge E$	\equiv	$p \wedge q \vdash p$
α_1	(4)	$q \wedge p$	2,3 $\wedge I$	\equiv	$p \wedge q \vdash q \wedge p$

column 1: assumption number *column 2:* line number

column 3: derivation *column 4:* how it was derived

- Note:** Each line represents a sequent! (Sequence of sequents.)

The 1-Step Rules (Based on Sequents): Derivation Rules

- Derivation Rules as considered so far:

$$\frac{A \wedge B}{A} \wedge E$$

$$\frac{A \wedge B}{B} \wedge E$$

$$\frac{A \quad B}{A \wedge B} \wedge I$$

- Re-written in terms of sequents:

$$\frac{X \vdash A \wedge B}{X \vdash A} \wedge E$$

$$\frac{X \vdash A \wedge B}{X \vdash B} \wedge E$$

$$\frac{X \vdash A \quad Y \vdash B}{X, Y \vdash A \wedge B} \wedge I$$

→ I.e., now we see how premises accumulate!

The 1-Step Rules (Based on Sequents): Accumulation of Assumptions, Example

$$p, q \vdash p \wedge q$$

$$\begin{array}{llllll}
 \alpha_1 & (1) & p & A & \equiv & p \vdash p \text{ (by assumption)} \\
 \alpha_2 & (2) & q & A & \equiv & q \vdash q \text{ (by assumption)} \\
 \alpha_1, \alpha_2 & (3) & p \wedge q & 1, 2 \wedge I & \equiv & p, q \vdash p \wedge q \quad (\wedge I)
 \end{array}$$

$$\boxed{
 \frac{X \vdash A \quad Y \vdash B}{X, Y \vdash A \wedge B} \wedge I
 }$$

Implication

The 1-Step Rules: Implication-Elimination and -Introduction

● Elimination rule:

$$\frac{A \rightarrow B \quad A}{B} \rightarrow E$$

● Introduction rule:

if we can derive
 B using A :

then we can derive
 $A \rightarrow B$ and discharge A :

$$\frac{\begin{array}{c} [A] \\ \vdots \\ B \end{array}}{A \rightarrow B} \rightarrow I \quad \equiv \quad \begin{array}{c} A \\ \vdots \\ B \end{array} \quad + \quad \frac{\begin{array}{c} [A] \\ \vdots \\ B \end{array}}{A \rightarrow B} \rightarrow I$$

The 1-Step Rules: Implication-Elimination and -Introduction (Based on Sequents)

- Derivation Rules as considered so far:

$$\frac{A \rightarrow B \quad A}{B} \rightarrow E$$

$$\frac{\begin{array}{c} [A] \\ \vdots \\ B \end{array}}{A \rightarrow B} \rightarrow I$$

- Re-written in terms of sequents:

$$\frac{X \vdash A \rightarrow B \quad Y \vdash A}{X, Y \vdash B} \rightarrow E$$

$$\frac{X, A \vdash B}{X \vdash A \rightarrow B} \rightarrow I$$

Has side effect of
removing the assumption A

- We say that A gets *discharged*, and annotate that in the proof.

The 1-Step Rules: Deduction Equivalence

$$\underbrace{X \vdash A \rightarrow B \quad \text{iff} \quad X, A \vdash B}$$

deduction equivalence
(or deduction theorem)

Why does this hold?

- If $X, A \vdash B$, then $X \vdash A \rightarrow B$:

$$\frac{X, A \vdash B}{X \vdash A \rightarrow B} \rightarrow I$$

- If $X \vdash A \rightarrow B$, then $X, A \vdash B$:

$$\frac{X \vdash A \rightarrow B \quad A \vdash A}{X, A \vdash B} \rightarrow E$$

(That's the $\rightarrow E$ rule with Y substituted by A)

The 1-Step Rules: Implication-Introduction and -Elimination, Example 1

$$p \rightarrow q \vdash (p \wedge r) \rightarrow q$$

α_1	(1)	$p \rightarrow q$	A
α_2	(2)	$p \wedge r$	A
α_2	(3)	p	2 $\wedge E$
α_1, α_2	(4)	q	1,3 $\rightarrow E$
α_1	(5)	$(p \wedge r) \rightarrow q$	4[α_2] $\rightarrow I$

$$\frac{X, A \vdash B}{X \vdash A \rightarrow B} \rightarrow I$$

$$\frac{X \vdash A \rightarrow B \quad Y \vdash A}{X, Y \vdash B} \rightarrow E$$

α_1, α_2	(n-1)	q	
α_1	(n)	$(p \wedge r) \rightarrow q$	(n-1)[α_2] $\rightarrow I$

- Assumption α_2 is a new one, which was not given in the original sequent, so we need to eliminate it later on.
- In the last step, we discharge assumption α_2 .

The 1-Step Rules: Implication-Introduction and -Elimination, Example 1 (cont'd)

The proof of $p \rightarrow q \vdash (p \wedge r) \rightarrow q$ in a tree-like structure:

$$\begin{array}{c}
 \frac{p \rightarrow q \quad \frac{[p \wedge r]^{(1)}}{p} \wedge E}{q} \rightarrow E \\
 \frac{q}{(p \wedge r) \rightarrow q} \rightarrow I(1)
 \end{array}$$

Here, we denote discharged assumptions by $[\dots]^{(n)}$, where we number each assumption so that they can be distinguished from each other, i.e., so that we know which rule discharged which assumption(s).

The 1-Step Rules: Implication-Introduction and -Elimination, Example 2

$$(p \wedge q) \rightarrow r \vdash p \rightarrow (q \rightarrow r)$$

α_1	(1)	$(p \wedge q) \rightarrow r$	A
α_2	(2)	p	A
α_3	(3)	q	A
α_2, α_3	(4)	$p \wedge q$	2,3 $\wedge I$
$\alpha_1, \alpha_2, \alpha_3$	(5)	r	1,4 $\rightarrow E$
α_1, α_2	(6)	$q \rightarrow r$	5[α_3] $\rightarrow I$
α_1	(7)	$p \rightarrow (q \rightarrow r)$	6[α_2] $\rightarrow I$

$$\frac{X, A \vdash B}{X \vdash A \rightarrow B} \rightarrow I$$

$$\frac{X \vdash A \rightarrow B \quad Y \vdash A}{X, Y \vdash B} \rightarrow E$$

$\alpha_1, \alpha_2, \alpha_3$	(n-2)	r	
α_1, α_2	(n-1)	$q \rightarrow r$	(n-2)[α_3] $\rightarrow I$
α_1	(n)	$p \rightarrow (q \rightarrow r)$	(n-1)[α_2] $\rightarrow I$

Vacuous Discharge: Discharging Non-existent Assumptions

- We can “discharge” assumptions that are not there; this happens if the conclusion does not depend on its assumption.

$$p \vdash q \rightarrow p$$

$$\begin{array}{llll}
 \alpha_1 & (1) & p & A \\
 \hline
 \alpha_2 & (2) & q & A \\
 \alpha_1 & (2) & q \rightarrow p & 1[] \rightarrow I
 \end{array}$$

$$\boxed{
 \frac{X, A \vdash B}{X \vdash A \rightarrow B} \rightarrow I
 }$$

$$\begin{array}{llll}
 \alpha_1, \cancel{\alpha_2} & (n-1) & p & \\
 \alpha_1 & (n) & q \rightarrow p & (n-1)[\cancel{\alpha_2}] \rightarrow I
 \end{array}$$

- We call such a discharge a *vacuous discharge*.
- I.e., whenever we “would remove” some assumption α from a set of assumptions X , but $\alpha \notin X$, we write $i[] \rightarrow I$ instead of $i[\alpha] \rightarrow I$

Excursion: \vdash vs. \rightarrow : An Often Asked Question in Previous Courses

- \vdash and \rightarrow seem to be of a very related nature:
E.g., compare $A, B \vdash C$ with $A \wedge B \rightarrow C$
- So what's the difference?
- \rightarrow is a logical connective, whereas \vdash is not.
- \vdash is a relation between formulae and cannot be used within a formula.
- They are linked by the deduction theorem: $X, A \vdash B$ if and only if $X \vdash A \rightarrow B$. In particular: $A \vdash B$ if and only if $\vdash A \rightarrow B$

Theorems

Unconditionally True Formulas

- Sequents that do not depend on anything are called *theorems*!
- Thus, A is a theorem if “ $\vdash A$ ”, e.g., $\vdash p \rightarrow (q \rightarrow p)$.
- Another (slightly more complex) example:

α_1	(1)	p	A
α_2	(2)	q	A
α_1, α_2	(3)	$p \wedge q$	1,2 $\wedge I$
α_1	(4)	$q \rightarrow (p \wedge q)$	3[α_2] $\rightarrow I$
	(5)	$p \rightarrow (q \rightarrow (p \wedge q))$	4[α_1] $\rightarrow I$

- Thus, we get $\vdash p \rightarrow (q \rightarrow (p \wedge q))$, so its formula is a theorem.
- Note that A in $\vdash A$ is a tautology!

Summary

Content of this Lecture

- Sequents and their semantics
 - What does $X \models A$ mean?
 - The most important basics of Natural Deduction!
 - How can proofs be written?
 - What does $X \vdash A$ mean?
 - Every logical connective comes with two 1-step rules:
Introduction and Elimination
 - What's a theorem?
- The Logic Notes sections:
- 3. *More about propositional logic: Truth Tables*
 - 2. *Propositional natural deduction: Conjunction*
 - 2. *Propositional natural deduction: Implication*
 - 2. *Propositional natural deduction: Counting assumptions*
(except *Contraction*, which you should look up!)