

Zircon oxybarometer - hygrometer

LIANG Xueyin
Precambrian-CUGB

China University of Geosciences, Beijing
School of Earth Sciences and Resources
Geology '26

July 30 th, 2025

Outline

1 Zircon oxybarometer

2 Zircon hygrometer

1. Zircon oxybarometer


Literature

JOURNAL OF
PETROLOGY

Journal of Petrology, 2020, Vol. 61, No. 3: egaa034
doi: 10.1093/petrology/egaa034
Advance Access Publication Date: 20 February 2020
Original Article

OXFORD

New Magmatic Oxybarometer Using Trace Elements in Zircon

Robert R. Loucks ^{1,2*}, **Marco L. Fiorentini**^{1,2} and
Gonzalo J. Henríquez^{1,2}

¹Centre for Exploration Targeting, School of Earth Sciences, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia; ²ARC Centre of Excellence for Core to Crust Fluid Systems, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

*Corresponding author. Telephone: +61 8 6488 6794. Fax: +61 8 6488 1178.
E-mail: robert.loucks@uwa.edu.au

Received December 31, 2018; Accepted February 13, 2020

Figure 1: Loucks et al., 2020

Click [here](#) to download the literature.

数据准备

- Zircon Trace elements
- 适用范围：
 - Fe–Ti oxide phenocrysts and hornblende phenocrysts quenched in eruptive I- and A-type dacites and rhyolites
 - slowly cooled plutonic rocks
 - detrital and xenocrystic zircons

Magma oxygen fugacity (Loucks et al., 2020)

Loucks et al., 2020

$$\Delta FMQ = \log fO_{2(\text{sample})} - \log fO_{2(\text{FMQ})} = 3.99 \times \log \frac{Ce}{\sqrt{U_i \times Ti}} + 2.28 \quad (1)$$

- U_i : age-corrected initial U content; $U_i = U \times e^{\lambda t}$
- 太古宙样品必须校正
- **independent estimates**

Magma oxygen fugacity (Smythe and Brennan, 2016)




Figure 2: Smythe and Brennan, 2016

- Principal: zircon-melt equilibrium

Smythe and Brennan, 2016

$$\ln \left[\frac{x_{\text{Ce}^{4+}}^{\text{melt}}}{x_{\text{Ce}^{3+}}^{\text{melt}}} \right] = \frac{1}{4} \ln f_{\text{O}_2} + \frac{13,136}{T} - 2.064 \frac{\text{NBO}}{T} - 8.878 \times x_{\text{H}_2\text{O}} - 8.955 \quad (2)$$

Crystallization Temperature (Ferry and Watson, 2007)

Ferry and Watson, 2007

$$\log (\text{Ti in Zircon}) + \log \alpha_{\text{SiO}_2} - \log \alpha_{\text{TiO}_2} = 5.711 - \frac{4800}{T} \quad (3)$$

- α_{SiO_2} :
 - ① 中酸性熔体 (石英饱和) $\alpha_{\text{SiO}_2} = 1$
 - ② 基性岩 $\alpha_{\text{SiO}_2} < 1$, 高级变质岩需考虑 P-T 影响, 需用热力学模拟计算
- $\alpha_{\text{TiO}_2} = 0.6$
 - ① 金红石饱和 \Rightarrow 出现金红石, $\alpha_{\text{TiO}_2} = 1$
 - ② 实验岩石学表明硅酸盐熔体中 $\alpha_{\text{TiO}_2} = 0.6 - 0.9$ (Watson et al., 2006; Hayden and Watson, 2007)

2. Zircon hygrometer

Article

Earth's early continental crust formed from wet and oxidizing arc magmas

<https://doi.org/10.1038/s41586-023-06552-0>

Received: 6 December 2022

Accepted: 17 August 2023

Published online: 27 September 2023

 Check for updates

Rong-Feng Ge^{1✉}, Simon A. Wilde², Wen-Bin Zhu¹ & Xiao-Lei Wang¹

Formation of continental crust has shaped the surface and interior of our planet and generated the land and mineral resources on which we rely. However, how the early continental crust of Earth formed is still debated^{1–7}. Modern continental crust is largely formed from wet and oxidizing arc magmas at subduction zones, in which

Figure 3: Ge et al., 2023

Click [here](#) to download the literature.

数据准备

- Zircon trace element data
- Whole-rock/melt inclusion data (parental magma composition)
- Filtered criteria ([Laurent et al., 2022](#)):
 - ① LREE-I ([Bell et al., 2016](#)): $\frac{Dy}{Sm} + \frac{Dy}{Nd} \geq 30$ decrease with increasing degree of hydrothermal alteration or contamination
 - ② $La \leq 0.5$ ppm (largest extent of modification during post-magmatic alteration)
 - ③ $Ca \leq 150$ ppm and $Fe \leq 150$ ppm (below the limit of detection)
 - ④ discordance of U-Pb ages within 5% and common Pb < 1%
- Calculate the median and 2SEM of the composition (Zircon and whole rock/melt)

Retrieve equilibrium magmatic H_2O content

Smythe and Brenan, 2016

$$\ln \left[\frac{x_{\text{Ce}^{4+}}^{\text{melt}}}{x_{\text{Ce}^{3+}}^{\text{melt}}} \right] = \frac{1}{4} \ln f_{\text{O}_2} + \frac{13,136}{T} - 2.064 \frac{\text{NBO}}{T} - 8.878 \times x_{\text{H}_2\text{O}} - 8.955 \quad (4)$$

Loucks et al., 2020

$$\Delta FMQ = 3.99 \times \log \frac{\text{Ce}}{\sqrt{U_i \times Ti}} + 2.28 \quad (5)$$

Ferry and Watson, 2007

$$\log (\text{Ti in Zircon}) + \log \alpha_{\text{SiO}_2} - \log \alpha_{\text{TiO}_2} = 5.711 - \frac{4800}{T} \quad (6)$$

Macro Excel

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O				
1	Zircon oxybarometer - hygrometer.xlsx is a spreadsheet attached to Ge et al (2023, Nature), which is modified from Smythe and Brenan (2016) and																		
2	Areas in yellow require input (median zircon and equilibrium melt/whole-rock compositions, as well as the $f_{\text{H}_2\text{O}}$ values given by the zircon Ce-U-Th																		
3	Unit	Zircon composition																	
4		Ti	Y	Zr	La	Ce	Pr	Nd	Sm	Eu	Gd	Tb	Dy	Ho	Er				
5	lg2E unit,	2.61	1901	473708	0.07	37.2	0.09	1.80	5.62	0.13	42.0	16.7	202	79.6	349				
6	2SEM	1.31	309	3376	0.06	5.61	0.02	0.27	0.91	0.02	4.02	1.44	17.4	7.13	32.0				
7	Calculation results																		
8	NBO/T		Ce ⁴⁺ /Ce ³⁺ Melt		T (K)		Δ FMQ		logfO ₂		H ₂ O (mol. %)			H ₂ O (wt. %)					
9	value	2SEM	value	2SEM	value	2SEM	value	2SEM	value	2SEM	value	2SEM	2SEM (to	value	2SEM				
10	0.03	0.02	0.00187	0.00015	946	46	1.22	0.28	-16.64	0.28	0.17	0.02	0.08	5.81	0.62				
11	Parent Melt Composition				0.05 (s)														
12	SiO ₂		TiO ₂		Al ₂ O ₃		MgO		MnO		FeO		Fe ₂ O ₃						
13	wt. %	2SEM	wt. %	2SEM	wt. %	2SEM	wt. %	2SEM	wt. %	2SEM	wt. %	2SEM	wt. %	2SEM					
14	77.8	0.70	0.07	0.01	12.5	0.40	0.02	0.01	0.00	0.00	0.62	0.03							
15																			
16	Element	Trace Element Concentration				a_{SiO_2}		a_{TiO_2}	Zr/Hf		11.2								
17		Melt		Zircon		1.000		0.600	Th/U		2.6								
18		ppm (s)		ppm (s)		(s)		(s)											
19		Y		1901		0.05		0.1											
20	La		15.9		0.30		0.07		0.06										

Figure 4: Macro Excel

Metamorphic disturbance

- Preferential U loss (high Th/U ratios of several TTG suites)
⇒ Overestimate magmatic H_2O content
- Assuming $Th/U = 5$ (the median composition of the global sodic TTGs)

Reference I

- Bell, E. A., Boehnke, P., & Harrison, T. M. (2016). Recovering the primary geochemistry of jack hills zircons through quantitative estimates of chemical alteration. *Geochimica et Cosmochimica Acta*, 191, 187–202.
- Ferry, J., & Watson, E. (2007). New thermodynamic models and revised calibrations for the ti-in-zircon and zr-in-rutile thermometers. *Contributions to Mineralogy and Petrology*, 154(4), 429–437.
- Ge, R.-F., Wilde, S. A., Zhu, W.-B., & Wang, X.-L. (2023). Earth' s early continental crust formed from wet and oxidizing arc magmas. *Nature*, 623(7986), 334–339.
- Hayden, L. A., & Watson, E. B. (2007). Rutile saturation in hydrous siliceous melts and its bearing on ti-thermometry of quartz and zircon. *Earth and Planetary Science Letters*, 258(3-4), 561–568.

Reference II

- Laurent, O., Moyen, J.-F., Wotzlaw, J.-F., Björnsen, J., & Bachmann, O. (2022). Early earth zircons formed in residual granitic melts produced by tonalite differentiation. *Geology*, 50(4), 437–441.
- Loucks, R. R., Fiorentini, M. L., & Henríquez, G. J. (2020). New magmatic oxybarometer using trace elements in zircon. *Journal of Petrology*, 61(3), egaa034.
- Smythe, D. J., & Brenan, J. M. (2016). Magmatic oxygen fugacity estimated using zircon-melt partitioning of cerium. *Earth and Planetary Science Letters*, 453, 260–266.
- Watson, E., Wark, D., & Thomas, J. (2006). Crystallization thermometers for zircon and rutile. *Contributions to Mineralogy and Petrology*, 151(4), 413–433.

Thank You!