

Exponential smoothing and non-negative data

Muhammad Akram
Rob J Hyndman
J Keith Ord

Business & Economic Forecasting Unit



MONASH University

Outline

- 1 **Exponential smoothing models**
- 2 **Problems with some of the models**
- 3 **A new model for positive data**
- 4 **Conclusions**

Outline

- 1 **Exponential smoothing models**
- 2 Problems with some of the models
- 3 A new model for positive data
- 4 Conclusions

Problem

- Most forecasting methods in business are based on exponential smoothing.

Problem

- Most forecasting methods in business are based on exponential smoothing.
- Most time series in business are inherently non-negative.


Problem

- Most forecasting methods in business are based on exponential smoothing.
- Most time series in business are inherently non-negative.
- But exponential smoothing models for non-negative data don't always work!

Problem

- Most forecasting methods in business are based on exponential smoothing.
- Most time series in business are inherently non-negative.
- But exponential smoothing models for non-negative data don't always work!


1 They can produce negative forecasts



Problem

- Most forecasting methods in business are based on exponential smoothing.
- Most time series in business are inherently non-negative.
- But exponential smoothing models for non-negative data don't always work!


- 1 They can produce negative forecasts
- 2 They can produce infinite forecast variance



Problem

- Most forecasting methods in business are based on exponential smoothing.
- Most time series in business are inherently non-negative.
- But exponential smoothing models for non-negative data don't always work!

- 1 They can produce negative forecasts
- 2 They can produce infinite forecast variance
- 3 They can converge almost surely to zero.



Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A_d	(Additive damped)	A_d,N	A_d,A	A_d,M
M	(Multiplicative)	M,N	M,A	M,M
M_d	(Multiplicative damped)	M_d,N	M_d,A	M_d,M

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(*Error,Trend,Seasonal*)

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(*Error,Trend,Seasonal*)
Exponential Smoothing

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(Error,Trend,Seasonal)
Exponential Smoothing

ETS(A,N,N): Simple exponential smoothing with additive errors

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(*Error, Trend, Seasonal*)
Exponential Smoothing

ETS(A,A,N): Holt's linear method with additive errors

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(*Error, Trend, Seasonal*)
Exponential Smoothing

ETS(A,A,A): Additive Holt-Winters' method with additive errors

Taxonomy of models

		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A _d	(Additive damped)	A _d ,N	A _d ,A	A _d ,M
M	(Multiplicative)	M,N	M,A	M,M
M _d	(Multiplicative damped)	M _d ,N	M _d ,A	M _d ,M

General notation **ETS**(Error,Trend,Seasonal)
Exponential Smoothing

ETS(M,A,M): Multiplicative Holt-Winters' method
with multiplicative errors

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A_d	(Additive damped)	A_d, N	A_d, A	A_d, M
M	(Multiplicative)	M,N	M,A	M,M
M_d	(Multiplicative damped)	M_d, N	M_d, A	M_d, M

General notation **ETS**(Error,Trend,Seasonal)
Exponential Smoothing

ETS(A, A_d ,N): Damped trend method with additive errors

Taxonomy of models

Trend Component		Seasonal Component		
		N (None)	A (Additive)	M (Multiplicative)
N	(None)	N,N	N,A	N,M
A	(Additive)	A,N	A,A	A,M
A_d	(Additive damped)	A_d,N	A_d,A	A_d,M
M	(Multiplicative)	M,N	M,A	M,M
M_d	(Multiplicative damped)	M_d,N	M_d,A	M_d,M

General notation **ETS**(*Error, Trend, Seasonal*)
Exponential Smoothing

There are 30 separate models in the ETS framework

Innovations state space model

**No trend or seasonality
and multiplicative errors**

Example: ETS(M,N,N)

$$y_t = \ell_{t-1}(1 + \varepsilon_t)$$

$$\ell_t = \alpha y_t + (1 - \alpha)\ell_{t-1}$$

$$0 \leq \alpha \leq 1$$

ε_t is white noise with mean zero.

Innovations state space model

**No trend or seasonality
and multiplicative errors**

Example: ETS(M,N,N)

$$y_t = l_{t-1}(1 + \varepsilon_t)$$

$$l_t = l_{t-1}(1 + \alpha\varepsilon_t)$$

$$0 \leq \alpha \leq 1$$

ε_t is white noise with mean zero.

Innovations state space model

**No trend or seasonality
and multiplicative errors**

Example: ETS(M,N,N)

$$y_t = l_{t-1}(1 + \varepsilon_t)$$

$$l_t = l_{t-1}(1 + \alpha\varepsilon_t)$$

$$0 \leq \alpha \leq 1$$

ε_t is white noise with mean zero.

**All exponential smoothing models can be
written using analogous state space
equations.**

New book!

Springer Series in Statistics

Rob J. Hyndman · Anne B. Koehler
J. Keith Ord · Ralph D. Snyder

Forecasting with Exponential Smoothing

The State Space Approach

 Springer

- State space modeling framework
- Prediction intervals
- Model selection
- Maximum likelihood estimation
- All the important research results in one place with consistent notation
- Many new results
- 375 pages but only US\$39.95.

New book!

Springer Series in Statistics

Rob J. Hyndman · Anne B. Koehler
J. Keith Ord · Ralph D. Snyder

Forecasting with Exponential Smoothing

The State Space Approach

 Springer

- State space modeling framework
- Prediction intervals
- Model selection
- Maximum likelihood estimation
- All the important research results in one place with consistent notation
- Many new results
- 375 pages but only US\$39.95.


www.exponentialsMOOTHING.net

Outline

- 1 Exponential smoothing models
- 2 Problems with some of the models**
- 3 A new model for positive data
- 4 Conclusions

Negative forecasts


Forecasts from ETS(A,A,N)



- Could solve by taking logs or some other Box-Cox transformation. However, this limits models to be additive in the transformed space.

Negative forecasts

Forecasts from ETS(A,A,N)



- Could solve by taking logs or some other Box-Cox transformation. However, this limits models to be additive in the transformed space.
- Could solve by only using multiplicative models. But these can have other problems.

Infinite forecast variances


ETS(A,M,N) model

$$y_t = \ell_{t-1} b_{t-1} + \varepsilon_t$$

$$\ell_t = \ell_{t-1} b_{t-1} + \alpha \varepsilon_t$$

$$b_t = b_{t-1} + \beta \varepsilon_t / \ell_{t-1}.$$

- $\ell_0 = 0.1$
- $b_0 = 1$
- $\alpha = 0.1$
- $\beta = 0.05$
- $\sigma = 1$



Infinite forecast variances


ETS(A,M,N) model

$$y_t = \ell_{t-1} b_{t-1} + \varepsilon_t$$

$$\ell_t = \ell_{t-1} b_{t-1} + \alpha \varepsilon_t$$

$$b_t = b_{t-1} + \beta \varepsilon_t / \ell_{t-1}.$$

- $\ell_0 = 0.1$
- $b_0 = 1$
- $\alpha = 0.1$
- $\beta = 0.05$
- $\sigma = 1$



Infinite forecast variances

Suppose ε_t has positive density at 0

For ETS models (A,M,N), (A,M,A), (A,M_d,N), (A,M_d,A), (A,M,M), (A,M_d,M), (M,M,A) and (M,M_d,A):

- $V(y_{n+h} \mid \mathbf{x}_n) = \infty$ for $h \geq 3$;
- $E(y_{n+h} \mid \mathbf{x}_n)$ is undefined for $h \geq 3$.

Infinite forecast variances

Suppose ε_t has positive density at 0

For ETS models (A,M,N), (A,M,A), (A,M_d,N), (A,M_d,A), (A,M,M), (A,M_d,M), (M,M,A) and (M,M_d,A):

- $V(y_{n+h} \mid \mathbf{x}_n) = \infty$ for $h \geq 3$;
- $E(y_{n+h} \mid \mathbf{x}_n)$ is undefined for $h \geq 3$.

For ETS models (A,N,M), (A,A,M) and (A,A_d,M):

- $V(y_{n+h} \mid \mathbf{x}_n) = \infty$ for $h \geq m + 2$;
- $E(y_{n+h} \mid \mathbf{x}_n)$ is undefined for $h \geq m + 2$.

Infinite forecast variances

Suppose ε_t has positive density at 0

For ETS models (A,M,N), (A,M,A), (A,M_d,N), (A,M_d,A), (A,M,M), (A,M_d,M), (M,M,A) and (M,M_d,A):

- $V(y_{n+h} \mid \mathbf{x}_n) = \infty$ for $h \geq 3$;
- $E(y_{n+h} \mid \mathbf{x}_n)$ is undefined for $h \geq 3$.

For ETS models (A,N,M), (A,A,M) and (A,A_d,M):

- $V(y_{n+h} \mid \mathbf{x}_n) = \infty$ for $h \geq m + 2$;
- $E(y_{n+h} \mid \mathbf{x}_n)$ is undefined for $h \geq m + 2$.

➡ **These problems occur regardless of the sample space of $\{y_t\}$.**




Convergence to zero

ETS(M,N,N) model

$$y_t = l_{t-1}(1 + \varepsilon_t)$$

$$l_t = l_{t-1}(1 + \alpha\varepsilon_t)$$

- $l_0 = 10$
- $\alpha = 0.3$
- $\sigma = 0.3$ with truncated Gaussian errors



Convergence to zero

ETS(M,N,N) model

$$y_t = \ell_{t-1}(1 + \varepsilon_t)$$
$$\ell_t = \ell_{t-1}(1 + \alpha\varepsilon_t),$$

• $0 < \alpha \leq 1$

Convergence to zero

ETS(M,N,N) model

$$y_t = l_{t-1}(1 + \varepsilon_t)$$
$$l_t = l_{t-1}(1 + \alpha\varepsilon_t),$$

- $0 < \alpha \leq 1$
- $\delta_t = 1 + \varepsilon_t$ has mean 1 and variance σ^2 .

Convergence to zero

ETS(M,N,N) model

$$y_t = \ell_{t-1}(1 + \varepsilon_t)$$
$$\ell_t = \ell_{t-1}(1 + \alpha\varepsilon_t),$$

- $0 < \alpha \leq 1$
- $\delta_t = 1 + \varepsilon_t$ has mean 1 and variance σ^2 .
- δ_t are iid with positive distribution such as truncated normal, lognormal, gamma, etc.

Convergence to zero

ETS(M,N,N) model

$$y_t = \ell_{t-1}(1 + \varepsilon_t)$$
$$\ell_t = \ell_{t-1}(1 + \alpha\varepsilon_t),$$

- $0 < \alpha \leq 1$
- $\delta_t = 1 + \varepsilon_t$ has mean 1 and variance σ^2 .
- δ_t are iid with positive distribution such as truncated normal, lognormal, gamma, etc.

Convergence to zero

ETS(M,N,N) model

$$y_t = \ell_{t-1}(1 + \varepsilon_t)$$
$$\ell_t = \ell_{t-1}(1 + \alpha\varepsilon_t),$$

- $0 < \alpha \leq 1$
- $\delta_t = 1 + \varepsilon_t$ has mean 1 and variance σ^2 .
- δ_t are iid with positive distribution such as truncated normal, lognormal, gamma, etc.

$$\ell_t = \ell_0(1 + \alpha\varepsilon_1)(1 + \alpha\varepsilon_2) \cdots (1 + \alpha\varepsilon_t) = \ell_0 U_t,$$

where $U_t = U_{t-1}(1 + \alpha\varepsilon_t)$ and $U_0 = 1$. Thus U_t is a non-negative product martingale.

Convergence to zero

- Kakutani's Theorem says that ℓ_t will converge to 0 almost surely if ε_t has mean zero and is non-degenerate.

Convergence to zero

- Kakutani's Theorem says that ℓ_t will converge to 0 almost surely if ε_t has mean zero and is non-degenerate.
- Consequently, all sample paths for y_t converge to 0 almost surely.

Convergence to zero

- Kakutani's Theorem says that ℓ_t will converge to 0 almost surely if ε_t has mean zero and is non-degenerate.
- Consequently, all sample paths for y_t converge to 0 almost surely.
- Similar results follow for all purely multiplicative models: (M,N,N) , (M,N,M) , (M,M,N) , (M,M,M) , (M,M_d,N) and (M,M_d,M) .

Four model classes

Class M: Purely multiplicative models: (M,N,N) , (M,N,M) , (M,M,N) , (M,M,M) , (M,M_d,N) and (M,M_d,M) .

Class A: Purely additive models: (A,N,N) , (A,N,A) , (A,A,N) , (A,A,A) , (A,A_d,N) and (A,A_d,A) .

Class X: Mixed models: $(A,M,*)$, $(A,M_d,*)$, $(A,*,M)$, (M,M,A) , (M,M_d,A) . (11 models)

Class Y: Mixed models: $(M,A,*)$, $(M,A_d,*)$ or (M,N,A) . (7 models)

Four model classes

Class M: Purely multiplicative models: (M,N,N) , (M,N,M) , (M,M,N) , (M,M,M) , (M,M_d,N) and (M,M_d,M) .

Class A: Purely additive models: (A,N,N) , (A,N,A) , (A,A,N) , (A,A,A) , (A,A_d,N) and (A,A_d,A) .

Class X: Mixed models: $(A,M,*)$, $(A,M_d,*)$, $(A,*,M)$, (M,M,A) , (M,M_d,A) . (11 models)

Class Y: Mixed models: $(M,A,*)$, $(M,A_d,*)$ or (M,N,A) . (7 models)

- Only Class M can guarantee a sample space restricted to the positive half-line.

Four model classes

Class M: Purely multiplicative models: (M,N,N) , (M,N,M) , (M,M,N) , (M,M,M) , (M,M_d,N) and (M,M_d,M) .

Class A: Purely additive models: (A,N,N) , (A,N,A) , (A,A,N) , (A,A,A) , (A,A_d,N) and (A,A_d,A) .

Class X: Mixed models: $(A,M,*)$, $(A,M_d,*)$, $(A,*,M)$, (M,M,A) , (M,M_d,A) . (11 models)

Class Y: Mixed models: $(M,A,*)$, $(M,A_d,*)$ or (M,N,A) . (7 models)

- Only Class M can guarantee a sample space restricted to the positive half-line.
- All Class M models converge to 0 if $E(\varepsilon) = 0$

Four model classes

Class M: Purely multiplicative models: (M,N,N) , (M,N,M) , (M,M,N) , (M,M,M) , (M,M_d,N) and (M,M_d,M) .

Class A: Purely additive models: (A,N,N) , (A,N,A) , (A,A,N) , (A,A,A) , (A,A_d,N) and (A,A_d,A) .

Class X: Mixed models: $(A,M,*)$, $(A,M_d,*)$, $(A,*,M)$, (M,M,A) , (M,M_d,A) . (11 models)

Class Y: Mixed models: $(M,A,*)$, $(M,A_d,*)$ or (M,N,A) . (7 models)

- Only Class M can guarantee a sample space restricted to the positive half-line.
- All Class M models converge to 0 if $E(\varepsilon) = 0$
- All Class X models have infinite forecast variance for $h \geq m + 2$ where m is the seasonal period.

Outline

- 1 Exponential smoothing models
- 2 Problems with some of the models
- 3 A new model for positive data**
- 4 Conclusions

New models

Let $\delta_t = (1 + \varepsilon_t)$ be a positive random variable.

METS(M,N,N) model

$$y_t = \ell_{t-1} \delta_t$$

$$\ell_t = \ell_{t-1} \delta_t^\alpha$$

New models

Let $\delta_t = (1 + \varepsilon_t)$ be a positive random variable.

METS(M,N,N) model

$$y_t = \ell_{t-1} \delta_t$$

$$\ell_t = \ell_{t-1} \delta_t^\alpha$$

$$\log(y_t) = \log(\ell_{t-1}) + \log(\delta_t)$$

$$\log(\ell_t) = \log(\ell_{t-1}) + \alpha \log(\delta_t)$$

New models

Let $\delta_t = (1 + \varepsilon_t)$ be a positive random variable.

METS(M,N,N) model

$$y_t = \ell_{t-1} \delta_t$$

$$\ell_t = \ell_{t-1} \delta_t^\alpha$$

$$\log(y_t) = \log(\ell_{t-1}) + \log(\delta_t)$$

$$\log(\ell_t) = \log(\ell_{t-1}) + \alpha \log(\delta_t)$$

Thus the log-transformed model is identical to Gaussian ETS(A,N,N) model if δ_t is logNormal with median 1.

Long term forecast behaviour

METS(M,N,N; LN) model

$$y_t = l_{t-1} \delta_t$$

$$l_t = l_{t-1} \delta_t^\alpha,$$

$$\delta_t \sim \text{logN}(\mu, \omega)$$

Long term forecast behaviour

METS(M,N,N; LN) model







$$y_t = l_{t-1} \delta_t$$

$$l_t = l_{t-1} \delta_t^\alpha,$$

$$\delta_t \sim \log N(\mu, \omega)$$

Range	$E(\delta_t^\alpha)$	$E(\delta_t^{\alpha/2})$	$E(y_h)$	$V(y_h)$
$\mu + \alpha\omega < 0$	< 1	< 1	Decreasing	Decreasing
$\mu + \alpha\omega = 0$	< 1	< 1	Decreasing	Finite
$-\alpha\omega < \mu < -\alpha\omega/2$	< 1	< 1	Decreasing	Increasing
$\mu + \alpha\omega/2 = 0$	$= 1$	< 1	Finite	Increasing
$-\alpha\omega/2 < \mu < -\alpha\omega/4$	> 1	< 1	Increasing	Increasing
$\mu + \alpha\omega/4 = 0$	> 1	$= 1$	Increasing	Increasing
$\mu + \alpha\omega/4 > 0$	> 1	> 1	Increasing	Increasing

Long term forecast behaviour



METS(M,N,N;LN)
 $\delta_t \sim \log N(\mu, \omega)$:

(a) $\mu = \alpha\omega/4$

(b) $\mu = 0$

(c) $\mu = -\alpha\omega/4$

(d) $\mu = -3\alpha\omega/8$

(e) $\mu = -\alpha\omega/2$

(f) $\mu = -3\alpha\omega/4$

Outline

- 1 Exponential smoothing models
- 2 Problems with some of the models
- 3 A new model for positive data
- 4 Conclusions**

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.
- 3 An alternative formulation has been proposed that avoids these problems.

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.
- 3 An alternative formulation has been proposed that avoids these problems.
- 4 The ergodic behaviour of the alternative formulation requires careful parameter choice.

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.
- 3 An alternative formulation has been proposed that avoids these problems.
- 4 The ergodic behaviour of the alternative formulation requires careful parameter choice.

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.
- 3 An alternative formulation has been proposed that avoids these problems.
- 4 The ergodic behaviour of the alternative formulation requires careful parameter choice.

Paper: **www.robhyndman.info**

Conclusions

- 1 The standard exponential smoothing state space models all have theoretical problems when constrained to the positive half line.
- 2 This is not a concern for most short-term forecasting.
- 3 An alternative formulation has been proposed that avoids these problems.
- 4 The ergodic behaviour of the alternative formulation requires careful parameter choice.

Paper: **www.robhyndman.info**

Book: **www.exponentialsMOOTHING.net**