



Rob J Hyndman

Forecasting: Principles and Practice




11. Hierarchical forecasting

[OTexts.com/fpp/9/4/](https://otexts.com/fpp/9/4/)

Outline

- 1 Hierarchical and grouped time series**
- 2 Forecasting framework
- 3 Optimal forecasts
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism
- 6 Application: Australian labour market
- 7 hts package for R


Introduction



Examples

- Manufacturing product hierarchies
- Net labour turnover
- Pharmaceutical sales
- Customer demand by region and purchase


Introduction



Examples

- Manufacturing product hierarchies
- Net labour turnover
- Pharmaceutical sales
- Tourism demand by region and purpose


Introduction



Examples

- Manufacturing product hierarchies
- Net labour turnover
- Pharmaceutical sales
- Tourism demand by region and purpose


Introduction



Examples

- Manufacturing product hierarchies
- Net labour turnover
- Pharmaceutical sales
- Tourism demand by region and purpose

Introduction



Examples

- Manufacturing product hierarchies
- Net labour turnover
- Pharmaceutical sales
- Tourism demand by region and purpose

Forecasting the PBS



ATC drug classification

- A Alimentary tract and metabolism
- B Blood and blood forming organs
- C Cardiovascular system
- D Dermatologicals
- G Genito-urinary system and sex hormones
- H Systemic hormonal preparations, excluding sex hormones and insulins
- J Anti-infectives for systemic use
- L Antineoplastic and immunomodulating agents
- M Musculo-skeletal system
- N Nervous system
- P Antiparasitic products, insecticides and repellents
- R Respiratory system
- S Sensory organs
- V Various

ATC drug classification

14 classes

A

Alimentary tract and metabolism

84 classes

A10

Drugs used in diabetes

A10B

Blood glucose lowering drugs


A10BA

Biguanides

A10BA02

Metformin


Australian tourism



Australian tourism

Also split by purpose of travel:

- Holiday
- Visits to friends and relatives
- Business
- Other



Hierarchical/grouped time series

- A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.

Example: Pharmaceutical products are organized in a hierarchy under the Anatomical Therapeutic Chemical (ATC) Classification System.

- A **grouped time series** is a collection of time series that are aggregated in a number of non-hierarchical ways.

Hierarchical/grouped time series

- A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.

Example: Pharmaceutical products are organized in a hierarchy under the Anatomical Therapeutic Chemical (ATC) Classification System.

- A **grouped time series** is a collection of time series that are aggregated in a number of non-hierarchical ways.

Example: Australian tourism demand is grouped by region and purpose of travel.

Hierarchical/grouped time series

- A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.

Example: Pharmaceutical products are organized in a hierarchy under the Anatomical Therapeutic Chemical (ATC) Classification System.

- A **grouped time series** is a collection of time series that are aggregated in a number of non-hierarchical ways.

Example: Australian tourism demand is grouped by region and purpose of travel.

Hierarchical/grouped time series


- A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.

Example: Pharmaceutical products are organized in a hierarchy under the Anatomical Therapeutic Chemical (ATC) Classification System.

- A **grouped time series** is a collection of time series that are aggregated in a number of non-hierarchical ways.

Example: Australian tourism demand is grouped by region and purpose of travel.

Hierarchical data




Y_t : observed aggregate of all series at time t .

$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

Hierarchical data




Y_t : observed aggregate of all series at time t .

$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

Hierarchical data




Y_t : observed aggregate of all series at time t .

$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

$$\mathbf{Y}_t = [Y_t, Y_{A,t}, Y_{B,t}, Y_{C,t}]' = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \end{pmatrix}$$

Hierarchical data




Y_t : observed aggregate of all series at time t .

$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

$$\mathbf{Y}_t = [Y_t, Y_{A,t}, Y_{B,t}, Y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_S \begin{pmatrix} Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \end{pmatrix}$$

Hierarchical data




Y_t : observed aggregate of all series at time t .

$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

$$\mathbf{Y}_t = [Y_t, Y_{A,t}, Y_{B,t}, Y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \end{pmatrix}}_{\mathbf{B}_t}$$

Hierarchical data



Y_t : observed aggregate of all series at time t .


$Y_{X,t}$: observation on series X at time t .

B_t : vector of all series at bottom level in time t .

$$\mathbf{Y}_t = [Y_t, Y_{A,t}, Y_{B,t}, Y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \end{pmatrix}}_{\mathbf{B}_t}$$


$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t$$

Hierarchical data




$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix}}_{\mathbf{B}_t}$$

Hierarchical data



$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix}}_{\mathbf{B}_t}$$


Hierarchical data



$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{C,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{AZ,t} \\ Y_{BX,t} \\ Y_{BY,t} \\ Y_{BZ,t} \\ Y_{CX,t} \\ Y_{CY,t} \\ Y_{CZ,t} \end{pmatrix}}_{\mathbf{B}_t}$$


$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t$$

Grouped data




$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{X,t} \\ Y_{Y,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix}}_{\mathbf{B}_t}$$

Grouped data



$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{X,t} \\ Y_{Y,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix}}_{\mathbf{B}_t}$$

Grouped data



$$\mathbf{Y}_t = \begin{pmatrix} Y_t \\ Y_{A,t} \\ Y_{B,t} \\ Y_{X,t} \\ Y_{Y,t} \\ Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} Y_{AX,t} \\ Y_{AY,t} \\ Y_{BX,t} \\ Y_{BY,t} \end{pmatrix}}_{\mathbf{B}_t}$$

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t$$

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework**
- 3 Optimal forecasts
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism
- 6 Application: Australian labour market
- 7 hts package for R

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{Y}}_n(h)$ to get bottom-level forecasts.
- Roll them up
- Revised (re-rolled) forecasts $\tilde{\mathbf{Y}}_n(h)$.

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{Y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up
- Revised reconciled forecasts: $\tilde{\mathbf{Y}}_n(h)$.

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{Y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up
- Revised reconciled forecasts: $\tilde{\mathbf{Y}}_n(h)$.

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{Y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up
- Revised reconciled forecasts: $\tilde{\mathbf{Y}}_n(h)$.

Forecasting notation

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t . (They may not add up.)

Hierarchical forecasting methods of the form:

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

for some matrix \mathbf{P} .

- \mathbf{P} extracts and combines base forecasts $\hat{\mathbf{Y}}_n(h)$ to get bottom-level forecasts.
- \mathbf{S} adds them up
- Revised reconciled forecasts: $\tilde{\mathbf{Y}}_n(h)$.

Bottom-up forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}],$$

where $\mathbf{0}$ is null matrix and \mathbf{I} is identity matrix.

- \mathbf{P} matrix extracts only bottom-level forecasts from $\hat{\mathbf{Y}}_n(h)$
- \mathbf{S} adds them up to give the bottom-up forecasts.

Bottom-up forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}],$$

where $\mathbf{0}$ is null matrix and \mathbf{I} is identity matrix.

- \mathbf{P} matrix extracts only bottom-level forecasts from $\hat{\mathbf{Y}}_n(h)$
- \mathbf{S} adds them up to give the bottom-up forecasts.

Bottom-up forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}],$$

where $\mathbf{0}$ is null matrix and \mathbf{I} is identity matrix.

- \mathbf{P} matrix extracts only bottom-level forecasts from $\hat{\mathbf{Y}}_n(h)$
- \mathbf{S} adds them up to give the bottom-up forecasts.

Top-down forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$ is a vector of proportions that sum to one.

- \mathbf{P} distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors \mathbf{p} .

Top-down forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$ is a vector of proportions that sum to one.

- \mathbf{P} distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors \mathbf{p} .

Top-down forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$ is a vector of proportions that sum to one.

- \mathbf{P} distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors \mathbf{p} .

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$E[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = E[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = E[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.

Then $E[\tilde{\mathbf{Y}}_n(h)] = \mathbf{S}\mathbf{D}_n(h)$.

We want the revised forecasts to be unbiased:

$$E[\tilde{\mathbf{Y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{D}_n(h) = \mathbf{S}\mathbf{D}_n(h)$$

Result will hold as long as $\mathbf{S}\mathbf{P} = \mathbf{I}$.

The \mathbf{P} matrix is the \mathbf{P} matrix from the bottom level forecasting model.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbb{E}[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

■ Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.

■ Then $\mathbb{E}[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.

We want the revised forecasts to be unbiased:

$$\mathbb{E}[\tilde{\mathbf{Y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{S}\beta_n(h) = \mathbf{S}\beta_n(h).$$

For this to hold we need $\mathbf{P}\mathbf{S} = \mathbf{I}$.

The following is a sufficient condition for $\mathbf{P}\mathbf{S} = \mathbf{I}$:
If $\mathbf{S} = \mathbf{S}_1\mathbf{S}_2\mathbf{S}_3$ and $\mathbf{P} = \mathbf{P}_1\mathbf{P}_2\mathbf{P}_3$ with $\mathbf{P}_1\mathbf{S}_1 = \mathbf{I}$, $\mathbf{P}_2\mathbf{S}_2 = \mathbf{I}$ and $\mathbf{P}_3\mathbf{S}_3 = \mathbf{I}$ then $\mathbf{P}\mathbf{S} = \mathbf{I}$.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$E[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = E[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = E[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $E[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased:
 $E[\tilde{\mathbf{Y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$\mathbb{E}[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbb{E}[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased:
 $\mathbb{E}[\tilde{\mathbf{Y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{S}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Result will hold provided $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$E[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = E[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = E[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $E[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased:
 $E[\tilde{\mathbf{Y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

Result will hold provided $\mathbf{SPS} = \mathbf{S}$.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:

$$E[\hat{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = E[\mathbf{Y}_{n+h}|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = E[\hat{\mathbf{B}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $E[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased:
 $E[\tilde{\mathbf{Y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.

■ Result will hold provided $\mathbf{SPS} = \mathbf{S}$.

■ True for bottom-up, but not for any top-down method or middle-out method.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:
$$\mathbb{E}[\hat{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbb{E}[\mathbf{Y}_{n+h} | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{B}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased:
$$\mathbb{E}[\tilde{\mathbf{Y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h).$$
- Result will hold provided **$\mathbf{SPS} = \mathbf{S}$** .
- True for bottom-up, but not for *any* top-down method or middle-out method.

General properties: bias

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Assume: base forecasts $\hat{\mathbf{Y}}_n(h)$ are unbiased:
$$\mathbb{E}[\hat{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbb{E}[\mathbf{Y}_{n+h} | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

- Let $\hat{\mathbf{B}}_n(h)$ be bottom level base forecasts with $\beta_n(h) = \mathbb{E}[\hat{\mathbf{B}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- Then $\mathbb{E}[\hat{\mathbf{Y}}_n(h)] = \mathbf{S}\beta_n(h)$.
- We want the revised forecasts to be unbiased: $\mathbb{E}[\tilde{\mathbf{Y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$.
- Result will hold provided $\mathbf{SPS} = \mathbf{S}$.
- True for bottom-up, but not for *any* top-down method or middle-out method.

General properties: variance

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Let variance of base forecasts $\hat{\mathbf{Y}}_n(h)$ be given by

$$\Sigma_h = \text{Var}[\hat{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

Then the variance of the revised forecasts is given by

$$\text{Var}[\tilde{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{SP}\Sigma_h\mathbf{P}'\mathbf{S}'.$$

This is a general result for all existing methods.

General properties: variance

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Let variance of base forecasts $\hat{\mathbf{Y}}_n(h)$ be given by

$$\Sigma_h = \text{Var}[\hat{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

Then the variance of the revised forecasts is given by

$$\text{Var}[\tilde{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{SP}\Sigma_h\mathbf{P}'\mathbf{S}'.$$

This is a general result for all existing methods.

General properties: variance

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{SP}\hat{\mathbf{Y}}_n(h)$$

Let variance of base forecasts $\hat{\mathbf{Y}}_n(h)$ be given by

$$\Sigma_h = \text{Var}[\hat{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n]$$

Then the variance of the revised forecasts is given by

$$\text{Var}[\tilde{\mathbf{Y}}_n(h) | \mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{SP}\Sigma_h\mathbf{P}'\mathbf{S}'.$$

This is a general result for all existing methods.

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework
- 3 Optimal forecasts**
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism
- 6 Application: Australian labour market
- 7 hts package for R

Forecasts

Key idea: forecast reconciliation

- ➔ Ignore structural constraints and forecast every series of interest independently.
- ➔ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

Forecasts

Key idea: forecast reconciliation

- ➡ Ignore structural constraints and forecast every series of interest independently.
- ➡ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

Forecasts

Key idea: forecast reconciliation

- ➡ Ignore structural constraints and forecast every series of interest independently.
- ➡ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

$$\boldsymbol{\beta}_n(h) = E[\mathbf{B}_{n+h} \mid \mathbf{Y}_1, \dots, \mathbf{Y}_n].$$

Forecast reconciliation

Forecasts

Key idea: forecast reconciliation

- ➔ Ignore structural constraints and forecast every series of interest independently.
- ➔ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

- $\boldsymbol{\beta}_n(h) = E[\mathbf{B}_{n+h} \mid \mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- $\boldsymbol{\varepsilon}_h$ has zero mean and covariance $\boldsymbol{\Sigma}_h$.
- Estimate $\boldsymbol{\beta}_n(h)$ using GLS?

Forecasts

Key idea: forecast reconciliation

- ➔ Ignore structural constraints and forecast every series of interest independently.
- ➔ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

- $\boldsymbol{\beta}_n(h) = \mathbf{E}[\mathbf{B}_{n+h} \mid \mathbf{Y}_1, \dots, \mathbf{Y}_n].$
- $\boldsymbol{\varepsilon}_h$ has zero mean and covariance $\boldsymbol{\Sigma}_h.$
- Estimate $\boldsymbol{\beta}_n(h)$ using GLS?

Forecasts

Key idea: forecast reconciliation

- ➔ Ignore structural constraints and forecast every series of interest independently.
- ➔ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

- $\boldsymbol{\beta}_n(h) = \mathbb{E}[\mathbf{B}_{n+h} \mid \mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- $\boldsymbol{\varepsilon}_h$ has zero mean and covariance $\boldsymbol{\Sigma}_h$.
- Estimate $\boldsymbol{\beta}_n(h)$ using GLS?

Forecasts

Key idea: forecast reconciliation

- ➔ Ignore structural constraints and forecast every series of interest independently.
- ➔ Adjust forecasts to impose constraints.

Let $\hat{\mathbf{Y}}_n(h)$ be vector of initial h -step forecasts, made at time n , stacked in same order as \mathbf{Y}_t .

$$\mathbf{Y}_t = \mathbf{S}\mathbf{B}_t. \quad \text{So } \hat{\mathbf{Y}}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h) + \boldsymbol{\varepsilon}_h.$$

- $\boldsymbol{\beta}_n(h) = \text{E}[\mathbf{B}_{n+h} \mid \mathbf{Y}_1, \dots, \mathbf{Y}_n]$.
- $\boldsymbol{\varepsilon}_h$ has zero mean and covariance $\boldsymbol{\Sigma}_h$.
- Estimate $\boldsymbol{\beta}_n(h)$ using GLS?

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\boldsymbol{\beta}}_n(h) = \mathbf{S}(\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\boldsymbol{\beta}}_n(h) = \mathbf{S}(\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Initial forecasts

• $\boldsymbol{\Sigma}_h^\dagger$ is generalized inverse of $\boldsymbol{\Sigma}_h$.

• $\boldsymbol{\Sigma}_h^\dagger$ is not unique, but $\mathbf{S}(\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\boldsymbol{\Sigma}_h^\dagger$ is.

• $\boldsymbol{\Sigma}_h^\dagger$ is hard to estimate.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Σ_h^\dagger is generalized inverse of Σ_h .
- $\text{Var}[\tilde{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'$
- **Problem:** Σ_h hard to estimate.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Σ_h^\dagger is generalized inverse of Σ_h .
- $\text{Var}[\tilde{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'$
- **Problem:** Σ_h hard to estimate.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Σ_h^\dagger is generalized inverse of Σ_h .
- $\text{Var}[\tilde{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'$
- **Problem:** Σ_h hard to estimate.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Σ_h^\dagger is generalized inverse of Σ_h .
- $\text{Var}[\tilde{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'$
- **Problem:** Σ_h hard to estimate.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Σ_h^\dagger is generalized inverse of Σ_h .
- $\text{Var}[\tilde{\mathbf{Y}}_n(h)|\mathbf{Y}_1, \dots, \mathbf{Y}_n] = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'$
- **Problem:** Σ_h hard to estimate.

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework
- 3 Optimal forecasts
- 4 Approximately optimal forecasts**
- 5 Application: Australian tourism
- 6 Application: Australian labour market
- 7 hts package for R

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\epsilon_h \approx \mathbf{S}\epsilon_{B,h}$ where $\epsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\epsilon_{B,h})$.
- If Moore-Penrose generalized inverse is used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger\hat{\mathbf{Y}}_n(h)$$

Revised forecasts


Base forecasts

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.
- Or assume $\varepsilon_h \approx \mathbf{S}\varepsilon_{B,h}$ where $\varepsilon_{B,h}$ is the forecast error at bottom level.
- Then $\Sigma_h \approx \mathbf{S}\Omega_h\mathbf{S}'$ where $\Omega_h = \text{Var}(\varepsilon_{B,h})$.
- If Moore-Penrose generalized inverse used, then $(\mathbf{S}'\Sigma_h^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^\dagger = (\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$.


$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts



$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts




$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

Weights:

$$\mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}' = \begin{bmatrix} 0.75 & 0.25 & 0.25 & 0.25 \\ 0.25 & 0.75 & -0.25 & -0.25 \\ 0.25 & -0.25 & 0.75 & -0.25 \\ 0.25 & -0.25 & -0.25 & 0.75 \end{bmatrix}$$


Optimal combination forecasts



Weights: $S(S'S)^{-1}S' =$

0.69	0.23	0.23	0.23	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
0.23	0.58	-0.17	-0.17	0.19	0.19	0.19	-0.06	-0.06	-0.06	-0.06	-0.06	-0.06
0.23	-0.17	0.58	-0.17	-0.06	-0.06	-0.06	0.19	0.19	0.19	-0.06	-0.06	-0.06
0.23	-0.17	-0.17	0.58	-0.06	-0.06	-0.06	-0.06	-0.06	-0.06	0.19	0.19	0.19
0.08	0.19	-0.06	-0.06	0.73	-0.27	-0.27	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	0.19	-0.06	-0.06	-0.27	0.73	-0.27	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	0.19	-0.06	-0.06	-0.27	-0.27	0.73	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	0.73	-0.27	-0.27	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	-0.27	0.73	-0.27	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	-0.27	-0.27	0.73	-0.02	-0.02	-0.02
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	0.73	-0.27	-0.27
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	-0.27	0.73	-0.27
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	-0.27	-0.27	0.73

Optimal combination forecasts



Weights: $S(S'S)^{-1}S' =$

0.69	0.23	0.23	0.23	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08	0.08
0.23	0.58	-0.17	-0.17	0.19	0.19	0.19	-0.06	-0.06	-0.06	-0.06	-0.06	-0.06
0.23	-0.17	0.58	-0.17	-0.06	-0.06	-0.06	0.19	0.19	0.19	-0.06	-0.06	-0.06
0.23	-0.17	-0.17	0.58	-0.06	-0.06	-0.06	-0.06	-0.06	-0.06	0.19	0.19	0.19
0.08	0.19	-0.06	-0.06	0.73	-0.27	-0.27	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	0.19	-0.06	-0.06	-0.27	0.73	-0.27	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	0.19	-0.06	-0.06	-0.27	-0.27	0.73	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	0.73	-0.27	-0.27	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	-0.27	0.73	-0.27	-0.02	-0.02	-0.02
0.08	-0.06	0.19	-0.06	-0.02	-0.02	-0.02	-0.27	-0.27	0.73	-0.02	-0.02	-0.02
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	0.73	-0.27	-0.27
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	-0.27	0.73	-0.27
0.08	-0.06	-0.06	0.19	-0.02	-0.02	-0.02	-0.02	-0.02	-0.02	-0.27	-0.27	0.73

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- **$SPS = S$** so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- Weights are independent of the data and of the covariance structure of the hierarchy.

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- **$SPS = S$** so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- Weights are independent of the data and of the covariance structure of the hierarchy.

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- $SPS = S$ so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- Weights are independent of the data and of the covariance structure of the hierarchy.

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- ***SPS* = *S*** so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- Weights are independent of the data and of the covariance structure of the hierarchy.

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- **$SPS = S$** so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- Weights are independent of the data and of the covariance structure of the hierarchy.

Features

- Covariates can be included in initial forecasts.
- Adjustments can be made to initial forecasts at any level.
- Very simple and flexible method. Can work with *any* hierarchical or grouped time series.
- **$SPS = S$** so reconciled forecasts are unbiased.
- Conceptually easy to implement: OLS on base forecasts.
- **Weights are independent of the data and of the covariance structure of the hierarchy.**

Challenges



$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

- Computational difficulties in big hierarchies due to size of the \mathbf{S} matrix and singular behavior of $(\mathbf{S}'\mathbf{S})$.
- Need to estimate covariance matrix to produce prediction intervals.
- Ignores covariance matrix in computing point forecasts.

Challenges



$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

- Computational difficulties in big hierarchies due to size of the \mathbf{S} matrix and singular behavior of $(\mathbf{S}'\mathbf{S})$.
- Need to estimate covariance matrix to produce prediction intervals.
- Ignores covariance matrix in computing point forecasts.

Challenges



$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'\hat{\mathbf{Y}}_n(h)$$

- Computational difficulties in big hierarchies due to size of the \mathbf{S} matrix and singular behavior of $(\mathbf{S}'\mathbf{S})$.
- Need to estimate covariance matrix to produce prediction intervals.
- Ignores covariance matrix in computing point forecasts.

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_1^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_1^\dagger\hat{\mathbf{Y}}_n(h)$$

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.

Solution 2: Rescaling

- Suppose we approximate Σ_1 by its diagonal.
- Let $\Lambda = [\text{diagonal}(\Sigma_1)]^{-1}$ contain inverse one-step forecast variances.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_1^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_1^\dagger\hat{\mathbf{Y}}_n(h)$$

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.

Solution 2: Rescaling

- Suppose we approximate Σ_1 by its diagonal.
- Let $\Lambda = [\text{diagonal}(\Sigma_1)]^{-1}$ contain inverse one-step forecast variances.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_1^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_1^\dagger\hat{\mathbf{Y}}_n(h)$$

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.

Solution 2: Rescaling

- Suppose we approximate Σ_1 by its diagonal.
- Let $\Lambda = [\text{diagonal}(\Sigma_1)]^{-1}$ contain inverse one-step forecast variances.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_1^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_1^\dagger\hat{\mathbf{Y}}_n(h)$$

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.

Solution 2: Rescaling

- Suppose we approximate Σ_1 by its diagonal.
- Let $\Lambda = [\text{diagonal}(\Sigma_1)]^{-1}$ contain inverse one-step forecast variances.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Optimal combination forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_1^\dagger\mathbf{S})^{-1}\mathbf{S}'\Sigma_1^\dagger\hat{\mathbf{Y}}_n(h)$$

Solution 1: OLS

- Approximate Σ_1^\dagger by $c\mathbf{I}$.

Solution 2: Rescaling

- Suppose we approximate Σ_1 by its diagonal.
- Let $\Lambda = [\text{diagonal}(\Sigma_1)]^{-1}$ contain inverse one-step forecast variances.

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Optimal reconciled forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{\Lambda}\mathbf{S})^{-1}\mathbf{S}'\mathbf{\Lambda}\hat{\mathbf{Y}}_n(h)$$

- Easy to estimate, and places weight where we have best forecasts.

■ ignores covariances

■ for large numbers of time series, we can

try to make the other optimality property

more likely to hold

Optimal reconciled forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{\Lambda}\mathbf{S})^{-1}\mathbf{S}'\mathbf{\Lambda}\hat{\mathbf{Y}}_n(h)$$

Initial forecasts

- Easy to estimate, and places weight where we have best forecasts.
- Ignores covariances.
- For large numbers of time series, we need to do calculation without explicitly forming \mathbf{S} or $(\mathbf{S}'\mathbf{\Lambda}\mathbf{S})^{-1}$ or $\mathbf{S}'\mathbf{\Lambda}$.

Optimal reconciled forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\mathbf{\Lambda}\mathbf{S})^{-1}\mathbf{S}'\mathbf{\Lambda}\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Easy to estimate, and places weight where we have best forecasts.
- Ignores covariances.
- For large numbers of time series, we need to do calculation without explicitly forming \mathbf{S} or $(\mathbf{S}'\mathbf{\Lambda}\mathbf{S})^{-1}$ or $\mathbf{S}'\mathbf{\Lambda}$.

Optimal reconciled forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Revised forecasts

Initial forecasts

- Easy to estimate, and places weight where we have best forecasts.
- Ignores covariances.
- For large numbers of time series, we need to do calculation without explicitly forming \mathbf{S} or $(\mathbf{S}'\Lambda\mathbf{S})^{-1}$ or $\mathbf{S}'\Lambda$.

Optimal reconciled forecasts

$$\tilde{\mathbf{Y}}_n(h) = \mathbf{S}\hat{\beta}_n(h) = \mathbf{S}(\mathbf{S}'\Lambda\mathbf{S})^{-1}\mathbf{S}'\Lambda\hat{\mathbf{Y}}_n(h)$$

Revised forecasts


Initial forecasts

- Easy to estimate, and places weight where we have best forecasts.
- Ignores covariances.
- For large numbers of time series, we need to do calculation without explicitly forming \mathbf{S} or $(\mathbf{S}'\Lambda\mathbf{S})^{-1}$ or $\mathbf{S}'\Lambda$.

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework
- 3 Optimal forecasts
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism**
- 6 Application: Australian labour market
- 7 hts package for R

Australian tourism




Australian tourism

Domestic visitor nights

Quarterly data: 1998 – 2006.


From: *National Visitor Survey*, based on annual interviews of 120,000 Australians aged 15+, collected by Tourism Research Australia.



Australian tourism

Hierarchy:

- States (7)
- Zones (27)
- Regions (82)



Australian tourism


Hierarchy:

- States (7)
- Zones (27)
- Regions (82)

Base forecasts


ETS (exponential smoothing)
models

- 0 to 3%
- > 3%




Base forecasts

Domestic tourism forecasts: Total




Base forecasts

Domestic tourism forecasts: NSW




Base forecasts

Domestic tourism forecasts: VIC




Base forecasts

Domestic tourism forecasts: Nth.Coast.NSW




Base forecasts

Domestic tourism forecasts: Metro.QLD




Base forecasts

Domestic tourism forecasts: Sth.WA




Base forecasts

Domestic tourism forecasts: X201.Melbourne




Base forecasts

Domestic tourism forecasts: X402.Murraylands




Base forecasts





Domestic tourism forecasts: X809.Daly











Reconciled forecasts



Reconciled forecasts



Reconciled forecasts



Forecast evaluation

- **Select models using all observations;**
- Re-estimate models using first 12 observations and generate 1- to 8-step-ahead forecasts;
- Increase sample size one observation at a time, re-estimate models, generate forecasts until the end of the sample;
- In total 24 1-step-ahead, 23 2-steps-ahead, up to 17 8-steps-ahead for forecast evaluation.

Forecast evaluation

- Select models using all observations;
- Re-estimate models using first 12 observations and generate 1- to 8-step-ahead forecasts;
- Increase sample size one observation at a time, re-estimate models, generate forecasts until the end of the sample;
- In total 24 1-step-ahead, 23 2-steps-ahead, up to 17 8-steps-ahead for forecast evaluation.

Forecast evaluation

- Select models using all observations;
- Re-estimate models using first 12 observations and generate 1- to 8-step-ahead forecasts;
- Increase sample size one observation at a time, re-estimate models, generate forecasts until the end of the sample;
- In total 24 1-step-ahead, 23 2-steps-ahead, up to 17 8-steps-ahead for forecast evaluation.

Forecast evaluation

- Select models using all observations;
- Re-estimate models using first 12 observations and generate 1- to 8-step-ahead forecasts;
- Increase sample size one observation at a time, re-estimate models, generate forecasts until the end of the sample;
- In total 24 1-step-ahead, 23 2-steps-ahead, up to 17 8-steps-ahead for forecast evaluation.

Hierarchy: states, zones, regions

MAPE	$h = 1$	$h = 2$	$h = 4$	$h = 6$	$h = 8$	Average
<i>Top Level: Australia</i>						
Bottom-up	3.79	3.58	4.01	4.55	4.24	4.06
OLS	3.83	3.66	3.88	4.19	4.25	3.94
Scaling (st. dev.)	3.68	3.56	3.97	4.57	4.25	4.04
<i>Level: States</i>						
Bottom-up	10.70	10.52	10.85	11.46	11.27	11.03
OLS	11.07	10.58	11.13	11.62	12.21	11.35
Scaling (st. dev.)	10.44	10.17	10.47	10.97	10.98	10.67
<i>Level: Zones</i>						
Bottom-up	14.99	14.97	14.98	15.69	15.65	15.32
OLS	15.16	15.06	15.27	15.74	16.15	15.48
Scaling (st. dev.)	14.63	14.62	14.68	15.17	15.25	14.94
<i>Bottom Level: Regions</i>						
Bottom-up	33.12	32.54	32.26	33.74	33.96	33.18
OLS	35.89	33.86	34.26	36.06	37.49	35.43
Scaling (st. dev.)	31.68	31.22	31.08	32.41	32.77	31.89

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework
- 3 Optimal forecasts
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism
- 6 Application: Australian labour market**
- 7 hts package for R

Australia and New Zealand Standard Classification of Occupations

- 8 major groups
 - 43 sub-major groups
 - 97 minor groups
 - 359 unit groups
 - * 1023 occupations

Example: statistician

2 Professionals

22 Business, Human Resource and Marketing Professionals

224 Information and Organisation Professionals

2241 Actuaries, Mathematicians and Statisticians

224113 Statistician

Australia and New Zealand Standard Classification of Occupations

- 8 major groups
 - 43 sub-major groups
 - 97 minor groups
 - 359 unit groups
 - * 1023 occupations

Example: statistician

2 Professionals


22 Business, Human Resource and Marketing Professionals

224 Information and Organisation Professionals


2241 Actuaries, Mathematicians and Statisticians

224113 Statistician

Australian Labour Market data




Australian Labour Market data




- Lower three panels show largest sub-groups at each level.

Australian Labour Market data




Forecasting: Principles and Practice




Application: Australian labour market

Australian Labour Market data



- Base forecasts from `auto.arima()`
- Largest changes shown for each level



Forecast evaluation (rolling origin)

RMSE	$h = 1$	$h = 2$	$h = 3$	$h = 4$	$h = 5$	$h = 6$	$h = 7$	$h = 8$	Average
<i>Top level</i>									
Bottom-up	74.71	102.02	121.70	131.17	147.08	157.12	169.60	178.93	135.29
OLS	52.20	77.77	101.50	119.03	138.27	150.75	160.04	166.38	120.74
WLS	61.77	86.32	107.26	119.33	137.01	146.88	156.71	162.38	122.21
<i>Level 1</i>									
Bottom-up	21.59	27.33	30.81	32.94	35.45	37.10	39.00	40.51	33.09
OLS	21.89	28.55	32.74	35.58	38.82	41.24	43.34	45.49	35.96
WLS	20.58	26.19	29.71	31.84	34.36	35.89	37.53	38.86	31.87
<i>Level 2</i>									
Bottom-up	8.78	10.72	11.79	12.42	13.13	13.61	14.14	14.65	12.40
OLS	9.02	11.19	12.34	13.04	13.92	14.56	15.17	15.77	13.13
WLS	8.58	10.48	11.54	12.15	12.88	13.36	13.87	14.36	12.15
<i>Level 3</i>									
Bottom-up	5.44	6.57	7.17	7.53	7.94	8.27	8.60	8.89	7.55
OLS	5.55	6.78	7.42	7.81	8.29	8.68	9.04	9.37	7.87
WLS	5.35	6.46	7.06	7.42	7.84	8.17	8.48	8.76	7.44
<i>Bottom Level</i>									
Bottom-up	2.35	2.79	3.02	3.15	3.29	3.42	3.54	3.65	3.15
OLS	2.40	2.86	3.10	3.24	3.41	3.55	3.68	3.80	3.25
WLS	2.34	2.77	2.99	3.12	3.27	3.40	3.52	3.63	3.13

Outline

- 1 Hierarchical and grouped time series
- 2 Forecasting framework
- 3 Optimal forecasts
- 4 Approximately optimal forecasts
- 5 Application: Australian tourism
- 6 Application: Australian labour market
- 7 hts package for R**

hts package for R



hts: Hierarchical and grouped time series

Methods for analysing and forecasting hierarchical and grouped time series

Version: 4.3

Depends: forecast (\geq 5.0)

Imports: SparseM, parallel, utils

Published: 2014-06-10

Author: Rob J Hyndman, Earo Wang and Alan Lee

Maintainer: Rob J Hyndman <Rob.Hyndman@monash.edu>

BugReports: <https://github.com/robjhyndman/hts/issues>

License: GPL (\geq 2)

Example using R


```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```

Example using R

```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))
```



Example using R

```
library(hts)
```

```
# bts is a matrix containing the bottom level time series  
# nodes describes the hierarchical structure  
y <- hts(bts, nodes=list(2, c(3,2)))  
  
# Forecast 10-step-ahead using WLS combination method  
# ETS used for each series by default  
fc <- forecast(y, h=10)
```

forecast.gts function

Usage

```
forecast(object, h,  
  method = c("comb", "bu", "mo", "tdgsf", "tdgsa", "tdfp"),  
  fmethod = c("ets", "rw", "arima"),  
  weights = c("sd", "none", "nseries"),  
  positive = FALSE,  
  parallel = FALSE, num.cores = 2, ...)
```

Arguments

object	Hierarchical time series object of class gts.
h	Forecast horizon
method	Method for distributing forecasts within the hierarchy.
fmethod	Forecasting method to use
positive	If TRUE, forecasts are forced to be strictly positive
weights	Weights used for optimal combination method. When weights = sd , it takes account of the standard deviation of forecasts.
parallel	If TRUE, allow parallel processing
num.cores	If parallel = TRUE, specify how many cores are going to be used