



**MONASH**  
University

MONASH  
BUSINESS  
SCHOOL

# Optimal Forecast Reconciliation

**Rob J Hyndman**

[robjhyndman.com](http://robjhyndman.com)

# Outline

**1 Hierarchical and grouped time series**

2 Forecast reconciliation

3 Fast computational tricks

4 Temporal hierarchies

# Labour market participation

## Australia and New Zealand Standard Classification of Occupations

- 8 major groups
  - 43 sub-major groups
    - 97 minor groups
      - 359 unit groups
        - \* 1023 occupations

### Example: statistician

- 2 Professionals
  - 22 Business, Human Resource and Marketing Professionals
    - 224 Information and Organisation Professionals
      - 2241 Actuaries, Mathematicians and Statisticians
        - 224113 Statistician

# Labour market participation


## Australia and New Zealand Standard Classification of Occupations

- 8 major groups
  - 43 sub-major groups
    - 97 minor groups
      - 359 unit groups
        - \* 1023 occupations

### Example: statistician

- 2 Professionals
  - 22 Business, Human Resource and Marketing Professionals
    - 224 Information and Organisation Professionals
      - 2241 Actuaries, Mathematicians and Statisticians
        - 224113 Statistician

# Australian tourism demand



## Australian tourism demand

- Quarterly data on visitor night from 1998:Q1 – 2013:Q4
- From: *National Visitor Survey*, based on annual interviews of 120,000 Australians aged 15+, collected by Tourism Research Australia.
- Split by 7 states, 27 zones and 76 regions (a geographical hierarchy)
- Also split by purpose of travel
  - Holiday
  - Visiting friends and relatives (VFR)
  - Business
  - Other
- 304 bottom-level series

☐ > 3%



### 3. PBS sales



# 3. PBS sales


## ATC drug classification

- A Alimentary tract and metabolism
- B Blood and blood forming organs
- C Cardiovascular system
- D Dermatologicals
- G Genito-urinary system and sex hormones
- H Systemic hormonal preparations, excluding sex hormones and insulins
- J Anti-infectives for systemic use
- L Antineoplastic and immunomodulating agents
- M Musculo-skeletal system
- N Nervous system
- P Antiparasitic products, insecticides and repellents
- R Respiratory system
- S Sensory organs
- V Various



# 3. PBS sales

## ATC drug classification



# Spectacle sales



- Monthly UK sales data from 2000 – 2014
- Provided by a large spectacle manufacturer
- Split by brand (26), gender (3), price range (6), materials (4), and stores (600)
- About 1 million bottom-level series

# Spectacle sales



- Monthly UK sales data from 2000 – 2014
- Provided by a large spectacle manufacturer
- Split by brand (26), gender (3), price range (6), materials (4), and stores (600)
- About 1 million bottom-level series

# Spectacle sales



- Monthly UK sales data from 2000 – 2014
- Provided by a large spectacle manufacturer
- Split by brand (26), gender (3), price range (6), materials (4), and stores (600)
- About 1 million bottom-level series


# Spectacle sales



- Monthly UK sales data from 2000 – 2014
- Provided by a large spectacle manufacturer
- Split by brand (26), gender (3), price range (6), materials (4), and stores (600)
- About 1 million bottom-level series

# Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




## Examples

- Labour turnover by occupation
- Pharmaceutical sales
- Tourism by state and region

# Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




## Examples

- Labour turnover by occupation
- Pharmaceutical sales
- Tourism by state and region

# Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




## Examples

- Labour turnover by occupation
- Pharmaceutical sales
- Tourism by state and region



# Hierarchical time series

A **hierarchical time series** is a collection of several time series that are linked together in a hierarchical structure.




## Examples

- Labour turnover by occupation
- Pharmaceutical sales
- **Tourism by state and region**

# Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.




## Examples

- Labour turnover by occupation and state
- Spectacle sales by brand, gender, stores, etc.
- Tourism by state and purpose of travel

# Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.




## Examples

- Labour turnover by occupation and state
- Spectacle sales by brand, gender, stores, etc.
- Tourism by state and purpose of travel

# Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.




## Examples

- Labour turnover by occupation and state
- Spectacle sales by brand, gender, stores, etc.
- Tourism by state and purpose of travel

# Grouped time series

A **grouped time series** is a collection of time series that can be grouped together in a number of non-hierarchical ways.



## Examples

- Labour turnover by occupation and state
- Spectacle sales by brand, gender, stores, etc.
- **Tourism by state and purpose of travel**

# The problem

- 1 How to forecast time series at all nodes such that the forecasts add up in the same way as the original data?
- 2 Can we exploit relationships between the series to improve the forecasts?

# The problem

- 1 How to forecast time series at all nodes such that the forecasts add up in the same way as the original data?
- 2 Can we exploit relationships between the series to improve the forecasts?

# The solution

- 1 Forecast all series at all levels of aggregation using an automatic forecasting algorithm (e.g., `ets`, `auto.arima`, ...)
- 2 Reconcile the resulting forecasts so they add up correctly using least squares optimization (i.e., find closest reconciled forecasts to the original forecasts).
- 3 This is all available in the `hts` package in R.




# The solution

- 1 Forecast all series at all levels of aggregation using an automatic forecasting algorithm (e.g., `ets`, `auto.arima`, ...)
- 2 Reconcile the resulting forecasts so they add up correctly using least squares optimization (i.e., find closest reconciled forecasts to the original forecasts).
- 3 This is all available in the `hts` package in R.

# The solution

- 1 Forecast all series at all levels of aggregation using an automatic forecasting algorithm (e.g., `ets`, `auto.arima`, ...)
- 2 Reconcile the resulting forecasts so they add up correctly using least squares optimization (i.e., find closest reconciled forecasts to the original forecasts).
- 3 This is all available in the **hts** package in R.

# Hierarchical time series




$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series  $X$  at time  $t$ .

$b_t$  : vector of all series at bottom level in time  $t$ .

# Hierarchical time series




$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series  $X$  at time  $t$ .

$b_t$  : vector of all series at bottom level in time  $t$ .

# Hierarchical time series




$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series X at time  $t$ .

$b_t$  : vector of all series at bottom level in time  $t$ .

$$\mathbf{y}_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}$$

# Hierarchical time series




$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series  $X$  at time  $t$ .

$b_t$  : vector of all series at bottom level in time  $t$ .

$$y_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_S \begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}$$

# Hierarchical time series




$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series  $X$  at time  $t$ .

$b_t$  : vector of all series at bottom level in time  $t$ .

$$y_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}}_{b_t}$$

# Hierarchical time series



$y_t$  : observed aggregate of all series at time  $t$ .

$y_{X,t}$  : observation on series X at time  $t$ .


$b_t$  : vector of all series at bottom level in time  $t$ .

$$y_t = [y_t, y_{A,t}, y_{B,t}, y_{C,t}]' = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}}_{b_t}$$

$$y_t = S b_t$$




# Hierarchical time series




$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

# Hierarchical time series



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$


# Hierarchical time series



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$


$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

# Grouped data




$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

# Grouped data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

# Grouped data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{BX,t} \\ y_{BY,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

# Hierarchical and grouped time series

Every collection of time series with aggregation constraints can be written as

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

where

- $\mathbf{y}_t$  is vector of all series at time  $t$
- $\mathbf{b}_t$  is vector of the most disaggregated series at time  $t$
- $\mathbf{S}$  is “summing matrix” containing the aggregation constraints.

# Outline

1 Hierarchical and grouped time series

**2 Forecast reconciliation**

3 Fast computational tricks

4 Temporal hierarchies



# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - **Top-down**
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Hierarchical/grouped time series

- Forecasts should be “coherent”, unbiased, minimum variance.
- Existing methods:
  - Bottom-up
  - Top-down
  - Middle-out
- How to compute forecast intervals?
- Most research is concerned about relative performance of existing methods.

# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals



# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals

# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals

# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals

# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals

# Top-down method

## Advantages

- Works well in presence of low counts.
- Single forecasting model easy to build
- Provides reliable forecasts for aggregate levels.

## Disadvantages

- Loss of information, especially individual series dynamics.
- Distribution of forecasts to lower levels can be difficult
- No prediction intervals

# Bottom-up method

## Advantages

- No loss of information.
- Better captures dynamics of individual series.

## Disadvantages

- Large number of series to be forecast.
- Constructing forecasting models is harder because of noisy data at bottom level.
- No prediction intervals

# Bottom-up method

## Advantages

- No loss of information.
- Better captures dynamics of individual series.

## Disadvantages

- Large number of series to be forecast.
- Constructing forecasting models is harder because of noisy data at bottom level.
- No prediction intervals

# Bottom-up method

## Advantages

- No loss of information.
- Better captures dynamics of individual series.

## Disadvantages

- Large number of series to be forecast.
- Constructing forecasting models is harder because of noisy data at bottom level.
- No prediction intervals



# Bottom-up method

## Advantages

- No loss of information.
- Better captures dynamics of individual series.

## Disadvantages

- Large number of series to be forecast.
- Constructing forecasting models is harder because of noisy data at bottom level.
- No prediction intervals

# Bottom-up method

## Advantages

- No loss of information.
- Better captures dynamics of individual series.

## Disadvantages

- Large number of series to be forecast.
- Constructing forecasting models is harder because of noisy data at bottom level.
- No prediction intervals

# Forecasting notation

Let  $\hat{\mathbf{y}}_n(h)$  be vector of initial  $h$ -step forecasts, made at time  $n$ , stacked in same order as  $\mathbf{y}_t$ .

(In general, they will not “add up”.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

for some matrix  $\mathbf{P}$ .

# Forecasting notation

Let  $\hat{\mathbf{y}}_n(h)$  be vector of initial  $h$ -step forecasts, made at time  $n$ , stacked in same order as  $\mathbf{y}_t$ .  
(In general, they will not “add up”.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

for some matrix  $\mathbf{P}$ .

- $\mathbf{P}$  extracts and combines base forecasts  $\hat{\mathbf{y}}_n(h)$  to get bottom-level forecasts.

# Forecasting notation

Let  $\hat{\mathbf{y}}_n(h)$  be vector of initial  $h$ -step forecasts, made at time  $n$ , stacked in same order as  $\mathbf{y}_t$ .  
(In general, they will not “add up”.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

for some matrix  $\mathbf{P}$ .

- $\mathbf{P}$  extracts and combines base forecasts  $\hat{\mathbf{y}}_n(h)$  to get bottom-level forecasts.
- $\mathbf{S}$  adds them up

# Forecasting notation

Let  $\hat{\mathbf{y}}_n(h)$  be vector of initial  $h$ -step forecasts, made at time  $n$ , stacked in same order as  $\mathbf{y}_t$ .  
(In general, they will not “add up”.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

for some matrix  $\mathbf{P}$ .

- $\mathbf{P}$  extracts and combines base forecasts  $\hat{\mathbf{y}}_n(h)$  to get bottom-level forecasts.
- $\mathbf{S}$  adds them up

# Forecasting notation

Let  $\hat{\mathbf{y}}_n(h)$  be vector of initial  $h$ -step forecasts, made at time  $n$ , stacked in same order as  $\mathbf{y}_t$ .  
(In general, they will not “add up”.)

Reconciled forecasts must be of the form:

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

for some matrix  $\mathbf{P}$ .

- $\mathbf{P}$  extracts and combines base forecasts  $\hat{\mathbf{y}}_n(h)$  to get bottom-level forecasts.
- $\mathbf{S}$  adds them up

# Bottom-up forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}] ,$$

where  $\mathbf{0}$  is null matrix and  $\mathbf{I}$  is identity matrix.

- $\mathbf{P}$  matrix extracts only bottom-level forecasts from  $\hat{\mathbf{y}}_n(h)$
- $\mathbf{S}$  adds them up to give the bottom-up forecasts.



# Bottom-up forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}] ,$$

where  $\mathbf{0}$  is null matrix and  $\mathbf{I}$  is identity matrix.

- $\mathbf{P}$  matrix extracts only bottom-level forecasts from  $\hat{\mathbf{y}}_n(h)$
- $\mathbf{S}$  adds them up to give the bottom-up forecasts.

# Bottom-up forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Bottom-up forecasts are obtained using

$$\mathbf{P} = [\mathbf{0} \mid \mathbf{I}] ,$$

where  $\mathbf{0}$  is null matrix and  $\mathbf{I}$  is identity matrix.

- $\mathbf{P}$  matrix extracts only bottom-level forecasts from  $\hat{\mathbf{y}}_n(h)$
- $\mathbf{S}$  adds them up to give the bottom-up forecasts.

# Top-down forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where  $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$  is a vector of proportions that sum to one.

- $\mathbf{P}$  distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors  $\mathbf{p}$ .

# Top-down forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where  $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$  is a vector of proportions that sum to one.

- $\mathbf{P}$  distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors  $\mathbf{p}$ .

# Top-down forecasts

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Top-down forecasts are obtained using

$$\mathbf{P} = [\mathbf{p} \mid \mathbf{0}]$$

where  $\mathbf{p} = [p_1, p_2, \dots, p_{m_K}]'$  is a vector of proportions that sum to one.

- $\mathbf{P}$  distributes forecasts of the aggregate to the lowest level series.
- Different methods of top-down forecasting lead to different proportionality vectors  $\mathbf{p}$ .

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{b}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{b}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .

Then  $\mathbb{E}[\hat{b}_n(h)] = \beta_n(h)$ .

We want the revised forecasts to be unbiased:

$$\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbb{E}[\mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h).$$

Reconciled forecasts are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{S}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecasts are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\boldsymbol{\beta}_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\boldsymbol{\beta}_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{S}\mathbf{P}\boldsymbol{\beta}_n(h) = \mathbf{S}\boldsymbol{\beta}_n(h)$ .

Reconciled forecasts are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.



# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{S}\mathbf{P}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecast are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{S}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecast are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecast are unbiased if and only if  $\mathbf{SPS} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{S}\mathbf{P}\mathbf{S}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecast are unbiased if and only if  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: bias

$$\tilde{\mathbf{y}}_n(h) = \mathbf{SP}\hat{\mathbf{y}}_n(h)$$

**Assume:** base forecasts  $\hat{\mathbf{y}}_n(h)$  are unbiased:

$$\mathbb{E}[\hat{\mathbf{y}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbb{E}[\mathbf{y}_{n+h}|\mathbf{y}_1, \dots, \mathbf{y}_n]$$

- Let  $\hat{\mathbf{b}}_n(h)$  be bottom level base forecasts with  $\beta_n(h) = \mathbb{E}[\hat{\mathbf{b}}_n(h)|\mathbf{y}_1, \dots, \mathbf{y}_n]$ .
- Then  $\mathbb{E}[\hat{\mathbf{y}}_n(h)] = \mathbf{S}\beta_n(h)$ .
- We want the revised forecasts to be unbiased:  
 $\mathbb{E}[\tilde{\mathbf{y}}_n(h)] = \mathbf{SPS}\beta_n(h) = \mathbf{S}\beta_n(h)$ .

Reconciled forecast are unbiased if and only if  $\mathbf{SPS} = \mathbf{S}$ .

True for bottom-up, but *not* top-down or middle-out.

# General properties: variance

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Let error variance of  $h$ -step base forecasts  $\hat{\mathbf{y}}_n(h)$  be

$$\Sigma_h = \text{Var}[\mathbf{y}_{n+h} - \hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

Then the error variance of the corresponding reconciled forecasts is

$$\text{Var}[\mathbf{y}_{n+h} - \tilde{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}'$$

This is a general result for all existing methods.

# General properties: variance

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

Let error variance of  $h$ -step base forecasts  $\hat{\mathbf{y}}_n(h)$  be

$$\Sigma_h = \text{Var}[\mathbf{y}_{n+h} - \hat{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$$

Then the error variance of the corresponding reconciled forecasts is

$$\text{Var}[\mathbf{y}_{n+h} - \tilde{\mathbf{y}}_n(h) \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}'$$

This is a general result for all existing methods.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

## Theorem: MinT Reconciliation

If  $\mathbf{P}$  satisfies  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ , then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}']$$

has solution  $\mathbf{P} = (\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

- Assume that  $\Sigma_h = k_h \Sigma_1$  to simplify computations.



# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

## Theorem: MinT Reconciliation

If  $\mathbf{P}$  satisfies  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ , then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}']$$

has solution  $\mathbf{P} = (\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

- Assume that  $\Sigma_h = k_h \Sigma_1$  to simplify computations.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

## Theorem: MinT Reconciliation

If  $\mathbf{P}$  satisfies  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ , then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}']$$

has solution  $\mathbf{P} = (\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

■ Assume that  $\Sigma_h = k_h\Sigma_1$  to simplify computations.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}\mathbf{P}\hat{\mathbf{y}}_n(h)$$

## Theorem: MinT Reconciliation

If  $\mathbf{P}$  satisfies  $\mathbf{S}\mathbf{P}\mathbf{S} = \mathbf{S}$ , then

$$\min_{\mathbf{P}} = \text{trace}[\mathbf{S}\mathbf{P}\Sigma_h\mathbf{P}'\mathbf{S}']$$

has solution  $\mathbf{P} = (\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{S}(\mathbf{S}'\Sigma_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

- Assume that  $\Sigma_h = k_h\Sigma_1$  to simplify computations.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kl$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well
- Still need to estimate covariance matrix to produce prediction intervals

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kI$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kI$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kI$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kI$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well.
- Still need to estimate covariance matrix to produce prediction intervals.



# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_1^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_1^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 1: OLS

- Assume  $\Sigma_1 \approx kI$ .

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\mathbf{s})^{-1}\mathbf{s}'\hat{\mathbf{y}}_n(h)$$

- Reconciliation does not depend on data
- Works surprisingly well.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 2: WLS

- Approximate  $\Sigma_1$  by its diagonal.
- Easy to estimate, and places weight where we have best forecasts.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 2: WLS

- Approximate  $\Sigma_1$  by its diagonal.
- Easy to estimate, and places weight where we have best forecasts.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 2: WLS

- Approximate  $\Sigma_1$  by its diagonal.
- Easy to estimate, and places weight where we have best forecasts.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 2: WLS

- Approximate  $\Sigma_1$  by its diagonal.
- Easy to estimate, and places weight where we have best forecasts.
- Still need to estimate covariance matrix to produce prediction intervals.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 3: GLS

- Estimate  $\Sigma_1$  using shrinkage to the diagonal.
- Allows for covariances.
- Difficult to compute for large numbers of time series.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(\textcolor{red}{h}) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(\textcolor{blue}{h})$$

Reconciled forecasts

Base forecasts

## Solution 3: GLS

- Estimate  $\Sigma_1$  using shrinkage to the diagonal.
- Allows for covariances.
- Difficult to compute for large numbers of time series.

# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$

Reconciled forecasts

Base forecasts

## Solution 3: GLS

- Estimate  $\Sigma_1$  using shrinkage to the diagonal.
- Allows for covariances.
- Difficult to compute for large numbers of time series.



# Optimal forecast reconciliation

$$\tilde{\mathbf{y}}_n(h) = \mathbf{s}(\mathbf{s}'\Sigma_h^{-1}\mathbf{s})^{-1}\mathbf{s}'\Sigma_h^{-1}\hat{\mathbf{y}}_n(h)$$


Reconciled forecasts

Base forecasts


## Solution 3: GLS

- Estimate  $\Sigma_1$  using shrinkage to the diagonal.
- Allows for covariances.
- Difficult to compute for large numbers of time series.


# Australian tourism



# Australian tourism




# Australian tourism




# Base forecasts

**Domestic tourism forecasts: Total**




# Base forecasts

**Domestic tourism forecasts: NSW**




# Base forecasts

Domestic tourism forecasts: VIC



# Base forecasts


**Domestic tourism forecasts: Nth.Coast.NSW**






# Base forecasts

Domestic tourism forecasts: Metro.QLD




# Base forecasts

**Domestic tourism forecasts: Sth.WA**




# Base forecasts

Domestic tourism forecasts: X201.Melbourne




# Base forecasts

Domestic tourism forecasts: X402.Murraylands



# Base forecasts


Domestic tourism forecasts: X809.Daly



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets

Test sets  $h = 1$






# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets

Test sets  $h = 1$






# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets

Test sets  $h = 1$






# Forecast evaluation

Training sets


Test sets  $h = 1$



# Forecast evaluation

Training sets


Test sets  $h = 2$



# Forecast evaluation

Training sets


Test sets  $h = 3$



# Forecast evaluation

Training sets


Test sets  $h = 4$



# Forecast evaluation

Training sets


Test sets  $h = 5$



# Forecast evaluation

Training sets

Test sets  $h = 6$



# Hierarchy: states, zones, regions

RMSE	Forecast horizon						Ave
	$h = 1$	$h = 2$	$h = 3$	$h = 4$	$h = 5$	$h = 6$	
Australia							
Base	1762.04	1770.29	1766.02	1818.82	1705.35	1721.17	1757.28
Bottom	1736.92	1742.69	1722.79	1752.74	1666.73	1687.43	1718.22
OLS	1747.60	1757.68	1751.77	1800.67	1686.00	1706.45	1741.69
WLS	1705.21	1715.87	1703.75	1729.56	1627.79	1661.24	1690.57
GLS	1704.64	1715.60	1705.31	1729.04	1626.36	1661.64	1690.43
States							
Base	399.77	404.16	401.92	407.26	395.38	401.17	401.61
Bottom	404.29	406.95	404.96	409.02	399.80	401.55	404.43
OLS	404.47	407.62	405.43	413.79	401.10	404.90	406.22
WLS	398.84	402.12	400.71	405.03	394.76	398.23	399.95
GLS	398.84	402.16	400.86	405.03	394.59	398.22	399.95
Regions							
Base	93.15	93.38	93.45	93.79	93.50	93.56	93.47
Bottom	93.15	93.38	93.45	93.79	93.50	93.56	93.47
OLS	93.28	93.53	93.64	94.17	93.78	93.88	93.71
WLS	93.02	93.32	93.38	93.72	93.39	93.53	93.39
GLS	92.98	93.27	93.34	93.66	93.34	93.46	93.34

# Outline

1 Hierarchical and grouped time series


2 Forecast reconciliation

**3 Fast computational tricks**

4 Temporal hierarchies




# Fast computation: hierarchical data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$


# Fast computation: hierarchical data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{B,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{C,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_S \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

# Fast computation: hierarchical data




$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{B,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{C,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

# Fast computation: hierarchies

Think of the hierarchy as a tree of trees:




Then the summing matrix contains  $k$  smaller summing matrices:

$$S = \begin{bmatrix} \mathbf{1}'_{n_1} & \mathbf{1}'_{n_2} & \cdots & \mathbf{1}'_{n_K} \\ S_1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & S_2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & S_K \end{bmatrix}$$

where  $\mathbf{1}_n$  is an  $n$ -vector of ones and tree  $T_i$  has  $n_i$  terminal nodes.

# Fast computation: hierarchies

Think of the hierarchy as a tree of trees:



Then the summing matrix contains  $k$  smaller summing matrices:

$$S = \begin{bmatrix} \mathbf{1}'_{n_1} & \mathbf{1}'_{n_2} & \cdots & \mathbf{1}'_{n_K} \\ S_1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & S_2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & S_K \end{bmatrix}$$

where  $\mathbf{1}_n$  is an  $n$ -vector of ones and tree  $T_i$  has  $n_i$  terminal nodes.

# Fast computation: hierarchies

$$S'\Lambda S = \begin{bmatrix} S'_1\Lambda_1S_1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & S'_2\Lambda_2S_2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & S'_K\Lambda_KS_K \end{bmatrix} + \lambda_0 \mathbf{J}_n$$

- $\lambda_0$  is the top left element of  $\Lambda$ ;
- $\Lambda_k$  is a block of  $\Lambda$ , corresponding to tree  $T_k$ ;
- $\mathbf{J}_n$  is a matrix of ones;
- $n = \sum_k n_k$ .

Now apply the Sherman-Morrison formula ...

# Fast computation: hierarchies

$$S'\Lambda S = \begin{bmatrix} S'_1\Lambda_1S_1 & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & S'_2\Lambda_2S_2 & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & S'_K\Lambda_KS_K \end{bmatrix} + \lambda_0 J_n$$

- $\lambda_0$  is the top left element of  $\Lambda$ ;
- $\Lambda_k$  is a block of  $\Lambda$ , corresponding to tree  $T_k$ ;
- $J_n$  is a matrix of ones;
- $n = \sum_k n_k$ .

Now apply the Sherman-Morrison formula ...

# Fast computation: hierarchies

$$(\mathbf{S}'\Lambda\mathbf{S})^{-1} = \begin{bmatrix} (\mathbf{S}'_1\Lambda_1\mathbf{S}_1)^{-1} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & (\mathbf{S}'_2\Lambda_2\mathbf{S}_2)^{-1} & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & (\mathbf{S}'_K\Lambda_K\mathbf{S}_K)^{-1} \end{bmatrix} - c\mathbf{S}_0$$

- $\mathbf{S}_0$  can be partitioned into  $K^2$  blocks, with the  $(k, \ell)$  block (of dimension  $n_k \times n_\ell$ ) being

$$(\mathbf{S}'_k\Lambda_k\mathbf{S}_k)^{-1}J_{n_k,n_\ell}(\mathbf{S}'_\ell\Lambda_\ell\mathbf{S}_\ell)^{-1}$$

- $J_{n_k,n_\ell}$  is a  $n_k \times n_\ell$  matrix of ones.
- $c^{-1} = \lambda_0^{-1} + \sum_k \mathbf{1}'_{n_k} (\mathbf{S}'_k\Lambda_k\mathbf{S}_k)^{-1} \mathbf{1}_{n_k}$ .
- Each  $\mathbf{S}'_k\Lambda_k\mathbf{S}_k$  can be inverted similarly.
- $\mathbf{S}'\Lambda\mathbf{y}$  can also be computed recursively.



# Fast computation: hierarchies

$$(S'\Lambda S)^{-1} = \begin{bmatrix} (S'_1\Lambda_1S_1)^{-1} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & (S'_2\Lambda_2S_2)^{-1} & \cdots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \cdots & (S'_K\Lambda_KS_K)^{-1} \end{bmatrix} - cS_0$$

- $S_0$  can be partitioned into  $K^2$  blocks, with the  $(k, \ell)$  block (of dimension  $n_k \times n_\ell$ ) being


The recursive calculations can be done in such a way that we never store any of the large matrices involved.

- $J_{n_k, n_\ell}$
- $c^{-1} = c_0^{-1} - \sum_k \frac{c_k^{-1} (c_k^{-1} \Lambda_k^{-1} c_k^{-1})}{c_k^{-1}}$



- Each  $S'_k\Lambda_kS_k$  can be inverted similarly.
- $S'\Lambda y$  can also be computed recursively.

# Fast computation: grouped data



$$\mathbf{y}_t = \begin{pmatrix} y_t \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \\ y_{X,t} \\ y_{Y,t} \\ y_{Z,t} \\ y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix} \underbrace{\begin{pmatrix} y_{AX,t} \\ y_{AY,t} \\ y_{AZ,t} \\ y_{BX,t} \\ y_{BY,t} \\ y_{BZ,t} \\ y_{CX,t} \\ y_{CY,t} \\ y_{CZ,t} \end{pmatrix}}_{\mathbf{b}_t}$$

$$\mathbf{y}_t = \mathbf{S} \mathbf{b}_t$$

# Fast computation: grouped data

$$\mathbf{S} = \begin{bmatrix} \mathbf{1}'_m \otimes \mathbf{1}'_n \\ \mathbf{1}'_m \otimes \mathbf{I}_n \\ \mathbf{I}_m \otimes \mathbf{1}'_n \\ \mathbf{I}_m \otimes \mathbf{I}_n \end{bmatrix}$$

$m$  = number of rows

$n$  = number of columns

$$\mathbf{S}'\mathbf{\Lambda}\mathbf{S} = \lambda_{00} \mathbf{J}_{mn} + (\mathbf{\Lambda}_R \otimes \mathbf{J}_n) + (\mathbf{J}_m \otimes \mathbf{\Lambda}_C) + \mathbf{\Lambda}_U$$

- $\mathbf{\Lambda}_R$ ,  $\mathbf{\Lambda}_C$  and  $\mathbf{\Lambda}_U$  are diagonal matrices corresponding to rows, columns and unaggregated series;
- $\lambda_{00}$  corresponds to aggregate.

# Fast computation: grouped data

$$\mathbf{S} = \begin{bmatrix} \mathbf{1}'_m \otimes \mathbf{1}'_n \\ \mathbf{1}'_m \otimes \mathbf{I}_n \\ \mathbf{I}_m \otimes \mathbf{1}'_n \\ \mathbf{I}_m \otimes \mathbf{I}_n \end{bmatrix}$$

$m$  = number of rows

$n$  = number of columns

$$\mathbf{S}'\mathbf{\Lambda}\mathbf{S} = \lambda_{00} \mathbf{J}_{mn} + (\mathbf{\Lambda}_R \otimes \mathbf{J}_n) + (\mathbf{J}_m \otimes \mathbf{\Lambda}_C) + \mathbf{\Lambda}_U$$

- $\mathbf{\Lambda}_R$ ,  $\mathbf{\Lambda}_C$  and  $\mathbf{\Lambda}_U$  are diagonal matrices corresponding to rows, columns and unaggregated series;
- $\lambda_{00}$  corresponds to aggregate.

# Fast computation: grouped data

$$(\mathbf{S}\mathbf{\Lambda}\mathbf{S})^{-1} = \mathbf{A} - \frac{\mathbf{A}\mathbf{1}_{mn}\mathbf{1}_{mn}'\mathbf{A}}{1/\lambda_{00} + \mathbf{1}_{mn}'\mathbf{A}\mathbf{1}_{mn}}$$

$$\mathbf{A} = \mathbf{\Lambda}_U^{-1} - \mathbf{\Lambda}_U^{-1}(\mathbf{J}_m \otimes \mathbf{D})\mathbf{\Lambda}_U^{-1} - \mathbf{E}\mathbf{M}^{-1}\mathbf{E}'.$$

$\mathbf{D}$  is diagonal with elements  $d_j = \lambda_{0j}/(1 + \lambda_{0j} \sum_i \lambda_{ij}^{-1})$ .

$\mathbf{E}$  has  $m \times m$  blocks where  $\mathbf{e}_{ij}$  has  $k$ th element

$$(\mathbf{e}_{ij})_k = \begin{cases} \lambda_{i0}^{1/2} \lambda_{ik}^{-1} - \lambda_{i0}^{1/2} \lambda_{ik}^{-2} d_k, & i = j, \\ -\lambda_{j0}^{1/2} \lambda_{ik}^{-1} \lambda_{jk}^{-1} d_k, & i \neq j. \end{cases}$$

$\mathbf{M}$  is  $m \times m$  with  $(i, j)$  element

$$(\mathbf{M})_{ij} = \begin{cases} 1 + \lambda_{i0} \sum_k \lambda_{ik}^{-1} - \lambda_{i0} \sum_k \lambda_{ik}^{-2} d_k, & i = j, \\ -\lambda_{i0}^{1/2} \lambda_{j0}^{1/2} \sum_k \lambda_{ik}^{-1} \lambda_{jk}^{-1} d_k, & i \neq j. \end{cases}$$

# Fast computation: grouped data

$$(\mathbf{S}\mathbf{\Lambda}\mathbf{S})^{-1} = \mathbf{A} - \frac{\mathbf{A}\mathbf{1}_{mn}\mathbf{1}'_{mn}\mathbf{A}}{1/\lambda_{00} + \mathbf{1}'_{mn}\mathbf{A}\mathbf{1}_{mn}}$$

$$\mathbf{A} = \mathbf{\Lambda}_U^{-1} - \mathbf{\Lambda}_U^{-1}(\mathbf{J}_m \otimes \mathbf{D})\mathbf{\Lambda}_U^{-1} - \mathbf{E}\mathbf{M}^{-1}\mathbf{E}'.$$

$\mathbf{D}$  is diagonal with elements  $d_j = \lambda_{0j}/(1 + \lambda_{0j} \sum_i \lambda_{ij}^{-1})$ .

$\mathbf{E}$  has  $m \times m$  blocks where  $\mathbf{e}_{ij}$  has  $k$ th element

$$(-\lambda_{i0}^{-1/2} + \lambda_{i0}^{-1/2} \sum_k \lambda_{ik}^{-1} \lambda_{jk}^{-1} d_k)$$

Again, the calculations can be done in such a way that we never store any of the large matrices involved.

$\mathbf{M}$  is  $m \times$

$$(\mathbf{M})_{ij} = \begin{cases} 1 + \lambda_{i0} \sum_k \lambda_{ik}^{-1} - \lambda_{i0} \sum_k \lambda_{ik}^{-2} d_k, & i = j, \\ -\lambda_{i0}^{1/2} \lambda_{j0}^{1/2} \sum_k \lambda_{ik}^{-1} \lambda_{jk}^{-1} d_k, & i \neq j. \end{cases}$$



# Outline


1 Hierarchical and grouped time series

2 Forecast reconciliation

3 Fast computational tricks

**4 Temporal hierarchies**

# Temporal hierarchies




## Basic idea:

- ➡ Forecast series at each available frequency.
- ➡ Optimally reconcile forecasts within the same year.




# Temporal hierarchies



## Basic idea:

- ➡ Forecast series at each available frequency.
- ➡ Optimally reconcile forecasts within the same year.

# Monthly series




■  $k = 2, 4, 12$  nodes

■  $k = 3, 6, 12$  nodes


■ Why not  $k = 2, 3, 4, 6, 12$  nodes?

# Monthly series



- $k = 2, 4, 12$  nodes
- $k = 3, 6, 12$  nodes
- Why not  $k = 2, 3, 4, 6, 12$  nodes?

# Monthly series



- $k = 2, 4, 12$  nodes
- $k = 3, 6, 12$  nodes
- Why not  $k = 2, 3, 4, 6, 12$  nodes?

# Monthly data

$$\underbrace{\begin{pmatrix} A \\ \text{Semi}A_1 \\ \text{Semi}A_2 \\ \text{Four}M_1 \\ \text{Four}M_2 \\ \text{Four}M_3 \\ Q_1 \\ \vdots \\ Q_4 \\ \text{Bi}M_1 \\ \vdots \\ \text{Bi}M_6 \\ M_1 \\ \vdots \\ M_{12} \end{pmatrix}}_{(28 \times 1)} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & \vdots & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ & & & & & & \vdots & & & & & \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 \end{pmatrix}}_{\substack{I_{12} \\ S}} \underbrace{\begin{pmatrix} M_1 \\ M_2 \\ M_3 \\ M_4 \\ M_5 \\ M_6 \\ M_7 \\ M_8 \\ M_9 \\ M_{10} \\ M_{11} \\ M_{12} \end{pmatrix}}_{B_t}$$

# In general

For a time series  $y_1, \dots, y_T$ , observed at frequency  $m$ , we generate aggregate series

$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$ .
- A single unique hierarchy is only possible when there are no coprime pairs in  $F(m)$ .
- $M_k = m/k$  is seasonal period of aggregated series.

# In general

For a time series  $y_1, \dots, y_T$ , observed at frequency  $m$ , we generate aggregate series

$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$ .
- A single unique hierarchy is only possible when there are no coprime pairs in  $F(m)$ .
- $M_k = m/k$  is seasonal period of aggregated series.

# In general

For a time series  $y_1, \dots, y_T$ , observed at frequency  $m$ , we generate aggregate series


$$y_j^{[k]} = \sum_{t=1+(j-1)k}^{jk} y_t, \quad \text{for } j = 1, \dots, \lfloor T/k \rfloor$$

- $k \in F(m) = \{\text{factors of } m\}$ .
- A single unique hierarchy is only possible when there are no coprime pairs in  $F(m)$ .
- $M_k = m/k$  is seasonal period of aggregated series.




# UK Accidents and Emergency Demand


Annual ( $k=52$ )




Semi-annual ( $k=26$ )




Quarterly ( $k=13$ )




Monthly ( $k=4$ )



Bi-weekly ( $k=2$ )



Weekly ( $k=1$ )



--- base

— reconciled

# UK Accidents and Emergency Demand

- 1 Type 1 Departments — Major A&E
- 2 Type 2 Departments — Single Specialty
- 3 Type 3 Departments — Other A&E/Minor Injury
- 4 Total Attendances
- 5 Type 1 Departments — Major A&E  $> 4$  hrs
- 6 Type 2 Departments — Single Specialty  $> 4$  hrs
- 7 Type 3 Departments — Other A&E/Minor Injury  $> 4$  hrs
- 8 Total Attendances  $> 4$  hrs
- 9 Emergency Admissions via Type 1 A&E
- 10 Total Emergency Admissions via A&E
- 11 Other Emergency Admissions (i.e., not via A&E)
- 12 Total Emergency Admissions
- 13 Number of patients spending  $> 4$  hrs from decision to admission

# UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	<i>h</i>	Base	Reconciled	Change
Weekly	1	1.6	1.3	−17.2%
Weekly	4	1.9	1.5	−18.6%
Weekly	13	2.3	1.9	−16.2%
Weekly	1–52	2.0	1.9	−5.0%
Annual	1	3.4	1.9	−42.9%

# UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- **Base forecasts using `auto.arima()`.**
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	$h$	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

# UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- **Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.**

Aggr. Level	$h$	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

# UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	$h$	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

# UK Accidents and Emergency Demand

- **Minimum training set:** all data except the last year
- Base forecasts using `auto.arima()`.
- Mean Absolute Scaled Errors for 1, 4 and 13 weeks ahead using a rolling origin.

Aggr. Level	$h$	Base	Reconciled	Change
Weekly	1	1.6	1.3	-17.2%
Weekly	4	1.9	1.5	-18.6%
Weekly	13	2.3	1.9	-16.2%
Weekly	1-52	2.0	1.9	-5.0%
Annual	1	3.4	1.9	-42.9%

# References



RJ Hyndman, RA Ahmed, G Athanasopoulos and HL Shang (2011). Optimal combination forecasts for hierarchical time series. *Computational Statistics & Data Analysis* 55(9), 2579–2589.



RJ Hyndman, A Lee and E Wang (2016). Fast computation of reconciled forecasts for hierarchical and grouped time series. *Computational Statistics & Data Analysis* 97, 16–32



SL Wickramasuriya, G Athanasopoulos and RJ Hyndman (2015). *Forecasting hierarchical and grouped time series through trace minimization*. Working paper. Dept Econometrics & Business Statistics, Monash University



G Athanasopoulos, RJ Hyndman, N Kourentzes and F Petropoulos (2017). Forecasting with temporal hierarchies. *European Journal of Operational Research* 262(1), 60–74



RJ Hyndman and G Athanasopoulos (2018). *Forecasting: principles and practice*. 2nd ed. Melbourne, Australia: OTexts. [OTexts.org/fpp2/](http://OTexts.org/fpp2/).



# R packages



<https://github.com/earowang/tsibble>



<http://pkg.earo.me/sugrrants>



<http://pkg.robjhyndman.com/forecast>



<http://pkg.earo.me/hts>



<http://pkg.robjhyndman.com/thief>