




Rob J Hyndman


Coherent mortality forecasting using functional time series models



Mortality rates


Mortality rates

Australia: male death rates (1970)




Mortality rates

Australia: male death rates (1990)




Mortality rates

Australia: male death rates (1921–2009)




Mortality rates

Australia: male death rates (1921–2009)



Mortality rates

Australia: mortality sex ratio (1921–2009)



Outline

- 1 Functional forecasting
- 2 Forecasting groups
- 3 Coherent cohort life expectancy forecasts
- 4 Conclusions

Outline

- 1 Functional forecasting
- 2 Forecasting groups
- 3 Coherent cohort life expectancy forecasts
- 4 Conclusions

Some notation

Let $y_{t,x}$ be the observed (smoothed) data in period t at age x , $t = 1, \dots, n$.

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- Estimate $f_t(x)$ using penalized regression splines.
- Estimate $\mu(x)$ as mean $f_t(x)$ across years.
- Estimate $\beta_{t,k}$ and $\phi_k(x)$ using functional principal components.

Some notation

Let $y_{t,x}$ be the observed (smoothed) data in period t at age x , $t = 1, \dots, n$.

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- Estimate $f_t(x)$ using penalized regression splines.
- Estimate $\mu(x)$ as mean $f_t(x)$ across years.
- Estimate $\beta_{t,k}$ and $\phi_k(x)$ using functional principal components.
- $\varepsilon_{t,x} \stackrel{\text{iid}}{\sim} N(0, 1)$ and $e_t(x) \stackrel{\text{iid}}{\sim} N(0, v(x))$.

Some notation

Let $y_{t,x}$ be the observed (smoothed) data in period t at age x , $t = 1, \dots, n$.

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- Estimate $f_t(x)$ using penalized regression splines.
- Estimate $\mu(x)$ as mean $f_t(x)$ across years.
- Estimate $\beta_{t,k}$ and $\phi_k(x)$ using functional principal components.
- $\varepsilon_{t,x} \stackrel{\text{iid}}{\sim} N(0, 1)$ and $e_t(x) \stackrel{\text{iid}}{\sim} N(0, v(x))$.

Some notation

Let $y_{t,x}$ be the observed (smoothed) data in period t at age x , $t = 1, \dots, n$.

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- Estimate $f_t(x)$ using penalized regression splines.
- Estimate $\mu(x)$ as mean $f_t(x)$ across years.
- Estimate $\beta_{t,k}$ and $\phi_k(x)$ using functional principal components.
- $\varepsilon_{t,x} \stackrel{\text{iid}}{\sim} N(0, 1)$ and $e_t(x) \stackrel{\text{iid}}{\sim} N(0, v(x))$.

Some notation


Let $y_{t,x}$ be the observed (smoothed) data in period t at age x , $t = 1, \dots, n$.

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$


$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- Estimate $f_t(x)$ using penalized regression splines.
- Estimate $\mu(x)$ as mean $f_t(x)$ across years.
- Estimate $\beta_{t,k}$ and $\phi_k(x)$ using functional principal components.
- $\varepsilon_{t,x} \stackrel{\text{iid}}{\sim} N(0, 1)$ and $e_t(x) \stackrel{\text{iid}}{\sim} N(0, v(x))$.

Australian male mortality model



Australian male mortality model



Functional time series model

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- The eigenfunctions $\phi_k(x)$ show the main regions of variation.
- The scores $\{\beta_{t,k}\}$ are uncorrelated by construction. So we can forecast each $\beta_{t,k}$ using a univariate time series model.
- Univariate ARIMA models can be used for forecasting.

Functional time series model

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- The eigenfunctions $\phi_k(x)$ show the main regions of variation.
- The scores $\{\beta_{t,k}\}$ are uncorrelated by construction. So we can forecast each $\beta_{t,k}$ using a univariate time series model.
- Univariate ARIMA models can be used for forecasting.

Functional time series model

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- The eigenfunctions $\phi_k(x)$ show the main regions of variation.
- The scores $\{\beta_{t,k}\}$ are uncorrelated by construction. So we can forecast each $\beta_{t,k}$ using a univariate time series model.
- Univariate ARIMA models can be used for forecasting.

Functional time series model

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- The eigenfunctions $\phi_k(x)$ show the main regions of variation.
- The scores $\{\beta_{t,k}\}$ are uncorrelated by construction. So we can forecast each $\beta_{t,k}$ using a univariate time series model.
- **Univariate ARIMA models can be used for forecasting.**

Forecasts

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

Forecasts

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$


$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$E[y_{n+h,x} | \mathbf{y}] = \hat{\mu}(x) + \sum_{k=1}^K \hat{\beta}_{n+h,k} \hat{\phi}_k(x)$$

$$\text{Var}[y_{n+h,x} | \mathbf{y}] = \hat{\sigma}_{\mu}^2(x) + \sum_{k=1}^K v_{n+h,k} \hat{\phi}_k^2(x) + \sigma_t^2(x) + v(x)$$


where $v_{n+h,k} = \text{Var}(\beta_{n+h,k} | \beta_{1,k}, \dots, \beta_{n,k})$
and $\mathbf{y} = [y_{1,x}, \dots, y_{n,x}]$.

Forecasting the PC scores




Forecasts of $f_t(x)$

Australia: male death rates (1921–2009)




Forecasts of $f_t(x)$

Australia: male death rates (1921–2009)




Forecasts of $f_t(x)$

Australia: male death rates forecasts (2010–2059)




Forecasts of $f_t(x)$

Australia: male death rates forecasts (2010 and 2059)




Forecasts of mortality sex ratio

Australia: mortality sex ratio data




Forecasts of mortality sex ratio

Australia: mortality sex ratio data




Forecasts of mortality sex ratio

Australia: mortality sex ratio forecasts



Forecasts of mortality sex ratio

Australia: mortality sex ratio forecasts



Outline

- 1 Functional forecasting
- 2 Forecasting groups**
- 3 Coherent cohort life expectancy forecasts
- 4 Conclusions

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

- Groups may be males and females.
- Groups may be states within a country.
- Expected that groups will behave similarly.
- Coherent forecasts do not diverge over time.
- Existing functional models do not impose coherence.

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

- Groups may be males and females.
- Groups may be states within a country.
- Expected that groups will behave similarly.
- Coherent forecasts do not diverge over time.
- Existing functional models do not impose coherence.

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

- Groups may be males and females.
- Groups may be states within a country.
- Expected that groups will behave similarly.
- Coherent forecasts do not diverge over time.
- Existing functional models do not impose coherence.

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

- Groups may be males and females.
- Groups may be states within a country.
- Expected that groups will behave similarly.
- **Coherent forecasts do not diverge over time.**
- Existing functional models do not impose coherence.

The problem

Let $f_{t,j}(x)$ be the smoothed mortality rate for age x in group j in year t .

- Groups may be males and females.
- Groups may be states within a country.
- Expected that groups will behave similarly.
- **Coherent** forecasts do not diverge over time.
- Existing functional models do not impose coherence.

Forecasting the coefficients

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- We use ARIMA models for each coefficient $\{\beta_{1,j,k}, \dots, \beta_{n,j,k}\}$.
- The ARIMA models are non-stationary for the first few coefficients ($k = 1, 2$)
- Non-stationary ARIMA forecasts will diverge. Hence the mortality forecasts are not coherent.

Forecasting the coefficients

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$

$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- We use ARIMA models for each coefficient $\{\beta_{1,j,k}, \dots, \beta_{n,j,k}\}$.
- The ARIMA models are non-stationary for the first few coefficients ($k = 1, 2$)
- Non-stationary ARIMA forecasts will diverge. Hence the mortality forecasts are not coherent.


Forecasting the coefficients

$$y_{t,x} = f_t(x) + \sigma_t(x)\varepsilon_{t,x}$$


$$f_t(x) = \mu(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

- We use ARIMA models for each coefficient $\{\beta_{1,j,k}, \dots, \beta_{n,j,k}\}$.
- The ARIMA models are non-stationary for the first few coefficients ($k = 1, 2$)
- Non-stationary ARIMA forecasts will diverge. Hence the mortality forecasts are not coherent.

Male fts model




Female fts model




Australian mortality forecasts

(a) Males



(b) Females



Mortality product and ratios

Key idea

Model the geometric mean and the mortality ratio instead of the individual rates for each sex separately.

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

- Product and ratio are approximately independent

Both should be stationary (for coherence) but product is better than ratio.

Mortality product and ratios

Key idea

Model the geometric mean and the mortality ratio instead of the individual rates for each sex separately.

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

- Product and ratio are approximately independent
- Ratio should be stationary (for coherence) but product can be non-stationary.

Mortality product and ratios

Key idea

Model the geometric mean and the mortality ratio instead of the individual rates for each sex separately.

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

- Product and ratio are approximately independent
- Ratio should be stationary (for coherence) but product can be non-stationary.

Mortality product and ratios

Key idea


Model the geometric mean and the mortality ratio instead of the individual rates for each sex separately.

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

- Product and ratio are approximately independent
- Ratio should be stationary (for coherence) but product can be non-stationary.


Product data

Australia: product data



Ratio data

Australia: mortality sex ratio (1921–2009)



Model product and ratios

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.
- Forecasts: $f_{n+h|n,M}(x) = p_{n+h|n}(x)r_{n+h|n}(x)$
 $f_{n+h|n,F}(x) = p_{n+h|n}(x)/r_{n+h|n}(x).$

Model product and ratios

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.
- Forecasts: $f_{n+h|n,M}(x) = p_{n+h|n}(x)r_{n+h|n}(x)$
 $f_{n+h|n,F}(x) = p_{n+h|n}(x)/r_{n+h|n}(x).$

Model product and ratios

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.
- Forecasts: $f_{n+h|n,M}(x) = p_{n+h|n}(x)r_{n+h|n}(x)$
 $f_{n+h|n,F}(x) = p_{n+h|n}(x)/r_{n+h|n}(x).$

Model product and ratios


$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$


$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.
- **Forecasts:** $f_{n+h|n,M}(x) = p_{n+h|n}(x)r_{n+h|n}(x)$
 $f_{n+h|n,F}(x) = p_{n+h|n}(x)/r_{n+h|n}(x).$


Product model




Ratio model



Product forecasts




Ratio forecasts




Coherent forecasts

(a) Males




(b) Females




Ratio forecasts

Independent forecasts




Coherent forecasts




Life expectancy forecasts

Life expectancy forecasts



Life expectancy difference: F-M



Coherent forecasts for J groups

$$p_t(x) = [f_{t,1}(x)f_{t,2}(x)\cdots f_{t,J}(x)]^{1/J}$$

and

$$r_{t,j}(x) = f_{t,j}(x)/p_t(x),$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_{t,j}(x)] = \mu_{r,j}(x) + \sum_{l=1}^L \gamma_{t,l,j} \psi_{l,j}(x) + w_{t,j}(x).$$

- $p_t(x)$ and all $r_{t,j}(x)$ are approximately independent.
- $p_t(x)$ satisfies policy constraint: $p_t(x)/p_0(x) = \pi_t(x) = 1$
- $\sum_{j=1}^J r_{t,j}(x) = 1$ for all x

Coherent forecasts for J groups

$$p_t(x) = [f_{t,1}(x)f_{t,2}(x)\cdots f_{t,J}(x)]^{1/J}$$

and

$$r_{t,j}(x) = f_{t,j}(x)/p_t(x),$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_{t,j}(x)] = \mu_{r,j}(x) + \sum_{l=1}^L \gamma_{t,l,j} \psi_{l,j}(x) + w_{t,j}(x).$$

■ $p_t(x)$ and all $r_{t,j}(x)$ are approximately independent.

■ Ratios satisfy constraint $r_{t,1}(x)r_{t,2}(x)\cdots r_{t,J}(x) = 1$.

$$\log[f_{t,j}(x)] = \log[p_t(x)r_{t,j}(x)]$$

Coherent forecasts for J groups

$$p_t(x) = [f_{t,1}(x)f_{t,2}(x)\cdots f_{t,J}(x)]^{1/J}$$

and

$$r_{t,j}(x) = f_{t,j}(x)/p_t(x),$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_{t,j}(x)] = \mu_{r,j}(x) + \sum_{l=1}^L \gamma_{t,l,j} \psi_{l,j}(x) + w_{t,j}(x).$$

- $p_t(x)$ and all $r_{t,j}(x)$ are approximately independent.

- Ratios satisfy constraint $r_{t,1}(x)r_{t,2}(x)\cdots r_{t,J}(x) = 1$.
- $\log[f_{t,j}(x)] = \log[p_t(x)r_{t,j}(x)]$

Coherent forecasts for J groups

$$p_t(x) = [f_{t,1}(x)f_{t,2}(x)\cdots f_{t,J}(x)]^{1/J}$$

and

$$r_{t,j}(x) = f_{t,j}(x)/p_t(x),$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_{t,j}(x)] = \mu_{r,j}(x) + \sum_{l=1}^L \gamma_{t,l,j} \psi_{l,j}(x) + w_{t,j}(x).$$

- $p_t(x)$ and all $r_{t,j}(x)$ are approximately independent.

- Ratios satisfy constraint $r_{t,1}(x)r_{t,2}(x)\cdots r_{t,J}(x) = 1$.

- $\log[f_{t,j}(x)] = \log[p_t(x)r_{t,j}(x)]$

Coherent forecasts for J groups

$$p_t(x) = [f_{t,1}(x)f_{t,2}(x)\cdots f_{t,J}(x)]^{1/J}$$

and

$$r_{t,j}(x) = f_{t,j}(x)/p_t(x),$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_{t,j}(x)] = \mu_{r,j}(x) + \sum_{l=1}^L \gamma_{t,l,j} \psi_{l,j}(x) + w_{t,j}(x).$$

- $p_t(x)$ and all $r_{t,j}(x)$ are approximately independent.
- Ratios satisfy constraint $r_{t,1}(x)r_{t,2}(x)\cdots r_{t,J}(x) = 1$.
- $\log[f_{t,j}(x)] = \log[p_t(x)r_{t,j}(x)]$

Coherent forecasts for J groups

$$\begin{aligned}\log[f_{t,j}(x)] &= \log[p_t(x)r_{t,j}(x)] = \log[p_t(x)] + \log[r_{t,j}] \\ &= \mu_j(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + \sum_{\ell=1}^L \gamma_{t,\ell,j} \psi_{\ell,j}(x) + z_{t,j}(x)\end{aligned}$$

- $\mu_j(x) = \mu_p(x) + \mu_{r,j}(x)$ is group mean
- $z_{t,j}(x) = e_t(x) + w_{t,j}(x)$ is error term.
- $\{\gamma_{t,\ell,j}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restriction on $\beta_{t,k}$

Coherent forecasts for J groups

$$\begin{aligned}\log[f_{t,j}(x)] &= \log[p_t(x)r_{t,j}(x)] = \log[p_t(x)] + \log[r_{t,j}] \\ &= \mu_j(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + \sum_{\ell=1}^L \gamma_{t,\ell,j} \psi_{\ell,j}(x) + z_{t,j}(x)\end{aligned}$$

- $\mu_j(x) = \mu_p(x) + \mu_{r,j}(x)$ is group mean
- $z_{t,j}(x) = e_t(x) + w_{t,j}(x)$ is error term.
- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.

Coherent forecasts for J groups

$$\begin{aligned}\log[f_{t,j}(x)] &= \log[p_t(x)r_{t,j}(x)] = \log[p_t(x)] + \log[r_{t,j}] \\ &= \mu_j(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + \sum_{\ell=1}^L \gamma_{t,\ell,j} \psi_{\ell,j}(x) + z_{t,j}(x)\end{aligned}$$

- $\mu_j(x) = \mu_p(x) + \mu_{r,j}(x)$ is group mean
- $z_{t,j}(x) = e_t(x) + w_{t,j}(x)$ is error term.
- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.

Coherent forecasts for J groups

$$\begin{aligned}\log[f_{t,j}(x)] &= \log[p_t(x)r_{t,j}(x)] = \log[p_t(x)] + \log[r_{t,j}] \\ &= \mu_j(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + \sum_{\ell=1}^L \gamma_{t,\ell,j} \psi_{\ell,j}(x) + z_{t,j}(x)\end{aligned}$$

- $\mu_j(x) = \mu_p(x) + \mu_{r,j}(x)$ is group mean
- $z_{t,j}(x) = e_t(x) + w_{t,j}(x)$ is error term.
- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.

Coherent forecasts for J groups

$$\begin{aligned}\log[f_{t,j}(x)] &= \log[p_t(x)r_{t,j}(x)] = \log[p_t(x)] + \log[r_{t,j}] \\ &= \mu_j(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + \sum_{\ell=1}^L \gamma_{t,\ell,j} \psi_{\ell,j}(x) + z_{t,j}(x)\end{aligned}$$

- $\mu_j(x) = \mu_p(x) + \mu_{r,j}(x)$ is group mean
- $z_{t,j}(x) = e_t(x) + w_{t,j}(x)$ is error term.
- $\{\gamma_{t,\ell}\}$ restricted to be stationary processes: either ARMA(p, q) or ARFIMA(p, d, q).
- No restrictions for $\beta_{t,1}, \dots, \beta_{t,K}$.

Li-Lee method

Li & Lee (*Demography*, 2005) method is a special case of our approach.

$$f_{t,j}(x) = \mu_j(x) + \beta_t \phi(x) + \gamma_{t,j} \psi_j(x) + e_{t,j}(x)$$

where f is *unsmoothed* log mortality rate, β_t is a random walk with drift and $\gamma_{t,j}$ is AR(1) process.

- **No smoothing.**

- Only one basis function for each part,
- Random walk with drift very limiting.
- AR(1) very limiting.
- The $\gamma_{t,j}$ coefficients will be highly correlated with each other, and so independent models are not appropriate.

Li-Lee method

Li & Lee (*Demography*, 2005) method is a special case of our approach.

$$f_{t,j}(x) = \mu_j(x) + \beta_t \phi(x) + \gamma_{t,j} \psi_j(x) + e_{t,j}(x)$$

where f is *unsmoothed* log mortality rate, β_t is a random walk with drift and $\gamma_{t,j}$ is AR(1) process.

- No smoothing.
- Only one basis function for each part,
- Random walk with drift very limiting.
- AR(1) very limiting.
- The $\gamma_{t,j}$ coefficients will be highly correlated with each other, and so independent models are not appropriate.

Li-Lee method

Li & Lee (*Demography*, 2005) method is a special case of our approach.

$$f_{t,j}(x) = \mu_j(x) + \beta_t \phi(x) + \gamma_{t,j} \psi_j(x) + e_{t,j}(x)$$

where f is *unsmoothed* log mortality rate, β_t is a random walk with drift and $\gamma_{t,j}$ is AR(1) process.

- No smoothing.
- Only one basis function for each part,
- Random walk with drift very limiting.
- AR(1) very limiting.
- The $\gamma_{t,j}$ coefficients will be highly correlated with each other, and so independent models are not appropriate.

Li-Lee method

Li & Lee (*Demography*, 2005) method is a special case of our approach.

$$f_{t,j}(x) = \mu_j(x) + \beta_t \phi(x) + \gamma_{t,j} \psi_j(x) + e_{t,j}(x)$$

where f is *unsmoothed* log mortality rate, β_t is a random walk with drift and $\gamma_{t,j}$ is AR(1) process.

- No smoothing.
- Only one basis function for each part,
- Random walk with drift very limiting.
- **AR(1) very limiting.**
- The $\gamma_{t,j}$ coefficients will be highly correlated with each other, and so independent models are not appropriate.

Li-Lee method

Li & Lee (*Demography*, 2005) method is a special case of our approach.

$$f_{t,j}(x) = \mu_j(x) + \beta_t \phi(x) + \gamma_{t,j} \psi_j(x) + e_{t,j}(x)$$

where f is *unsmoothed* log mortality rate, β_t is a random walk with drift and $\gamma_{t,j}$ is AR(1) process.

- No smoothing.
- Only one basis function for each part,
- Random walk with drift very limiting.
- AR(1) very limiting.
- The $\gamma_{t,j}$ coefficients will be highly correlated with each other, and so independent models are not appropriate.

Outline

- 1 Functional forecasting
- 2 Forecasting groups
- 3 Coherent cohort life expectancy forecasts**
- 4 Conclusions

Life expectancy calculation

Using standard life table calculations:

$$\begin{aligned}\text{For } x = 0, 1, \dots, \omega - 1: \quad & q_x = m_x / (1 + (1 - a_x)m_x) \\ & \ell_{x+1} = \ell_x(1 - q_x) \\ & L_x = \ell_x[1 - q_x(1 - a_x)] \\ & T_x = L_x + L_{x+1} + \dots + L_{\omega-1} + L_{\omega+} \\ & e_x = T_x / L_x\end{aligned}$$

where $a_x = 0.5$ for $x \geq 1$ and a_0 taken from Coale et al (1983).
 $q_{\omega+} = 1$, $L_{\omega+} = \ell_{\omega}/m_{\omega}$, and $T_{\omega+} = L_{\omega+}$.

■ Period life expectancy: let $m_x = m_{x,t}$ for some year t .

■ Cohort life expectancy: let $m_x = m_{x,t+x}$ for birth cohort in year t .

Life expectancy calculation

Using standard life table calculations:

$$\begin{aligned}\text{For } x = 0, 1, \dots, \omega - 1: \quad & q_x = m_x / (1 + (1 - a_x)m_x) \\ & \ell_{x+1} = \ell_x(1 - q_x) \\ & L_x = \ell_x[1 - q_x(1 - a_x)] \\ & T_x = L_x + L_{x+1} + \dots + L_{\omega-1} + L_{\omega+} \\ & e_x = T_x / L_x\end{aligned}$$

where $a_x = 0.5$ for $x \geq 1$ and a_0 taken from Coale et al (1983).
 $q_{\omega+} = 1$, $L_{\omega+} = \ell_{\omega}/m_{\omega}$, and $T_{\omega+} = L_{\omega+}$.

- Period life expectancy: let $m_x = m_{x,t}$ for some year t .
- Cohort life expectancy: let $m_x = m_{x,t+x}$ for birth cohort in year t .

Life expectancy calculation

Using standard life table calculations:

$$\begin{aligned}\text{For } x = 0, 1, \dots, \omega - 1: \quad & q_x = m_x / (1 + (1 - a_x)m_x) \\ & \ell_{x+1} = \ell_x(1 - q_x) \\ & L_x = \ell_x[1 - q_x(1 - a_x)] \\ & T_x = L_x + L_{x+1} + \dots + L_{\omega-1} + L_{\omega+} \\ & e_x = T_x / L_x\end{aligned}$$

where $a_x = 0.5$ for $x \geq 1$ and a_0 taken from Coale et al (1983).
 $q_{\omega+} = 1$, $L_{\omega+} = \ell_{\omega}/m_{\omega}$, and $T_{\omega+} = L_{\omega+}$.

- **Period life expectancy:** let $m_x = m_{x,t}$ for some year t .
- Cohort life expectancy: let $m_x = m_{x,t+x}$ for birth cohort in year t .

Life expectancy calculation

Using standard life table calculations:

$$\begin{aligned}\text{For } x = 0, 1, \dots, \omega - 1: \quad & q_x = m_x / (1 + (1 - a_x)m_x) \\ & \ell_{x+1} = \ell_x(1 - q_x) \\ & L_x = \ell_x[1 - q_x(1 - a_x)] \\ & T_x = L_x + L_{x+1} + \dots + L_{\omega-1} + L_{\omega+} \\ & e_x = T_x / L_x\end{aligned}$$

where $a_x = 0.5$ for $x \geq 1$ and a_0 taken from Coale et al (1983).
 $q_{\omega+} = 1$, $L_{\omega+} = \ell_{\omega}/m_{\omega}$, and $T_{\omega+} = L_{\omega+}$.

- Period life expectancy: let $m_x = m_{x,t}$ for some year t .
- Cohort life expectancy: let $m_x = m_{x,t+x}$ for birth cohort in year t .


Cohort life expectancy

- Because we can forecast $m_{x,t}$ we can estimate the mortality rates for each birth cohort (using actual values when they are available).
- We can simulate future $m_{x,t}$ in order to estimate the uncertainty associated with e_x .

Cohort life expectancy

- Because we can forecast $m_{x,t}$ we can estimate the mortality rates for each birth cohort (using actual values when they are available).
- We can simulate future $m_{x,t}$ in order to estimate the uncertainty associated with e_x .

Cohort life expectancy



Simulate future mortality rates

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ and $\{\beta_{t,k}\}$ simulated.
- $\{e_t(x)\}$ and $\{w_t(x)\}$ bootstrapped.
- Generate many future sample paths for $f_{t,M}(x)$ and $f_{t,F}(x)$ to estimate uncertainty in e_t .

Simulate future mortality rates

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ and $\{\beta_{t,k}\}$ simulated.
- $\{e_t(x)\}$ and $\{w_t(x)\}$ bootstrapped.
- Generate many future sample paths for $f_{t,M}(x)$ and $f_{t,F}(x)$ to estimate uncertainty in e_x .

Simulate future mortality rates

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$

$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ and $\{\beta_{t,k}\}$ simulated.
- $\{e_t(x)\}$ and $\{w_t(x)\}$ bootstrapped.
- Generate many future sample paths for $f_{t,M}(x)$ and $f_{t,F}(x)$ to estimate uncertainty in e_x .

Simulate future mortality rates

$$p_t(x) = \sqrt{f_{t,M}(x)f_{t,F}(x)} \quad \text{and} \quad r_t(x) = \sqrt{f_{t,M}(x)/f_{t,F}(x)}.$$


$$\log[p_t(x)] = \mu_p(x) + \sum_{k=1}^K \beta_{t,k} \phi_k(x) + e_t(x)$$

$$\log[r_t(x)] = \mu_r(x) + \sum_{\ell=1}^L \gamma_{t,\ell} \psi_{\ell}(x) + w_t(x).$$

- $\{\gamma_{t,\ell}\}$ and $\{\beta_{t,k}\}$ simulated.
- $\{e_t(x)\}$ and $\{w_t(x)\}$ bootstrapped.
- Generate many future sample paths for $f_{t,M}(x)$ and $f_{t,F}(x)$ to estimate uncertainty in e_x .

Cohort life expectancy

Australia: cohort life expectancy at age 50



Complete code

```
library(demography)

# Read data
aus <- hmd.mx("AUS", "username", "password", "Australia")

# Smooth data
aus.sm <- smooth.demogdata(aus)


# Fit model
aus.pr <- coherentfdm(aus.sm)

# Forecast
aus.pr.fc <- forecast(aus.pr, h=100)

# Compute life expectancies
e50.m.aus.fc <- flife.expectancy(aus.pr.fc, series="male",
  age=50, PI=TRUE, nsim=1000, type="cohort")
e50.f.aus.fc <- flife.expectancy(aus.pr.fc, series="female",
  age=50, PI=TRUE, nsim=1000, type="cohort")
```

Forecast accuracy evaluation

Australian female cohort e50



Forecast accuracy evaluation

Forecast accuracy evaluation

- Compute age 50 remaining cohort life expectancy with a rolling forecast origin beginning in 1921.
- Compare against actual cohort life expectancy where available.
- Compute 80% prediction interval actual coverage.


Forecast accuracy evaluation

- Compute age 50 remaining cohort life expectancy with a rolling forecast origin beginning in 1921.
- Compare against actual cohort life expectancy where available.
- Compute 80% prediction interval actual coverage.


Forecast accuracy evaluation

- Compute age 50 remaining cohort life expectancy with a rolling forecast origin beginning in 1921.
- Compare against actual cohort life expectancy where available.
- Compute 80% prediction interval actual coverage.

Forecast accuracy evaluation



Forecast accuracy evaluation



Outline

- 1 Functional forecasting
- 2 Forecasting groups
- 3 Coherent cohort life expectancy forecasts
- 4 Conclusions**

Some conclusions

- New, automatic, flexible method for coherent forecasting of groups of functional time series.
- Suitable for age-specific mortality.
- Based on geometric means and ratios, so interpretable results.
- More general and flexible than existing methods.
- Easy to compute prediction intervals for any computable statistics.

Some conclusions

- New, automatic, flexible method for coherent forecasting of groups of functional time series.
- Suitable for age-specific mortality.
- Based on geometric means and ratios, so interpretable results.
- More general and flexible than existing methods.
- Easy to compute prediction intervals for any computable statistics.

Some conclusions

- New, automatic, flexible method for coherent forecasting of groups of functional time series.
- Suitable for age-specific mortality.
- Based on geometric means and ratios, so interpretable results.
- More general and flexible than existing methods.
- Easy to compute prediction intervals for any computable statistics.

Some conclusions

- New, automatic, flexible method for coherent forecasting of groups of functional time series.
- Suitable for age-specific mortality.
- Based on geometric means and ratios, so interpretable results.
- More general and flexible than existing methods.
- Easy to compute prediction intervals for any computable statistics.

Some conclusions

- New, automatic, flexible method for coherent forecasting of groups of functional time series.
- Suitable for age-specific mortality.
- Based on geometric means and ratios, so interpretable results.
- More general and flexible than existing methods.
- Easy to compute prediction intervals for any computable statistics.

Selected references

- Hyndman, Ullah (2007). “Robust forecasting of mortality and fertility rates: A functional data approach”. *Computational Statistics and Data Analysis* **51**(10), 4942–4956
- Hyndman, Shang (2009). “Forecasting functional time series (with discussion)”. *Journal of the Korean Statistical Society* **38**(3), 199–221
- Hyndman, Booth, Yasmeen (2013). “Coherent mortality forecasting: the product-ratio method with functional time series models”. *Demography* **50**(1), 261–283
- Booth, Hyndman, Tickle (2013). “Prospective Life Tables”. *Computational Actuarial Science, with R.* ed. by Charpentier. Chapman & Hall/CRC, 323–348
- Hyndman (2013). *demography: Forecasting mortality, fertility, migration and population data.* v1.16. cran.r-project.org/package=demography

Selected references

- Hyndman, Ullah (2007). “Robust forecasting of mortality and fertility rates: A functional data approach”. *Computational Statistics and Data Analysis* **51**(10), 4942–4956
- Hyndman, Shang (2009). “Forecasting functional time series (with discussion)”. *Journal of the Korean Statistical Society* **38**(3), 199–221
- Hyndman, Booth, Yasmineen (2013). “Coherent mortality forecasting: the product-ratio method with functional time series models”. *Demography* **50**(1), 261–283

➞ Papers and R code:

robjhyndman.com

➞ Email: Rob.Hyndman@monash.edu

cran.r-project.org/package=demography