
a
From per-sample LIANA to focus axes
Per-sample LIANA rows
(within shared CM top-10)
11,210,000
rank05 events
(magnitude & specificity ≤ 0.05)
382,000
Consensus
(CM, sender, receiver, LR)
29,605
Research priorities
≥10 eligible, ≥3 GSE, ≥3 cancers
139
Focus axes
(CM01 / 07 / 08)
12
0
2
4
6
log10 count
c
Collagen–DDR pairs present
(only COL1A1 in resource)
COL1A1–DDR2 → myCAF
34 GSE
COL1A1–DDR1 → Malignant epi
28 GSE
COL1A1–DDR1 → Bcell (unres.)
5 GSE
COL1A1–DDR1 → Plasma (unres.)
4 GSE
COL1A1–DDR1 → Granulocyte (unres.)
3 GSE
COL1A1–DDR2 → Endo (blood)
2 GSE
COL1A1–DDR1 → Macrophage (unres.)
2 GSE
COL1A1–DDR1 → Infl. Mono
2 GSE
0
50
100
150
200
Supporting samples
b
Breadth vs threshold robustness
Strict / standard retention
1.0
0.9
0.8
0.7
0.6
0.5
0.4
lower-left: threshold-sensitive
0.5
0.6
0.7
0.8
0.9
1.0
Support fraction (rank ≤ 0.05)
CM01 C1QA–CD93 (SPP1 Mac→Endo (blood))
CM01 S100A9–CD68 (FCN1 Mono→SPP1 Mac)
CM08 S100A9–CD68 (Infl. Mono→Infl. Mac)
CM07 COL1A1–CD44 (myCAF→Infl. Mono)
CM08 S100A8–CD69 (Infl. Mono→CD8 cytotoxic)
CM01 APP–CD74 (Endo (blood)→SPP1 Mac)
CM08 MYL9–CD69 (myCAF→CD8 cytotoxic)
CM08 COL1A1–CD44 (myCAF→CD8 cytotoxic)
CM07 COL1A1–ITGA1_ITGB1 (myCAF→myCAF)
CM07 COL1A1–DDR1 (myCAF→Malignant epi)
CM07 COL1A1–DDR2 (myCAF→myCAF)
CM08 LGALS1–PTPRC (myCAF→CD8 cytotoxic)