



Sharpening Filters





Sharpening (change detection)

1. First derivative

- a) must be zero in areas of constant intensity
- b) must be nonzero at the onset of an intensity step or ramp
- c) must be nonzero along ramps

2. Second derivative

- a) must be zero in constant areas
- b) must be nonzero at the onset and end of an intensity step or ramp
- c) must be zero along ramps of constant slope

Sharpening (change detection)

1. First derivative

$$\frac{\partial f}{\partial x} = f(x + 1) - f(x)$$


2. Second derivative

$$\frac{\partial^2 f}{\partial x^2} = f(x + 1) + f(x - 1) - 2f(x)$$

Performance of derivatives

a
b
c

FIGURE 3.36
Illustration of the first and second derivatives of a 1-D digital function representing a section of a horizontal intensity profile from an image. In (a) and (c) data points are joined by dashed lines as a visualization aid.



Gradient (first derivative)

$$\nabla f \equiv \text{grad}(f) \equiv \begin{bmatrix} g_x \\ g_y \end{bmatrix} = \begin{bmatrix} \frac{\partial f}{\partial x} \\ \frac{\partial f}{\partial y} \end{bmatrix}$$

$$M(x, y) = \text{mag}(\nabla f) = \sqrt{g_x^2 + g_y^2}$$

$$M(x, y) \approx |g_x| + |g_y|$$

$$\alpha(x, y) = \tan^{-1} \left[\frac{g_y}{g_x} \right]$$

Gradient operators

$$g_x = \frac{\partial f(x, y)}{\partial x} = f(x + 1, y) - f(x, y)$$

$$g_y = \frac{\partial f(x, y)}{\partial y} = f(x, y + 1) - f(x, y)$$

Gradient operators

z_1	z_2	z_3
z_4	z_5	z_6
z_7	z_8	z_9

-1	0	0	-1
0	1	1	0

Roberts

-1	-1	-1	-1	0	1
0	0	0	-1	0	1
1	1	1	-1	0	1

Prewitt

-1	-2	-1	-1	0	1
0	0	0	-2	0	2
1	2	1	-1	0	1

Sobel

a
b c
d e
f g

FIGURE 10.14
A 3×3 region of an image (the z 's are intensity values) and various masks used to compute the gradient at the point labeled z_5 .

Gradient operators

a	b
c	d

FIGURE 10.15
Prewitt and Sobel masks for detecting diagonal edges.

0	1	1
-1	0	1
-1	-1	0

-1	-1	0
-1	0	1
0	1	1

Prewitt

0	1	2
-1	0	1
-2	-1	0

-2	-1	0
-1	0	1
0	1	2

Sobel

Gradient operators



a	b
c	d


FIGURE 10.16

(a) Original image of size 834×1114 pixels, with intensity values scaled to the range $[0, 1]$.
(b) $|g_x|$, the component of the gradient in the x -direction, obtained using the Sobel mask in Fig. 10.14(f) to filter the image.
(c) $|g_y|$, obtained using the mask in Fig. 10.14(g).
(d) The gradient image, $|g_x| + |g_y|$.

FIGURE 10.17

Gradient angle image computed using Eq. (10.2-11). Areas of constant intensity in this image indicate that the direction of the gradient vector is the same at all the pixel locations in those regions.

Gradient operators



a b

FIGURE 3.42

(a) Optical image of contact lens (note defects on the boundary at 4 and 5 o'clock).

(b) Sobel gradient.

(Original image courtesy of Pete Sites, Perceptics Corporation.)

Sobel Gradient operator in OpenCV

Sobel(src, dst, ddepth, dx, dy)

Sobel(input, horizontal_derivative, CV_32F, 1, 0);

Sobel(input, vertical_derivative, CV_32F, 0, 1);