

# MM2090

## Assignment 4

### Authors

Akshat Rakesh Garhwal  
me20b020@smail.iitm.ac.in  
Archish S  
me20b032@smail.iitm.ac.in  
Prabhat Bedida  
me20b132@smail.iitm.ac.in  
Albin George  
mm20b005@smail.iitm.ac.in  
Prithviraj Pratap Bhosle  
mm20b049@smail.iitm.ac.in

This is the submission file for  
Group 7

Indian Institute Of Technology Madras  
June 30, 2021

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Akshat Rakesh Garhwal me20b020</b>	<b>2</b>
2.1	Snell's Law . . . . .	2
2.2	Snell's Law equation . . . . .	2
2.3	Terms in Snell's Law equation. . . . .	2
2.4	Importance of Snell's Law . . . . .	2
<b>3</b>	<b>Archish S me20b032</b>	<b>4</b>
3.1	The Value Function . . . . .	5
3.2	The Optimal Policy . . . . .	5
<b>4</b>	<b>Prabhat Bedida me20b132</b>	<b>6</b>
4.1	Terms used in the equation . . . . .	6
<b>5</b>	<b>Albin George mm20b005</b>	<b>7</b>
5.1	The Equation . . . . .	7
5.2	Analysis . . . . .	7

## 1 Introduction

This is the submission file for Assignment 4.

## 2 Akshat Rakesh Garhwal me20b020

### 2.1 Snell's Law

Snell's law is a formula used to explain the change in angle of movement of light when it passes through a surface between two different mediums. While the law is named after Willebrord Snellius, it was first accurately described by Ibn Sahl over 600 years before Willebrord Snellius. It states that the sines of angles of incidence and refraction is equivalent to the ratio of velocity in medium or equivalent to reciprocal of the indices of refraction.

### 2.2 Snell's Law equation

$$\frac{\sin(\theta_2)}{\sin(\theta_1)} = \frac{\nu_2}{\nu_1} = \frac{n_1}{n_2} \quad (1)$$

### 2.3 Terms in Snell's Law equation.

The terms in the equation are

- $\theta_1$  - The angle of incidence (angle measured from the normal to the incident light)
- $\theta_2$  - The angle of refraction (angle measured from the normal to the refracted ray)
- $\nu_1$  - Velocity of light in the incident medium
- $\nu_2$  - Velocity of light in the refraction medium
- $n_1$  - Refractive index of incident medium
- $n_2$  - Refractive index of refraction medium

### 2.4 Importance of Snell's Law




Figure 1: Light changes its direction of propagation as it passes through an interface of two different medium

Snell's Law serves as backbone for Optics and related studies, with every application of refractive optics relying on this formula. It can predict the movement of light through surfaces

(refer fig.(1)) and this has led to countless applications. It is used while making lens for spectacles and experimental purposes.

### 3 Archish S me20b032

A Markov decision process [?] can be described as a tuple  $\langle S, A, T, R \rangle$ , where

- $S$  is a finite set of states of the world;
- $A$  is a finite set of actions;
- $T : S \times A \rightarrow \Pi(S)$  is the *state-transition function*, giving for each world state and agent action, a probability distribution over world states (we write  $T(s, a, s')$  for the probability of ending in state  $s'$ , given that the agent starts in state  $s$  and takes action  $a$ );
- $R : S \times A \rightarrow \mathbb{R}$  is the reward function, giving the expected immediate reward gained by the agent for taking each action in each state (we write  $R(s, a)$  for the expected reward for taking action  $a$  in state  $s$ );
- A stationary policy,  $\pi : S \rightarrow A$ , is a situation-action mapping that specifies, for each state, an action to be taken.
- $V_\pi(s)$  is the expected discounted sum of future reward for starting in state  $s$  and executing policy  $\pi$ .

In this model, as described by figure 2, the next state and the expected reward depend only on the previous state and the action taken; even if we were to condition on additional previous states, the transition probabilities and the expected rewards would remain the same. This is known as the Markov property.




Figure 2: An MDP models the synchronous interaction between agent and world

### 3.1 The Value Function

$$V_{\pi}(s) = R(s, \pi(s)) + \gamma \sum_{s' \in S} T(s, \pi(s), s') V_{\pi}(s') \quad (2)$$

### 3.2 The Optimal Policy

Given the Value Function 2 a greedy policy with respect to that value function,  $\pi_V$ , is defined as

$$\pi_V(s) = \operatorname{argmax}_a \left[ R(s, a) + \gamma \sum_{s' \in S} T(s, a, s') V(s') \right] \quad (3)$$

## 4 Prabhat Bedida me20b132

$$\nabla \times \vec{B} = \mu_0 \vec{J} + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t} \quad (4)$$

Eqn. 4 is commonly known as Maxwell's fourth electromagnetic equation. It is an expansion on Ampère's Circuital Law and is hence also known as the Ampère-Maxwell Law. Currently a more generalized form is used, which takes into account the behaviour and interference of material substances in electric and magnetic fields [?].

### 4.1 Terms used in the equation

- $\nabla \times$  - Curl Operator
- $\vec{B}$  - Magnetic Field Vector
- $\vec{J}$  - Current Density Vector
- $\vec{E}$  - Electric Field Vector
- $\mu_0$  (Constant) - Permeability of Free Space
- $\epsilon_0$  (Constant) - Permittivity of Free Space

It is an important equation as it links two phenomena, which were previously thought to be separate, magnetism and electricity. It also finally proved that light is an electromagnetic wave, (as seen in Fig. 3 taken from a paper by J. Gratus, M.W.McCall and P.Kinsler [?]) leading to a whole new area and direction of research.




Figure 3: Propagation of Light with respect to magnetic and electric fields

## 5 Albin George mm20b005

### 5.1 The Equation

The Coulombs Law :

$$F = \frac{K q_1 q_2}{r^2} \quad (5)$$

### 5.2 Analysis

Following contains a brief explanation of the variables and the importance of the equation :

The above given equation ?? has terms **F** ,**K** , **q1,q2** and **r**.

Here,

**F** represents the force exerted by m1 on m2 or vice versa

**K** represents the Coulombs Constant

**q1** represents the Charge of entity 1

**q2** represents the Charge of entity 2

**r** represents the distance between entity 1 and entity 2

Coulomb's law, or Coulomb's inverse-square law, is an experimental law[1] of physics that quantifies the amount of force between two stationary, electrically charged particles. The electric force between charged bodies at rest is conventionally called electrostatic force or Coulomb force. The law was first discovered in 1785 by French physicist Charles-Augustin de Coulomb, hence the name. Coulomb's law was essential to the development of the theory of electromagnetism, maybe even its starting point, as it made it possible to discuss the quantity of electric charge in a meaningful way.

Webpage Links [?]






Figure 4: The Coulombs Law[?]