

# Algebraic Number Theory

## Problem sheet 3

1. (3 points) Verify the Chinese Remainder Theorem for Dedekind domains: If  $A \triangleleft \mathcal{O}$  is an ideal in the Dedekind domain  $\mathcal{O}$  with decomposition  $A = P_1^{\nu_1} \dots P_t^{\nu_t}$  as a product of prime ideals then we have

$$\mathcal{O}/A \cong \bigoplus_{i=1}^t \mathcal{O}/P_i^{\nu_i}.$$

2. (3 points) Show that a lattice  $\Gamma \subset V$  is complete if and only if  $V/\Gamma$  is compact.
3. (2 points) Let  $L_i(x_1, \dots, x_n) = \sum_{j=1}^n a_{ij}x_j$  ( $i = 1, \dots, n$ ) be real homogeneous linear polynomials such that  $\det((a_{ij}))_{ij} \neq 0$  and let  $c_1, \dots, c_n$  be positive real numbers with  $c_1 \dots c_n > |\det((a_{ij}))_{ij}|$ . Verify that there exist not all zero integers  $m_1, \dots, m_n \in \mathbb{Z}$  such that  $|L_i(m_1, \dots, m_n)| < c_i$  ( $i = 1, \dots, n$ ).
4. (2+2 points) Show that the map

$$\begin{aligned} \mathbb{C} \otimes_{\mathbb{Q}} K &\rightarrow K_{\mathbb{C}} \\ z \otimes \alpha &\mapsto z \cdot (j\alpha) \end{aligned}$$

is an isomorphism of rings. Moreover, prove that its restriction to  $\mathbb{R} \otimes_{\mathbb{Q}} K$  also induces an isomorphism between  $K \otimes_{\mathbb{Q}} \mathbb{R}$  and  $K_{\mathbb{R}}$ .

5. (1 point for each) Let  $K = \mathbb{Q}[\sqrt{d}]$  be an **imaginary** quadratic field, where  $d < 0$  is squarefree. Show that  $\mathcal{O}_K$  is a PID if  $d$  is one of the following integers:

$$-1, -2, -3, -7, -11, -19, -43, -67, -163.$$

**Remark.** Historically, Gauss proved the above result and conjectured that there was no other such  $d$ , which was finally proved by Baker, Heegner and Stark in the middle of the 20th century, after people has worked on it for more than 150 years.

6. (1 point for each) Let  $K = \mathbb{Q}[\sqrt{d}]$  be a **real** quadratic field, where  $d > 0$  is squarefree. Show that  $\mathcal{O}_K$  is a PID if  $d$  is one of the following integers:

$$2, 3, 5, 6, 7, 11, 13, 14, 17, 19, 21, 22, 23, 29, 31, 33, 37, 38, 41, 43, \\ 46, 47, 53, 57, 59, 61, 62, 67, 69, 71, 73, 77, 83, 86, 89, 93, 94, 97$$

**Remark.** It is not known whether there are infinitely many such  $d$  for real quadratic fields.

7. (a) (4 points) Let  $X_t := \{z \in K_{\mathbb{R}} \mid \sum_{\tau} |z_{\tau}| < t\}$  for any positive real number  $t$ . Verify  $\text{vol}(X_t) = 2^r \pi^s \frac{t^n}{n!}$ .
- (b) (4 points) Assume  $|K/\mathbb{Q}| = n$ . Show  $|d_K|^{1/2} \geq \frac{n^n}{n!} \left(\frac{\pi}{4}\right)^{n/2}$ . (Hint: choose  $t$  so that  $\text{vol}(X_t) > 2^n \text{vol}(\Gamma)$  where  $\Gamma = j(\mathcal{O}_K) \subset K_{\mathbb{R}}$ . Apply the inequality between geometric and arithmetic mean on the numbers  $|\tau(\alpha)|$  where  $0 \neq \alpha \in \mathcal{O}_K$  is the element with  $j(\alpha) \in X_t$  guaranteed by Minkowski's lattice point theorem. Finally, note that  $1 \leq |N_{K/\mathbb{Q}}(\alpha)|$ .)
- (c) (1 point) Show that whenever the degree  $|K : \mathbb{Q}|$  goes to infinity, so does the discriminant  $d_K$ . Moreover, we have  $d_K > 1$  for all extensions  $K \neq \mathbb{Q}$ .

8. (a) (3 points) Let  $A \triangleleft \mathcal{O}_K$  be an ideal whose class has order  $m$  in the class group (ie.  $m$  is the smallest integer such that  $A^m = (\alpha)$  is a principal ideal). Prove that  $A\mathcal{O}_L = (\beta)$  where  $L = K(\beta)$  and  $\beta^m = \alpha$ .
- (b) (1 point) Verify that for each number field  $K$  there is a finite extension  $L/K$  in which all ideals of  $\mathcal{O}_K$  become principal (ie. all ideals of  $\mathcal{O}_K$  *capitulate* in  $\mathcal{O}_L$ ).
- (c) (2 points) Verify that the ring  $\Omega$  of all algebraic integers (in  $\mathbb{C}$ ) is a *Bézout domain*, ie. all finitely generated ideals are principal (but not noetherian, so not a PID).
9. (a) (3 points) Show that over a Bézout domain every finitely generated torsion-free module is free.
- (b) (2 points) Verify that  $\bigcup_{n=1}^{\infty} \mathbb{C}[[x^{1/n}]]$  is a Bézout domain.
10. In this exercise we compute the rank of the unit group  $\mathcal{O}_K^\times$  as an abelian group using a multiplicative version of Minkowski's theory.
- (a) (3 points) Let  $K_{\mathbb{C}}^\times = \prod_{\tau} \mathbb{C}^\times$  be the multiplicative group of invertible elements in the ring  $K_{\mathbb{C}}$  and let  $N: K_{\mathbb{C}}^\times \rightarrow \mathbb{C}^\times$  be the group homomorphism defined as the product of coordinates. Further define the homomorphism  $l := \log |\cdot|: K_{\mathbb{C}}^\times \rightarrow \prod_{\tau} \mathbb{R}$  coordinatewise. Show that the kernel of  $l \circ j: \mathcal{O}_K^\times \rightarrow \prod_{\tau} \mathbb{R}$  is the torsion subgroup in  $\mathcal{O}_K^\times$ , ie. the group  $\mu(K)$  of roots of unity in  $K$ .
- (b) (2 points) Verify that for each  $\alpha \in \mathcal{O}_K^\times$  the element  $l \circ j(\alpha) \in \prod_{\tau} \mathbb{R}$  is fixed by the complex conjugation (permuting  $\tau \in \text{Hom}_{\mathbb{Q}}(K, \mathbb{C})$ ). Further show that the sum of coordinates of  $l \circ j(\alpha)$  equals 0.
- (c) (5 points) Show that  $l \circ j(\mathcal{O}_K^\times)$  is a full lattice in the subspace  $H$  where

$$H = \left\{ x_{\tau} \in \prod_{\tau} \mathbb{R} \mid \sum_{\tau} x_{\tau} = 0 \text{ and } x_{\sigma_k} = x_{\overline{\sigma_k}} \ (k = 1, \dots, s) \right\} .$$

In particular,  $\mathcal{O}_K^\times \cong \mu(K) \times \mathbb{Z}^{r+s-1}$  (as an abelian group).