

# Contents

<b>2</b>	<b>Bayesian decision theory</b>	<b>3</b>
2.1	Introduction . . . . .	3
2.2	Bayesian Decision Theory – Continuous Features . . . . .	7
2.2.1	Two-Category Classification . . . . .	8
2.3	Minimum-Error-Rate Classification . . . . .	9
2.3.1	*Minimax Criterion . . . . .	10
2.3.2	*Neyman-Pearson Criterion . . . . .	12
2.4	Classifiers, Discriminants and Decision Surfaces . . . . .	13
2.4.1	The Multi-Category Case . . . . .	13
2.4.2	The Two-Category Case . . . . .	14
2.5	The Normal Density . . . . .	15
2.5.1	Univariate Density . . . . .	16
2.5.2	Multivariate Density . . . . .	17
2.6	Discriminant Functions for the Normal Density . . . . .	19
2.6.1	Case 1: $\Sigma_i = \sigma^2 \mathbf{I}$ . . . . .	19
2.6.2	Case 2: $\Sigma_i = \Sigma$ . . . . .	23
2.6.3	Case 3: $\Sigma_i = \text{arbitrary}$ . . . . .	25
	<i>Example 1: Decisions for Gaussian data</i> . . . . .	29
2.7	*Error Probabilities and Integrals . . . . .	30
2.8	*Error Bounds for Normal Densities . . . . .	31
2.8.1	Chernoff Bound . . . . .	31
2.8.2	Bhattacharyya Bound . . . . .	32
	<i>Example 2: Error bounds</i> . . . . .	33
2.8.3	Signal Detection Theory and Operating Characteristics . . . . .	33
2.9	Bayes Decision Theory — Discrete Features . . . . .	36
2.9.1	Independent Binary Features . . . . .	36
	<i>Example 3: Bayes decisions for binary data</i> . . . . .	38
2.10	*Missing and Noisy Features . . . . .	39
2.10.1	Missing Features . . . . .	39
2.10.2	Noisy Features . . . . .	40
2.11	*Compound Bayes Decision Theory and Context . . . . .	41
	Summary . . . . .	42
	Bibliographical and Historical Remarks . . . . .	43
	Problems . . . . .	44
	Computer exercises . . . . .	59
	Bibliography . . . . .	61
	Index . . . . .	65

## Chapter 2

# Bayesian decision theory

---

### 2.1 Introduction

Bayesian decision theory is a fundamental statistical approach to the problem of pattern classification. This approach is based on quantifying the tradeoffs between various classification decisions using probability and the costs that accompany such decisions. It makes the assumption that the decision problem is posed in probabilistic terms, and that all of the relevant probability values are known. In this chapter we develop the fundamentals of this theory, and show how it can be viewed as being simply a formalization of common-sense procedures; in subsequent chapters we will consider the problems that arise when the probabilistic structure is not completely known.

While we will give a quite general, abstract development of Bayesian decision theory in Sect. ??, we begin our discussion with a specific example. Let us reconsider the hypothetical problem posed in Chap. ?? of designing a classifier to separate two kinds of fish: sea bass and salmon. Suppose that an observer watching fish arrive along the conveyor belt finds it hard to predict what type will emerge next and that the sequence of types of fish appears to be random. In decision-theoretic terminology we would say that as each fish emerges nature is in one or the other of the two possible states: either the fish is a sea bass or the fish is a salmon. We let  $\omega$  denote the *state of nature*, with  $\omega = \omega_1$  for sea bass and  $\omega = \omega_2$  for salmon. Because the state of nature is so unpredictable, we consider  $\omega$  to be a variable that must be described probabilistically.

STATE OF  
NATURE

If the catch produced as much sea bass as salmon, we would say that the next fish is equally likely to be sea bass or salmon. More generally, we assume that there is some *a priori probability* (or simply *prior*)  $P(\omega_1)$  that the next fish is sea bass, and some prior probability  $P(\omega_2)$  that it is salmon. If we assume there are no other types of fish relevant here, then  $P(\omega_1)$  and  $P(\omega_2)$  sum to one. These prior probabilities reflect our prior knowledge of how likely we are to get a sea bass or salmon before the fish actually appears. It might, for instance, depend upon the time of year or the choice of fishing area.

PRIOR

Suppose for a moment that we were forced to make a decision about the type of fish that will appear next without being allowed to see it. For the moment, we shall

DECISION  
RULE

assume that any incorrect classification entails the same cost or consequence, and that the only information we are allowed to use is the value of the prior probabilities. If a decision must be made with so little information, it seems logical to use the following *decision rule*: Decide  $\omega_1$  if  $P(\omega_1) > P(\omega_2)$ ; otherwise decide  $\omega_2$ .

This rule makes sense if we are to judge just one fish, but if we are to judge many fish, using this rule repeatedly may seem a bit strange. After all, we would always make the same decision even though we know that *both* types of fish will appear. How well it works depends upon the values of the prior probabilities. If  $P(\omega_1)$  is very much greater than  $P(\omega_2)$ , our decision in favor of  $\omega_1$  will be right most of the time. If  $P(\omega_1) = P(\omega_2)$ , we have only a fifty-fifty chance of being right. In general, the probability of error is the smaller of  $P(\omega_1)$  and  $P(\omega_2)$ , and we shall see later that under these conditions no other decision rule can yield a larger probability of being right.

In most circumstances we are not asked to make decisions with so little information. In our example, we might for instance use a lightness measurement  $x$  to improve our classifier. Different fish will yield different lightness readings and we express this variability in probabilistic terms; we consider  $x$  to be a continuous random variable whose distribution depends on the state of nature, and is expressed as  $p(x|\omega_1)$ .<sup>\*</sup> This is the *class-conditional probability density* function. Strictly speaking, the probability density function  $p(x|\omega_1)$  should be written as  $p_X(x|\omega_1)$  to indicate that we are speaking about a particular density function for the random variable  $X$ . This more elaborate subscripted notation makes it clear that  $p_X(\cdot)$  and  $p_Y(\cdot)$  denote two different functions, a fact that is obscured when writing  $p(x)$  and  $p(y)$ . Since this potential confusion rarely arises in practice, we have elected to adopt the simpler notation. Readers who are unsure of our notation or who would like to review probability theory should see Appendix ??). This is the probability density function for  $x$  given that the state of nature is  $\omega_1$ . (It is also sometimes called state-conditional probability density.) Then the difference between  $p(x|\omega_1)$  and  $p(x|\omega_2)$  describes the difference in lightness between populations of sea bass and salmon (Fig. 2.1).

Suppose that we know both the prior probabilities  $P(\omega_j)$  and the conditional densities  $p(x|\omega_j)$ . Suppose further that we measure the lightness of a fish and discover that its value is  $x$ . How does this measurement influence our attitude concerning the true state of nature — that is, the category of the fish? We note first that the (joint) probability density of finding a pattern that is in category  $\omega_j$  *and* has feature value  $x$  can be written two ways:  $p(\omega_j, x) = P(\omega_j|x)p(x) = p(x|\omega_j)P(\omega_j)$ . Rearranging these leads us to the answer to our question, which is called *Bayes' formula*:

$$\boxed{P(\omega_j|x) = \frac{p(x|\omega_j)P(\omega_j)}{p(x)}}, \quad (1)$$

where in this case of two categories

$$p(x) = \sum_{j=1}^2 p(x|\omega_j)P(\omega_j). \quad (2)$$

Bayes' formula can be expressed informally in English by saying that

$$\text{posterior} = \frac{\text{likelihood} \times \text{prior}}{\text{evidence}}. \quad (3)$$

---

<sup>\*</sup> We generally use an upper-case  $P(\cdot)$  to denote a probability mass function and a lower-case  $p(\cdot)$  to denote a probability density function.

Bayes' formula shows that by observing the value of  $x$  we can convert the prior probability  $P(\omega_j)$  to the *a posteriori* probability (or *posterior*) probability  $P(\omega_j|x)$  — the probability of the state of nature being  $\omega_j$  given that feature value  $x$  has been measured. We call  $p(x|\omega_j)$  the *likelihood* of  $\omega_j$  with respect to  $x$  (a term chosen to indicate that, other things being equal, the category  $\omega_j$  for which  $p(x|\omega_j)$  is large is more “likely” to be the true category). Notice that it is the product of the likelihood and the prior probability that is most important in determining the posterior probability; the evidence factor,  $p(x)$ , can be viewed as merely a scale factor that guarantees that the posterior probabilities sum to one, as all good probabilities must. The variation of  $P(\omega_j|x)$  with  $x$  is illustrated in Fig. 2.2 for the case  $P(\omega_1) = 2/3$  and  $P(\omega_2) = 1/3$ .

POSTERIOR  
LIKELIHOOD




Figure 2.1: Hypothetical class-conditional probability density functions show the probability density of measuring a particular feature value  $x$  given the pattern is in category  $\omega_i$ . If  $x$  represents the length of a fish, the two curves might describe the difference in length of populations of two types of fish. Density functions are normalized, and thus the area under each curve is 1.0.

If we have an observation  $x$  for which  $P(\omega_1|x)$  is greater than  $P(\omega_2|x)$ , we would naturally be inclined to decide that the true state of nature is  $\omega_1$ . Similarly, if  $P(\omega_2|x)$  is greater than  $P(\omega_1|x)$ , we would be inclined to choose  $\omega_2$ . To justify this decision procedure, let us calculate the probability of error whenever we make a decision. Whenever we observe a particular  $x$ ,

$$P(\text{error}|x) = \begin{cases} P(\omega_1|x) & \text{if we decide } \omega_2 \\ P(\omega_2|x) & \text{if we decide } \omega_1. \end{cases} \quad (4)$$

Clearly, for a given  $x$  we can minimize the probability of error by deciding  $\omega_1$  if  $P(\omega_1|x) > P(\omega_2|x)$  and  $\omega_2$  otherwise. Of course, we may never observe exactly the same value of  $x$  twice. Will this rule minimize the average probability of error? Yes, because the average probability of error is given by




Figure 2.2: Posterior probabilities for the particular priors  $P(\omega_1) = 2/3$  and  $P(\omega_2) = 1/3$  for the class-conditional probability densities shown in Fig. 2.1. Thus in this case, given that a pattern is measured to have feature value  $x = 14$ , the probability it is in category  $\omega_2$  is roughly 0.08, and that it is in  $\omega_1$  is 0.92. At every  $x$ , the posteriors sum to 1.0.

$$P(\text{error}) = \int_{-\infty}^{\infty} P(\text{error}, x) dx = \int_{-\infty}^{\infty} P(\text{error}|x)p(x) dx \quad (5)$$

and if for every  $x$  we insure that  $P(\text{error}|x)$  is as small as possible, then the integral must be as small as possible. Thus we have justified the following *Bayes' decision rule* for minimizing the probability of error:

BAYES'  
DECISION  
RULE

$$\text{Decide } \omega_1 \text{ if } P(\omega_1|x) > P(\omega_2|x); \text{ otherwise decide } \omega_2, \quad (6)$$

and under this rule Eq. 4 becomes

$$P(\text{error}|x) = \min [P(\omega_1|x), P(\omega_2|x)]. \quad (7)$$

This form of the decision rule emphasizes the role of the posterior probabilities. By using Eq. 1, we can instead express the rule in terms of the conditional and prior probabilities. First note that the *evidence*,  $p(x)$ , in Eq. 1 is unimportant as far as making a decision is concerned. It is basically just a scale factor that states how frequently we will actually measure a pattern with feature value  $x$ ; its presence in Eq. 1 assures us that  $P(\omega_1|x) + P(\omega_2|x) = 1$ . By eliminating this scale factor, we obtain the following completely equivalent decision rule:

EVIDENCE

$$\text{Decide } \omega_1 \text{ if } p(x|\omega_1)P(\omega_1) > p(x|\omega_2)P(\omega_2); \text{ otherwise decide } \omega_2. \quad (8)$$

Some additional insight can be obtained by considering a few special cases. If for some  $x$  we have  $p(x|\omega_1) = p(x|\omega_2)$ , then that particular observation gives us no

information about the state of nature; in this case, the decision hinges entirely on the prior probabilities. On the other hand, if  $P(\omega_1) = P(\omega_2)$ , then the states of nature are equally probable; in this case the decision is based entirely on the likelihoods  $p(x|\omega_j)$ . In general, both of these factors are important in making a decision, and the Bayes decision rule combines them to achieve the minimum probability of error.

## 2.2 Bayesian Decision Theory – Continuous Features

We shall now formalize the ideas just considered, and generalize them in four ways:

- by allowing the use of more than one feature
- by allowing more than two states of nature
- by allowing actions other than merely deciding the state of nature
- by introducing a loss function more general than the probability of error.

These generalizations and their attendant notational complexities should not obscure the central points illustrated in our simple example. Allowing the use of more than one feature merely requires replacing the scalar  $x$  by the *feature vector*  $\mathbf{x}$ , where  $\mathbf{x}$  is in a  $d$ -dimensional Euclidean space  $\mathbf{R}^d$ , called the *feature space*. Allowing more than two states of nature provides us with a useful generalization for a small notational expense. Allowing actions other than classification primarily allows the possibility of rejection, i.e., of refusing to make a decision in close cases; this is a useful option if being indecisive is not too costly. Formally, the *loss function* states exactly how costly each action is, and is used to convert a probability determination into a decision. Cost functions let us treat situations in which some kinds of classification mistakes are more costly than others, although we often discuss the simplest case, where all errors are equally costly. With this as a preamble, let us begin the more formal treatment.

FEATURE  
SPACE  
  
LOSS  
FUNCTION

Let  $\omega_1, \dots, \omega_c$  be the finite set of  $c$  states of nature (“categories”) and  $\alpha_1, \dots, \alpha_a$  be the finite set of  $a$  possible actions. The loss function  $\lambda(\alpha_i|\omega_j)$  describes the loss incurred for taking action  $\alpha_i$  when the state of nature is  $\omega_j$ . Let the feature vector  $\mathbf{x}$  be a  $d$ -component vector-valued random variable, and let  $p(\mathbf{x}|\omega_j)$  be the state-conditional probability density function for  $\mathbf{x}$  — the probability density function for  $\mathbf{x}$  conditioned on  $\omega_j$  being the true state of nature. As before,  $P(\omega_j)$  describes the prior probability that nature is in state  $\omega_j$ . Then the posterior probability  $P(\omega_j|\mathbf{x})$  can be computed from  $p(\mathbf{x}|\omega_j)$  by Bayes’ formula:

$$P(\omega_j|\mathbf{x}) = \frac{p(\mathbf{x}|\omega_j)P(\omega_j)}{p(\mathbf{x})}, \quad (9)$$

where the evidence is now

$$p(\mathbf{x}) = \sum_{j=1}^c p(\mathbf{x}|\omega_j)P(\omega_j). \quad (10)$$

Suppose that we observe a particular  $\mathbf{x}$  and that we contemplate taking action  $\alpha_i$ . If the true state of nature is  $\omega_j$ , by definition we will incur the loss  $\lambda(\alpha_i|\omega_j)$ . Since  $P(\omega_j|\mathbf{x})$  is the probability that the true state of nature is  $\omega_j$ , the expected loss associated with taking action  $\alpha_i$  is merely

$$R(\alpha_i|\mathbf{x}) = \sum_{j=1}^c \lambda(\alpha_i|\omega_j)P(\omega_j|\mathbf{x}). \quad (11)$$

RISK

In decision-theoretic terminology, an expected loss is called a *risk*, and  $R(\alpha_i|\mathbf{x})$  is called the *conditional risk*. Whenever we encounter a particular observation  $\mathbf{x}$ , we can minimize our expected loss by selecting the action that minimizes the conditional risk. We shall now show that this *Bayes decision procedure* actually provides the optimal performance on an overall risk.

DECISION

RULE

Stated formally, our problem is to find a decision rule against  $P(\omega_j)$  that minimizes the overall risk. A general *decision rule* is a function  $\alpha(\mathbf{x})$  that tells us which action to take for every possible observation. To be more specific, for every  $\mathbf{x}$  the *decision function*  $\alpha(\mathbf{x})$  assumes one of the  $a$  values  $\alpha_1, \dots, \alpha_a$ . The overall risk  $R$  is the expected loss associated with a given decision rule. Since  $R(\alpha_i|\mathbf{x})$  is the conditional risk associated with action  $\alpha_i$ , and since the decision rule specifies the action, the overall risk is given by

$$R = \int R(\alpha(\mathbf{x})|\mathbf{x})p(\mathbf{x}) d\mathbf{x}, \quad (12)$$

where  $d\mathbf{x}$  is our notation for a  $d$ -space volume element, and where the integral extends over the entire feature space. Clearly, if  $\alpha(\mathbf{x})$  is chosen so that  $R(\alpha_i(\mathbf{x}))$  is as small as possible for every  $\mathbf{x}$ , then the overall risk will be minimized. This justifies the following statement of the *Bayes decision rule*: To minimize the overall risk, compute the conditional risk

$$R(\alpha_i|\mathbf{x}) = \sum_{j=1}^c \lambda(\alpha_i|\omega_j)P(\omega_j|\mathbf{x}) \quad (13)$$

BAYES RISK

for  $i = 1, \dots, a$  and select the action  $\alpha_i$  for which  $R(\alpha_i|\mathbf{x})$  is minimum.\* The resulting minimum overall risk is called the *Bayes risk*, denoted  $R^*$ , and is the best performance that can be achieved.

### 2.2.1 Two-Category Classification

Let us consider these results when applied to the special case of two-category classification problems. Here action  $\alpha_1$  corresponds to deciding that the true state of nature is  $\omega_1$ , and action  $\alpha_2$  corresponds to deciding that it is  $\omega_2$ . For notational simplicity, let  $\lambda_{ij} = \lambda(\alpha_i|\omega_j)$  be the loss incurred for deciding  $\omega_i$  when the true state of nature is  $\omega_j$ . If we write out the conditional risk given by Eq. 13, we obtain

$$\begin{aligned} R(\alpha_1|\mathbf{x}) &= \lambda_{11}P(\omega_1|\mathbf{x}) + \lambda_{12}P(\omega_2|\mathbf{x}) \quad \text{and} \\ R(\alpha_2|\mathbf{x}) &= \lambda_{21}P(\omega_1|\mathbf{x}) + \lambda_{22}P(\omega_2|\mathbf{x}). \end{aligned} \quad (14)$$

There are a variety of ways of expressing the minimum-risk decision rule, each having its own minor advantages. The fundamental rule is to decide  $\omega_1$  if  $R(\alpha_1|\mathbf{x}) < R(\alpha_2|\mathbf{x})$ . In terms of the posterior probabilities, we decide  $\omega_1$  if

$$(\lambda_{21} - \lambda_{11})P(\omega_1|\mathbf{x}) > (\lambda_{12} - \lambda_{22})P(\omega_2|\mathbf{x}). \quad (15)$$

---

\* Note that if more than one action minimizes  $R(\alpha|\mathbf{x})$ , it does not matter which of these actions is taken, and any convenient tie-breaking rule can be used.

Ordinarily, the loss incurred for making an error is greater than the loss incurred for being correct, and both of the factors  $\lambda_{21} - \lambda_{11}$  and  $\lambda_{12} - \lambda_{22}$  are positive. Thus in practice, our decision is generally determined by the more likely state of nature, although we must scale the posterior probabilities by the loss differences. By employing Bayes' formula, we can replace the posterior probabilities by the prior probabilities and the conditional densities. This results in the equivalent rule, to decide  $\omega_1$  if

$$(\lambda_{21} - \lambda_{11})p(\mathbf{x}|\omega_1)P(\omega_1) > (\lambda_{12} - \lambda_{22})p(\mathbf{x}|\omega_2)P(\omega_2), \quad (16)$$

and  $\omega_2$  otherwise.

Another alternative, which follows at once under the reasonable assumption that  $\lambda_{21} > \lambda_{11}$ , is to decide  $\omega_1$  if

$$\frac{p(\mathbf{x}|\omega_1)}{p(\mathbf{x}|\omega_2)} > \frac{\lambda_{12} - \lambda_{22}}{\lambda_{21} - \lambda_{11}} \frac{P(\omega_2)}{P(\omega_1)}. \quad (17)$$

This form of the decision rule focuses on the  $\mathbf{x}$ -dependence of the probability densities. We can consider  $p(\mathbf{x}|\omega_j)$  a function of  $\omega_j$  (i.e., the likelihood function), and then form the *likelihood ratio*  $p(\mathbf{x}|\omega_1)/p(\mathbf{x}|\omega_2)$ . Thus the Bayes decision rule can be interpreted as calling for deciding  $\omega_1$  if the likelihood ratio exceeds a threshold value that is independent of the observation  $\mathbf{x}$ .

LIKELIHOOD  
RATIO

## 2.3 Minimum-Error-Rate Classification

In classification problems, each state of nature is usually associated with a different one of the  $c$  classes, and the action  $\alpha_i$  is usually interpreted as the decision that the true state of nature is  $\omega_i$ . If action  $\alpha_i$  is taken and the true state of nature is  $\omega_j$ , then the decision is correct if  $i = j$ , and in error if  $i \neq j$ . If errors are to be avoided, it is natural to seek a decision rule that minimizes the probability of error, i.e., the *error rate*.

The loss function of interest for this case is hence the so-called *symmetrical* or *zero-one* loss function,

ZERO-ONE  
LOSS

$$\lambda(\alpha_i|\omega_j) = \begin{cases} 0 & i = j \\ 1 & i \neq j \end{cases} \quad i, j = 1, \dots, c. \quad (18)$$

This loss function assigns no loss to a correct decision, and assigns a unit loss to any error; thus, all errors are equally costly.\* The risk corresponding to this loss function is precisely the average probability of error, since the conditional risk is

$$\begin{aligned} R(\alpha_i|\mathbf{x}) &= \sum_{j=1}^c \lambda(\alpha_i|\omega_j)P(\omega_j|\mathbf{x}) \\ &= \sum_{j \neq i} P(\omega_j|\mathbf{x}) \\ &= 1 - P(\omega_i|\mathbf{x}) \end{aligned} \quad (19)$$

---

\* We note that other loss functions, such as quadratic and linear-difference, find greater use in regression tasks, where there is a natural ordering on the predictions and we can meaningfully penalize predictions that are "more wrong" than others.



and  $P(\omega_i|\mathbf{x})$  is the conditional probability that action  $\alpha_i$  is correct. The Bayes decision rule to minimize risk calls for selecting the action that minimizes the conditional risk. Thus, to minimize the average probability of error, we should select the  $i$  that *maximizes* the posterior probability  $P(\omega_i|\mathbf{x})$ . In other words, for *minimum error rate*:

$$\text{Decide } \omega_i \text{ if } P(\omega_i|\mathbf{x}) > P(\omega_j|\mathbf{x}) \quad \text{for all } j \neq i. \quad (20)$$

This is the same rule as in Eq. 6.

We saw in Fig. 2.2 some class-conditional probability densities and the posterior probabilities; Fig. 2.3 shows the likelihood ratio  $p(x|\omega_1)/p(x|\omega_2)$  for the same case. In general, this ratio can range between zero and infinity. The threshold value  $\theta_a$  marked is from the same prior probabilities but with zero-one loss function. Notice that this leads to the same decision boundaries as in Fig. 2.2, as it must. If we penalize mistakes in classifying  $\omega_1$  patterns as  $\omega_2$  more than the converse (i.e.,  $\lambda_{21} > \lambda_{12}$ ), then Eq. 17 leads to the threshold  $\theta_b$  marked. Note that the range of  $x$  values for which we classify a pattern as  $\omega_1$  gets smaller, as it should.




Figure 2.3: The likelihood ratio  $p(x|\omega_1)/p(x|\omega_2)$  for the distributions shown in Fig. 2.1. If we employ a zero-one or classification loss, our decision boundaries are determined by the threshold  $\theta_a$ . If our loss function penalizes miscategorizing  $\omega_2$  as  $\omega_1$  patterns more than the converse, (i.e.,  $\lambda_{12} > \lambda_{21}$ ), we get the larger threshold  $\theta_b$ , and hence  $\mathcal{R}_1$  becomes smaller.

### 2.3.1 \*Minimax Criterion

Sometimes we must design our classifier to perform well over a *range* of prior probabilities. For instance, in our fish categorization problem we can imagine that whereas the physical properties of lightness and width of each type of fish remain constant, the prior probabilities might vary widely and in an unpredictable way, or alternatively we want to use the classifier in a different plant where we do not know the prior probabilities. A reasonable approach is then to design our classifier so that the *worst* overall risk for any value of the priors is as small as possible — that is, minimize the maximum possible overall risk.

In order to understand this, we let  $\mathcal{R}_1$  denote that (as yet unknown) region in feature space where the classifier decides  $\omega_1$  and likewise for  $\mathcal{R}_2$  and  $\omega_2$ , and then write our overall risk Eq. 12 in terms of conditional risks:

$$\begin{aligned} R &= \int_{\mathcal{R}_1} [\lambda_{11}P(\omega_1) p(\mathbf{x}|\omega_1) + \lambda_{12}P(\omega_2) p(\mathbf{x}|\omega_2)] d\mathbf{x} \\ &+ \int_{\mathcal{R}_2} [\lambda_{21}P(\omega_1) p(\mathbf{x}|\omega_1) + \lambda_{22}P(\omega_2) p(\mathbf{x}|\omega_2)] d\mathbf{x}. \end{aligned} \quad (21)$$

We use the fact that  $P(\omega_2) = 1 - P(\omega_1)$  and that  $\int_{\mathcal{R}_1} p(\mathbf{x}|\omega_1) d\mathbf{x} = 1 - \int_{\mathcal{R}_2} p(\mathbf{x}|\omega_1) d\mathbf{x}$  to rewrite the risk as:

$$\begin{aligned} R(P(\omega_1)) &= \overbrace{\lambda_{22} + (\lambda_{12} - \lambda_{22}) \int_{\mathcal{R}_1} p(\mathbf{x}|\omega_2) d\mathbf{x}}^{= R_{mm}, \text{ minimax risk}} \\ &+ P(\omega_1) \underbrace{\left[ (\lambda_{11} - \lambda_{22}) - (\lambda_{21} - \lambda_{11}) \int_{\mathcal{R}_2} p(\mathbf{x}|\omega_1) d\mathbf{x} - (\lambda_{12} - \lambda_{22}) \int_{\mathcal{R}_1} p(\mathbf{x}|\omega_2) d\mathbf{x} \right]}_{= 0 \text{ for minimax solution}}. \end{aligned} \quad (22)$$

This equation shows that once the decision boundary is set (i.e.,  $\mathcal{R}_1$  and  $\mathcal{R}_2$  determined), the overall risk is linear in  $P(\omega_1)$ . If we can find a boundary such that the constant of proportionality is 0, then the risk is independent of priors. This is the *minimax solution*, and the *minimax risk*,  $R_{mm}$ , can be read from Eq. 22:

MINIMAX  
RISK

$$\begin{aligned} R_{mm} &= \lambda_{22} + (\lambda_{12} - \lambda_{22}) \int_{\mathcal{R}_1} p(\mathbf{x}|\omega_2) d\mathbf{x} \\ &= \lambda_{11} + (\lambda_{21} - \lambda_{11}) \int_{\mathcal{R}_2} p(\mathbf{x}|\omega_1) d\mathbf{x}. \end{aligned} \quad (23)$$

Figure 2.4 illustrates the approach. Briefly stated, we search for the prior for which the Bayes risk is *maximum*, the corresponding decision boundary gives the minimax solution. The value of the minimax risk,  $R_{mm}$ , is hence equal to the worst Bayes risk. In practice, finding the decision boundary for minimax risk may be difficult, particularly when distributions are complicated. Nevertheless, in some cases the boundary can be determined analytically (Problem 3).

The minimax criterion finds greater use in game theory than it does in traditional pattern recognition. In game theory, you have a hostile opponent who can be expected to take an action maximally detrimental to you. Thus it makes great sense for you to take an action (e.g., make a classification) where your costs — due to your opponent's subsequent actions — are minimized.




Figure 2.4: The curve at the bottom shows the minimum (Bayes) error as a function of prior probability  $P(\omega_1)$  in a two-category classification problem of fixed distributions. For each value of the priors (e.g.,  $P(\omega_1) = 0.25$ ) there is a corresponding optimal decision boundary and associated Bayes error rate. For any (fixed) such boundary, if the priors are then changed, the probability of error will change as a linear function of  $P(\omega_1)$  (shown by the dashed line). The maximum such error will occur at an extreme value of the prior, here at  $P(\omega_1) = 1$ . To minimize the maximum of such error, we should design our decision boundary for the maximum Bayes error (here  $P(\omega_1) = 0.6$ ), and thus the error will not change as a function of prior, as shown by the solid red horizontal line.

### 2.3.2 \*Neyman-Pearson Criterion

In some problems, we may wish to minimize the overall risk subject to a constraint; for instance, we might wish to minimize the total risk subject to the constraint  $\int R(\alpha_i|\mathbf{x}) d\mathbf{x} < \text{constant}$  for some particular  $i$ . Such a constraint might arise when there is a fixed resource that accompanies one particular action  $\alpha_i$ , or when we must not misclassify pattern from a particular state of nature  $\omega_i$  at more than some limited frequency. For instance, in our fish example, there might be some government regulation that we must not misclassify more than 1% of salmon as sea bass. We might then seek a decision that minimizes the chance of classifying a sea bass as a salmon subject to this condition.

We generally satisfy such a *Neyman-Pearson criterion* by adjusting decision boundaries numerically. However, for Gaussian and some other distributions, Neyman-Pearson solutions can be found analytically (Problems 5 & 6). We shall have cause to mention Neyman-Pearson criteria again in Sect. 2.8.3 on operating characteristics.

## 2.4 Classifiers, Discriminant Functions and Decision Surfaces

### 2.4.1 The Multi-Category Case

There are many different ways to represent pattern classifiers. One of the most useful is in terms of a set of *discriminant functions*  $g_i(\mathbf{x})$ ,  $i = 1, \dots, c$ . The classifier is said to assign a feature vector  $\mathbf{x}$  to class  $\omega_i$  if

$$g_i(\mathbf{x}) > g_j(\mathbf{x}) \quad \text{for all } j \neq i. \quad (24)$$

Thus, the classifier is viewed as a network or machine that computes  $c$  discriminant functions and selects the category corresponding to the largest discriminant. A network representation of a classifier is illustrated in Fig. 2.5.




Figure 2.5: The functional structure of a general statistical pattern classifier which includes  $d$  inputs and  $c$  discriminant functions  $g_i(\mathbf{x})$ . A subsequent step determines which of the discriminant values is the maximum, and categorizes the input pattern accordingly. The arrows show the direction of the flow of information, though frequently the arrows are omitted when the direction of flow is self-evident.

A Bayes classifier is easily and naturally represented in this way. For the general case with risks, we can let  $g_i(\mathbf{x}) = -R(\alpha_i|\mathbf{x})$ , since the maximum discriminant function will then correspond to the minimum conditional risk. For the minimum-error-rate case, we can simplify things further by taking  $g_i(\mathbf{x}) = P(\omega_i|\mathbf{x})$ , so that the maximum discriminant function corresponds to the maximum posterior probability.

Clearly, the choice of discriminant functions is not unique. We can always multiply all the discriminant functions by the same positive constant or shift them by the same additive constant without influencing the decision. More generally, if we replace every  $g_i(\mathbf{x})$  by  $f(g_i(\mathbf{x}))$ , where  $f(\cdot)$  is a monotonically increasing function, the resulting classification is unchanged. This observation can lead to significant analytical and computational simplifications. In particular, for minimum-error-rate classification, any of the following choices gives identical classification results, but some can be much simpler to understand or to compute than others:

$$g_i(\mathbf{x}) = P(\omega_i|\mathbf{x}) = \frac{p(\mathbf{x}|\omega_i)P(\omega_i)}{\sum_{j=1}^c p(\mathbf{x}|\omega_j)P(\omega_j)} \quad (25)$$

$$g_i(\mathbf{x}) = p(\mathbf{x}|\omega_i)P(\omega_i) \quad (26)$$

$$g_i(\mathbf{x}) = \ln p(\mathbf{x}|\omega_i) + \ln P(\omega_i), \quad (27)$$

where  $\ln$  denotes natural logarithm.

DECISION  
REGION

Even though the discriminant functions can be written in a variety of forms, the decision rules are equivalent. The effect of any decision rule is to divide the feature space into  $c$  *decision regions*,  $\mathcal{R}_1, \dots, \mathcal{R}_c$ . If  $g_i(\mathbf{x}) > g_j(\mathbf{x})$  for all  $j \neq i$ , then  $\mathbf{x}$  is in  $\mathcal{R}_i$ , and the decision rule calls for us to assign  $\mathbf{x}$  to  $\omega_i$ . The regions are separated by *decision boundaries*, surfaces in feature space where ties occur among the largest discriminant functions (Fig. 2.6).




Figure 2.6: In this two-dimensional two-category classifier, the probability densities are Gaussian (with  $1/e$  ellipses shown), the decision boundary consists of two hyperbolas, and thus the decision region  $\mathcal{R}_2$  is not simply connected.

### 2.4.2 The Two-Category Case

While the two-category case is just a special instance of the multicategory case, it has traditionally received separate treatment. Indeed, a classifier that places a pattern in

one of only two categories has a special name — a *dichotomizer*.<sup>\*</sup> Instead of using two discriminant functions  $g_1$  and  $g_2$  and assigning  $\mathbf{x}$  to  $\omega_1$  if  $g_1 > g_2$ , it is more common to define a single discriminant function

DICHOTOMIZER

$$g(\mathbf{x}) \equiv g_1(\mathbf{x}) - g_2(\mathbf{x}), \quad (28)$$

and to use the following decision rule: Decide  $\omega_1$  if  $g(\mathbf{x}) > 0$ ; otherwise decide  $\omega_2$ . Thus, a dichotomizer can be viewed as a machine that computes a single discriminant function  $g(\mathbf{x})$ , and classifies  $\mathbf{x}$  according to the algebraic sign of the result. Of the various forms in which the minimum-error-rate discriminant function can be written, the following two (derived from Eqs. 25 & 27) are particularly convenient:

$$g(\mathbf{x}) = P(\omega_1|\mathbf{x}) - P(\omega_2|\mathbf{x}) \quad (29)$$

$$g(\mathbf{x}) = \ln \frac{p(\mathbf{x}|\omega_1)}{p(\mathbf{x}|\omega_2)} + \ln \frac{P(\omega_1)}{P(\omega_2)}. \quad (30)$$

## 2.5 The Normal Density

The structure of a Bayes classifier is determined by the conditional densities  $p(\mathbf{x}|\omega_i)$  as well as by the prior probabilities. Of the various density functions that have been investigated, none has received more attention than the multivariate normal or Gaussian density. To a large extent this attention is due to its analytical tractability. However the multivariate normal density is also an appropriate model for an important situation, viz., the case where the feature vectors  $\mathbf{x}$  for a given class  $\omega_i$  are continuous valued, randomly corrupted versions of a single typical or prototype vector  $\boldsymbol{\mu}_i$ . In this section we provide a brief exposition of the multivariate normal density, focusing on the properties of greatest interest for classification problems.

First, recall the definition of the *expected value* of a scalar function  $f(x)$ , defined for some density  $p(x)$ :

EXPECTATION

$$\mathcal{E}[f(x)] \equiv \int_{-\infty}^{\infty} f(x)p(x)dx. \quad (31)$$

If we have samples in a set  $\mathcal{D}$  from a discrete distribution, we must sum over all samples as

$$\mathcal{E}[f(x)] = \sum_{x \in \mathcal{D}} f(x)P(x), \quad (32)$$

where  $P(x)$  is the probability mass at  $x$ . We shall often have call to calculate expected values — by these and analogous equations defined in higher dimensions (see Appendix Secs. ??, ?? & ??).<sup>\*</sup>

<sup>\*</sup> A classifier for more than two categories is called a polychotomizer.

<sup>\*</sup> We will often use somewhat loose engineering terminology and refer to a single point as a “sample.” Statisticians, though, always refer to a sample as a *collection* of points, and discuss “a sample of size  $n$ .” When taken in context, there are rarely ambiguities in such usage.

### 2.5.1 Univariate Density

We begin with the continuous univariate normal or Gaussian density,

$$p(x) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left[ -\frac{1}{2} \left( \frac{x - \mu}{\sigma} \right)^2 \right], \quad (33)$$

for which the *expected value* of  $x$  (an average, here taken over the feature space) is

$$\mu \equiv \mathcal{E}[x] = \int_{-\infty}^{\infty} xp(x) dx, \quad (34)$$

VARIANCE and where the expected squared deviation or *variance* is

$$\sigma^2 \equiv \mathcal{E}[(x - \mu)^2] = \int_{-\infty}^{\infty} (x - \mu)^2 p(x) dx. \quad (35)$$

MEAN

The univariate normal density is completely specified by two parameters: its mean  $\mu$  and variance  $\sigma^2$ . For simplicity, we often abbreviate Eq. 33 by writing  $p(x) \sim N(\mu, \sigma^2)$  to say that  $x$  is distributed normally with mean  $\mu$  and variance  $\sigma^2$ . Samples from normal distributions tend to cluster about the mean, with a spread related to the standard deviation  $\sigma$  (Fig. 2.7).




Figure 2.7: A univariate normal distribution has roughly 95% of its area in the range  $|x - \mu| \leq 2\sigma$ , as shown. The peak of the distribution has value  $p(\mu) = 1/\sqrt{2\pi}\sigma$ .

ENTROPY

There is a deep relationship between the normal distribution and *entropy*. We shall consider entropy in greater detail in Chap. ??, but for now we merely state that the entropy of a distribution is given by

$$H(p(x)) = - \int p(x) \ln p(x) dx, \quad (36)$$

NAT

and measured in *nats*. If a  $\log_2$  is used instead, the unit is the *bit*. The entropy is a non-negative quantity that describes the fundamental uncertainty in the values of points

BIT

selected randomly from a distribution. It can be shown that the normal distribution has the maximum entropy of all distributions having a given mean and variance (Problem 20). Moreover, as stated by the *Central Limit Theorem*, the aggregate effect of a large number of small, independent random disturbances will lead to a Gaussian distribution (Computer exercise ??). Because many patterns — from fish to handwritten characters to some speech sounds — can be viewed as some ideal or prototype pattern corrupted by a large number of random processes, the Gaussian is often a good model for the actual probability distribution.

CENTRAL  
LIMIT  
THEOREM

## 2.5.2 Multivariate Density

The general multivariate normal density in  $d$  dimensions is written as

$$p(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} |\mathbf{\Sigma}|^{1/2}} \exp \left[ -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^t \mathbf{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right], \quad (37)$$

where  $\mathbf{x}$  is a  $d$ -component column vector,  $\boldsymbol{\mu}$  is the  $d$ -component *mean vector*,  $\mathbf{\Sigma}$  is the  $d$ -by- $d$  *covariance matrix*,  $|\mathbf{\Sigma}|$  and  $\mathbf{\Sigma}^{-1}$  are its determinant and inverse, respectively, and  $(\mathbf{x} - \boldsymbol{\mu})^t$  is the transpose of  $\mathbf{x} - \boldsymbol{\mu}$ .<sup>\*</sup> Our notation for the *inner product* is

COVARIANCE  
MATRIX

$$\mathbf{a}^t \mathbf{b} = \sum_{i=1}^d a_i b_i, \quad (38)$$

INNER  
PRODUCT

and often called a *dot product*.

For simplicity, we often abbreviate Eq. 37 as  $p(\mathbf{x}) \sim N(\boldsymbol{\mu}, \mathbf{\Sigma})$ . Formally, we have

$$\boldsymbol{\mu} \equiv \mathcal{E}[\mathbf{x}] = \int \mathbf{x} p(\mathbf{x}) d\mathbf{x} \quad (39)$$

and

$$\mathbf{\Sigma} \equiv \mathcal{E}[(\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^t] = \int (\mathbf{x} - \boldsymbol{\mu})(\mathbf{x} - \boldsymbol{\mu})^t p(\mathbf{x}) d\mathbf{x}, \quad (40)$$

where the expected value of a vector or a matrix is found by taking the expected values of its components. In other words, if  $x_i$  is the  $i$ th component of  $\mathbf{x}$ ,  $\mu_i$  the  $i$ th component of  $\boldsymbol{\mu}$ , and  $\sigma_{ij}$  the  $ij$ th component of  $\mathbf{\Sigma}$ , then

$$\mu_i = \mathcal{E}[x_i] \quad (41)$$

and

$$\sigma_{ij} = \mathcal{E}[(x_i - \mu_i)(x_j - \mu_j)]. \quad (42)$$

The covariance matrix  $\mathbf{\Sigma}$  is always symmetric and positive semidefinite. We shall restrict our attention to the case in which  $\mathbf{\Sigma}$  is positive definite, so that the determinant of  $\mathbf{\Sigma}$  is strictly positive.<sup>†</sup> The diagonal elements  $\sigma_{ii}$  are the variances of the respective  $x_i$  (i.e.,  $\sigma_i^2$ ), and the off-diagonal elements  $\sigma_{ij}$  are the *covariances* of  $x_i$  and  $x_j$ . We would expect a positive covariance for the length and weight features of a population of fish, for instance. If  $x_i$  and  $x_j$  are *statistically independent*,  $\sigma_{ij} = 0$ . If

COVARIANCE

STATISTICAL  
INDEPENDENCE

<sup>\*</sup> The mathematical expressions for the multivariate normal density are greatly simplified by employing the concepts and notation of linear algebra. Readers who are unsure of our notation or who would like to review linear algebra should see Appendix ??.

<sup>†</sup> If sample vectors are drawn from a linear subspace,  $|\mathbf{\Sigma}| = 0$  and  $p(\mathbf{x})$  is degenerate. This occurs, for example, when one component of  $\mathbf{x}$  has zero variance, or when two components are identical or multiples of one another.



all the off-diagonal elements are zero,  $p(\mathbf{x})$  reduces to the product of the univariate normal densities for the components of  $\mathbf{x}$ .

Linear combinations of jointly normally distributed random variables, independent or not, are normally distributed. In particular, if  $\mathbf{A}$  is a  $d$ -by- $k$  matrix and  $\mathbf{y} = \mathbf{A}^t \mathbf{x}$  is a  $k$ -component vector, then  $p(\mathbf{y}) \sim N(\mathbf{A}^t \boldsymbol{\mu}, \mathbf{A}^t \boldsymbol{\Sigma} \mathbf{A})$ , as illustrated in Fig. 2.8. In the special case where  $k = 1$  and  $\mathbf{A}$  is a unit-length vector  $\mathbf{a}$ ,  $y = \mathbf{a}^t \mathbf{x}$  is a scalar that represents the projection of  $\mathbf{x}$  onto a line in the direction of  $\mathbf{a}$ ; in that case  $\mathbf{a}^t \boldsymbol{\Sigma} \mathbf{a}$  is the variance of the projection of  $\mathbf{x}$  onto  $\mathbf{a}$ . In general then, knowledge of the covariance matrix allows us to calculate the dispersion of the data in any direction, or in any subspace.

WHITENING  
TRANSFORM

It is sometimes convenient to perform a coordinate transformation that converts an arbitrary multivariate normal distribution into a spherical one, i.e., one having a covariance matrix proportional to the identity matrix  $\mathbf{I}$ . If we define  $\Phi$  to be the matrix whose columns are the orthonormal eigenvectors of  $\boldsymbol{\Sigma}$ , and  $\Lambda$  the diagonal matrix of the corresponding eigenvalues, then the transformation  $\mathbf{A}_w = \Phi \Lambda^{-1/2}$  applied to the coordinates insures that the transformed distribution has covariance matrix equal to the identity matrix. In signal processing, the transform  $\mathbf{A}_w$  is called a *whitening* transformation, since it makes the spectrum of eigenvectors of the transformed distribution uniform.

The multivariate normal density is completely specified by  $d + d(d + 1)/2$  parameters — the elements of the mean vector  $\boldsymbol{\mu}$  and the independent elements of the covariance matrix  $\boldsymbol{\Sigma}$ . Samples drawn from a normal population tend to fall in a single cloud or cluster (Fig. 2.9); the center of the cluster is determined by the mean vector, and the shape of the cluster is determined by the covariance matrix. It follows from Eq. 37 that the loci of points of constant density are hyperellipsoids for which the quadratic form  $(\mathbf{x} - \boldsymbol{\mu})^t \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu})$  is constant. The principal axes of these hyperellipsoids are given by the eigenvectors of  $\boldsymbol{\Sigma}$  (described by  $\Phi$ ); the eigenvalues (described by  $\Lambda$ ) determine the lengths of these axes. The quantity

$$r^2 = (\mathbf{x} - \boldsymbol{\mu})^t \boldsymbol{\Sigma}^{-1} (\mathbf{x} - \boldsymbol{\mu}) \quad (43)$$

MAHALANOBIS  
DISTANCE

is sometimes called the squared *Mahalanobis distance* from  $\mathbf{x}$  to  $\boldsymbol{\mu}$ . Thus, the contours of constant density are hyperellipsoids of constant Mahalanobis distance to  $\boldsymbol{\mu}$  and the volume of these hyperellipsoids measures the scatter of the samples about the mean. It can be shown (Problems 15 & 16) that the volume of the hyperellipsoid corresponding to a Mahalanobis distance  $r$  is given by

$$V = V_d |\boldsymbol{\Sigma}|^{1/2} r^d, \quad (44)$$

where  $V_d$  is the volume of a  $d$ -dimensional unit hypersphere:

$$V_d = \begin{cases} \pi^{d/2} / (d/2)! & d \text{ even} \\ 2^d \pi^{(d-1)/2} (\frac{d-1}{2})! / (d)! & d \text{ odd.} \end{cases} \quad (45)$$

Thus, for a given dimensionality, the scatter of the samples varies directly with  $|\boldsymbol{\Sigma}|^{1/2}$  (Problem 17).




Figure 2.8: The action of a linear transformation on the feature space will convert an arbitrary normal distribution into another normal distribution. One transformation,  $\mathbf{A}$ , takes the source distribution into distribution  $N(\mathbf{A}^t \boldsymbol{\mu}, \mathbf{A}^t \boldsymbol{\Sigma} \mathbf{A})$ . Another linear transformation — a projection  $\mathbf{P}$  onto line  $\mathbf{a}$  — leads to  $N(\mu, \sigma^2)$  measured along  $\mathbf{a}$ . While the transforms yield distributions in a different space, we show them superimposed on the original  $x_1 - x_2$  space. A whitening transform leads to a circularly symmetric Gaussian, here shown displaced.

## 2.6 Discriminant Functions for the Normal Density

In Sect. 2.4.1 we saw that the minimum-error-rate classification can be achieved by use of the discriminant functions

$$g_i(\mathbf{x}) = \ln p(\mathbf{x}|\omega_i) + \ln P(\omega_i). \quad (46)$$

This expression can be readily evaluated if the densities  $p(\mathbf{x}|\omega_i)$  are multivariate normal, i.e., if  $p(\mathbf{x}|\omega_i) \sim N(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$ . In this case, then, from Eq. 37 we have

$$g_i(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_i)^t \boldsymbol{\Sigma}_i^{-1} (\mathbf{x} - \boldsymbol{\mu}_i) - \frac{d}{2} \ln 2\pi - \frac{1}{2} \ln |\boldsymbol{\Sigma}_i| + \ln P(\omega_i). \quad (47)$$

Let us examine the discriminant function and resulting classification for a number of special cases.

### 2.6.1 Case 1: $\boldsymbol{\Sigma}_i = \sigma^2 \mathbf{I}$

The simplest case occurs when the features are statistically independent, and when each feature has the same variance,  $\sigma^2$ . In this case the covariance matrix is diagonal,




Figure 2.9: Samples drawn from a two-dimensional Gaussian lie in a cloud centered on the mean  $\boldsymbol{\mu}$ . The red ellipses show lines of equal probability density of the Gaussian.

being merely  $\sigma^2$  times the identity matrix  $\mathbf{I}$ . Geometrically, this corresponds to the situation in which the samples fall in equal-size hyperspherical clusters, the cluster for the  $i$ th class being centered about the mean vector  $\boldsymbol{\mu}_i$ . The computation of the determinant and the inverse of  $\boldsymbol{\Sigma}_i$  is particularly easy:  $|\boldsymbol{\Sigma}_i| = \sigma^{2d}$  and  $\boldsymbol{\Sigma}_i^{-1} = (1/\sigma^2)\mathbf{I}$ . Since both  $|\boldsymbol{\Sigma}_i|$  and the  $(d/2) \ln 2\pi$  term in Eq. 47 are independent of  $i$ , they are unimportant additive constants that can be ignored. Thus we obtain the simple discriminant functions

$$g_i(\mathbf{x}) = -\frac{\|\mathbf{x} - \boldsymbol{\mu}_i\|^2}{2\sigma^2} + \ln P(\omega_i), \quad (48)$$

EUCLIDEAN  
NORM

where  $\|\cdot\|$  is the *Euclidean norm*, that is,

$$\|\mathbf{x} - \boldsymbol{\mu}_i\|^2 = (\mathbf{x} - \boldsymbol{\mu}_i)^t(\mathbf{x} - \boldsymbol{\mu}_i). \quad (49)$$

If the prior probabilities are not equal, then Eq. 48 shows that the squared distance  $\|\mathbf{x} - \boldsymbol{\mu}\|^2$  must be normalized by the variance  $\sigma^2$  and offset by adding  $\ln P(\omega_i)$ ; thus, if  $\mathbf{x}$  is equally near two different mean vectors, the optimal decision will favor the a priori more likely category.

Regardless of whether the prior probabilities are equal or not, it is not actually necessary to compute distances. Expansion of the quadratic form  $(\mathbf{x} - \boldsymbol{\mu}_i)^t(\mathbf{x} - \boldsymbol{\mu}_i)$  yields

$$g_i(\mathbf{x}) = -\frac{1}{2\sigma^2}[\mathbf{x}^t\mathbf{x} - 2\boldsymbol{\mu}_i^t\mathbf{x} + \boldsymbol{\mu}_i^t\boldsymbol{\mu}_i] + \ln P(\omega_i), \quad (50)$$

which appears to be a quadratic function of  $\mathbf{x}$ . However, the quadratic term  $\mathbf{x}^t\mathbf{x}$  is the same for all  $i$ , making it an ignorable additive constant. Thus, we obtain the equivalent *linear discriminant functions*

LINEAR  
DISCRIMINANT

$$g_i(\mathbf{x}) = \mathbf{w}_i^t\mathbf{x} + w_{i0}, \quad (51)$$

where




Figure 2.10: If the covariances of two distributions are equal and proportional to the identity matrix, then the distributions are spherical in  $d$  dimensions, and the boundary is a generalized hyperplane of  $d - 1$  dimensions, perpendicular to the line separating the means. In these 1-, 2-, and 3-dimensional examples, we indicate  $p(\mathbf{x}|\omega_i)$  and the boundaries for the case  $P(\omega_1) = P(\omega_2)$ . In the 3-dimensional case, the grid plane separates  $\mathcal{R}_1$  from  $\mathcal{R}_2$ .

$$\mathbf{w}_i = \frac{1}{\sigma^2} \boldsymbol{\mu}_i \quad (52)$$

and

$$w_{i0} = \frac{-1}{2\sigma^2} \boldsymbol{\mu}_i^t \boldsymbol{\mu}_i + \ln P(\omega_i). \quad (53)$$

We call  $w_{i0}$  the *threshold* or *bias* in the  $i$ th direction.

A classifier that uses linear discriminant functions is called a *linear machine*. This kind of classifier has many interesting theoretical properties, some of which will be discussed in detail in Chap. ???. At this point we merely note that the decision surfaces for a linear machine are pieces of hyperplanes defined by the linear equations  $g_i(\mathbf{x}) = g_j(\mathbf{x})$  for the two categories with the highest posterior probabilities. For our particular case, this equation can be written as

$$\mathbf{w}^t(\mathbf{x} - \mathbf{x}_0) = 0, \quad (54)$$

where

$$\mathbf{w} = \boldsymbol{\mu}_i - \boldsymbol{\mu}_j \quad (55)$$

and

$$\mathbf{x}_0 = \frac{1}{2}(\boldsymbol{\mu}_i + \boldsymbol{\mu}_j) - \frac{\sigma^2}{\|\boldsymbol{\mu}_i - \boldsymbol{\mu}_j\|^2} \ln \frac{P(\omega_i)}{P(\omega_j)} (\boldsymbol{\mu}_i - \boldsymbol{\mu}_j). \quad (56)$$

This equation defines a hyperplane through the point  $\mathbf{x}_0$  and orthogonal to the vector  $\mathbf{w}$ . Since  $\mathbf{w} = \boldsymbol{\mu}_i - \boldsymbol{\mu}_j$ , the hyperplane separating  $\mathcal{R}_i$  and  $\mathcal{R}_j$  is orthogonal to the line linking the means. If  $P(\omega_i) = P(\omega_j)$ , the second term on the right of Eq. 56 vanishes, and thus the point  $\mathbf{x}_0$  is halfway between the means, and the hyperplane is the perpendicular bisector of the line between the means (Fig. 2.11). If  $P(\omega_i) \neq P(\omega_j)$ , the point  $\mathbf{x}_0$  shifts away from the more likely mean. Note, however, that if the variance

THRESHOLD

BIAS

LINEAR  
MACHINE




Figure 2.11: As the priors are changed, the decision boundary shifts; for sufficiently disparate priors the boundary will not lie between the means of these 1-, 2- and 3-dimensional spherical Gaussian distributions.

$\sigma^2$  is small relative to the squared distance  $\|\mu_i - \mu_j\|$ , then the position of the decision boundary is relatively insensitive to the exact values of the prior probabilities.

If the prior probabilities  $P(\omega_i)$  are the same for all  $c$  classes, then the  $\ln P(\omega_i)$  term becomes another unimportant additive constant that can be ignored. When this happens, the optimum decision rule can be stated very simply: to classify a feature vector  $\mathbf{x}$ , measure the Euclidean distance  $\|\mathbf{x} - \mu_i\|$  from each  $\mathbf{x}$  to each of the  $c$  mean vectors, and assign  $\mathbf{x}$  to the category of the nearest mean. Such a classifier is called a *minimum distance classifier*. If each mean vector is thought of as being an ideal prototype or template for patterns in its class, then this is essentially a *template-matching* procedure (Fig. 2.10), a technique we will consider again in Chap. ?? Sect. ?? on the nearest-neighbor algorithm.

MINIMUM  
DISTANCE  
CLASSIFIER

TEMPLATE-  
MATCHING

### 2.6.2 Case 2: $\Sigma_i = \Sigma$

Another simple case arises when the covariance matrices for all of the classes are identical but otherwise arbitrary. Geometrically, this corresponds to the situation in which the samples fall in hyperellipsoidal clusters of equal size and shape, the cluster for the  $i$ th class being centered about the mean vector  $\boldsymbol{\mu}_i$ . Since both  $|\Sigma_i|$  and the  $(d/2) \ln 2\pi$  term in Eq. 47 are independent of  $i$ , they can be ignored as superfluous additive constants. This simplification leads to the discriminant functions

$$g_i(\mathbf{x}) = -\frac{1}{2}(\mathbf{x} - \boldsymbol{\mu}_i)^t \Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}_i) + \ln P(\omega_i). \quad (57)$$

If the prior probabilities  $P(\omega_i)$  are the same for all  $c$  classes, then the  $\ln P(\omega_i)$  term can be ignored. In this case, the optimal decision rule can once again be stated very simply: to classify a feature vector  $\mathbf{x}$ , measure the squared Mahalanobis distance  $(\mathbf{x} - \boldsymbol{\mu}_i)^t \Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}_i)$  from  $\mathbf{x}$  to each of the  $c$  mean vectors, and assign  $\mathbf{x}$  to the category of the nearest mean. As before, unequal prior probabilities bias the decision in favor of the a priori more likely category.

Expansion of the quadratic form  $(\mathbf{x} - \boldsymbol{\mu}_i)^t \Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}_i)$  results in a sum involving a quadratic term  $\mathbf{x}^t \Sigma^{-1} \mathbf{x}$  which here is independent of  $i$ . After this term is dropped from Eq. 57, the resulting discriminant functions are again linear:

$$g_i(\mathbf{x}) = \mathbf{w}_i^t \mathbf{x} + w_{i0}, \quad (58)$$

where

$$\mathbf{w}_i = \Sigma^{-1} \boldsymbol{\mu}_i \quad (59)$$

and

$$w_{i0} = -\frac{1}{2} \boldsymbol{\mu}_i^t \Sigma^{-1} \boldsymbol{\mu}_i + \ln P(\omega_i). \quad (60)$$

Since the discriminants are linear, the resulting decision boundaries are again hyperplanes (Fig. 2.10). If  $\mathcal{R}_i$  and  $\mathcal{R}_j$  are contiguous, the boundary between them has the equation

$$\mathbf{w}^t (\mathbf{x} - \mathbf{x}_0) = 0, \quad (61)$$

where

$$\mathbf{w} = \Sigma^{-1}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j) \quad (62)$$

and

$$\mathbf{x}_0 = \frac{1}{2}(\boldsymbol{\mu}_i + \boldsymbol{\mu}_j) - \frac{\ln [P(\omega_i)/P(\omega_j)]}{(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)^t \Sigma^{-1}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j). \quad (63)$$

Since  $\mathbf{w} = \Sigma^{-1}(\boldsymbol{\mu}_i - \boldsymbol{\mu}_j)$  is generally not in the direction of  $\boldsymbol{\mu}_i - \boldsymbol{\mu}_j$ , the hyperplane separating  $\mathcal{R}_i$  and  $\mathcal{R}_j$  is generally not orthogonal to the line between the means. However, it does intersect that line at the point  $\mathbf{x}_0$  which is halfway between the means if the prior probabilities are equal. If the prior probabilities are not equal, the optimal boundary hyperplane is shifted away from the more likely mean (Fig. 2.12). As before, with sufficient bias the decision plane need not lie between the two mean vectors.




Figure 2.12: Probability densities (indicated by the surfaces in two dimensions and ellipsoidal surfaces in three dimensions) and decision regions for equal but asymmetric Gaussian distributions. The decision hyperplanes need not be perpendicular to the line connecting the means.

### 2.6.3 Case 3: $\Sigma_i = \text{arbitrary}$

In the general multivariate normal case, the covariance matrices are different for each category. The only term that can be dropped from Eq. 47 is the  $(d/2) \ln 2\pi$  term, and the resulting discriminant functions are inherently quadratic:

$$g_i(\mathbf{x}) = \mathbf{x}^t \mathbf{W}_i \mathbf{x} + \mathbf{w}_i^t \mathbf{x} + w_{i0}, \quad (64)$$

where

$$\mathbf{W}_i = -\frac{1}{2} \Sigma_i^{-1}, \quad (65)$$

$$\mathbf{w}_i = \Sigma_i^{-1} \boldsymbol{\mu}_i \quad (66)$$

and

$$w_{i0} = -\frac{1}{2} \boldsymbol{\mu}_i^t \Sigma_i^{-1} \boldsymbol{\mu}_i - \frac{1}{2} \ln |\Sigma_i| + \ln P(\omega_i). \quad (67)$$

The decision surfaces are *hyperquadrics*, and can assume any of the general forms — hyperplanes, pairs of hyperplanes, hyperspheres, hyperellipsoids, hyperparaboloids, and hyperhyperboloids of various types (Problem 29). Even in one dimension, for arbitrary covariance the decision regions need not be simply connected (Fig. 2.13). The two- and three-dimensional examples in Fig. 2.14 & 2.15 indicate how these different forms can arise. These variances are indicated by the contours of constant probability density.

HYPER-  
QUADRIC

The extension of these results to more than two categories is straightforward though we need to keep clear which two of the total  $c$  categories are responsible for any boundary segment. Figure 2.16 shows the decision surfaces for a four-category case made up of Gaussian distributions. Of course, if the distributions are more complicated, the decision regions can be even more complex, though the same underlying theory holds there too.




Figure 2.13: Non-simply connected decision regions can arise in one dimensions for Gaussians having unequal variance.






Figure 2.14: Arbitrary Gaussian distributions lead to Bayes decision boundaries that are general hyperquadrics. Conversely, given any hyperquadratic, one can find two Gaussian distributions whose Bayes decision boundary is that hyperquadratic.




Figure 2.15: Arbitrary three-dimensional Gaussian distributions yield Bayes decision boundaries that are two-dimensional hyperquadrics. There are even degenerate cases in which the decision boundary is a line.




Figure 2.16: The decision regions for four normal distributions. Even with such a low number of categories, the shapes of the boundary regions can be rather complex.

**Example 1: Decision regions for two-dimensional Gaussian data**

To clarify these ideas, we explicitly calculate the decision boundary for the two-category two-dimensional data in the Example figure. Let  $\omega_1$  be the set of the four black points, and  $\omega_2$  the red points. Although we will spend much of the next chapter understanding how to estimate the parameters of our distributions, for now we simply assume that we need merely calculate the means and covariances by the discrete versions of Eqs. 39 & 40; they are found to be:

$$\boldsymbol{\mu}_1 = \begin{bmatrix} 3 \\ 6 \end{bmatrix}; \quad \boldsymbol{\Sigma}_1 = \begin{pmatrix} 1/2 & 0 \\ 0 & 2 \end{pmatrix} \text{ and } \boldsymbol{\mu}_2 = \begin{bmatrix} 3 \\ -2 \end{bmatrix}; \quad \boldsymbol{\Sigma}_2 = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}.$$


The inverse matrices are then,

$$\boldsymbol{\Sigma}_1^{-1} = \begin{pmatrix} 2 & 0 \\ 0 & 1/2 \end{pmatrix} \quad \text{and} \quad \boldsymbol{\Sigma}_2^{-1} = \begin{pmatrix} 1/2 & 0 \\ 0 & 1/2 \end{pmatrix}.$$

We assume equal prior probabilities,  $P(\omega_1) = P(\omega_2) = 0.5$ , and substitute these into the general form for a discriminant, Eqs. 64 – 67, setting  $g_1(\mathbf{x}) = g_2(\mathbf{x})$  to obtain the decision boundary:

$$x_2 = 3.514 - 1.125x_1 + 0.1875x_1^2.$$

This equation describes a parabola with vertex at  $(\frac{3}{1.83})$ . Note that despite the fact that the variance in the data along the  $x_2$  direction for both distributions is the same, the decision boundary does not pass through the point  $(\frac{3}{2})$ , midway between the means, as we might have naively guessed. This is because for the  $\omega_1$  distribution, the probability distribution is “squeezed” in the  $x_1$ -direction more so than for the  $\omega_2$  distribution. Because the overall prior probabilities are the same (i.e., the integral over space of the probability density), the distribution is increased along the  $x_2$  direction (relative to that for the  $\omega_2$  distribution). Thus the decision boundary lies slightly lower than the point midway between the two means, as can be seen in the decision boundary.



The computed Bayes decision boundary for two Gaussian distributions, each based on four data points.

## 2.7 Error Probabilities and Integrals

We can obtain additional insight into the operation of a general classifier — Bayes or otherwise — if we consider the sources of its error. Consider first the two-category case, and suppose the dichotomizer has divided the space into two regions  $\mathcal{R}_1$  and  $\mathcal{R}_2$  in a possibly non-optimal way. There are two ways in which a classification error can occur; either an observation  $\mathbf{x}$  falls in  $\mathcal{R}_2$  and the true state of nature is  $\omega_1$ , or  $\mathbf{x}$  falls in  $\mathcal{R}_1$  and the true state of nature is  $\omega_2$ . Since these events are mutually exclusive and exhaustive, the probability of error is

$$\begin{aligned} P(\text{error}) &= P(\mathbf{x} \in \mathcal{R}_2, \omega_1) + P(\mathbf{x} \in \mathcal{R}_1, \omega_2) \\ &= P(\mathbf{x} \in \mathcal{R}_2 | \omega_1)P(\omega_1) + P(\mathbf{x} \in \mathcal{R}_1 | \omega_2)P(\omega_2) \\ &= \int_{\mathcal{R}_2} p(\mathbf{x} | \omega_1)P(\omega_1) d\mathbf{x} + \int_{\mathcal{R}_1} p(\mathbf{x} | \omega_2)P(\omega_2) d\mathbf{x}. \end{aligned} \quad (68)$$

This result is illustrated in the one-dimensional case in Fig. 2.17. The two integrals in Eq. 68 represent the pink and the gray areas in the tails of the functions  $p(\mathbf{x} | \omega_i)P(\omega_i)$ . Because the decision point  $x^*$  (and hence the regions  $\mathcal{R}_1$  and  $\mathcal{R}_2$ ) were chosen arbitrarily for that figure, the probability of error is not as small as it might be. In particular, the triangular area marked “reducible error” can be eliminated if the decision boundary is moved to  $x_B$ . This is the Bayes optimal decision boundary and gives the lowest probability of error. In general, if  $p(\mathbf{x} | \omega_1)P(\omega_1) > p(\mathbf{x} | \omega_2)P(\omega_2)$ , it is advantageous to classify  $\mathbf{x}$  as in  $\mathcal{R}_1$  so that the smaller quantity will contribute to the error integral; this is exactly what the Bayes decision rule achieves.




Figure 2.17: Components of the probability of error for equal priors and (non-optimal) decision point  $x^*$ . The pink area corresponds to the probability of errors for deciding  $\omega_1$  when the state of nature is in fact  $\omega_2$ ; the gray area represents the converse, as given in Eq. 68. If the decision boundary is instead at the point of equal posterior probabilities,  $x_B$ , then this reducible error is eliminated and the total shaded area is the minimum possible — this is the Bayes decision and gives the Bayes error rate.

In the multicategory case, there are more ways to be wrong than to be right, and it is simpler to compute the probability of being correct. Clearly

$$P(\text{correct}) = \sum_{i=1}^c P(\mathbf{x} \in \mathcal{R}_i, \omega_i)$$

$$\begin{aligned}
&= \sum_{i=1}^c P(\mathbf{x} \in \mathcal{R}_i | \omega_i) P(\omega_i) \\
&= \sum_{i=1}^c \int_{\mathcal{R}_i} p(\mathbf{x} | \omega_i) P(\omega_i) d\mathbf{x}.
\end{aligned} \tag{69}$$

The general result of Eq. 69 depends neither on how the feature space is partitioned into decision regions nor on the form of the underlying distributions. The Bayes classifier maximizes this probability by choosing the regions so that the integrand is maximal for all  $\mathbf{x}$ ; no other partitioning can yield a smaller probability of error.

## 2.8 Error Bounds for Normal Densities

The Bayes decision rule guarantees the lowest average error rate, and we have seen how to calculate the decision boundaries for normal densities. However, these results do not tell us what the probability of error actually *is*. The full calculation of the error for the Gaussian case would be quite difficult, especially in high dimensions, because of the discontinuous nature of the decision regions in the integral in Eq. 69. However, in the two-category case the general error integral of Eq. 5 can be approximated analytically to give us an upper bound on the error.

### 2.8.1 Chernoff Bound

To derive a bound for the error, we need the following inequality:

$$\min[a, b] \leq a^\beta b^{1-\beta} \quad \text{for } a, b \geq 0 \text{ and } 0 \leq \beta \leq 1. \tag{70}$$

To understand this inequality we can, without loss of generality, assume  $a \geq b$ . Thus we need only show that  $b \leq a^\beta b^{1-\beta} = (\frac{a}{b})^\beta b$ . But this inequality is manifestly valid, since  $(\frac{a}{b})^\beta \geq 1$ . Using Eqs. 7 & 1, we apply this inequality to Eq. 5 and get the bound:

$$P(\text{error}) \leq P^\beta(\omega_1) P^{1-\beta}(\omega_2) \int p^\beta(\mathbf{x} | \omega_1) p^{1-\beta}(\mathbf{x} | \omega_2) d\mathbf{x} \quad \text{for } 0 \leq \beta \leq 1. \tag{71}$$

Note especially that this integral is over *all* feature space — we do not need to impose integration limits corresponding to decision boundaries.

If the conditional probabilities are normal, the integral in Eq. 71 can be evaluated analytically (Problem 35), yielding:

$$\int p^\beta(\mathbf{x} | \omega_1) p^{1-\beta}(\mathbf{x} | \omega_2) d\mathbf{x} = e^{-k(\beta)} \tag{72}$$

where

$$\begin{aligned}
k(\beta) &= \frac{\beta(1-\beta)}{2} (\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1)^t [\beta \boldsymbol{\Sigma}_1 + (1-\beta) \boldsymbol{\Sigma}_2]^{-1} (\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1) + \\
&\quad \frac{1}{2} \ln \frac{|\beta \boldsymbol{\Sigma}_1 + (1-\beta) \boldsymbol{\Sigma}_2|}{|\boldsymbol{\Sigma}_1|^\beta |\boldsymbol{\Sigma}_2|^{1-\beta}}.
\end{aligned} \tag{73}$$

The graph in Fig. 2.18 shows a typical example of how  $e^{-k(\beta)}$  varies with  $\beta$ . The *Chernoff bound*, on  $P(\text{error})$  is found by analytically or numerically finding the value

of  $\beta$  that minimizes  $e^{-k(\beta)}$ , and substituting the results in Eq. 71. The key benefit here is that this optimization is in the one-dimensional  $\beta$  space, despite the fact that the distributions themselves might be in a space of arbitrarily high dimension.




Figure 2.18: The Chernoff error bound is never looser than the Bhattacharyya bound. For this example, the Chernoff bound happens to be at  $\beta^* = 0.66$ , and is slightly tighter than the Bhattacharyya bound ( $\beta = 0.5$ ).

### 2.8.2 Bhattacharyya Bound

The general dependence of the Chernoff bound upon  $\beta$  shown in Fig. 2.18 is typical of a wide range of problems — the bound is loose for extreme values (i.e.,  $\beta \rightarrow 1$  and  $\beta \rightarrow 0$ ), and tighter for intermediate ones. While the precise value of the optimal  $\beta$  depends upon the parameters of the distributions and the prior probabilities, a computationally simpler, but slightly less tight bound can be derived by merely using the results for  $\beta = 1/2$ . This result is the so-called *Bhattacharyya bound* on the error, where Eq. 71 then has the form

$$\begin{aligned} P(\text{error}) &\leq \sqrt{P(\omega_1)P(\omega_2)} \int \sqrt{p(\mathbf{x}|\omega_1)p(\mathbf{x}|\omega_2)} d\mathbf{x} \\ &= \sqrt{P(\omega_1)P(\omega_2)} e^{-k(1/2)}, \end{aligned} \quad (74)$$

where by Eq. 73 we have for the Gaussian case:

$$\begin{aligned} k(1/2) &= 1/8(\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1)^t \left[ \frac{\boldsymbol{\Sigma}_1 + \boldsymbol{\Sigma}_2}{2} \right]^{-1} (\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1) + \\ &\quad \frac{1}{2} \ln \frac{\left| \frac{\boldsymbol{\Sigma}_1 + \boldsymbol{\Sigma}_2}{2} \right|}{\sqrt{|\boldsymbol{\Sigma}_1||\boldsymbol{\Sigma}_2|}}. \end{aligned} \quad (75)$$

The Chernoff and Bhattacharyya bounds may still be used even if the underlying distributions are not Gaussian. However, for distributions that deviate markedly from a Gaussian, the bounds will not be informative (Problem 32).

---

**Example 2: Error bounds for Gaussian distributions.**


---

It is a straightforward matter to calculate the Bhattacharyya bound for the two-dimensional data sets of Example 1. Substituting the means and covariances of Example 1 into Eq. 75 we find  $k(1/2) = 4.11$  and thus by Eqs. 74 & 75 the Bhattacharyya bound on the error is  $P(\text{error}) \leq 0.016382$ .

A tighter bound on the error can be approximated by searching numerically for the Chernoff bound of Eq. 73, which for this problem gives 0.016380. One can get the best estimate by numerically integrating the error rate directly Eq. 5, which gives 0.0021, and thus the bounds here are not particularly tight. Such numerical integration is often impractical for Gaussians in higher than two or three dimensions.

---

### 2.8.3 Signal Detection Theory and Operating Characteristics

Another measure of distance between two Gaussian distributions has found great use in experimental psychology, radar detection and other fields. Suppose we are interested in detecting a single weak pulse, such as a dim flash of light or a weak radar reflection. Our model is, then, that at some point in the detector there is an internal signal (such as a voltage)  $x$ , whose value has mean  $\mu_2$  when the external signal (pulse) is present, and mean  $\mu_1$  when it is not present. Because of random noise — within and outside the detector itself — the actual value is a random variable. We assume the distributions are normal with different means but the same variance, i.e.,  $p(x|\omega_i) \sim N(\mu_i, \sigma^2)$ , as shown in Fig. 2.19.




Figure 2.19: During any instant when no external pulse is present, the probability density for an internal signal is normal, i.e.,  $p(x|\omega_1) \sim N(\mu_1, \sigma^2)$ ; when the external signal is present, the density is  $p(x|\omega_2) \sim N(\mu_2, \sigma^2)$ . Any decision threshold  $x^*$  will determine the probability of a hit (the red area under the  $\omega_2$  curve, above  $x^*$ ) and of a false alarm (the black area under the  $\omega_1$  curve, above  $x^*$ ).

The detector (classifier) employs a threshold value  $x^*$  for determining whether the external pulse is present, but suppose we, as experimenters, do not have access to this value (nor to the means and standard deviations of the distributions). We seek to find some measure of the ease of discriminating whether the pulse is present or not, in a form independent of the choice of  $x^*$ . Such a measure is the *discriminability*, which describes the inherent and unchangeable properties due to noise and the strength of the external signal, but not on the decision strategy (i.e., the actual choice of  $x^*$ ). This discriminability is defined as

DISCRIMIN-  
ABILITY



$$d' = \frac{|\mu_2 - \mu_1|}{\sigma}. \quad (76)$$

A high  $d'$  is of course desirable.

While we do not know  $\mu_1$ ,  $\mu_2$ ,  $\sigma$  nor  $x^*$ , we assume here that we know the state of nature and the decision of the system. Such information allows us to find  $d'$ . To this end, we consider the following four probabilities:

- $P(x > x^* | x \in \omega_2)$ : a *hit* — the probability that the internal signal is above  $x^*$  given that the external signal is present
- $P(x > x^* | x \in \omega_1)$ : a *false alarm* — the probability that the internal signal is above  $x^*$  despite there being no external signal is present
- $P(x < x^* | x \in \omega_2)$ : a *miss* — the probability that the internal signal is below  $x^*$  given that the external signal is present
- $P(x < x^* | x \in \omega_1)$ : a *correct rejection* — the probability that the internal signal is below  $x^*$  given that the external signal is not present.

If we have a large number of trials (and we can assume  $x^*$  is fixed, albeit at an unknown value), we can determine these probabilities experimentally, in particular the hit and false alarm rates. We plot a point representing these rates on a two-dimensional graph. If the densities are fixed but the threshold  $x^*$  is changed, then our hit and false alarm rates will also change. Thus we see that for a given discriminability  $d'$ , our point will move along a smooth curve — a *receiver operating characteristic* or ROC curve (Fig. 2.20).

RECEIVER  
OPERATING  
CHARACTER-  
ISTIC




Figure 2.20: In a receiver operating characteristic (ROC) curve, the abscissa is the probability of false alarm,  $P(x > x^* | x \in \omega_1)$ , and the ordinate the probability of hit,  $P(x > x^* | x \in \omega_2)$ . From the measured hit and false alarm rates (here corresponding to  $x^*$  in Fig. 2.19, and shown as the red dot), we can deduce that  $d' = 3$ .

The great benefit of this signal detection framework is that we can distinguish operationally between *discriminability* and *decision bias* — while the former is an inherent property of the detector system, the latter is due to the receiver's implied but changeable loss matrix. Through any pair of hit and false alarm rates passes one and only one ROC curve; thus, so long as neither rate is exactly 0 or 1, we can determine the discriminability from these rates (Problem 38). Moreover, if the Gaussian assumption holds, a determination of the discriminability (from an arbitrary  $x^*$ ) allows us to calculate the Bayes error rate — the most important property of any

classifier. If the actual error rate differs from the Bayes rate inferred in this way, we should alter the threshold  $x^*$  accordingly.

It is a simple matter to generalize the above discussion and apply it to two categories having arbitrary multidimensional distributions, Gaussian or not. Suppose we have two distributions  $p(\mathbf{x}|\omega_1)$  and  $p(\mathbf{x}|\omega_2)$  which overlap, and thus have non-zero Bayes classification error. Just as we saw above, any pattern actually from  $\omega_2$  could be properly classified as  $\omega_2$  (a “hit”) or misclassified as  $\omega_1$  (a “false alarm”). Unlike the one-dimensional case above, however, there may be *many* decision boundaries that give a particular hit rate, each with a different false alarm rate. Clearly here we cannot determine a fundamental measure of discriminability without knowing more about the underlying decision rule than just the hit and false alarm rates.

In a rarely attainable ideal, we can imagine that our measured hit and false alarm rates are *optimal*, for example that of all the decision rules giving the measured hit rate, the rule that is actually used is the one having the minimum false alarm rate. If we constructed a multidimensional classifier — regardless of the distributions used — we might try to characterize the problem in this way, though it would probably require great computational resources to search for such optimal hit and false alarm rates.

In practice, instead we eschew optimality, and simply vary a single parameter controlling the decision rule and plot the resulting hit and false alarm rates — a curve called merely an *operating characteristic*. Such a control parameter might be the bias or nonlinearity in a discriminant function. It is traditional to choose a control parameter that can yield, at extreme values, either a vanishing false alarm or a vanishing hit rate, just as can be achieved with a very large or a very small  $x^*$  in an ROC curve. We should note that since the distributions can be arbitrary, the operating characteristic need not be symmetric (Fig. 2.21); in rare cases it need not even be concave down at all points.

OPERATING  
CHARACTER-  
ISTIC




Figure 2.21: In a general operating characteristic curve, the abscissa is the probability of false alarm,  $P(x \in \mathcal{R}_2|x \in \omega_1)$ , and the ordinate the probability of hit,  $P(x \in \mathcal{R}_2|x \in \omega_2)$ . As illustrated here, operating characteristic curves are generally not symmetric, as shown at the right.

Classifier operating curves are of value for problems where the loss matrix  $\lambda_{ij}$  might be changed. If the operating characteristic has been determined as a function of the control parameter ahead of time, it is a simple matter, when faced with a new loss function, to deduce the control parameter setting that will minimize the expected risk (Problem 38).

## 2.9 Bayes Decision Theory — Discrete Features

Until now we have assumed that the feature vector  $\mathbf{x}$  could be any point in a  $d$ -dimensional Euclidean space,  $\mathbf{R}^d$ . However, in many practical applications the components of  $\mathbf{x}$  are binary-, ternary-, or higher integer valued, so that  $\mathbf{x}$  can assume only one of  $m$  discrete values  $\mathbf{v}_1, \dots, \mathbf{v}_m$ . In such cases, the probability density function  $p(\mathbf{x}|\omega_j)$  becomes singular; integrals of the form

$$\int p(\mathbf{x}|\omega_j) d\mathbf{x} \quad (77)$$

must then be replaced by corresponding sums, such as

$$\sum_{\mathbf{x}} P(\mathbf{x}|\omega_j), \quad (78)$$

where we understand that the summation is over all values of  $\mathbf{x}$  in the discrete distribution.\* Bayes' formula then involves probabilities, rather than probability densities:

$$\boxed{P(\omega_j|\mathbf{x}) = \frac{P(\mathbf{x}|\omega_j)P(\omega_j)}{P(\mathbf{x})}}, \quad (79)$$

where

$$P(\mathbf{x}) = \sum_{j=1}^c P(\mathbf{x}|\omega_j)P(\omega_j). \quad (80)$$

The definition of the conditional risk  $R(\alpha|\mathbf{x})$  is unchanged, and the fundamental Bayes decision rule remains the same: To minimize the overall risk, select the action  $\alpha_i$  for which  $R(\alpha_i|\mathbf{x})$  is minimum, or stated formally,

$$\alpha^* = \arg \max_i R(\alpha_i|\mathbf{x}). \quad (81)$$

The basic rule to minimize the error-rate by maximizing the posterior probability is also unchanged as are the discriminant functions of Eqs. 25 – 27, given the obvious replacement of densities  $p(\cdot)$  by probabilities  $P(\cdot)$ .

### 2.9.1 Independent Binary Features

As an example of a classification involving discrete features, consider the two-category problem in which the components of the feature vector are binary-valued and conditionally independent. To be more specific we let  $\mathbf{x} = (x_1, \dots, x_d)^t$ , where the components  $x_i$  are either 0 or 1, with

$$p_i = \text{Prob}(x_i = 1|\omega_1) \quad (82)$$

and

$$q_i = \text{Prob}(x_i = 1|\omega_2). \quad (83)$$

---

\* Technically speaking, Eq. 78 should be written as  $\sum_{\mathbf{v}_k} P(\mathbf{v}_k|\omega_j)$  where  $P(\mathbf{v}_k|\omega_j)$  is the conditional probability that  $\mathbf{x} = \mathbf{v}_k$  given that the state of nature is  $\omega_j$ .

This is a model of a classification problem in which each feature gives us a yes/no answer about the pattern. If  $p_i > q_i$ , we expect the  $i$ th feature to give a “yes” answer more frequently when the state of nature is  $\omega_1$  than when it is  $\omega_2$ . (As an example, consider two factories each making the same automobile, each of whose  $d$  components could be functional or defective. If it was known how the factories differed in their reliabilities for making each component, then this model could be used to judge which factory manufactured a given automobile based on the knowledge of which features are functional and which defective.) By assuming conditional independence we can write  $P(\mathbf{x}|\omega_i)$  as the product of the probabilities for the components of  $\mathbf{x}$ . Given this assumption, a particularly convenient way of writing the class-conditional probabilities is as follows:

$$P(\mathbf{x}|\omega_1) = \prod_{i=1}^d p_i^{x_i} (1 - p_i)^{1-x_i} \quad (84)$$

and

$$P(\mathbf{x}|\omega_2) = \prod_{i=1}^d q_i^{x_i} (1 - q_i)^{1-x_i}. \quad (85)$$

Then the likelihood ratio is given by

$$\frac{P(\mathbf{x}|\omega_1)}{P(\mathbf{x}|\omega_2)} = \prod_{i=1}^d \left( \frac{p_i}{q_i} \right)^{x_i} \left( \frac{1-p_i}{1-q_i} \right)^{1-x_i} \quad (86)$$

and consequently Eq. 30 yields the discriminant function

$$g(\mathbf{x}) = \sum_{i=1}^d \left[ x_i \ln \frac{p_i}{q_i} + (1 - x_i) \ln \frac{1-p_i}{1-q_i} \right] + \ln \frac{P(\omega_1)}{P(\omega_2)}. \quad (87)$$

We note especially that this discriminant function is linear in the  $x_i$  and thus we can write

$$g(\mathbf{x}) = \sum_{i=1}^d w_i x_i + w_0, \quad (88)$$

where

$$w_i = \ln \frac{p_i(1-q_i)}{q_i(1-p_i)} \quad i = 1, \dots, d \quad (89)$$

and

$$w_0 = \sum_{i=1}^d \ln \frac{1-p_i}{1-q_i} + \ln \frac{P(\omega_1)}{P(\omega_2)}. \quad (90)$$

Let us examine these results to see what insight they can give. Recall first that we decide  $\omega_1$  if  $g(\mathbf{x}) > 0$  and  $\omega_2$  if  $g(\mathbf{x}) \leq 0$ . We have seen that  $g(\mathbf{x})$  is a weighted combination of the components of  $\mathbf{x}$ . The magnitude of the weight  $w_i$  indicates the relevance of a “yes” answer for  $x_i$  in determining the classification. If  $p_i = q_i$ ,  $x_i$  gives us no information about the state of nature, and  $w_i = 0$ , just as we might expect. If  $p_i > q_i$ , then  $1 - p_i < 1 - q_i$  and  $w_i$  is positive. Thus in this case a “yes” answer

for  $x_i$  contributes  $w_i$  votes for  $\omega_1$ . Furthermore, for any fixed  $q_i < 1$ ,  $w_i$  gets larger as  $p_i$  gets larger. On the other hand, if  $p_i < q_i$ ,  $w_i$  is negative and a “yes” answer contributes  $|w_i|$  votes for  $\omega_2$ .

The condition of feature independence leads to a very simple (linear) classifier; of course if the features were not independent, a more complicated classifier would be needed. We shall come across this again for systems with continuous features in Chap. ??, but note here that the more independent we can make the features, the simpler the classifier can be.

The prior probabilities  $P(\omega_i)$  appear in the discriminant only through the threshold weight  $w_0$ . Increasing  $P(\omega_1)$  increases  $w_0$  and biases the decision in favor of  $\omega_1$ , whereas decreasing  $P(\omega_1)$  has the opposite effect. Geometrically, the possible values for  $\mathbf{x}$  appear as the vertices of a  $d$ -dimensional hypercube; the decision surface defined by  $g(\mathbf{x}) = 0$  is a hyperplane that separates  $\omega_1$  vertices from  $\omega_2$  vertices.

---

**Example 3: Bayesian decisions for three-dimensional binary features**

---

Suppose two categories consist of independent binary features in three dimensions with known feature probabilities. Let us construct the Bayesian decision boundary if  $P(\omega_1) = P(\omega_2) = 0.5$  and the individual components obey:


$$\begin{cases} p_i = 0.8 \\ q_i = 0.5 \end{cases} \quad i = 1, 2, 3.$$

By Eqs. 89 & 90 we have that the weights are

$$w_i = \ln \frac{.8(1 - .5)}{.5(1 - .8)} = 1.3863$$

and the bias value is

$$w_0 = \sum_{i=1}^3 \ln \frac{1 - .8}{1 - .5} + \ln \frac{.5}{.5} = 1.2.$$



The decision boundary for the Example involving three-dimensional binary features. On the left we show the case  $p_i = .8$  and  $q_i = .5$ . On the right we use the same values except  $p_3 = q_3$ , which leads to  $w_3 = 0$  and a decision surface parallel to the  $x_3$  axis.

The surface  $g(\mathbf{x}) = 0$  from Eq. 88 is shown on the left of the figure. Indeed, as we might have expected, the boundary places points with two or more “yes” answers into category  $\omega_1$ , since that category has a higher probability of having any feature have value 1.

Suppose instead that while the prior probabilities remained the same, our individual components obeyed:

$$\begin{cases} p_1 = p_2 = 0.8, & p_3 = 0.5 \\ q_1 = q_2 = q_3 = 0.5 \end{cases}$$

In this case feature  $x_3$  gives us no predictive information about the categories, and hence the decision boundary is parallel to the  $x_3$  axis. Note that in this discrete case there is a large range in positions of the decision boundary that leaves the categorization unchanged, as is particularly clear in the figure on the right.

## 2.10 Missing and Noisy Features

If we know the full probability structure of a problem, we can construct the (optimal) Bayes decision rule. Suppose we develop a Bayes classifier using uncorrupted data, but our input (test) data are then corrupted in particular known ways. How can we classify such corrupted inputs to obtain a minimum error now?

There are two analytically solvable cases of particular interest: when some of the features are *missing*, and when they are corrupted by a *noise source* with known properties. In each case our basic approach is to recover as much information about the underlying distribution as possible and use the Bayes decision rule.

### 2.10.1 Missing Features

Suppose we have a Bayesian (or other) recognizer for a problem using two features, but that for a particular pattern to be classified, one of the features is missing.\* For example, we can easily imagine that the lightness can be measured from a portion of a fish, but the width cannot because of occlusion by another fish.

We can illustrate with four categories a somewhat more general case (Fig. 2.22). Suppose for a particular test pattern the feature  $x_1$  is missing, and the measured value of  $x_2$  is  $\hat{x}_2$ . Clearly if we assume the missing value is the *mean* of all the  $x_1$  values, i.e.,  $\bar{x}_1$ , we will classify the pattern as  $\omega_3$ . However, if the priors are equal,  $\omega_2$  would be a better decision, since the figure implies that  $p(\hat{x}_2|\omega_2)$  is the largest of the four likelihoods.

To clarify our derivation we let  $\mathbf{x} = [\mathbf{x}_g, \mathbf{x}_b]$ , where  $\mathbf{x}_g$  represents the known or “good” features and  $\mathbf{x}_b$  represents the “bad” ones, i.e., either unknown or missing. We seek the Bayes rule given the good features, and for that the posterior probabilities are needed. In terms of the good features the posteriors are

$$P(\omega_i|\mathbf{x}_g) = \frac{p(\omega_i, \mathbf{x}_g)}{p(\mathbf{x}_g)} = \frac{\int p(\omega_i, \mathbf{x}_g, \mathbf{x}_b) d\mathbf{x}_b}{p(\mathbf{x}_g)}$$

\* In practice, just determining that the feature is in fact *missing* rather than having a value of zero (or the mean value) can be difficult in itself.




Figure 2.22: Four categories have equal priors and the class-conditional distributions shown. If a test point is presented in which one feature is missing (here,  $x_1$ ) and the other is measured to have value  $\hat{x}_2$  (red dashed line), we want our classifier to classify the pattern as category  $\omega_2$ , because  $p(\hat{x}_2|\omega_2)$  is the largest of the four likelihoods.

$$\begin{aligned}
 &= \frac{\int P(\omega_i|\mathbf{x}_g, \mathbf{x}_b) p(\mathbf{x}_g, \mathbf{x}_b) d\mathbf{x}_b}{p(\mathbf{x}_g)} \\
 &= \frac{\int g_i(\mathbf{x}) p(\mathbf{x}) d\mathbf{x}_b}{\int p(\mathbf{x}) d\mathbf{x}_b}, \tag{91}
 \end{aligned}$$

where  $g_i(\mathbf{x}) = g_i(\mathbf{x}_g, \mathbf{x}_b) = P(\omega_i|\mathbf{x}_g, \mathbf{x}_b)$  is one form of our discriminant function.

MARGINAL

We refer to  $\int p(\omega_i, \mathbf{x}_g, \mathbf{x}_b) d\mathbf{x}_b$ , as a *marginal distribution*; we say the full joint distribution is marginalized over the variable  $\mathbf{x}_b$ . In short, Eq. 91 shows that we must integrate (marginalize) the posterior probability over the bad features. Finally we use the Bayes decision rule on the resulting posterior probabilities, i.e., choose  $\omega_i$  if  $P(\omega_i|\mathbf{x}_g) > P(\omega_j|\mathbf{x}_g)$  for all  $i$  and  $j$ . We shall consider the Expectation-Maximization (EM) algorithm in Chap. ??, which addresses a related problem involving missing features.

### 2.10.2 Noisy Features

It is a simple matter to generalize the results of Eq. 91 to the case where a particular feature has been corrupted by statistically independent noise.\* For instance, in our fish classification example, we might have a reliable measurement of the length, while variability of the light source might degrade the measurement of the lightness. We assume we have uncorrupted (good) features  $\mathbf{x}_g$ , as before, and a *noise model*, expressed as  $p(\mathbf{x}_b|\mathbf{x}_t)$ . Here we let  $\mathbf{x}_t$  denote the true value of the observed  $\mathbf{x}_b$  features, i.e., without the noise present; that is, the  $\mathbf{x}_b$  are observed instead of the true  $\mathbf{x}_t$ . We assume that if  $\mathbf{x}_t$  were known,  $\mathbf{x}_b$  would be independent of  $\omega_i$  and  $\mathbf{x}_g$ . From such an assumption we get:

$$P(\omega_i|\mathbf{x}_g, \mathbf{x}_b) = \frac{\int p(\omega_i, \mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t) d\mathbf{x}_t}{p(\mathbf{x}_g, \mathbf{x}_b)}. \tag{92}$$

\* Of course, to tell the classifier that a feature value is missing, the feature extractor must be designed to provide more than just a numerical value for each feature.

Now  $p(\omega_i, \mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t) = P(\omega_i | \mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t) p(\mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t)$ , but by our independence assumption, if we know  $\mathbf{x}_t$ , then  $\mathbf{x}_b$  does not provide any additional information about  $\omega_i$ . Thus we have  $P(\omega_i | \mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t) = P(\omega_i | \mathbf{x}_g, \mathbf{x}_t)$ . Similarly, we have  $p(\mathbf{x}_g, \mathbf{x}_b, \mathbf{x}_t) = p(\mathbf{x}_b | \mathbf{x}_g, \mathbf{x}_t) p(\mathbf{x}_g, \mathbf{x}_t)$ , and  $p(\mathbf{x}_b | \mathbf{x}_g, \mathbf{x}_t) = p(\mathbf{x}_b | \mathbf{x}_t)$ . We put these together and thereby obtain

$$\begin{aligned} P(\omega_i | \mathbf{x}_g, \mathbf{x}_b) &= \frac{\int P(\omega_i | \mathbf{x}_g, \mathbf{x}_t) p(\mathbf{x}_g, \mathbf{x}_t) p(\mathbf{x}_b | \mathbf{x}_t) d\mathbf{x}_t}{\int p(\mathbf{x}_g, \mathbf{x}_t) p(\mathbf{x}_b | \mathbf{x}_t) d\mathbf{x}_t} \\ &= \frac{\int g_i(\mathbf{x}) p(\mathbf{x}) p(\mathbf{x}_b | \mathbf{x}_t) d\mathbf{x}_t}{\int p(\mathbf{x}) p(\mathbf{x}_b | \mathbf{x}_t) d\mathbf{x}_t}, \end{aligned} \quad (93)$$

which we use as discriminant functions for classification in the manner dictated by Bayes.

Equation 93 differs from Eq. 91 solely by the fact that the integral is weighted by the noise model. In the extreme case where  $p(\mathbf{x}_b | \mathbf{x}_t)$  is uniform over the entire space (and hence provides no predictive information for categorization), the equation reduces to the case of missing features — a satisfying result.

## 2.11 Compound Bayesian Decision Theory and Context

Let us reconsider our introductory example of designing a classifier to sort two types of fish. Our original assumption was that the sequence of types of fish was so unpredictable that the state of nature looked like a random variable. Without abandoning this attitude, let us consider the possibility that the consecutive states of nature might not be statistically independent. We should be able to exploit such statistical dependence to gain improved performance. This is one example of the use of *context* to aid decision making.

The way in which we exploit such context information is somewhat different when we can wait for  $n$  fish to emerge and then make all  $n$  decisions jointly than when we must decide as each fish emerges. The first problem is a *compound decision problem*, and the second is a *sequential compound decision problem*. The former case is conceptually simpler, and is the one we shall examine here.

To state the general problem, let  $\boldsymbol{\omega} = (\omega(1), \dots, \omega(n))^t$  be a vector denoting the  $n$  states of nature, with  $\omega(i)$  taking on one of the  $c$  values  $\omega_1, \dots, \omega_c$ . Let  $P(\boldsymbol{\omega})$  be the prior probability for the  $n$  states of nature. Let  $X = (\mathbf{x}_1, \dots, \mathbf{x}_n)$  be a matrix giving the  $n$  observed feature vectors, with  $\mathbf{x}_i$  being the feature vector obtained when the state of nature was  $\omega(i)$ . Finally, let  $p(X | \boldsymbol{\omega})$  be the conditional probability density function for  $X$  given the true set of states of nature  $\boldsymbol{\omega}$ . Using this notation we see that the posterior probability of  $\boldsymbol{\omega}$  is given by

$$P(\boldsymbol{\omega} | X) = \frac{p(X | \boldsymbol{\omega}) P(\boldsymbol{\omega})}{p(X)} = \frac{p(X | \boldsymbol{\omega}) P(\boldsymbol{\omega})}{\sum_{\boldsymbol{\omega}} p(X | \boldsymbol{\omega}) P(\boldsymbol{\omega})}. \quad (94)$$

In general, one can define a loss matrix for the compound decision problem and seek a decision rule that minimizes the compound risk. The development of this theory parallels our discussion for the simple decision problem, and concludes that the optimal procedure is to minimize the compound conditional risk. In particular, if there is no loss for being correct, and if all errors are equally costly, then the procedure



reduces to computing  $P(\omega|X)$  for all  $\omega$  and selecting the  $\omega$  for which this posterior probability is maximum.

While this provides the theoretical solution, in practice the computation of  $P(\omega|X)$  can easily prove to be an enormous task. If each component  $\omega(i)$  can have one of  $c$  values, there are  $c^n$  possible values of  $\omega$  to consider. Some simplification can be obtained if the distribution of the feature vector  $\mathbf{x}_i$  depends only on the corresponding state of nature  $\omega(i)$ , not on the values of the other feature vectors or the other states of nature. In this case the joint density  $p(X|\omega)$  is merely the product of the component densities  $p(\mathbf{x}_i|\omega(i))$ :

$$p(X|\omega) = \prod_{i=1}^n p(\mathbf{x}_i|\omega(i)). \quad (95)$$

While this simplifies the problem of computing  $p(X|\omega)$ , there is still the problem of computing the prior probabilities  $P(\omega)$ . This joint probability is central to the compound Bayes decision problem, since it reflects the interdependence of the states of nature. Thus it is unacceptable to simplify the problem of calculating  $P(\omega)$  by assuming that the states of nature are independent. In addition, practical applications usually require some method of avoiding the computation of  $P(\omega|X)$  for all  $c^n$  possible values of  $\omega$ . We shall find some solutions to this problem in Chap. ??.

## Summary

The basic ideas underlying Bayes decision theory are very simple. To minimize the overall risk, one should always choose the action that minimizes the conditional risk  $R(\alpha|\mathbf{x})$ . In particular, to minimize the probability of error in a classification problem, one should always choose the state of nature that maximizes the posterior probability  $P(\omega_j|\mathbf{x})$ . Bayes' formula allows us to calculate such probabilities from the prior probabilities  $P(\omega_j)$  and the conditional densities  $p(\mathbf{x}|\omega_j)$ . If there are different penalties for misclassifying patterns from  $\omega_i$  as if from  $\omega_j$ , the posteriors must be first weighted according to such penalties before taking action.

If the underlying distributions are multivariate Gaussian, the decision boundaries will be hyperquadrics, whose form and position depends upon the prior probabilities, means and covariances of the distributions in question. The true expected error can be bounded above by the Chernoff and computationally simpler Bhattacharyya bounds. If an input (test) pattern has missing or corrupted features, we should form the marginal distributions by integrating over such features, and then using Bayes decision procedure on the resulting distributions. Receiver operating characteristic curves describe the inherent and unchangeable properties of a classifier and can be used, for example, to determine the Bayes rate.

For many pattern classification applications, the chief problem in applying these results is that the conditional densities  $p(\mathbf{x}|\omega_j)$  are not known. In some cases we may know the form these densities assume, but may not know characterizing parameter values. The classic case occurs when the densities are known to be, or can assumed to be multivariate normal, but the values of the mean vectors and the covariance matrices are not known. More commonly even less is known about the conditional densities, and procedures that are less sensitive to specific assumptions about the densities must be used. Most of the remainder of this book will be devoted to various procedures that have been developed to attack such problems.

## Bibliographical and Historical Remarks

The power, coherence and elegance of Bayesian theory in pattern recognition make it among the most beautiful formalisms in science. Its foundations go back to Bayes himself, of course [3], but he stated his theorem (Eq. 1) for the case of uniform priors. It was Laplace [25] who first stated it for the more general (but discrete) case. There are several modern and clear descriptions of the ideas — in pattern recognition and general decision theory — that can be recommended [7, 6, 26, 15, 13, 20, 27]. Since Bayesian theory rests on an axiomatic foundation, it is guaranteed to have quantitative coherence; some other classification methods do not. Wald presents a non-Bayesian perspective on these topics that can be highly recommended [36], and the philosophical foundations of Bayesian and non-Bayesian methods are explored in [16]. Neyman and Pearson provided some of the most important pioneering work in hypothesis testing, and used the probability of error as the criterion [28]; Wald extended this work by introducing the notions of loss and risk [35]. Certain conceptual problems have always attended the use of loss functions and prior probabilities. In fact, the Bayesian approach is avoided by many statisticians, partly because there are problems for which a decision is made only once, and partly because there may be no reasonable way to determine the prior probabilities. Neither of these difficulties seems to present a serious drawback in typical pattern recognition applications: for nearly all critical pattern recognition problems we *will* have training data; we will use our recognizer more than once. For these reasons, the Bayesian approach will continue to be of great use in pattern recognition. The single most important drawback of the Bayesian approach is its assumption that the true probability distributions for the problem can be represented by the classifier, for instance the true distributions are Gaussian, and all that is unknown are parameters describing these Gaussians. This is a strong assumption that is not always fulfilled and we shall later consider other approaches that do not have this requirement.

Chow[10] was among the earliest to use Bayesian decision theory for pattern recognition, and he later established fundamental relations between error and reject rate [11]. Error rates for Gaussians have been explored by [18], and the Chernoff and Bhattacharyya bounds were first presented in [9, 8], respectively and are explored in a number of statistics texts, such as [17]. Computational approximations for bounding integrals for Bayesian probability of error (the source for one of the homework problems) appears in [2]. Neyman and Pearson also worked on classification given constraints [28], and the analysis of minimax estimators for multivariate normals is presented in [5, 4, 14]. Signal detection theory and receiver operating characteristics are fully explored in [21]; a brief overview, targetting experimental psychologists, is [34]. Our discussion of the missing feature problem follows closely the work of [1] while the definitive book on missing features, including a great deal beyond our discussion here, can be found in [30].

Entropy was the central concept in the foundation of information theory [31] and the relation of Gaussians to entropy is explored in [33]. Readers requiring a review of information theory [12], linear algebra [24, 23], calculus and continuous mathematics, [38, 32] probability [29] calculus of variations and Lagrange multipliers [19] should consult these texts and those listed in our Appendix.

## Problems

### ⊕ Section 2.1

1. In the two-category case, under the Bayes' decision rule the conditional error is given by Eq. 7. Even if the posterior densities are continuous, this form of the conditional error virtually always leads to a discontinuous integrand when calculating the full error by Eq. 5.

- (a) Show that for arbitrary densities, we can replace Eq. 7 by  $P(\text{error}|x) = 2P(\omega_1|x)P(\omega_2|x)$  in the integral and get an upper bound on the full error.
- (b) Show that if we use  $P(\text{error}|x) = \alpha P(\omega_1|x)P(\omega_2|x)$  for  $\alpha < 2$ , then we are not guaranteed that the integral gives an upper bound on the error.
- (c) Analogously, show that we can use instead  $P(\text{error}|x) = P(\omega_1|x)P(\omega_2|x)$  and get a lower bound on the full error.
- (d) Show that if we use  $P(\text{error}|x) = \beta P(\omega_1|x)P(\omega_2|x)$  for  $\beta > 1$ , then we are not guaranteed that the integral gives an lower bound on the error.

### ⊕ Section 2.2

2. Consider minimax criterion for the zero-one loss function, i.e.,  $\lambda_{11} = \lambda_{22} = 0$  and  $\lambda_{12} = \lambda_{21} = 1$ .

- (a) Prove that in this case the decision regions will satisfy

$$\int_{\mathcal{R}_2} p(\mathbf{x}|\omega_1) d\mathbf{x} = \int_{\mathcal{R}_1} p(\mathbf{x}|\omega_2) d\mathbf{x}$$

- (b) Is this solution always unique? If not, construct a simple counterexample.

3. Consider the minimax criterion for a two-category classification problem.

- (a) Fill in the steps of the derivation of Eq. 22.
- (b) Explain why the overall Bayes risk must be concave down as a function of the prior  $P(\omega_1)$ , as shown in Fig. 2.4.
- (c) Assume we have one-dimensional Gaussian distributions  $p(x|\omega_i) \sim N(\mu_i, \sigma_i^2)$ ,  $i = 1, 2$  but completely unknown prior probabilities. Use the minimax criterion to find the optimal decision point  $x^*$  in terms of  $\mu_i$  and  $\sigma_i$  under a zero-one risk.
- (d) For the decision point  $x^*$  you found in (c), what is the overall minimax risk? Express this risk in terms of an error function  $\text{erf}(\cdot)$ .
- (e) Assume  $p(x|\omega_1) \sim N(0, 1)$  and  $p(x|\omega_2) \sim N(1/2, 1/4)$ , under a zero-one loss. Find  $x^*$  and the overall minimax loss.
- (f) Assume  $p(x|\omega_1) \sim N(5, 1)$  and  $p(x|\omega_2) \sim N(6, 1)$ . Without performing any explicit calculations, determine  $x^*$  for the minimax criterion. Explain your reasoning.

4. Generalize the minimax decision rule in order to classify patterns from three categories having triangle densities as follows:

$$p(x|\omega_i) = T(\mu_i, \delta_i) \equiv \begin{cases} (\delta_i - |x - \mu_i|)/\delta_i^2 & \text{for } |x - \mu_i| < \delta_i \\ 0 & \text{otherwise,} \end{cases}$$

where  $\delta_i > 0$  is the half-width of the distribution ( $i = 1, 2, 3$ ). Assume for convenience that  $\mu_1 < \mu_2 < \mu_3$ , and make some minor simplifying assumptions about the  $\delta_i$ 's as needed, to answer the following:

- (a) In terms of the priors  $P(\omega_i)$ , means and half-widths, find the optimal decision points  $x_1^*$  and  $x_2^*$  under a zero-one (categorization) loss.
- (b) Generalize the minimax decision rule to *two* decision points,  $x_1^*$  and  $x_2^*$  for such triangular distributions.
- (c) Let  $\{\mu_i, \delta_i\} = \{0, 1\}, \{.5, .5\}$ , and  $\{1, 1\}$ . Find the minimax decision rule (i.e.,  $x_1^*$  and  $x_2^*$ ) for this case.
- (d) What is the minimax risk?

5. Consider the Neyman-Pearson criterion for two univariate normal distributions:  $p(x|\omega_i) \sim N(\mu_i, \sigma_i^2)$  and  $P(\omega_i) = 1/2$  for  $i = 1, 2$ . Assume a zero-one error loss, and for convenience  $\mu_2 > \mu_1$ .

- (a) Suppose the maximum acceptable error rate for classifying a pattern that is actually in  $\omega_1$  as if it were in  $\omega_2$  is  $E_1$ . Determine the decision boundary in terms of the variables given.
- (b) For this boundary, what is the error rate for classifying  $\omega_2$  as  $\omega_1$ ?
- (c) What is the overall error rate under zero-one loss?
- (d) Apply your results to the specific case  $p(x|\omega_1) \sim N(-1, 1)$  and  $p(x|\omega_2) \sim N(1, 1)$  and  $E_1 = 0.05$ .
- (e) Compare your result to the Bayes error rate (i.e., without the Neyman-Pearson conditions).

6. Consider Neyman-Pearson criteria for two Cauchy distributions in one dimension

$$p(x|\omega_i) = \frac{1}{\pi b} \cdot \frac{1}{1 + \left(\frac{x-a_i}{b}\right)^2}, \quad i = 1, 2.$$

Assume a zero-one error loss, and for simplicity  $a_2 > a_1$ , the same “width”  $b$ , and equal priors.

- (a) Suppose the maximum acceptable error rate for classifying a pattern that is actually in  $\omega_1$  as if it were in  $\omega_2$  is  $E_1$ . Determine the decision boundary in terms of the variables given.
- (b) For this boundary, what is the error rate for classifying  $\omega_2$  as  $\omega_1$ ?
- (c) What is the overall error rate under zero-one loss?
- (d) Apply your results to the specific case  $b = 1$  and  $a_1 = -1$ ,  $a_2 = 1$  and  $E_1 = 0.1$ .

- (e) Compare your result to the Bayes error rate (i.e., without the Neyman-Pearson conditions).

⊕ *Section 2.4*

**7.** Let the conditional densities for a two-category one-dimensional problem be given by the Cauchy distribution described in Problem 6.

- (a) By explicit integration, check that the distributions are indeed normalized.
- (b) Assuming  $P(\omega_1) = P(\omega_2)$ , show that  $P(\omega_1|x) = P(\omega_2|x)$  if  $x = (a_1 + a_2)/2$ , i.e., the minimum error decision boundary is a point midway between the peaks of the two distributions, regardless of  $b$ .
- (c) Plot  $P(\omega_1|x)$  for the case  $a_1 = 3$ ,  $a_2 = 5$  and  $b = 1$ .
- (d) How do  $P(\omega_1|x)$  and  $P(\omega_2|x)$  behave as  $x \rightarrow -\infty$ ?  $x \rightarrow +\infty$ ? Explain.

**8.** Use the conditional densities given in Problem 6, and assume equal prior probabilities for the categories.

- (a) Show that the minimum probability of error is given by

$$P(\text{error}) = \frac{1}{2} - \frac{1}{\pi} \tan^{-1} \left| \frac{a_2 - a_1}{2b} \right|.$$

- (b) Plot this as a function of  $|a_2 - a_1|/b$ .
- (c) What is the maximum value of  $P(\text{error})$  and under which conditions can this occur? Explain.

**9.** Consider the following decision rule for a two-category one-dimensional problem: Decide  $\omega_1$  if  $x > \theta$ ; otherwise decide  $\omega_2$ .

- (a) Show that the probability of error for this rule is given by

$$P(\text{error}) = P(\omega_1) \int_{-\infty}^{\theta} p(x|\omega_1) dx + P(\omega_2) \int_{\theta}^{\infty} p(x|\omega_2) dx.$$

- (b) By differentiating, show that a necessary condition to minimize  $P(\text{error})$  is that  $\theta$  satisfy

$$p(\theta|\omega_1)P(\omega_1) = p(\theta|\omega_2)P(\omega_2).$$

- (c) Does this equation define  $\theta$  uniquely?
- (d) Give an example where a value of  $\theta$  satisfying the equation actually *maximizes* the probability of error.

**10.** Consider

- (a) True or false: In a two-category one-dimensional problem with continuous feature  $x$ , a monotonic transformation of  $x$  leave the Bayes error rate unchanged.

- (b) True or false: In a two-category two-dimensional problem with continuous feature  $\mathbf{x}$ , monotonic transformations of both  $x_1$  and  $x_2$  leave the Bayes error rate unchanged.

**11.** Suppose that we replace the deterministic decision function  $\alpha(\mathbf{x})$  with a *randomized rule*, viz., the probability  $P(\alpha_i|\mathbf{x})$  of taking action  $\alpha_i$  upon observing  $\mathbf{x}$ .

- (a) Show that the resulting risk is given by

$$R = \int \left[ \sum_{i=1}^a R(\alpha_i|\mathbf{x}) P(\alpha_i|\mathbf{x}) \right] p(\mathbf{x}) d\mathbf{x}.$$

- (b) In addition, show that  $R$  is minimized by choosing  $P(\alpha_i|\mathbf{x}) = 1$  for the action  $\alpha_i$  associated with the minimum conditional risk  $R(\alpha_i|\mathbf{x})$ , thereby showing that no benefit can be gained from randomizing the best decision rule.

- (c) Can we benefit from randomizing a suboptimal rule? Explain.

**12.** Let  $\omega_{max}(\mathbf{x})$  be the state of nature for which  $P(\omega_{max}|\mathbf{x}) \geq P(\omega_i|\mathbf{x})$  for all  $i$ ,  $i = 1, \dots, c$ .

- (a) Show that  $P(\omega_{max}|\mathbf{x}) \geq 1/c$ .
- (b) Show that for the minimum-error-rate decision rule the average probability of error is given by

$$P(\text{error}) = 1 - \int P(\omega_{max}|\mathbf{x}) p(\mathbf{x}) d\mathbf{x}.$$

- (c) Use these two results to show that  $P(\text{error}) \leq (c-1)/c$ .
- (d) Describe a situation for which  $P(\text{error}) = (c-1)/c$ .

**13.** In many pattern classification problems one has the option either to assign the pattern to one of  $c$  classes, or to *reject* it as being unrecognizable. If the cost for rejects is not too high, rejection may be a desirable action. Let

$$\lambda(\alpha_i|\omega_j) = \begin{cases} 0 & i = j \\ \lambda_r & i = c+1 \\ \lambda_s & \text{otherwise,} \end{cases} \quad i, j = 1, \dots, c$$

where  $\lambda_r$  is the loss incurred for choosing the  $(c+1)$ th action, rejection, and  $\lambda_s$  is the loss incurred for making a substitution error. Show that the minimum risk is obtained if we decide  $\omega_i$  if  $P(\omega_i|\mathbf{x}) \geq P(\omega_j|\mathbf{x})$  for all  $j$  and if  $P(\omega_i|\mathbf{x}) \geq 1 - \lambda_r/\lambda_s$ , and reject otherwise. What happens if  $\lambda_r = 0$ ? What happens if  $\lambda_r > \lambda_s$ ?

**14.** Consider the classification problem with rejection option.

- (a) Use the results of Problem 13 to show that the following discriminant functions are optimal for such problems:

$$g_i(\mathbf{x}) = \begin{cases} p(\mathbf{x}|\omega_i)P(\omega_i) & i = 1, \dots, c \\ \frac{\lambda_s - \lambda_r}{\lambda_s} \sum_{j=1}^c p(\mathbf{x}|\omega_j)P(\omega_j) & i = c+1. \end{cases}$$

- (b) Plot these discriminant functions and the decision regions for the two-category one-dimensional case having
- $p(x|\omega_1) \sim N(1, 1)$ ,
  - $p(x|\omega_2) \sim N(-1, 1)$ ,
  - $P(\omega_1) = P(\omega_2) = 1/2$ , and
  - $\lambda_r/\lambda_s = 1/4$ .
- (c) Describe qualitatively what happens as  $\lambda_r/\lambda_s$  is increased from 0 to 1.
- (d) Repeat for the case having
- $p(x|\omega_1) \sim N(1, 1)$ ,
  - $p(x|\omega_2) \sim N(0, 1/4)$ ,
  - $P(\omega_1) = 1/3, P(\omega_2) = 2/3$ , and
  - $\lambda_r/\lambda_s = 1/2$ .

### ⊕ Section 2.5

**15.** Confirm Eq. 45 for the volume of a  $d$ -dimensional hypersphere as follows:

- (a) Verify that the equation is correct for a line ( $d = 1$ ).
- (b) Verify that the equation is correct for a disk ( $d = 2$ ).
- (c) Integrate the volume of a line over appropriate limits to obtain the volume of a disk.
- (d) Consider a general  $d$ -dimensional hypersphere. Integrate its volume to obtain a formula (involving the ratio of gamma functions,  $\Gamma(\cdot)$ ) for the volume of a  $(d + 1)$ -dimensional hypersphere.
- (e) Apply your formula to find the volume of a hypersphere in an odd-dimensional space by integrating the volume of a hypersphere in the lower even-dimensional space, and thereby confirm Eq. 45 for odd dimensions.
- (f) Repeat the above but for finding the volume of a hypersphere in even dimensions.

**16.** Derive the formula for the volume of a  $d$ -dimensional hypersphere in Eq. 45 as follows:

- (a) State by inspection the formula for  $V_1$ .
- (b) Follow the general procedure outlined in Problem 15 and integrate twice to find  $V_{d+2}$  as a function of  $V_d$ .
- (c) Assume that the functional form of  $V_d$  is the same for all odd dimensions (and likewise for all even dimensions). Use your integration results to determine the formula for  $V_d$  for  $d$  odd.
- (d) Use your intermediate integration results to determine  $V_d$  for  $d$  even.
- (e) Explain why we should expect the functional form of  $V_d$  to be different in even and in odd dimensions.

**17.** Derive the formula (Eq. 44) for the volume  $V$  of a hyperellipsoid of constant Mahalanobis distance  $r$  (Eq. 43) for a Gaussian distribution having covariance  $\Sigma$ .

**18.** Consider two normal distributions in one dimension:  $N(\mu_1, \sigma_1^2)$  and  $N(\mu_2, \sigma_2^2)$ . Imagine that we choose two random samples  $x_1$  and  $x_2$ , one from each of the normal distributions and calculate their sum  $x_3 = x_1 + x_2$ . Suppose we do this repeatedly.

- (a) Consider the resulting distribution of the values of  $x_3$ . Show from first principles that this is also a normal distribution.
- (b) What is the mean,  $\mu_3$ , of your new distribution?
- (c) What is the variance,  $\sigma_3^2$ ?
- (d) Repeat the above with two distributions in a multi-dimensional space, i.e.,  $N(\mu_1, \Sigma_1)$  and  $N(\mu_2, \Sigma_2)$ .

**19.** Starting from the definition of entropy (Eq. 36), derive the general equation for the maximum-entropy distribution given constraints expressed in the general form

$$\int b_k(x)p(x) dx = a_k, \quad k = 1, 2, \dots, q$$

as follows:

- (a) Use Lagrange undetermined multipliers  $\lambda_1, \lambda_2, \dots, \lambda_q$  and derive the synthetic function:

$$H_s = - \int p(x) \left[ \ln p(x) - \sum_{k=1}^q \lambda_k b_k(x) \right] dx - \sum_{k=1}^q \lambda_k a_k.$$

State why we know  $a_0 = 1$  and  $b_0(x) = 1$  for all  $x$ .

- (b) Take the derivative of  $H_s$  with respect to  $p(x)$ . Equate the integrand to zero, and thereby prove that the minimum-entropy distribution obeys

$$p(x) = \exp \left[ \sum_{k=1}^q \lambda_k b_k(x) - 1 \right],$$

where the  $q + 1$  parameters are determined by the constraint equation.

**20.** Use the final result from Problem 19 for the following.

- (a) Suppose we know only that a distribution is non-zero in the range  $x_l \leq x \leq x_u$ . Prove that the maximum entropy distribution is uniform in that range, i.e.,

$$p(x) \sim U(x_l, x_u) = \begin{cases} 1/|x_u - x_l| & x_l \leq x \leq x_u \\ 0 & \text{otherwise.} \end{cases}$$



- (b) Suppose we know only that a distribution is non-zero for  $x \geq 0$  and that its mean is  $\mu$ . Prove that the maximum entropy distribution is

$$p(x) = \begin{cases} \frac{1}{\mu} e^{-x/\mu} & \text{for } x \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$

- (c) Now suppose we know solely that the distribution is normalized, has mean  $\mu$ , and standard deviation  $\sigma^2$ , and thus from Problem 19 our maximum entropy distribution must be of the form

$$p(x) = \exp[\lambda_0 - 1 + \lambda_1 x + \lambda_2 x^2].$$

Write out the three constraints and solve for  $\lambda_0, \lambda_1$ , and  $\lambda_2$  and thereby prove that the maximum entropy solution is a Gaussian, i.e.,

$$p(x) = \frac{1}{\sqrt{2\pi}} \exp \left[ \frac{-(x - \mu)^2}{2\sigma^2} \right].$$

**21.** Three distributions — a Gaussian, a uniform distribution, and a triangle distribution (cf., Problem 4) — each have mean zero and standard deviation  $\sigma^2$ . Use Eq. 36 to calculate and compare their entropies.

**22.** Calculate the entropy of a multidimensional Gaussian  $p(\mathbf{x}) \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$ .

$\oplus$  Section 2.6

**23.** Consider the three-dimensional normal distribution  $p(\mathbf{x}|\omega) \sim N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  where  $\boldsymbol{\mu} = \begin{pmatrix} 1 \\ 2 \\ 2 \end{pmatrix}$  and  $\boldsymbol{\Sigma} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 5 & 2 \\ 0 & 2 & 5 \end{pmatrix}$ .

- Find the probability density at the point  $\mathbf{x}_0 = (.5, 0, 1)^t$ .
- Construct the whitening transformation  $\mathbf{A}_w$ . Show your  $\boldsymbol{\Lambda}$  and  $\boldsymbol{\Phi}$  matrices. Next, convert the distribution to one centered on the origin with covariance matrix equal to the identity matrix,  $p(\mathbf{x}|\omega) \sim N(\mathbf{0}, \mathbf{I})$ .
- Apply the same overall transformation to  $\mathbf{x}_0$  to yield a transformed point  $\mathbf{x}_w$ .
- By explicit calculation, confirm that the Mahalanobis distance from  $\mathbf{x}_0$  to the mean  $\boldsymbol{\mu}$  in the original distribution is the same as for  $\mathbf{x}_w$  to  $\mathbf{0}$  in the transformed distribution.
- Does the probability density remain unchanged under a general linear transformation? In other words, is  $p(\mathbf{x}_0|N(\boldsymbol{\mu}, \boldsymbol{\Sigma})) = p(\mathbf{T}^t \mathbf{x}_0|N(\mathbf{T}^t \boldsymbol{\mu}, \mathbf{T}^t \boldsymbol{\Sigma} \mathbf{T}))$  for some linear transform  $\mathbf{T}$ ? Explain.
- Prove that a general whitening transform  $\mathbf{A}_w = \boldsymbol{\Phi} \boldsymbol{\Lambda}^{-1/2}$  when applied to a Gaussian distribution insures that the final distribution has covariance proportional to the identity matrix  $\mathbf{I}$ . Check whether normalization is preserved by the transformation.

**24.** Consider the multivariate normal density for which  $\sigma_{ij} = 0$  and  $\sigma_{ii} = \sigma_i^2$ , i.e.,  $\Sigma = \text{diag}(\sigma_1^2, \sigma_2^2, \dots, \sigma_d^2)$ .

- (a) Show that the evidence is

$$p(\mathbf{x}) = \frac{1}{\prod_{i=1}^d \sqrt{2\pi}\sigma_i} \exp \left[ -\frac{1}{2} \sum_{i=1}^d \left( \frac{x_i - \mu_i}{\sigma_i} \right)^2 \right].$$

- (b) Plot and describe the contours of constant density.

- (c) Write an expression for the Mahalanobis distance from  $\mathbf{x}$  to  $\boldsymbol{\mu}$ .

**25.** Fill in the steps in the derivation from Eq. 57 to Eqs. 58–63.

**26.** Let  $p(\mathbf{x}|\omega_i) \sim N(\boldsymbol{\mu}_i, \Sigma)$  for a two-category  $d$ -dimensional problem with the same covariances but arbitrary means and prior probabilities. Consider the squared Mahalanobis distance

$$r_i^2 = (\mathbf{x} - \boldsymbol{\mu}_i)^t \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}_i).$$

- (a) Show that the gradient of  $r_i^2$  is given by

$$\nabla r_i^2 = 2\Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu}_i).$$

- (b) Show that at any position on a given line through  $\boldsymbol{\mu}_i$  the gradient  $\nabla r_i^2$  points in the same direction. Must this direction be parallel to that line?

- (c) Show that  $\nabla r_1^2$  and  $\nabla r_2^2$  point in opposite directions along the line from  $\boldsymbol{\mu}_1$  to  $\boldsymbol{\mu}_2$ .

- (d) Show that the optimal separating hyperplane is tangent to the constant probability density hyperellipsoids at the point that the separating hyperplane cuts the line from  $\boldsymbol{\mu}_1$  to  $\boldsymbol{\mu}_2$ .

- (e) True or False: For a two-category problem involving normal densities with arbitrary means and covariances, and  $P(\omega_1) = P(\omega_2) = 1/2$ , the Bayes decision boundary consists of the set of points of equal Mahalanobis distance from the respective sample means. Explain.

**27.** Suppose we have two normal distributions with the same covariances but different means:  $N(\boldsymbol{\mu}_1, \Sigma)$  and  $N(\boldsymbol{\mu}_2, \Sigma)$ . In terms of their prior probabilities  $P(\omega_1)$  and  $P(\omega_2)$ , state the condition that the Bayes decision boundary *not* pass between the two means.

**28.** Two random variables  $\mathbf{x}$  and  $\mathbf{y}$  are called “statistically independent” if  $p(\mathbf{x}, \mathbf{y}|\omega) = p(\mathbf{x}|\omega)p(\mathbf{y}|\omega)$ .

- (a) Prove that if  $x_i - \mu_i$  and  $x_j - \mu_j$  are statistically independent (for  $i \neq j$ ) then  $\sigma_{ij}$  as defined in Eq. 42 is 0.

- (b) Prove that the converse is true for the Gaussian case.

- (c) Show by counterexample that this converse is *not* true in the general case.

**29.** Consider the Bayes decision boundary for two-category classification in  $d$  dimensions.

- (a) Prove that for any arbitrary hyperquadratic in  $d$  dimensions, there exist normal distributions  $p(x|\omega_i) \sim N(\boldsymbol{\mu}_i, \boldsymbol{\Sigma}_i)$  and priors  $P(\omega_i)$ ,  $i = 1, 2$ , that possess this hyperquadratic as their Bayes decision boundary.
- (b) Is the above also true if the priors are held fixed and non-zero, e.g.,  $P(\omega_1) = P(\omega_2) = 1/2$ ?

$\oplus$  Section 2.7

**30.** Let  $p(x|\omega_i) \sim N(\mu_i, \sigma^2)$  for a two-category one-dimensional problem with  $P(\omega_1) = P(\omega_2) = 1/2$ .

- (a) Show that the minimum probability of error is given by

$$P_e = \frac{1}{\sqrt{2\pi}} \int_a^\infty e^{-u^2/2} du,$$

where  $a = |\mu_2 - \mu_1|/(2\sigma)$ .

- (b) Use the inequality

$$P_e = \frac{1}{\sqrt{2\pi}} \int_a^\infty e^{-t^2/2} dt \leq \frac{1}{\sqrt{2\pi}a} e^{-a^2/2}$$

to show that  $P_e$  goes to zero as  $|\mu_2 - \mu_1|/\sigma$  goes to infinity.

**31.** Let  $p(\mathbf{x}|\omega_i) \sim N(\boldsymbol{\mu}_i, \sigma^2 \mathbf{I})$  for a two-category  $d$ -dimensional problem with  $P(\omega_1) = P(\omega_2) = 1/2$ .

- (a) Show that the minimum probability of error is given by

$$P_e = \frac{1}{\sqrt{2\pi}} \int_a^\infty e^{-u^2/2} du,$$

where  $a = \|\boldsymbol{\mu}_2 - \boldsymbol{\mu}_1\|/(2\sigma)$ .

- (b) Let  $\boldsymbol{\mu}_1 = \mathbf{0}$  and  $\boldsymbol{\mu} = (\mu_1, \dots, \mu_d)^t$ . Use the inequality from Problem 30 to show that  $P_e$  approaches zero as the dimension  $d$  approaches infinity.
- (c) Express the meaning of this result in words.

**32.** Show that if the densities in a two-category classification problem differ markedly from Gaussian, the Chernoff and Bhattacharyya bounds are not likely to be informative by considering the following one-dimensional examples. Consider a number of problems in which the mean and variance are the same (and thus the Chernoff bound and the Bhattacharyya bound remain the same), but nevertheless have a wide range in Bayes error. For definiteness, assume the distributions have means at  $\mu_1 = -\mu$  and  $\mu_2 = +\mu$ , and  $\sigma_1^2 = \sigma_2^2 = \mu^2$ .

- (a) Use the equations in the text to calculate the Chernoff and the Bhattacharyya bounds on the error.
- (b) Suppose the distributions are both Gaussian. Calculate explicitly the Bayes error. Express it in terms of an error function  $\text{erf}(\cdot)$  and as a numerical value.
- (c) Now consider a another case, in which half the density for  $\omega_1$  is concentrated at a point  $x = -2\mu$  and half at  $x = 0$ ; likewise (symmetrically) the density for  $\omega_2$  has half its mass at  $x = +2\mu$  and half at  $x = 0$ . Show that the means and variance remain as desired, but that now the Bayes error is 0.5.
- (d) Now consider yet another case, in which half the density for  $\omega_1$  is concentrated near  $x = -2$  and half at  $x = -\epsilon$ , where  $\epsilon$  is an infinitesimally small positive distance; likewise (symmetrically) the density for  $\omega_2$  has half its mass near  $x = +2\mu$  and half at  $+\epsilon$ . Show that by making  $\epsilon$  sufficiently small, the means and variances can be made arbitrarily close to  $\mu$  and  $\mu^2$ , respectively. Show, too, that now the Bayes error is zero.
- (e) Compare your errors in (b), (c) and (d) to your Chernoff and Bhattacharyya bounds of (a) and explain in words why those bounds are unlikely to be of much use if the distributions differ markedly from Gaussians.

**33.** Suppose we know exactly two arbitrary distributions  $p(\mathbf{x}|\omega_i)$  and priors  $P(\omega_i)$  in a  $d$ -dimensional feature space.

- (a) Prove that the true error cannot decrease if we first project the distributions to a lower dimensional space and then classify them.
- (b) Despite this fact, suggest why in an actual pattern recognition application we might not want to include an arbitrarily high number of feature dimensions.

### ⊕ Section 2.8

**34.** Show for non-pathological cases that if we include more feature dimensions in a Bayesian classifier for multidimensional Gaussian distributions then the Bhattacharyya bound decreases. Do this as follows: Let  $P_d(P(\omega_1), \boldsymbol{\mu}_1, \boldsymbol{\Sigma}_1, P(\omega_2), \boldsymbol{\mu}_2, \boldsymbol{\Sigma}_2)$ , or simply  $P_d$ , be the Bhattacharyya bound if we consider the distributions restricted to  $d$  dimensions.

- (a) Using general properties of a covariance matrix, prove that  $k(1/2)$  of Eq. 75 must increase as we increase from  $d$  to  $d + 1$  dimensions, and hence the error bound must decrease.
- (b) Explain why this general result does or does not depend upon *which* dimension is added.
- (c) What is a “pathological” case in which the error bound does *not* decrease, i.e., for which  $P_{d+1} = P_d$ ?
- (d) Is it ever possible that the *true* error could *increase* as we go to higher dimension?
- (e) Prove that as  $d \rightarrow \infty$ ,  $P_d \rightarrow 0$  for non-pathological distributions. Describe pathological distributions for which this infinite limit does not hold.

- (f) Given that the Bhattacharyya bound decreases for the inclusion of a particular dimension, does this guarantee that the *true* error will decrease? Explain.

**35.** Derive Eqs. 72 & 73 from Eq. 71 by the following steps:

- (a) Substitute the normal distributions into the integral and gather the terms dependent upon  $\mathbf{x}$  and those that are not dependent upon  $\mathbf{x}$ .
- (b) Factor the term independent of  $\mathbf{x}$  from the integral.
- (c) Integrate explicitly the term dependent upon  $\mathbf{x}$ .

**36.** Consider a two-category classification problem in two dimensions with  $p(\mathbf{x}|\omega_1) \sim N(\mathbf{0}, \mathbf{I})$ ,  $p(\mathbf{x}|\omega_2) \sim N\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{I}\right)$ , and  $P(\omega_1) = P(\omega_2) = 1/2$ .

- (a) Calculate the Bayes decision boundary.
- (b) Calculate the Bhattacharyya error bound.
- (c) Repeat the above for the same prior probabilities, but  $p(\mathbf{x}|\omega_1) \sim N\left(\mathbf{0}, \begin{pmatrix} 2 & .5 \\ .5 & 2 \end{pmatrix}\right)$  and  $p(\mathbf{x}|\omega_2) \sim N\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 5 & 4 \\ 4 & 5 \end{pmatrix}\right)$ .

**37.** Derive the Bhattacharyya error bound without the need for first examining the Chernoff bound. Do this as follows:

- (a) If  $a$  and  $b$  are nonnegative numbers, show directly that  $\min[a, b] \leq \sqrt{ab}$ .
- (b) Use this to show that the error rate for a two-category Bayes classifier must satisfy

$$P(\text{error}) \leq \sqrt{P(\omega_1)P(\omega_2)} \rho \leq \rho/2,$$

where  $\rho$  is the so-called *Bhattacharyya coefficient*

$$\rho = \int \sqrt{p(\mathbf{x}|\omega_1) p(\mathbf{x}|\omega_2)} d\mathbf{x}.$$

**38.** Use the signal detection theory, the notation and basic Gaussian assumptions described in the text to address the following.

- (a) Prove that  $P(x > x^*|x \in \omega_2)$  and  $P(x < x^*|x \in \omega_2)$ , taken together, uniquely determine the discriminability  $d'$ .
- (b) Use error functions  $\text{erf}(\cdot)$  to express  $d'$  in terms of the hit and false alarm rates. Estimate  $d'$  if  $P(x > x^*|x \in \omega_2) = 0.8$  and  $P(x < x^*|x \in \omega_2) = 0.3$ . Repeat for  $P(x > x^*|x \in \omega_2) = 0.7$  and  $P(x < x^*|x \in \omega_2) = 0.4$ .
- (c) Given that the Gaussian assumption is valid, calculate the Bayes error for both the cases in (b).
- (d) Determine by a trivial one-line computation, which case has the higher  $d'$ :

**case A:**  $P(x > x^*|x \in \omega_2) = 0.8, P(x < x^*|x \in \omega_2) = 0.3$  or

**case B:**  $P(x > x^*|x \in \omega_2) = 0.9, P(x < x^*|x \in \omega_2) = 0.7$ .

Explain your logic.

**39.** Suppose in our signal detection framework we had two Gaussians, but with different variances (cf., Fig. 2.20), that is,  $p(x|\omega_1) \sim N(\mu_1, \sigma_1^2)$  and  $p(x|\omega_2) \sim N(\mu_2, \sigma_2^2)$  for  $\mu_2 > \mu_1$  and  $\sigma_2^2 \neq \sigma_1^2$ . In that case the resulting ROC curve would no longer be symmetric.

- Suppose in this asymmetric case we modified the definition of the discriminability to be  $d'_a = |\mu_2 - \mu_1|/\sqrt{\sigma_1\sigma_2}$ . Show by non-trivial counterexample or analysis that one cannot determine  $d'_a$  uniquely based on a single pair of hit and false alarm rates.
- Assume we measure the hit and false alarm rates for two different, but unknown, values of the threshold  $x^*$ . Derive a formula for  $d'_a$  based on measurements.
- State and explain all pathological values for which your formula does not give a meaningful value for  $d'_a$ .
- Plot several ROC curves for the case  $p(x|\omega_1) \sim N(0, 1)$  and  $p(x|\omega_2) \sim N(1, 2)$ .

**40.** Consider two one-dimensional triangle distributions having different means, but the same width:

$$p(x|\omega_i) = T(\mu_i, \delta) = \begin{cases} (\delta - |x - \mu_i|)/\delta^2 & \text{for } |x - \mu_i| < \delta \\ 0 & \text{otherwise,} \end{cases}$$

with  $\mu_2 > \mu_1$ . We define a new discriminability here as  $d'_T = (\mu_2 - \mu_1)/\delta_i$ .

- Write an analytic function, parameterized by  $d'_T$ , for the operating characteristic curves.
- Plot these novel operating characteristic curves for  $d'_T = \{.1, .2, \dots, 1.0\}$ . Interpret your answer for the case  $d'_T = 1.0$ .
- Suppose we measure  $P(x > x^*|x \in \omega_2) = .4$  and  $P(x > x^*|x \in \omega_1) = .2$ . What is  $d'_T$ ? What is the Bayes error rate?
- Infer the decision rule. That is, express  $x^*$  in terms of the variables given in the problem.
- Suppose we measure  $P(x > x^*|x \in \omega_2) = .9$  and  $(x > x^*|x \in \omega_1) = .3$ . What is  $d'_T$ ? What is the Bayes error rate?
- Infer the decision rule. That is, express  $x^*$  in terms of the variables given in the problem.

**41.** Equation 70 can be used to obtain an upper bound on the error. One can also derive tighter analytic bounds in the two-category case — both upper and lower bounds — analogous to Eq. 71 for general distributions. If we let  $p \equiv p(x|\omega_1)$ , then we seek tighter bounds on  $\text{Min}[p, 1 - p]$  (which has discontinuous derivative).

- Prove that

$$b_L(p) = \frac{1}{\beta} \ln \left[ \frac{1 + e^{-\beta}}{e^{-\beta p} + e^{-\beta(1-p)}} \right]$$

for any  $\beta > 0$  is a lower bound on  $\text{Min}[p, 1 - p]$ .

- (b) Prove that one can choose  $\beta$  in (a) to give an arbitrarily tight lower bound.  
 (c) Repeat (a) and (b) for the upper bound given by

$$b_U(p) = b_L(p) + [1 - 2g_L(0.5)]b_G(p)$$

where  $b_G(p)$  is any upper bound that obeys

$$\begin{aligned} b_G(p) &\geq \text{Min}[p, 1 - p] \\ b_G(p) &= b_G(1 - p) \\ b_G(0) &= b_G(1) = 0 \\ b_G(0.5) &= 0.5. \end{aligned}$$

- (d) Confirm that  $b_G(p) = 1/2 \sin[\pi p]$  obeys the conditions in (c).  
 (e) Let  $b_G(p) = 1/2 \sin[\pi p]$ , and plot your upper and lower bounds as a function of  $p$ , for  $0 \leq p \leq 1$  and  $\beta = 1, 10, 50$ .

### ⊕ Section 2.9

**42.** Let the components of the vector  $\mathbf{x} = (x_1, \dots, x_d)^t$  be binary valued (0 or 1) and  $P(\omega_j)$  be the prior probability for the state of nature  $\omega_j$  and  $j = 1, \dots, c$ . Now define

$$p_{ij} = \text{Prob}(x_i = 1 | \omega_j) \quad \begin{array}{l} i = 1, \dots, d \\ j = 1, \dots, c, \end{array}$$

with the components of  $x_i$  being statistically independent for all  $\mathbf{x}$  in  $\omega_j$ .

- (a) Interpret in words the meaning of  $p_{ij}$ .  
 (b) Show that the minimum probability of error is achieved by the following decision rule: Decide  $\omega_k$  if  $g_k(\mathbf{x}) \geq g_j(\mathbf{x})$  for all  $j$  and  $k$ , where

$$g_j(\mathbf{x}) = \sum_{i=1}^d x_i \ln \frac{p_{ij}}{1 - p_{ij}} + \sum_{i=1}^d \ln (1 - p_{ij}) + \ln P(\omega_j).$$

**43.** Let the components of the vector  $\mathbf{x} = (x_1, \dots, x_d)^t$  be ternary valued (1, 0 or -1), with

$$\begin{aligned} p_{ij} &= \text{Prob}(x_i = 1 | \omega_j) \\ q_{ij} &= \text{Prob}(x_i = 0 | \omega_j) \\ r_{ij} &= \text{Prob}(x_i = -1 | \omega_j), \end{aligned}$$

and with the components of  $x_i$  being statistically independent for all  $\mathbf{x}$  in  $\omega_j$ .

- (a) Show that a minimum probability of error decision rule can be derived that involves discriminant functions  $g_j(\mathbf{x})$  that are quadratic function of the components  $x_i$ .  
 (b) Suggest a generalization to more categories of your answers to this and Problem 42.

**44.** Let  $\mathbf{x}$  be distributed as in Problem 42 with  $c = 2$ ,  $d$  odd, and

$$\begin{aligned} p_{i1} &= p > 1/2 & i = 1, \dots, d \\ p_{i2} &= 1 - p & i = 1, \dots, d \end{aligned}$$

and  $P(\omega_1) = P(\omega_2) = 1/2$ .

(a) Show that the minimum-error-rate decision rule becomes:

$$\text{Decide } \omega_1 \text{ if } \sum_{i=1}^d x_i > d/2 \text{ and } \omega_2 \text{ otherwise.}$$

(b) Show that the minimum probability of error is given by

$$P_e(d, p) = \sum_{k=0}^{(d-1)/2} \binom{d}{k} p^k (1-p)^{d-k}.$$

where  $\binom{d}{k} = d!/(k!(d-k)!)$  is the binomial coefficient.

(c) What is the limiting value of  $P_e(d, p)$  as  $p \rightarrow 1/2$ ? Explain.

(d) Show that  $P_e(d, p)$  approaches zero as  $d \rightarrow \infty$ . Explain.

**45.** Under the natural assumption concerning losses, i.e., that  $\lambda_{21} > \lambda_{11}$  and  $\lambda_{12} > \lambda_{22}$ , show that the general minimum risk discriminant function for the independent binary case described in Sect. 2.9.1 is given by  $g(\mathbf{x}) = \mathbf{w}^t \mathbf{x} + w_0$ , where  $\mathbf{w}$  is unchanged, and

$$w_0 = \sum_{i=1}^d \ln \frac{1-p_i}{1-q_i} + \ln \frac{P(\omega_1)}{P(\omega_2)} + \ln \frac{\lambda_{21} - \lambda_{11}}{\lambda_{12} - \lambda_{22}}.$$

**46.** The Poisson distribution for a discrete variable  $x = 0, 1, 2, \dots$  and real parameter  $\lambda$  is

$$P(x|\lambda) = e^{-\lambda} \frac{\lambda^x}{x!}.$$

(a) Prove that the mean of such a distribution is  $\mathcal{E}[x] = \lambda$ .

(b) Prove that the variance of such a distribution is  $\mathcal{E}[x - \bar{x}] = \lambda$ .

(c) The *mode* of a distribution is the value of  $x$  that has the maximum probability. Prove that the mode of a Poisson distribution is the greatest integer that does not exceed  $\lambda$ , i.e., the mode is  $\lceil \lambda \rceil$ . (If  $\lambda$  is an integer, then both  $\lambda$  and  $\lambda - 1$  are modes.)

(d) Consider two equally probable categories having Poisson distributions but with differing parameters; assume for definiteness  $\lambda_1 > \lambda_2$ . What is the Bayes classification decision?

(e) What is the Bayes error rate?



⊕ Section 2.10

47. Suppose we have three categories in two dimensions with the following underlying distributions:

- $p(\mathbf{x}|\omega_1) \sim N(\mathbf{0}, \mathbf{I})$
- $p(\mathbf{x}|\omega_2) \sim N\left(\begin{pmatrix} 1 \\ 1 \end{pmatrix}, \mathbf{I}\right)$
- $p(\mathbf{x}|\omega_3) \sim \frac{1}{2}N\left(\begin{pmatrix} .5 \\ .5 \end{pmatrix}, \mathbf{I}\right) + \frac{1}{2}N\left(\begin{pmatrix} -.5 \\ .5 \end{pmatrix}, \mathbf{I}\right)$

with  $P(\omega_i) = 1/3$ ,  $i = 1, 2, 3$ .

- (a) By explicit calculation of posterior probabilities, classify the point  $\mathbf{x} = \begin{pmatrix} .3 \\ .3 \end{pmatrix}$  for minimum probability of error.
- (b) Suppose that for a particular test point the first feature is missing. That is, classify  $\mathbf{x} = \begin{pmatrix} * \\ .3 \end{pmatrix}$ .
- (c) Suppose that for a particular test point the second feature is missing. That is, classify  $\mathbf{x} = \begin{pmatrix} .3 \\ * \end{pmatrix}$ .
- (d) Repeat all of the above for  $\mathbf{x} = \begin{pmatrix} .2 \\ .6 \end{pmatrix}$ .

48. Show that Eq. 93 reduces to Bayes rule when the true feature is  $\boldsymbol{\mu}_i$  and  $p(\mathbf{x}_b|\mathbf{x}_t) \sim N(\mathbf{x}_t, \boldsymbol{\Sigma})$ . Interpret this answer in words.


⊕ Section 2.11

49. Suppose we have three categories with  $P(\omega_1) = 1/2$ ,  $P(\omega_2) = P(\omega_3) = 1/4$  and the following distributions

- $p(x|\omega_1) \sim N(0, 1)$
- $p(x|\omega_2) \sim N(.5, 1)$
- $p(x|\omega_3) \sim N(1, 1)$  ,

and that we sample the following four points:  $x = 0.6, 0.1, 0.9, 1.1$ .

- (a) Calculate explicitly the probability that the sequence actually came from  $\omega_1, \omega_3, \omega_3, \omega_2$ . Be careful to consider normalization.
- (b) Repeat for the sequence  $\omega_1, \omega_2, \omega_2, \omega_3$ .
- (c) Find the sequence having the maximum probability.



## Computer exercises

Several of the computer exercises will rely on the following data.

sample	$\omega_1$			$\omega_2$			$\omega_3$		
	$x_1$	$x_2$	$x_3$	$x_1$	$x_2$	$x_3$	$x_1$	$x_2$	$x_3$
1	-5.01	-8.12	-3.68	-0.91	-0.18	-0.05	5.35	2.26	8.13
2	-5.43	-3.48	-3.54	1.30	-2.06	-3.53	5.12	3.22	-2.66
3	1.08	-5.52	1.66	-7.75	-4.54	-0.95	-1.34	-5.31	-9.87
4	0.86	-3.78	-4.11	-5.47	0.50	3.92	4.48	3.42	5.19
5	-2.67	0.63	7.39	6.14	5.72	-4.85	7.11	2.39	9.21
6	4.94	3.29	2.08	3.60	1.26	4.36	7.17	4.33	-0.98
7	-2.51	2.09	-2.59	5.37	-4.63	-3.65	5.75	3.97	6.65
8	-2.25	-2.13	-6.94	7.18	1.46	-6.66	0.77	0.27	2.41
9	5.56	2.86	-2.26	-7.39	1.17	6.30	0.90	-0.43	-8.71
10	1.03	-3.33	4.33	-7.50	-6.32	-0.31	3.52	-0.36	6.43

### ⊕ Section 2.2

1. You may need the following procedures for several exercises below.
  - (a) Write a procedure to generate random samples according to a normal distribution  $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$  in  $d$  dimensions.
  - (b) Write a procedure to calculate the discriminant function (of the form given in Eq. 47) for a given normal distribution and prior probability  $P(\omega_i)$ .
  - (c) Write a procedure to calculate the Euclidean distance between two arbitrary points.
  - (d) Write a procedure to calculate the Mahalanobis distance between the mean  $\boldsymbol{\mu}$  and an arbitrary point  $\mathbf{x}$ , given the covariance matrix  $\boldsymbol{\Sigma}$ .

### ⊕ Section 2.5

2. Use your classifier from Problem ?? to classify the following 10 samples from the table above in the following way. Assume that the underlying distributions are normal.
  - (a) Assume that the prior probabilities for the first two categories are equal ( $P(\omega_1) = P(\omega_2) = 1/2$  and  $P(\omega_3) = 0$ ) and design a dichotomizer for those two categories using only the  $x_1$  feature value.
  - (b) Determine the empirical training error on your samples, i.e., the percentage of points misclassified.
  - (c) Use the Bhattacharyya bound to bound the error you will get on novel patterns drawn from the distributions.
  - (d) Repeat all of the above, but now use *two* feature values,  $x_1$ , and  $x_2$ .
  - (e) Repeat, but use all *three* feature values.
  - (f) Discuss your results. In particular, is it ever possible for a finite set of data that the empirical error might be *larger* for more data dimensions?

3. Repeat Computer exercise 2 but for categories  $\omega_1$  and  $\omega_3$ .
4. Repeat Computer exercise 2 but for categories  $\omega_2$  and  $\omega_3$ .
5. Consider the three categories in Computer exercise 2, and assume  $P(\omega_i) = 1/3$ .
  - (a) What is the Mahalanobis distance between each of the following test points and each of the category means in Computer exercise 2:  $(1, 2, 1)^t$ ,  $(5, 3, 2)^t$ ,  $(0, 0, 0)^t$ ,  $(1, 0, 0)^t$ .
  - (b) Classify those points.
  - (c) Assume instead that  $P(\omega_1) = 0.8$ , and  $P(\omega_2) = P(\omega_3) = 0.1$  and classify the test points again.
6. Illustrate the fact that the average of a large number of independent random variables will approximate a Gaussian by the following:
  - (a) Write a program to generate  $n$  random integers from a uniform distribution  $U(x_l, x_u)$ . (Some computer systems include this as a single, compiled function call.)
  - (b) Now write a routine to choose  $x_l$  and  $x_u$  randomly, in the range  $-100 \leq x_l < x_u \leq +100$ , and  $n$  (the number of samples) randomly in the range  $0 < n \leq 1000$ .
  - (c) Generate and plot a histogram of the accumulation of  $10^4$  points sampled as just described.
  - (d) Calculate the mean and standard deviation of your histogram, and plot it
  - (e) Repeat the above for  $10^5$  and for  $10^6$ . Discuss your results.

### ⊕ Section 2.8

7. Explore how the empirical error does or does not approach the Bhattacharyya bound as follows:
  - (a) Write a procedure to generate sample points in  $d$  dimensions with a normal distribution having mean  $\boldsymbol{\mu}$  and covariance matrix  $\boldsymbol{\Sigma}$ .
  - (b) Consider  $p(\mathbf{x}|\omega_1) \sim N\left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \mathbf{I}\right)$  and  $p(\mathbf{x}|\omega_2) \sim N\left(\begin{pmatrix} -1 \\ 0 \end{pmatrix}, \mathbf{I}\right)$  with  $P(\omega_1) = P(\omega_2) = 1/2$ . By inspection, state the Bayes decision boundary.
  - (c) Generate  $n = 100$  points (50 for  $\omega_1$  and 50 for  $\omega_2$ ) and calculate the empirical error.
  - (d) Repeat for increasing values of  $n$ ,  $100 \leq n \leq 1000$ , in steps of 100 and plot your empirical error.
  - (e) Discuss your results. In particular, is it ever possible that the empirical error is greater than the Bhattacharyya or Chernoff bound?
8. Consider two one-dimensional normal distributions  $p(x|\omega_1) \sim N(-.5, 1)$  and  $p(x|\omega_2) \sim N(+.5, 1)$  and  $P(\omega_1) = P(\omega_2) = 0.5$ .
  - (a) Calculate the Bhattacharyya bound for the error of a Bayesian classifier.
  - (b) Express the true error rate in terms of an error function,  $\text{erf}(\cdot)$ .

- (c) Evaluate this true error to four significant figures by numerical integration (or other routine).
- (d) Generate 10 points each for the two categories and determine the empirical error using your Bayesian classifier. (You should recalculate the decision boundary for each of your data sets.)
- (e) Plot the empirical error as a function of the number of points from either distribution by repeating the previous part for 50, 100, 200, 500 and 1000 sample points from each distribution. Compare your asymptotic empirical error to the true error and the Bhattacharyya error bound.

**9.** Repeat Computer exercise 8 with the following conditions:

- (a)  $p(x|\omega_1) \sim N(-.5, 2)$  and  $p(x|\omega_2) \sim N(.5, 2)$ ,  $P(\omega_1) = 2/3$  and  $P(\omega_2) = 1/3$ .
- (b)  $p(x|\omega_1) \sim N(-.5, 2)$  and  $p(x|\omega_2) \sim N(.5, 2)$  and  $P(\omega_1) = P(\omega_2) = 1/2$ .
- (c)  $p(x|\omega_1) \sim N(-.5, 3)$  and  $p(x|\omega_2) \sim N(.5, 1)$  and  $P(\omega_1) = P(\omega_2) = 1/2$ .



# Bibliography

- [1] Subutai Ahmad and Volker Tresp. Some solutions to the missing feature problem in vision. In Stephen J. Hanson, Jack D. Cowan, and C. Lee Giles, editors, *Neural Information Processing Systems*, volume 5, pages 393–400, San Mateo, CA, 1993. Morgan Kaufmann.
- [2] Hadar Avi-Itzhak and Thanh Diep. Arbitrarily tight upper and lower bounds on the Bayesian probability of error. *IEEE Transaction on Pattern Analysis and Machine Intelligence*, PAMI-18(1):89–91, 1996.
- [3] Thomas Bayes. An essay towards solving a problem in the doctrine of chances. *Philosophical Transactions of the Royal Society (London)*, 53:370–418, 1763.
- [4] James O. Berger. Minimax estimation of a multivariate normal mean under arbitrary quadratic loss. *Journal of Multivariate Analysis*, 6:256–264, 1976.
- [5] James O. Berger. Selecting a minimax estimator of a multivariate normal mean. *Annals of Statistics*, 10:81–92, 1982.
- [6] James O. Berger. *Statistical Decision Theory and Bayesian Analysis*. Springer-Verlag, New York, NY, 2nd edition, 1985.
- [7] José M. Bernardo and Adrian F. M. Smith. *Bayesian Theory*. John Wiley, New York, NY, 1996.
- [8] Anil Bhattacharyya. On a measure of divergence between two statistical populations defined by their probability distributions. *Bulletin of the Calcutta Mathematical Society*, 35:99–110, 1943.
- [9] Herman Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *Annals of Mathematical Statistics*, 23:493–507, 1952.
- [10] Chao K. Chow. An optimum character recognition system using decision functions. *IRE Transactions*, pages 247–254, 1957.
- [11] Chao K. Chow. On optimum recognition error and reject tradeoff. *IEEE Transactions on Information Theory*, IT-16:41–46, 1970.
- [12] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. Wiley Interscience, New York, NY, 1991.
- [13] Morris H. DeGroot. *Optimal Statistical Decisions*. McGraw Hill, New York, NY, 1970.

- [14] Bradley Efron and Carl Morris. Families of minimax estimators of the mean of a multivariate normal distribution. *Annals of Statistics*, 4:11–21, 1976.
- [15] Thomas S. Ferguson. *Mathematical Statistics: A Decision Theoretic Approach*. Academic Press, New York, NY, 1967.
- [16] Simon French. *Decision Theory: An introduction to the mathematics of rationality*. Halsted Press, New York, NY, 1986.
- [17] Keinosuke Fukunaga. *Introduction to Statistical Pattern Recognition*. Academic Press, New York, NY, 2nd edition, 1990.
- [18] Keinosuke Fukunaga and Thomas F. Krile. Calculation of Bayes recognition error for two multivariate Gaussian distributions. *IEEE Transactions on Computers*, C-18:220–229, 1969.
- [19] Izrail M. Gelfand and Sergei Vasilevich Fomin. *Calculus of Variations*. Prentice-Hall, Englewood Cliffs, NJ, translated from the Russian by Richard A. Silverman edition, 1963.
- [20] Andrew Gelman, John B. Carlin, Hal S. Stern, and Donald B. Rubin. *Bayesian Data Analysis*. Chapman & Hall, New York, NY, 1995.
- [21] David M. Green and John A. Swets. *Signal Detection Theory and Psychophysics*. Wiley, New York, NY, 1974.
- [22] David J. Hand. *Construction and Assessment of Classification Rules*. Wiley, New York, NY, 1997.
- [23] Thomas Kailath. *Linear Systems*. Prentice-Hall, Englewood Cliffs, NJ, 1980.
- [24] Bernard Kolman. *Elementary Linear Algebra*. MacMillan College Division, New York, NY, fifth edition, 1991.
- [25] Pierre Simon Laplace. *Théorie Analytique des Probabilités*. Courcier, Paris, France, 1812.
- [26] Peter M Lee. *Bayesian Statistics: An Introduction*. Edward Arnold, London, UK, 1989. there is no period after the M in his name.
- [27] Dennis V. Lindley. *Making Decisions*. Wiley, New York, NY, 1991.
- [28] Jerzy Neyman and Egon S. Pearson. On the problem of the most efficient tests of statistical hypotheses. *Philosophical Transactions of the Royal Society, London*, 231:289–337, 1928.
- [29] Sheldon M. Ross. *Introduction to Probability and Statistics for Engineers*. John Wiley and Sons, New York, NY, 1987.
- [30] Donald B. Rubin and Roderick J. A. Little. *Statistical Analysis with Missing Data*. John Wiley, New York, NY, 1987.
- [31] Claude E. Shannon. A mathematical theory of communication. *Bell Systems Technical Journal*, 6:379–423, 623–656, 1948.
- [32] George B. Thomas, Jr. and Ross L. Finney. *Calculus and Analytic Geometry*. Addison-Wesley, New York, NY, ninth edition, 1996.

- [33] Julius T. Tou and Rafael C. Gonzalez. *Pattern Recognition Principles*. Addison-Wesley, New York, NY, 1974.
- [34] William R. Uttal. *The psychobiology of sensory coding*. HarperCollins, New York, NY, 1973.
- [35] Abraham Wald. Contributions to the theory of statistical estimation and testing of hypotheses. *Annals of Mathematical Statistics*, 10:299–326, 1939.
- [36] Abraham Wald. *Statistical Decision Functions*. John Wiley, New York, NY, 1950.
- [37] C. T. Wolverton and T. J. Wagner. Asymptotically optimal discriminant functions for pattern classifiers. *IEEE Transactions on Information Theory*, IT-15:258–265, 1969.
- [38] C. Ray Wylie and Louis C. Barrett. *Advanced Engineering Mathematics*. McGraw Hill, New York, NY, sixth edition, 1995.