

# Problem Solving I: Mathematical Techniques

For the basics of dimensional analysis and limiting cases, see chapter 1 of Morin or chapter 2 of Order of Magnitude Physics. Many more examples are featured in The Art of Insight; some particularly relevant sections are 2.1, 5.5, 6.3, 8.2, and 8.3. Other sections will be mentioned throughout the course. There is a total of **82** points.

## 1 Dimensional Analysis

### Idea 1

The dimensions of all physical equations should match on both sides. Sometimes, this constraint alone can determine the answers to physical questions.

Even a statement such as “the speed  $v$  is slow” does not make sense, since we need to compare it to something else with dimensions of speed. A more meaningful statement would be “ $v$  is slow compared to the speed of light”, which would correspond to  $v/c \ll 1$ .

### Example 1: $F = ma$ 2018 B11

A circle of rope is spinning in outer space with an angular velocity  $\omega_0$ . Transverse waves on the rope have speed  $v_0$ , as measured in a rotating reference frame where the rope is at rest. If the angular velocity of the rope is doubled, what is the new speed of transverse waves?

### Solution

To solve this problem by dimensional analysis, we reason about what could possibly affect the speed of transverse waves. The result could definitely depend on the rope’s length  $L$ , mass per length  $\lambda$ , and angular velocity  $\omega_0$ . It could also depend on the tension, but since this tension balances the centrifugal force, it is determined by all of the other quantities. Thus the quantities we have are

$$[L] = \text{m}, \quad [\lambda] = \text{kg/m}, \quad [\omega_0] = 1/\text{s}.$$

Since  $\lambda$  is the only thing with dimensions of mass, it can’t affect the speed, because there is nothing that could cancel out the mass dimension. So the only possible answer is

$$v_0 \sim L\omega_0$$

where the  $\sim$  indicates equality up to a dimensionless constant, which cannot be found by dimensional analysis alone. In practice, the constant usually won’t be too big or too small, so  $L\omega_0$  is a decent estimate of  $v_0$ . But even if it isn’t, the dimensional analysis tells us the scaling: if  $\omega_0$  is doubled, the new speed is  $2v_0$ .

### Example 2

Find the dimensions of the magnetic field.

**Solution**

To do this, we just think of some simple equation involving  $B$ , then solve for its dimensions. For example, we know that  $\mathbf{F} = q(\mathbf{v} \times \mathbf{B})$ , so

$$[B] = \frac{[F]}{[q][v]} = \frac{\text{kg} \cdot \text{m}}{\text{s}^2} \frac{1}{\text{C}} \frac{1}{\text{m/s}} = \frac{\text{kg}}{\text{C} \cdot \text{s}}.$$

[2] **Problem 1.** Find the dimensions of power, the gravitational constant  $G$ , the permittivity of free space  $\epsilon_0$ , and the ideal gas constant  $R$ .

[2] **Problem 2.** Some questions about vibrations.

- (a) The typical frequency  $f$  of a vibrating star depends only on its radius  $R$ , density  $\rho$ , and the gravitational constant  $G$ . Use dimensional analysis to find an expression for  $f$ , up to a dimensionless constant. Then estimate  $f$  for the Sun, looking up any numbers you need.
- (b) The typical frequency  $f$  of a small water droplet freely vibrating in zero gravity could depend on its radius  $R$ , density  $\rho$ , surface tension  $\gamma$ , and the gravitational constant  $G$ . Argue that at least one of these parameters doesn't matter, and find an expression for  $f$  up to a dimensionless constant.

[3] **Problem 3.** Some questions about the speed of waves. For all estimates, you can look up any numbers you need.

- (a) The speed of sound in an ideal gas depends on its pressure  $p$  and density  $\rho$ . Explain why we don't have to use the temperature  $T$  or ideal gas constant  $R$  in the dimensional analysis, and then estimate the speed of sound in air.
- (b) The speed of sound in a fluid depends only on its density  $\rho$  and bulk modulus  $B = -V dP/dV$ . Estimate the speed of sound in water, which has  $B = 2.1 \text{ GPa}$ .

The speed of waves on top of the surface of water can depend on the water depth  $h$ , the wavelength  $\lambda$ , the density  $\rho$ , the surface tension  $\gamma$ , and the gravitational acceleration  $g$ .

- (c) Find the speed of capillary waves, i.e. water waves of very short wavelength, up to a dimensionless constant.
- (d) Find the speed of long-wavelength waves in very deep water, up to a dimensionless constant.

[3] **Problem 4** (Morin 1.5). A particle with mass  $m$  and initial speed  $v$  is subject to a velocity-dependent damping force of the form  $bv^n$ .

- (a) For  $n = 0, 1, 2, \dots$ , determine how the stopping time and stopping distance depend on  $m$ ,  $v$ , and  $b$ .
- (b) Check that these results actually make sense as  $m$ ,  $v$ , and  $b$  are changed, for a few values of  $n$ . You should find something puzzling going on. (Hint: to resolve the problem, it may be useful to find the stopping time explicitly in a few examples.)

**Idea 2**

Dimensional analysis applies everywhere. The argument of any function that is not a monomial, such as  $\sin x$ , must have no dimensions. The derivative  $d/dx$  has the opposite dimensions to  $x$ , and the  $dx$  in an integral has the same dimensions as  $x$ .

**Example 3**

We are given the integral

$$\int_{-\infty}^{\infty} e^{-x^2} dx = \sqrt{\pi}.$$

Find the value of the integral

$$\int_{-\infty}^{\infty} e^{-ax^2} dx.$$

**Solution**

In the first equation,  $x$  must be dimensionless, so both sides are dimensionless. The second equation would also be consistent if both  $x$  and  $a$  were dimensionless, but we can do better. Suppose we arbitrarily assign  $x$  dimensions of length,  $[x] = \text{m}$ . Then to make the argument of the exponential dimensionless,  $a$  must have dimensions  $[a] = \text{m}^{-2}$ . The dimensions of the left-hand side are  $[dx] = \text{m}$ . In order to make the dimensions work out on the right-hand side, we must have

$$\int_{-\infty}^{\infty} e^{-ax^2} dx \propto \frac{1}{\sqrt{a}}.$$

To find the value of the constant, treat  $x$  and  $a$  as dimensionless again. Then we know the answer must reduce to  $\sqrt{\pi}$  when  $a = 1$ , so

$$\int_{-\infty}^{\infty} e^{-ax^2} dx = \sqrt{\frac{\pi}{a}}.$$

This procedure is completely equivalent to using  $u$ -substitution to nondimensionalize everything, but may be faster to see. In general, for all integrals except for the simplest ones, you should use either dimensional analysis or  $u$ -substitution to reduce the integral to a dimensionless one.

**Remark**

Consider the value of the definite integral

$$\int_{-\infty}^y e^{-x^2} dx.$$

You can try all day to compute the value of this integral, using all the integration tricks you know, but nothing will work. The function  $e^{-x^2}$  simply doesn't have an antiderivative in terms of the functions you already know, i.e. in terms of polynomials, exponents and logarithms, and trigonometric functions (for more discussion, see [here](#)).

If you ask a computer algebra system like Mathematica, it'll spit out something like  $\text{erf}(x)$ , which is defined by being an antiderivative of  $e^{-x^2}$ . But is this really an “analytic” solution? Isn't that just saying “the integral of  $e^{-x^2}$  is equal to the integral of  $e^{-x^2}$ ”? Well, like many things in math, it depends on what the meaning of the word “is” is.


The fact is, the set of functions we regard as “elementary” is arbitrary; we just choose a set that's big enough to solve most of the problems we want, and small enough to attain fluency with. (Back in the days before calculators, it just meant all the functions whose values were tabulated in the references on hand.) If you're uncomfortable with  $\text{erf}(x)$ , note that a similar thing would happen if a middle schooler asked you what the ratio of the opposite to adjacent sides of a right triangle is. You'd say  $\tan(x)$ , but they could say it's tautological, because the only way to define  $\tan(x)$  at the middle school level is the ratio of opposite to adjacent sides. Similarly,  $1/x$  has no elementary antiderivative – unless you count  $\log(x)$  as elementary, but  $\log(x)$  is simply *defined* to be such an antiderivative. It's all tautology, but it's still useful.

- [1] **Problem 5** (SFM 1.11). Find the integral

$$\int \frac{dx}{x^2 + 1}$$

using a trigonometric substitution. Using dimensional analysis, find the integral

$$\int \frac{dx}{x^2 + a^2}.$$

- [2] **Problem 6** (SFM 1.15). Derive Kepler's Third Law using only dimensional analysis. Why do you think people didn't figure out this argument 2000 years ago?
- [2] **Problem 7**. In particle physics, it is conventional to set  $\hbar = c = 1$ , resulting in equations with seemingly incorrect units. For example, it is said that the mass of the Higgs boson is about 125 GeV, where 1 eV is the energy gained by an electron accelerated through a voltage difference of 1 V. Energy doesn't have the same units as mass, but the units can be fixed by adding appropriate factors of  $\hbar$  and  $c$ . Do this, and find the mass of the Higgs boson in kilograms.
- [3] **Problem 8**.  USAPhO 2002, problem A3.

#### Example 4

The Schrodinger equation for an electron in the electric field of a proton is

$$-\frac{\hbar^2}{2m}\nabla^2\psi - \frac{e^2}{4\pi\epsilon_0 r}\psi = E\psi.$$

Estimate the size of the hydrogen atom.

**Solution**

This is yet another dimensional analysis problem: there is only one way to form a length using the quantities given above. We have

$$[m] = \text{kg}, \quad [\hbar] = \text{J} \cdot \text{s} = \text{kg m}^2 \text{s}^{-1}, \quad [e^2/4\pi\epsilon_0] = \text{J} \cdot \text{m} = \text{kg m}^3 \text{s}^{-2}.$$

Doing dimensional analysis, the only length scale is the Bohr radius,

$$a_0 = \frac{4\pi\hbar^2\epsilon_0}{me^2} \sim 10^{-10} \text{ m}.$$

I've thrown in a  $4\pi$  above because  $\epsilon_0$  always appears in the equations as  $4\pi\epsilon_0$ . The dimensional analysis would be valid without this factor, but as you'll see in problem 11, if you don't include it then annoying compensating factors of  $4\pi$  will appear elsewhere.

Classically (i.e. without  $\hbar$ ), there is no way to form a length, and hence there should be no classically stable radius for the atom. (This was one of the arguments used by Bohr to motivate quantum mechanics; it appears in the beginning of his paper introducing the Bohr model.) Once we introduce  $\hbar$ , there are three dimensionful parameters in the problem, as listed above. And there are exactly three fundamental dimensions. So there is only one way to create a length, which we found above, one way to create a time, one way to create an energy, and so on. This means that the solutions to the Schrodinger equation above look qualitatively the same no matter what these parameters are; all that changes are the overall length, time, and energy scales. In problem 11, you'll investigate how this conclusion changes when we add more dimensionful parameters.

Dimensional analysis is especially helpful with scaling relations. For example, a question might ask you how the radius of the hydrogen atom would change in a world where the electron mass was twice as large. You would solve this problem in the exact same way as the example above, using dimensional analysis to show that  $a_0 \propto 1/m$ .

- [3] **Problem 9.** In this problem we'll continue the dimensional analysis of the Schrodinger equation.
- Estimate the typical energy scale of quantum states of the hydrogen atom, as well as the typical "velocity" of the electron, using dimensional analysis.
  - Do the same for one-electron helium, the system consisting of a helium nucleus (containing two protons) and one electron.
  - Estimate the electric field needed to rip the electron off the hydrogen atom.

**Idea 3: Buckingham Pi Theorem**

Dimensional analysis can't always pin down the form of the answer. If one has  $N$  quantities with  $D$  independent dimensions, then one can form  $N - D$  independent dimensionless quantities. Dimensional analysis can't say how the answer depends on them.

A familiar but somewhat trivial example is the pendulum: its period depends on  $L$ ,  $g$ , and the amplitude  $\theta_0$ , three quantities which contain two dimensions (length and time). Hence we can form one dimensionless group, which is clearly just  $\theta_0$  itself. The period of a pendulum is  $T = f(\theta_0)\sqrt{L/g}$ .

**Example 5:  $F = ma$  2014 12**

A paper helicopter with rotor radius  $r$  and weight  $W$  is dropped from a height  $h$  in air with a density of  $\rho$ . Assuming the helicopter quickly reaches terminal velocity, use dimensional analysis to analyze the total flight time  $T$ .

**Solution**

The answer can only depend on the parameters  $r$ ,  $W$ ,  $h$ , and  $\rho$ . There are four quantities in total, but three dimensions (mass, length, and time), so by the Buckingham Pi theorem we can form one independent dimensionless quantity. In this case, it's clearly  $r/h$ . Continuing with routine dimensional analysis, we find

$$T = f(r/h) h^2 \sqrt{\frac{\rho}{W}}.$$

The form of this expression is a bit arbitrary; for instance, we could also have written  $f(r/h)r^2$  in front, or even  $f(r/h)r^{37}h^{-35}$ . These adjustments just correspond to pulling factors of  $r/h$  out of  $f$ , not to changing the actual result.

This is as far as we can get with dimensional analysis alone, but we can go further using physical reasoning. If the helicopter quickly reaches terminal velocity, then it travels at a constant speed. So we must have  $T \propto h$ , which means that  $f(x) \propto x$ , and

$$T \propto rh \sqrt{\frac{\rho}{W}}.$$

**Example 6**

An hourglass is constructed with sand of density  $\rho$  and an orifice of diameter  $d$ . When the sand level above the orifice is  $h$ , what is the mass flow rate  $\mu$ ?

**Solution**

The answer can only depend on  $\rho$ ,  $d$ ,  $h$ , and  $g$ . The Buckingham Pi theorem gives

$$\mu = f(h/d) \rho \sqrt{gd^5}.$$

That's as far as we can get with dimensional analysis; to go further we need to know more about sand. If we were dealing with an ideal fluid, then the flow speed would be  $v = \sqrt{2gh}$  by Torricelli's law, which means the flow rate has to be proportional to  $\sqrt{h}$ . Then  $f(x) \propto \sqrt{x}$ , giving the result  $\mu \propto \rho d^2 \sqrt{gh}$ . This is a good estimate as long as the orifice isn't so small that viscosity starts to dominate.

But this isn't how sand works. Sand is a granular material, whose motion is dominated by the friction between sand grains. So the higher pressure doesn't actually propagate to the orifice, and the flow rate is independent of  $h$ , which is apparent from watching an hourglass run. Then  $f(x)$  is a constant, giving  $\mu \propto \rho \sqrt{gd^5}$ , which is [experimentally verified](#).

**Remark**

One has to be a little careful with the Buckingham Pi theorem. For example, if all we had were 3 speeds  $v_i$ , we can form two dimensionless quantities:  $v_1/v_2$  and  $v_1/v_3$ . (The quantity  $v_2/v_3$  is not independent, since it is the quotient of these two.) But there are 3 quantities with 2 dimensions (length and time), so we expect only 1 dimensionless quantity.

The problem is that the two dimensions really aren't independent: for any quantity built from the  $v_i$ , a power of length always comes with an inverse power of time, so there's only one independent dimension. These considerations can be put on a more rigorous footing in linear algebra, where the Buckingham Pi theorem is merely a special case of the rank-nullity theorem. If you're ever in doubt, you can just forget about the theorem and play with the equations directly.

**Remark**

Dimensional analysis is an incredibly common tool in Olympiad physics because it lets you say a lot even without much advanced knowledge. If a problem ever says to find some quantity "up to a constant/dimensionless factor", or how that quantity scales as another quantity changes, or what that quantity is proportional to, it's almost certainly asking you to do dimensional analysis. Another giveaway is if the problem *looks* extremely technical and advanced, because they can't actually be.

[3] **Problem 10** (Insight). In this problem we'll do one of the most famous dimensional analyses of all time: estimating the yield of the first atomic bomb blast. Such a blast will create a shockwave of air, which reaches a radius  $R$  at time  $t$  after the blast. The air density is  $\rho$ , and we want to estimate the blast energy  $E$ .

- (a) Declassified photographs of the blast indicate that  $R \approx 100$  m at time  $t \approx 15$  ms. The density of air is  $\rho \approx 1$  kg/m<sup>3</sup>. Estimate the blast energy  $E$ .
- (b) How much mass-energy (in grams) was used up in this blast?
- (c) If we measure the entire function  $R(t)$ , what general form would we expect it to have, if this dimensional analysis argument is correct?

**Remark**

The British physicist G. I. Taylor performed the dimensional analysis in problem 10 upon seeing a picture of the first atomic blast in a magazine. The result was so good that the physicists at the Manhattan project thought their security had been breached!

During World War II, the exact value of the critical mass needed to set off a nuclear explosion was important and nontrivial information. The Nazi effort to make a bomb had been stopped by Werner Heisenberg's huge overestimation of this quantity, and after the war, the specific value was kept a closely guarded secret. That is, it was until 1947, when a Chinese physicist [got the answer](#) using a rough estimate that took four lines of algebra.

[5] **Problem 11.** We now consider the Schrodinger equation for the hydrogen atom in greater depth. We begin by switching to dimensionless variables, which is useful for the same reason that writing integrals in terms of dimensionless variables is: it highlights what is independent of unit choices.

- (a) Define a dimensionless length variable  $\tilde{r} = r/a_0$ , where  $a_0$  is the length scale found in example 4. The  $\nabla^2$  term in the Schrodinger equation is a second derivative, the 3D generalization of  $d^2/dx^2$ . Using the chain rule, argue that

$$\tilde{\nabla}^2 = a_0^2 \nabla^2$$

where  $\tilde{\nabla}$  is the gradient with respect to  $\tilde{r}$ .

- (b) Similarly define a dimensionless energy  $\tilde{E} = E/E_0$ , using the energy scale  $E_0$  found in problem 9. Show that the Schrodinger equation can be written in a form like

$$-\tilde{\nabla}^2 \psi - \frac{1}{\tilde{r}} \psi = \tilde{E} \psi$$

Here I've suppressed all dimensionless constants, like factors of 2, because they depend on how you choose to define  $E_0$  and don't really matter at this level of precision.

The result of this part confirms what we concluded above: solutions to the Schrodinger equation don't qualitatively depend on the values of the parameters, because they all come from scaling a solution to this one dimensionless equation appropriately.

- (c) This is no longer true in relativity, where the total energy is

$$E = \sqrt{p^2 c^2 + m^2 c^4}.$$

Assuming  $p \ll mc$ , perform a Taylor expansion to show that the next term is  $Ap^4$ , and find the coefficient  $A$ . (If you don't know how to do this, work through the next section first.)

- (d) In quantum mechanics, the momentum is represented by a gradient,  $p \rightarrow -i\hbar\nabla$ . (We will see why in **X1**.) Show that the Schrodinger equation with the first relativistic correction is

$$-\frac{\hbar^2}{2m} \nabla^2 \psi - \frac{e^2}{4\pi\epsilon_0 r} \psi + \hbar^4 A \nabla^4 \psi = E \psi.$$

- (e) Since there is now one more dimensionful quantity in the game, it is possible to combine the quantities to form a dimensionless one. Create a dimensionless quantity  $\alpha$  that is proportional to  $e^2/4\pi$ , then numerically evaluate it. This is called the fine structure constant. It serves as an "objective" measure of the strength of the electromagnetic force, because it is dimensionless, and hence its value doesn't depend on an arbitrary unit system.
- (f) As the number of protons in the nucleus increases, the relativistic correction becomes more important. Estimate the atomic number  $Z$  where the correction becomes very important.

You probably won't see any differential equations as complex as the ones in the above problem anywhere in Olympiad physics, but the key idea of using dimensionless quantities to simplify and clarify the physics can be used everywhere.

[5] **Problem 12.**  IPhO 2007, problem "blue". This problem applies thermodynamics and dimensional analysis in some exotic contexts.



**Example 7**

Estimate the Young's modulus for a material with interatomic separation  $a$  and typical atomic bond energy  $E_b$ . Use this to estimate the spring constant of a rod of area  $A$  and length  $L$ , as well as the speed of sound, if each atom has mass  $m$ .

**Solution**

This example is to get you comfortable with the Young's modulus  $Y$ , which occasionally comes up. It is defined in terms of how much a material stretches as it is pulled apart,

$$Y = \frac{\text{stress}}{\text{strain}} = \frac{\text{restoring force/cross-sectional area}}{\text{change in length/length}}.$$

The Young's modulus is a useful way to characterize materials, because unlike the spring constant, it doesn't depend on the shape of the material. For example, putting two identical springs side-by-side doubles the spring constant, because they both contribute to the force. But since the stress is the force per area, it's unchanged. Similarly, putting two identical springs end-to-end halves the spring constants, because they both stretch, but since the strain is change in length per length, it's unchanged. So you would quote a material's Young's modulus instead of its spring constant, for the same reason you would quote a material's resistivity instead of its resistance.

We note that  $Y$  has the dimensions of energy per length cubed, so

$$Y \sim \frac{E_b}{a^3}$$

solely by dimensional analysis. (Of course, for this dimensional analysis to work, one has to understand why  $E_b$  and  $a$  are the only relevant quantities. It's because  $Y$ , or equivalently the spring constant  $k$ , determines the energy stored in a stretched spring. But microscopically this comes from the energy stored in interatomic bonds when they're stretched. So the relevant energy scale is the bond energy  $E_b$ , and the relevant distance scale is  $a$ , because that determines how many bonds get stretched, and by how much.)

To relate  $Y$  to the spring constant of a rod, note that

$$Y = \frac{F/A}{\Delta L/L} = \frac{L}{A} \frac{F}{\Delta L} = k \frac{L}{A}$$

for a rod, giving the estimate  $k \sim AE_b/La^3$ . This is correct to within an order of magnitude!

To relate  $Y$  to the speed of sound, note that the sound speed, like most wave speeds, depends on the material's inertia and its restoring force against distortions. Since the speed of sound doesn't depend on the extrinsic features of a metal object, such as a length, both of these should be measured intrinsically. The intrinsic measure of inertia is the mass density  $\rho \sim m/a^3$ , while the intrinsic measure of restoring force is just  $Y$ . By dimensional analysis,

$$v \sim \sqrt{\frac{Y}{\rho}} \sim \sqrt{\frac{E_b/a^3}{m/a^3}} \sim \sqrt{\frac{E_b}{m}}.$$

This is also reasonably accurate. For example, in diamond,  $E_b \sim 1 \text{ eV}$  (a typical atomic energy scale), while a carbon nucleus contains 12 nucleons, so to the nearest order of magnitude,  $m \sim 10m_p$ , where a useful fact is  $m_p \sim 1 \text{ GeV}/c^2$ . Thus,

$$v \sim \sqrt{\frac{1 \text{ eV}}{10^{10} \text{ eV}}} c \sim 10^{-5} c \sim 3 \text{ km/s}$$

which is roughly right. (The true answer is 12 km/s.)

Amazingly, we can get an even rougher estimate of  $v$  for any solid in terms of nothing besides fundamental constants. To be very rough, the binding energy is on the order of that of hydrogen. As you found in problem 9, this is, by dimensional analysis,

$$E_b \sim \frac{1}{4\pi\epsilon_0} \frac{e^2}{a_0} \sim m_e \left( \frac{e^2}{4\pi\epsilon_0 \hbar} \right)^2.$$

We take the nuclear mass to be very roughly the proton mass  $m_p$ , which gives

$$\frac{v}{c} \sim \sqrt{\frac{m_e}{m_p} \left( \frac{e^2}{4\pi\epsilon_0 \hbar c} \right)^2} \sim \alpha \sqrt{\frac{m_e}{m_p}}$$

where  $\alpha$  is as found in problem 11. This expresses the speed of sound in terms of the dimensionless strength of electromagnetism  $\alpha$ , the electron to proton mass ratio, and the speed of light. Of course, the approximations we have made here have been so rough that now the answer is off by *two* orders of magnitude, but now we know how the answer would change if the fundamental constants did.

Estimates as simple as these can be surprising to even seasoned physicists: in 2020, the simple estimate above was rediscovered and [published](#) in one of the top journals in science. If you want to learn how to do more of these estimates, [this paper](#) is a good starting point.

### Remark

A warning: from these examples, you could get the idea that dimensional analysis gives you nearly godlike powers, and the ability to write down the answer to most physics problems instantly. In reality, it only works if you're pretty sure your physical system depends on only about 3 or 4 variables – and the hard part is often finding *which* variables matter. For example, as we saw above, you can't get Kepler's third law for free because that requires knowing the dimensions of  $G$ , which require knowing that gravity is an inverse square law in the first place, a luxury Kepler didn't have. And as another example, we couldn't have figured out  $E = mc^2$  long before Einstein, as who would have thought that the speed of light had anything to do with the energy of a lump of matter? Without the framework of relativity, it seems as irrelevant as the speed of sound or the speed of water waves. To illustrate this point, we consider two contrasting examples below.

In reality, dimensional analysis is best for problems where it's easy to see which variables

matter, problems where you're explicitly told what variables matter, and for checking your work, which you should get in the habit of doing all the time!

### Example 8

Archaeologists and biologists have discovered that the famed T. Rex was essentially a gigantic chicken. Suppose that a T. Rex is about  $N = 20$  times larger in scale than a chicken. Compared to a chicken, how much larger is its weight, cross-sectional area of bone, walking speed, and maximum jump height?

### Solution

These kinds of biological scaling questions are fun to think about, though the accuracy of the results is questionable. Here's the typical reasoning:

- Since the densities should match, the weight should scale with the volume, so as  $N^3$ .
- Since the maximum compressive pressure that bone can take should be the same, the bone area should scale with the weight, so also as  $N^3$ . That is, the width of the bones scales as  $N^{3/2}$ , while their length  $L$  scales only as  $N$ . This is one of the reasons it's impractical to have huge land animals; the biggest animals now are all whales.
- As a very crude model of walking, we can think of the legs as swinging like a free pendulum. The length of one step is proportional to  $L$ , while the period of the steps is proportional to  $\sqrt{L}$ . Thus, the walking speed scales as  $\sqrt{L} \propto \sqrt{N}$ .
- The energy stored in the muscle cells scales with the volume, but the mass also scales with the volume. Since the jump height satisfies  $E = mgh$  where  $E$  is the energy stored by the muscles,  $h \propto N^0$ . And indeed, a flea can jump nearly as high as a human!

### Example 9

A person with density  $\rho$  and total energy  $E$  stored in their muscles can jump to a height  $h$  in gravity  $g$ . How high would they be able to jump in gravity  $10g$ ?

### Solution

By dimensional analysis, the only possible answer is

$$h \propto \left( \frac{E}{\rho g} \right)^{1/4}$$

which means that in gravity  $10g$ , they can jump to height  $h/10^{1/4}$ .

But this is completely wrong! In gravity  $10g$ , a person wouldn't be able to jump at all; they'd be so crushed by their own weight that they wouldn't even be able to stand. The actual answer depends on details of the biomechanics of muscles and bone, which involve more dimensionful quantities than just the total energy  $E$ . So, as remarked above, you can't

solve literally any problem by just listing a few relevant quantities and doing dimensional analysis – you need to make sure those are the *only* relevant quantities.

## 2 Approximations

### Idea 4: Taylor Series

For small  $x$ , a function  $f(x)$  may be approximated as

$$f(x) = f(0) + xf'(0) + \frac{x^2}{2}f''(0) + \dots + \frac{x^n}{n!}f^{(n)}(0) + O(x^{n+1})$$

where  $O(x^{n+1})$  stands for an error term which grows at most as fast as  $x^{n+1}$ .

There are a few Taylor series that are essential to know. The most important are

$$\exp x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + O(x^4), \quad \log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - O(x^4)$$

and the small angle approximations

$$\sin x = x - \frac{x^3}{6} + O(x^5), \quad \cos x = 1 - \frac{x^2}{2} + O(x^4).$$

Another Taylor series you learned long before calculus class is

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + O(x^4)$$

Usually you'll only need the first one or two terms, but for practice we'll do an example with more. If any of these results aren't familiar, you should rederive them!

### Example 10

Find the Taylor series for  $\tan x$  up to, and including the fourth order term.

### Solution

By the fourth order term, we mean the term proportional to  $x^4$ . (Not the fourth nonzero term, which would be  $O(x^7)$ .) Of course,  $\tan x$  is an odd function, so the  $O(x^4)$  term is zero, which means we only need to expand up to  $O(x^3)$ . That means we can neglect  $O(x^4)$  terms and higher everywhere in the computation, subject to some caveats we'll point out later.

By definition, we have

$$\tan x = \frac{\sin x}{\cos x} = \frac{x - x^3/6 + O(x^5)}{1 - x^2/2 + O(x^4)}.$$

However, it's a little tricky because we have a Taylor series in a denominator. There are two ways to deal with this. We could multiply both sides by  $\cos x$ , and expand  $\tan x$  in a Taylor

series with unknown coefficients. Then we would get a system of equations that will allow us to solve for the coefficients recursively.

A faster method is to use the Taylor series for  $1/(1-x)$ . We have

$$\frac{1}{1-u} = 1 + u + O(u^2)$$

and substituting  $u = x^2/2 - O(x^4)$  gives

$$\frac{1}{\cos x} = 1 + \frac{x^2}{2} + O(x^4).$$

Therefore, we conclude

$$\tan x = (x - x^3/6 + O(x^5))(1 + x^2/2 + O(x^4)) = x + x^3/3 + O(x^5).$$

Here I was fairly careful with writing out all the error terms and intermediate steps, but as you get better at this process, you'll be able to do it faster. (Of course, one could also have done this example by just directly computing the Taylor series of  $\tan x$  from its derivatives. This is possible, but for more complicated situations it's generally not a good idea, because computing high derivatives of a complex expression tends to get very messy. It's better to just Taylor expand the individual pieces and combine the results, as we did here.)

### Remark

Finding series up to a given order can be subtle. For example, if you want to compute an  $O(x^4)$  term, it is *not* always enough to expand everything up to  $O(x^4)$ , because powers of  $x$  might cancel. To illustrate this, the last step here is wrong:

$$\tan x = \frac{x^3 \sin x}{x^3 \cos x} = \frac{x^4 + O(x^6)}{x^3 + O(x^5)} = x + O(x^5).$$

- [2] **Problem 13.** Find the Taylor series for  $1/\cos x$  up to and including the fourth order ( $O(x^4)$ ) term.
- [2] **Problem 14.** Extend the computation above to get the  $x^5$  term in the Taylor series for  $\tan x$ .
- [3] **Problem 15.** For small  $x$ , approximate the quantity

$$\frac{x^2 e^x}{(e^x - 1)^2} - 1$$

to lowest order. That is, find the first nonzero term in the Taylor series. If you don't take enough terms in the Taylor series to begin with, you'll get an answer of zero, indicating you approximated too loosely. But if you take too many, the computation will get extremely messy. Your final answer should contain a 12, which is actually the same 12 as in the classic result  $1 + 2 + 3 + \dots = -1/12$ .

- [2] **Problem 16.** The function  $\cos^{-1}(1-x)$  does not have a Taylor series about  $x=0$ . However, it does have a series expansion about  $x=0$  in a different variable. What is this variable, and what's the first term in the series? (Optionally, can you find higher order terms in the series?)

**Idea 5: Binomial Theorem**

When the quantity  $xn$  is small, it is useful to use the binomial theorem,

$$(1+x)^n = 1 + xn + O(x^2n^2).$$

It applies even when  $n$  is not an integer. In particular,  $n$  can be very large, very small, or even negative. The extra terms will be small as long as  $xn$  is small. If desired, one can find higher terms using binomial coefficients,

$$(1+x)^n = \sum_{m=0}^{\infty} \binom{n}{m} x^m$$

where the definition of the binomial coefficient is formally extended to arbitrary real  $n$ .

The binomial theorem is one of the most common approximations in physics. It's really just taking the first two terms in the Taylor series of  $(1+x)^n$ , but we give it a name because it's so useful.

- [1] **Problem 17.** Suppose the period of a pendulum is one second, and recall that

$$T = 2\pi\sqrt{\frac{L}{g}}.$$

If the length is increased by 3% and  $g$  is increased by 1%, use the binomial theorem to estimate how much the period changes. This kind of thinking is extremely useful when doing experimental physics, and you should be able to do it in your head.

- [2] **Problem 18.** Consider an electric charge  $q$  placed at  $x = 0$  and a charge  $-q$  placed at  $x = d$ . The electric field along the  $x$  axis is then

$$E(x) = \frac{q}{4\pi\epsilon_0} \left( \frac{1}{x^2} - \frac{1}{(x-d)^2} \right).$$

For large  $x$ , use the binomial theorem to approximate the field.

- [3] **Problem 19.** Some exercises involving square roots.

- Manually find the Taylor series for  $\sqrt{1+x}$  up to second order, and verify they agree with the binomial theorem.
- Approximate  $\sqrt{1+2x+x^2}$  for small  $x$  using the binomial theorem. Does the result match what you expect? If not, how can you correct it?

**Example 11: Birthday Paradox**

If you have  $n$  people in a room, around how large does  $n$  have to be for there to be at least a 50% chance of two people sharing the same birthday?

**Solution**

Imagine adding people one at a time. The second person has a  $1/365$  chance of sharing a birthday with the first. If they don't share a birthday, the third person has a  $2/365$  chance of sharing a birthday with either, and so on. So a decent estimate for  $n$  is the  $n$  where

$$\left(1 - \frac{1}{365}\right) \left(1 - \frac{2}{365}\right) \cdots \left(1 - \frac{n}{365}\right) \approx \frac{1}{2}.$$

The surprising point of the birthday paradox is that  $n \ll 365$ . So we can use the binomial theorem in reverse, approximating the left-hand side as

$$\left(1 - \frac{1}{365}\right) \left(1 - \frac{1}{365}\right)^2 \cdots \left(1 - \frac{1}{365}\right)^n \approx \left(1 - \frac{1}{365}\right)^{n^2/2}$$

which is valid since  $n/365$  is small. It's tempting to use the binomial theorem again to write

$$\left(1 - \frac{1}{365}\right)^{n^2/2} \approx 1 - \frac{n^2}{2 \cdot 365} = \frac{1}{2}$$

which gives  $n = 19$ . However, this is a bad approximation, because the binomial theorem only works if  $(n^2/2)(1/365)$  is very small, but here we've set it to  $1/2$ , which isn't particularly small. Since the series expansion variable is  $1/2$ , each term in the series expansion is *roughly*  $1/2$  as big as the last (ignoring numerical coefficients), so we expect to be off by about  $(1/2)^2 = 25\%$ .

The binomial theorem is an expansion for  $(1+x)^y$  which works when  $xy$  is small. Here  $xy$  isn't small, and we instead want an approximation that works when only  $x$  is small. One trick to dealing with an annoying exponent is to take the logarithm, since that just turns it into a multiplicative factor. Note that

$$\log((1+x)^y) = y \log(1+x) \approx yx$$

by Taylor series, which implies that

$$(1+x)^y \approx e^{yx}$$

when  $x$  is small, an important fact which you should remember. So we have

$$\left(1 - \frac{1}{365}\right)^{n^2/2} \approx e^{-n^2/2(365)} = \frac{1}{2}$$

and solving gives  $n = 22.5$ . We should round up since  $n$  is actually an integer, giving  $n = 23$ , which is indeed the exact answer.

**Remark**

Precisely how accurate is the approximation  $(1+x)^y \approx e^{yx}$ ? Note that the only approximate step used to derive it was taking  $\log(1+x) \approx x$ , which means we can get the corrections by

expanding to higher order. If we take the next term,  $\log(1+x) \approx x - x^2/2$ , then we find

$$(1+x)^y \approx e^{yx} e^{-x^2 y/2}.$$

Note that because we are approximating the logarithm of the quantity we want, the next correction is multiplicative rather than additive; we'll see a similar situation with Stirling's approximation in **T2**. So, while we motivated the approximation by imagining  $x \ll 1$ , the real condition we need to have a good approximation is  $x^2 y \ll 1$ . For instance, in the previous example we had  $x^2 y/2 = (22.5/365)^2/4 = 0.1\%$ , so our answer was quite accurate.

- [2] **Problem 20.** Find a series approximation for  $x^y$ , given that  $y$  is small and  $x$  is neither small nor exponentially huge. (Hint: to check if you have it right, you can try concrete numbers, such as  $y = 0.01$  and  $x = 10$ . The series expansion variable may look a bit unusual.)
- [3] **Problem 21.** Consider the equation  $\epsilon x^3 - x^2 + 1 = 0$ , where  $\epsilon$  is small. Find approximate expressions for all three roots of this equation, up to and including terms of order  $\epsilon$ .
- [3] **Problem 22.** Sometimes, transcendental equations can be solved numerically by “iteration”. This is especially useful in Olympiad contexts where only basic calculators are allowed.
- Take a scientific calculator (in radians), put in any number, and press the “cos” button many times. Convince yourself that the final number you get is the unique solution to  $x = \cos x$ .
  - What are the key features of the graphs of  $x$  and  $\cos x$  that made this work? For example, why doesn't pressing  $\cos^{-1}$  repeatedly give the same result? As another example, since  $x = \sin x$  has a unique solution, why does repeatedly pressing sin not work so well?
  - Find a nonzero solution for  $x = \tan(x/2)$ .
  - Find a nonzero solution for  $e^x - 1 = 2x$ .
  - Find a positive solution for  $x^x = e$ .

If you ever have trouble getting iteration to work, another useful technique is ordinary binary search. It's not as flashy or fast as iteration, but it's easy and reliable. Numeric solutions like these will appear in future problem sets.

### Example 12

In units where  $c = 1$ , the Lorentz factor is defined as

$$\gamma = \frac{1}{\sqrt{1-v^2}}.$$

Suppose that a particle traveling very close to the speed of light has  $\gamma = 10^8$ . Find the difference  $\Delta v$  between its speed and the speed of light.



**Solution**

This problem looks easy; by some trivial algebra we find

$$\Delta v = 1 - \sqrt{1 - 1/\gamma^2}.$$

But when you plug this into a cheap scientific calculator, you get *zero*, or something that's quite far from the right result. The problem is that we are trying to find a small quantity  $\Delta v$  by subtracting two nearby, much larger quantities. But the calculator has limited precision, and it ends up rounding  $1 - 1/\gamma^2 = 1 - 10^{-16}$  a bit, giving a completely wrong answer!

Instead, we can apply the binomial theorem to find

$$\Delta v \approx \frac{1}{2\gamma^2} + O(1/\gamma^4).$$

This is no longer the exact answer, but it's a great approximation, because the error term is around  $1/\gamma^2 \sim 10^{-16}$  times as small as the answer, and it's easy for a calculator to evaluate. The lesson is that it's better to be accurate in practice than to be precise in theory.

**Remark**

If these questions seem complicated, rest assured that 90% of approximations on the USAPhO and IPhO boil down to using

$$\sin x \approx x, \quad \cos x \approx 1 - x^2/2, \quad (1+x)^n \approx 1 + nx, \quad e^x \approx 1 + x, \quad \log(1+x) \approx x.$$

I've given you a lot of subtle situations above, but it's these that you have to know by heart. Almost all situations where you will use these will look like problem 17 or problem 18.

**3 Limiting Cases****Idea 6**

Limiting cases can be used to infer how the answer to a physical problem depends on its parameters. It is primarily useful for remembering the forms of formulas, but can also be powerful enough to solve multiple choice questions by itself.

**Example 13**

What is the horizontal range of a rock thrown with speed  $v$  at an angle  $\theta$  to the horizontal?

**Solution**

This result is easy to derive, but dimensional analysis and extreme cases can be used to recover the result too. The answer can only depend on  $v$ ,  $g$ , and  $\theta$ , so by dimensional analysis it is proportional to  $v^2/g$ . This is sensible, since the range increases with  $v$  and decreases

with  $g$ . Now, the range is zero in the extreme cases  $\theta = 0$  and  $\theta = \pi/2$ , but not anywhere in between, so if we remember the range contains a simple trigonometric function, it must be  $\sin(2\theta)$ , so

$$R \propto \frac{v^2}{g} \sin(2\theta).$$

We can also get the prefactor by a simple limiting case, the case  $\theta \ll 1$ . In this case, by the small angle approximation,

$$v_x \approx v, \quad v_y \approx v\theta.$$

The time taken is  $t = 2v_y/g$ , so the range is

$$R \approx v_x t = \frac{2v^2}{g} \theta.$$

Thus there is no proportionality constant; the answer is

$$R = \frac{v^2}{g} \sin(2\theta).$$

In reality, it's probably faster to go through the full derivation than all of this reasoning, but if you're just not sure about whether it's a sine or a cosine, or what the prefactor is, then limiting cases can be quickly used to recover that piece. Also note that the approximations we used above are frequently useful for evaluating limiting cases.

### Example 14

Consider an [Atwood's machine](#) with masses  $m$  and  $M$ , and a massless pulley. Find the tension in the string.

### Solution

Since the equations involved are all linear equations, we expect the answer should also be simple. It can only depend on  $g$ ,  $m$ , and  $M$ , so by dimensional analysis, it must be proportional to  $g$ . By dimensional analysis, this must be multiplied by something with one net power of mass. Since the answer remains the same if we switch the masses, it should be symmetric in  $m$  and  $M$ .

Given all of this, the simplest possible answer would be

$$T \propto g(m + M).$$

To test this, we consider some limiting cases. If  $M \gg m$ , the mass  $M$  is essentially in freefall, so the mass  $m$  accelerates upward with acceleration  $g$ . Then the tension is approximately  $2mg$ . Similarly, in the case  $M \ll m$ , the tension is approximately  $2Mg$ . These can't be satisfied by the form above.

The next simplest option is a quadratic divided by a linear expression. Both of these must

be symmetric, so the most general possibility is

$$T = g \frac{A(m^2 + M^2) + BmM}{m + M}.$$

Then the limiting cases can be satisfied if  $A = 0$  and  $B = 2$ , giving

$$T = \frac{2gmM}{m + M}.$$

- [1] **Problem 23.** Find the perimeter of a regular  $N$ -gon, if  $L$  is the distance from the center to any of the sides. By considering a limiting case, use this to derive the circumference of a circle.
- [1] **Problem 24.** Use similar reasoning to find the acceleration of the Atwood's machine. (We will show an even easier way to do this, using “generalized coordinates”, in **M4**.)
- [2] **Problem 25** (Morin 1.6). A person throws a ball (at an angle of her choosing, to achieve the maximum distance) with speed  $v$  from the edge of a cliff of height  $h$ . Which of the below could be an expression for the maximal range?

$$\frac{gh^2}{v^2}, \quad \frac{v^2}{g}, \quad \sqrt{\frac{v^2 h}{g}}, \quad \frac{v^2}{g} \sqrt{1 + \frac{2gh}{v^2}}, \quad \frac{v^2}{g} \left(1 + \frac{2gh}{v^2}\right), \quad \frac{v^2/g}{1 - 2gh/v^2}.$$

If desired, try Morin problems 1.13, 1.14, and 1.15 for additional practice.

- [2] **Problem 26.** Consider a triangle with side lengths  $a$ ,  $b$ , and  $c$ . It turns out the area of its incircle can be expressed purely by multiplying and dividing combinations of these lengths. Moreover, the answer is the simplest possible one consistent with limiting cases, scaling, and symmetry. Guess it!

While we won't have more questions that are explicitly about dimensional analysis or limiting cases, these are not techniques but ways of life. For all future problems you solve, you should be constantly checking the dimensions and limiting cases to make sure everything makes sense.

## 4 Manipulating Differentials

You might have been taught in math class that manipulating differentials like they're just small, finite quantities, and treating derivatives like fractions is “illegal”. But it's also very useful.

### Idea 7

Derivatives can be treated like fractions, if all functions have a single argument.

The reason is simply the chain rule. The motion of a single particle only depends on a single parameter, so the chain rule is just the same as fraction cancellation. For example,

$$\frac{dv}{dt} = \frac{d}{dt}v(x(t)) = \frac{dv}{dx} \frac{dx}{dt}$$

which show that “canceling a  $dx$ ” is valid. Similarly, you can show that

$$\frac{dy}{dx} \frac{dx}{dy} = 1$$

by considering the derivative with respect to  $x$  of the function  $x(y(x)) = x$ .

### Remark

Math students tend to get very upset about the above idea – you can read thousands of rants online from people who have spent whole days mad at it, and at physics in general. The crux of their objection is that we shouldn’t use convenient notation if it hides what’s “really” going on. But that attitude is unhelpful.

For example, the “rigorous” definition of the number 2 is  $S(1)$ , where  $S$  is the successor function. And 4 is just a shorthand for  $S(S(S(1)))$ . So if you want to do arithmetic the “right” way, you shouldn’t even write  $2 + 2 = 4$ , because that hides the true definition of the natural numbers. You should at minimum expand it out to

$$S(1) + S(1) = S(S(S(1))).$$

Or really, if you want to go all the way to the set-theoretic foundations, you should use the “real” definition of the successor function in terms of sets, where for example,

$$S(S(S(1))) = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}, \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}.$$

People have actually advocated for 1<sup>st</sup> grade math to be taught this way – to a certain type of mathematician, an equation becomes more elegant, and more true, the longer it is. For them, inventing and then following many steps of elaborate logic is inherently enjoyable, even if they know it won’t lead to anything they didn’t already know. But in physics we have no time to luxuriate in such rigor, because we want to figure out how specific things work. The point of notation is to help us do that by suppressing mathematical clutter, not inventing more of it.

However, the mathematicians are right in one sense: this trick won’t work for functions with multiple arguments. For example, for a function  $f(x(t), y(t))$ , the chain rule says

$$\frac{df}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

where there are two terms, representing the change in  $f$  from changes only in  $x$ , and only in  $y$ . Luckily, in mechanics we’ll usually consider functions with one argument. We’ll come back to the multivariable chain rule in **T1**.

### Example 15

Derive the work-kinetic energy theorem,  $dW = F dx$ .

**Solution**

Canceling the mass from both sides, we wish to show

$$\frac{1}{2}d(v^2) = a \, dx.$$

To do this, note that

$$\frac{1}{2}d(v^2) = v \, dv = \frac{dx}{dt} dv = \frac{dv}{dt} dx = a \, dx$$

as desired. If you're not satisfied with this derivation, because of the bare differentials floating around, we can equivalently prove that  $F = dW/dx$ , by noting

$$\frac{dW}{dx} = mv \frac{dv}{dx} = mv \frac{dv}{dt} \frac{dt}{dx} = m \frac{dv}{dt} = F.$$

[2] **Problem 27.** Some more about power.

- (a) Use similar reasoning to derive  $P = Fv$ .
- (b) An electric train has a power line that can deliver power  $P(x)$ , where  $x$  is the distance along the track. If the train starts at rest at  $x = 0$ , find its speed at point  $x_0$  in terms of an integral of  $P(x)$ . (Hint: try to get rid of the  $dt$ 's to avoid having to think about the time-dependence.)

[2] **Problem 28** (Kalda). The deceleration of a boat in water due to drag is given by a function  $a(v)$ . Given an initial velocity  $v_0$ , write the total distance the boat travels as a single integral.

[5] **Problem 29.** A particle in a potential well.

- (a) Consider a particle of mass  $m$  and energy  $E$  with potential energy  $V(x)$ , which performs periodic motion. Write the period of the motion in terms of a single integral over  $x$ .
- (b) Suppose the potential well has the form  $V(x) = V_0(x/a)^n$  for  $n > 0$ . If the period of the motion is  $T_0$  when it has amplitude  $A_0$ , find the period when the amplitude is  $A$ , by considering how the integral you found in part (a) scales with  $A$ .
- (c) Find a special case where you can check your answer to part (b). (In fact, there are two more special cases you can check, one which requires negative  $n$  and negative  $V_0$ , and one which requires  $V(x)$  to be replaced with its absolute value.)
- (d) Using a similar method to part (a), write down an integral over  $\theta$  giving the period of a pendulum with length  $L$  in gravity  $g$ , without the small angle approximation. Using this, compute the period of the pendulum with amplitude  $\theta_0$ , up to and including order  $\theta_0^2$ .

## 5 Multiple Integrals

It's also useful to know how to set up multiple integrals. This is fairly straightforward, though technically an "advanced" topic, so we'll demonstrate it by example. For further examples, see chapter 2 of Wang and Ricardo, volume 1, or [MIT OCW 18.02](#), lectures 16, 17, 25, and 26.

**Idea 8**

In most Olympiad problems, multiple integrals can be reduced to single integrals by symmetry.

**Example 16**

Calculate the area of a circle of radius  $R$ .

**Solution**

The area  $A$  is the integral of  $dA$ , i.e. the sum of the infinitesimal areas of pieces we break the circle into. As a first example, let's consider using Cartesian coordinates. Then the pieces will be the rectangular regions centered at  $(x, y)$  with sides  $(dx, dy)$ , which have area  $dx dy$ . The area is thus

$$A = \int dA = \int dx \int dy.$$

The only tricky thing about setting up the integral is writing down the bounds. The inner integral is done first, so its bounds depend on the value of  $x$ . Since the boundary of the circle is  $x^2 + y^2 = R^2$ , the bounds are  $y = \pm\sqrt{R^2 - x^2}$ . Thus we have

$$A = \int_{-R}^R dx \int_{-\sqrt{R^2-x^2}}^{\sqrt{R^2-x^2}} dy.$$

We then just do the integrals one at a time, from the inside out, like regular integrals,

$$A = \int_{-R}^R 2\sqrt{R^2 - x^2} dx = 2R^2 \int_{-1}^1 \sqrt{1 - u^2} du = 2R^2 \int_{-\pi/2}^{\pi/2} \cos^2 \theta d\theta = \pi R^2$$

where we nondimensionalized the integral by letting  $u = x/R$ , and then did the trigonometric substitution  $u = \sin \theta$ . (To do the final integral trivially, notice that the average value of  $\cos^2 \theta$  along any of its periods is  $1/2$ .)

We can also use polar coordinates. We break the circle into regions bounded by radii  $r$  and  $r + dr$ , and angles  $\theta$  and  $\theta + d\theta$ . These regions are rectangular, with side lengths of  $dr$  and  $r d\theta$ , so the area element is  $dA = r dr d\theta$ . Then we have

$$A = \int_0^R r dr \int_0^{2\pi} d\theta = 2\pi \int_0^R r dr = \pi R^2$$

which is quite a bit easier. In fact, it's so much easier that we didn't even need to use double integrals at all. We could have decomposed the circle into a bunch of thin circular shells, argued that each shell contributed area  $(2\pi r) dr$ , then integrated over them,

$$A = \int_0^R 2\pi r dr = \pi R^2.$$

In Olympiad physics, there's usually a method like this, that allows you to get the answer without explicitly writing down any multiple integrals.

**Example 17**

Calculate the moment of inertia of the circle above, about the  $y$  axis, if it has total mass  $M$  and uniform density.

**Solution**

The moment of inertia of a small piece of the circle is

$$dI = x^2 dm = x^2 \sigma dA = \frac{x^2 M}{\pi R^2} dA$$

where  $x^2$  appears because  $x$  is the distance to the rotation axis, and  $\sigma$  is the mass density per unit area. Using Cartesian coordinates, we have

$$I = \frac{M}{\pi R^2} \int_{-R}^R dx \int_{-\sqrt{R^2-x^2}}^{\sqrt{R^2-x^2}} x^2 dy.$$

The inner integral is still trivial; the  $x^2$  doesn't change anything, because from the perspective of the  $dy$  integral,  $x$  is just some constant. However, the remaining integral becomes a bit nasty. In general, when this happens, we can try flipping the order of integration, giving

$$I = \frac{M}{\pi R^2} \int_{-R}^R dy \int_{-\sqrt{R^2-y^2}}^{\sqrt{R^2-y^2}} x^2 dx.$$

Unfortunately, this is also nasty. Both of these integrals can be done using trigonometric substitutions, but in this case there's a clever symmetry argument.

Notice that  $I$  is also equal to the moment of inertia about the  $x$  axis, by symmetry. So if we add them together, we get

$$2I = \int (x^2 + y^2) dm = \int r^2 dm.$$

The  $r^2$  factor has no dependence on  $\theta$  at all, so the angular integral in polar coordinates is trivial. We end up with

$$2I = \frac{M}{\pi R^2} \int_0^R 2\pi r r^2 dr = \frac{1}{2} MR^2$$

which gives an answer of  $I = MR^2/4$ , as expected.

[2] **Problem 30.** Evaluate  $I$  in the previous example using Cartesian coordinates.

[3] **Problem 31.** In this problem we'll generalize some of the ideas above to three dimensions, where we need triple integrals. Consider a ball of radius  $R$ .

- (a) In Cartesian coordinates, the volume element is  $dV = dx dy dz$ . Set up an appropriate triple integral for the volume.

- (b) The inner two integrals might look a bit nasty, but we already have essentially done them. Using the result we already know, perform the inner two integrals in a single step, and then perform the remaining integral to derive the volume of a sphere.
- (c) In cylindrical coordinates, the volume element is  $dV = r dr d\theta dz$ . Set up a triple integral for the volume, and perform it. (Hint: this can either be hard, or a trivial extension of part (b), depending on what order of integration you choose.)
- (d) In spherical coordinates, the volume element is  $dV = r^2 dr \sin \phi d\phi d\theta$ . Set up a triple integral for the volume, and perform it.
- (e) Let the ball have uniform density and total mass  $M$ . Compute its moment of inertia about the  $z$ -axis. (Hint: this can be reduced to a single integral if you use an appropriate trick.)
- [2] **Problem 32.** Consider a spherical cap that is formed by slicing a sphere of radius  $R$  by a plane, so that the altitude from the vertex to the base is  $h$ . Find the area of its curved surface using an appropriate integral.
- [3] **Problem 33.** Find the average distance (in 3D space) between two random points on a unit sphere.

### Remark

The funny thing about problem 33 is that it's pretty popular, and therefore provides an illustrative comparison of the reliability of different websites.

- On [Math StackExchange](#), there are 4 answers, and all but the last are detailed and correct.
- On [Reddit](#), there are 5 top-level comments, all of which say they can't solve it.
- On [Yahoo Answers](#), there are 9 answers. The top answer is correct, while the rest are confident but wrong.
- On [Quora](#), there are 11 answers, almost all of which are completely wrong, or just irrelevant. The single correct answer is buried down the list.

When I started college, Quora was one of the hottest new tech companies. They were growing fast with an endless pile of cash, and seemingly everybody wanted to intern with them. Quora's press releases declared they were revolutionizing human knowledge. A Stanford particle physicist quit his professorship, declaring that our field was dead and Quora was the future. Brilliant minds devoted all their energy to maximizing the site's growth, using every cheap trick in the book.

Unfortunately, along the way they forgot that engagement numbers and SEO are only worth something if your site actually has content worth reading. Now it's a sad pile of trash – though it's also a really *big* pile of trash, with a \$2 billion valuation, which should allow them to continue playing this game for years. The lesson you can take away is that the sites holding the top Google search results are often there because they hire engineers to optimize their rankings, not because they're any good. Be cautious with search results. Despite all the promise of the internet, it's still not as reliable as the average library.