

# **Adv Abstract Algebra: AAA #HW03**

Due on 2022 at 11:59PM

*Prof. Peter Hermann Spr 2022*

**Xianzhi**

2023

Homework set 3

**Problem 1**

Suppose that a group  $G$  has order 312. Prove that  $G$  has a proper normal subgroup.

**Solution:**

$$312 = 2^3 \times 3 \times 13.$$

The number of Sylow  $p = 2$  subgroup,  $n$ , has several possibilities

$$1, 3, 13, 39 \equiv 1 \pmod{p = 2} \tag{1}$$

and they divide  $m = 3 \times 13$ .

The number of Sylow  $p = 3$  subgroup,  $n$  has several possibilities.

$$1, 4, 13 \equiv 1 \pmod{p = 3} \tag{2}$$

$$\tag{3}$$

and they divide  $m = 8 \cdot 13$ .

However, the number of Sylow  $p = 13$  subgroup is one, since 1 is the only number  $\equiv 1 \pmod{p = 13}$  and divide  $m = 2^3 \cdot 3$  at the same time.

By Sylow's theorem, (and corollary)  $G$  has a proper normal subgroup of order 13.

The unique Sylow 13 subgroup.

## Problem 2

Suppose that a group  $G$  has order 1960. Prove that  $G$  has a proper normal subgroup.

### Solution:

Suppose a group has order  $1960 = 2^3 \cdot 5 \cdot 7^2$ , the number of Sylow  $p = 2$  subgroup has several possibilities

$$1, 5, 7 \equiv 1 \pmod{p = 2} \quad (4)$$

and divide  $m = 5 \cdot 7 \cdot 7$ .

The number of Sylow  $p = 5$  subgroup has several possibilities, for example,  $1, 56, 196 \equiv 1 \pmod{p = 5}$  and divide  $m = 2^3 \cdot 7^2$ .

But the number  $n$  of Sylow  $p = 7$  subgroup has 2 possibilities

$$1, 8 \equiv 1 \pmod{p = 7} \quad (5)$$

$$(6)$$

and divide  $m = 2^3 \cdot 5 = 40$ .

Suppose  $n = 8$ . (If  $n = 1$ , then we are done)

Since  $G$  acts on  $Syl_7(G)$  by conjugation and the action is transitive,  $G$  is essentially permuting the 8  $Syl_7(G)$  subgroups.

Thus, we could define a homomorphism

$$G \xrightarrow{\psi} S_8 \quad (7)$$

( $\psi$  is indeed a homomorphism because of the definition of group action)

$\ker \psi$  cannot be the whole group  $G$ , since  $G$  acts transitively on the  $Syl_7(G)$  groups, and we are assuming there is 8 of them, so  $\psi$  cannot map everything in  $G$  to the identity permutation.

**Problem 3**

For  $A \leq G$ ,  $|G : A|$  finite and  $A$  abelian, let  $\tau_{G \rightarrow A}$  denote the transfer homomorphism from  $G$  to  $A$ . Let  $g \in G$  and  $b \in N_G(A)$ . Show that  $\tau_{G \rightarrow A}(g)$  commutes with  $b$ .

Hint: If  $h_1, \dots, h_n$  is a set of right coset representatives of  $A$  then show that  $bh_1, \dots, bh_n$  is also a set of right coset representatives of  $A$ .