

Lec 5: Matrix Decomposition

Ailin Zhang

2022-09-25

Agenda

- Matrix Decomposition (QR)
- Gram-Schmidt Process
- Householder Reflection

Matrix Decomposition

QR decomposition decomposes a matrix \mathbf{X} into a product $\mathbf{X} = \mathbf{Q}\mathbf{R}$ where \mathbf{Q} is an orthogonal matrix and \mathbf{R} is an upper triangular matrix.

Matrix Decomposition

QR decomposition decomposes a matrix \mathbf{X} into a product $\mathbf{X} = \mathbf{Q}\mathbf{R}$ where \mathbf{Q} is an orthogonal matrix and \mathbf{R} is an upper triangular matrix. It does not require the computation of the cross-product matrix such as $\mathbf{X}^\top \mathbf{X}$.

Matrix Decomposition

QR decomposition decomposes a matrix \mathbf{X} into a product $\mathbf{X} = QR$ where Q is an orthogonal matrix and R is an upper triangular matrix. It does not require the computation of the cross-product matrix such as $\mathbf{X}^\top \mathbf{X}$.

Let $Q = (q_1, q_2, \dots, q_n)$ be an orthogonal matrix, $Q^\top Q = QQ^\top = I$, then Q forms an orthogonal basis:

- (1) For each vector q_i , $\|q_i\| = 1$.
- (2) For any two different vectors q_i and q_j , $\langle q_i, q_j \rangle = 0$, i.e., $q_i \perp q_j$.

Matrix Decomposition

QR decomposition decomposes a matrix \mathbf{X} into a product $\mathbf{X} = \mathbf{Q}\mathbf{R}$ where \mathbf{Q} is an orthogonal matrix and \mathbf{R} is an upper triangular matrix. It does not require the computation of the cross-product matrix such as $\mathbf{X}^\top \mathbf{X}$.

Let $\mathbf{Q} = (q_1, q_2, \dots, q_n)$ be an orthogonal matrix, $\mathbf{Q}^\top \mathbf{Q} = \mathbf{Q}\mathbf{Q}^\top = \mathbf{I}$, then \mathbf{Q} forms an orthogonal basis:

- (1) For each vector q_i , $\|q_i\| = 1$.
- (2) For any two different vectors q_i and q_j , $\langle q_i, q_j \rangle = 0$, i.e., $q_i \perp q_j$.

For any vector v , we have

- (1) Analysis: $u_i = \langle v, q_i \rangle = q_i^\top v$ is the coordinate of v on the axis q_i , for $i = 1, \dots, n$, i.e., $u = \mathbf{Q}^\top v$.
- (2) Synthesis: $v = \sum_{i=1}^n q_i u_i = \mathbf{Q}u$.

From (1) and (2), \mathbf{Q} and \mathbf{Q}^\top are inverse of each other.

Gram-Schmidt Process

Sketch it!

Gram-Schmidt Process

Sketch it!

$$\begin{aligned}
 u_1 &= a_1, & q_1 &= \frac{u_1}{\|u_1\|_2}, & \tilde{r}_{11} &= \frac{1}{\|u_1\|_2} \\
 u_2 &= a_2 - (q_1^T a_2)q_1, & q_2 &= \frac{u_2}{\|u_2\|_2}, & \tilde{r}_{12} &= -\frac{q_1^T a_2}{\|u_2\|_2}, & \tilde{r}_{22} &= \frac{1}{\|u_2\|_2}
 \end{aligned}$$

\vdots

$$u_n = a_n - \sum_{j=1}^{n-1} (q_j^T a_n)q_j, \quad q_n = \frac{u_n}{\|u_n\|_2}, \quad \tilde{r}_{jn} = -\frac{q_j^T a_n}{\|u_n\|_2}, \quad \tilde{r}_{nn} = \frac{1}{\|u_n\|_2}$$

$$A \begin{bmatrix} \tilde{r}_{11} & & & \\ & 1 & & \\ & & \ddots & \\ & & & 1 \end{bmatrix} \begin{bmatrix} 1 & \tilde{r}_{12} & & \\ & \tilde{r}_{22} & & \\ & & \ddots & \\ & & & 1 \end{bmatrix} \cdots \begin{bmatrix} 1 & & & \tilde{r}_{1n} \\ & 1 & & \tilde{r}_{2n} \\ & & \ddots & \vdots \\ & & & \tilde{r}_{n-1,n} \\ & & & & 1 \end{bmatrix} = Q$$

Projection Matrix

In Gram-Schmidt, we can treat each vector q_j as a product of projection of A_j , where the projection matrix is $P_j = I - Q_{j-1}Q_{j-1}^T$ and it satisfies the following:

- ① $P^2 = P$
- ② $I - P$ is also a projection matrix

$$Q_j = \begin{bmatrix} | & | & \cdots & | \\ q_1 & q_2 & \cdots & q_j \\ | & | & \cdots & | \end{bmatrix}$$

$$q_1 = \frac{P_1 a_1}{\|P_1 a_1\|} \qquad P_1 = I$$

$$q_2 = \frac{P_2 a_2}{\|P_2 a_2\|}$$

$$\vdots$$

Householder reflection

To obtain a QR decomposition, we can apply the Householder reflections repeatedly. Given an $n \times p$ matrix \mathbf{X} , as the first step, we want to find an orthogonal transformation H_1 such that only the first element in the first column is non-zero after the transformation:

$$\begin{bmatrix} x_{11} & x_{12} & \dots & y_1 \\ x_{21} & x_{22} & \dots & y_2 \\ \dots & \dots & \dots & \dots \\ x_{n1} & x_{n2} & \dots & y_n \end{bmatrix} \xrightarrow{H_1} \begin{bmatrix} x_{11}^* & x_{12}^* & \dots & y_1^* \\ 0 & x_{22}^* & \dots & y_2^* \\ \dots & \dots & \dots & \dots \\ 0 & x_{n2}^* & \dots & y_n^* \end{bmatrix}$$

Graphical Representation

Since the orthogonal transformation preserves the length of vectors, we know

$$|\mathbf{X}_1^*| = |\mathbf{X}_1| = \sqrt{x_{11}^2 + x_{12}^2 + \dots + x_{1n}^2},$$

which means the value of x_{11}^* is determined by

$$x_{11}^* = \pm |\mathbf{X}_1|.$$

The sign of x_{11}^* is chosen as the opposite of x_{11} for numerical stability.




Figure 1: Householder Reflection

Discussion: Choice of X_{11}^*

The sign of x_{11}^* is chosen as the opposite of x_{11} for numerical stability.

Discussion: Recursion

$$\begin{bmatrix} x_{11}^* & x_{12}^* & \cdots & y_1^* \\ 0 & x_{22}^* & \cdots & y_2^* \\ \cdots & \cdots & \cdots & \cdots \\ 0 & x_{n2}^* & \cdots & y_n^* \end{bmatrix} \xrightarrow{H_2} \begin{bmatrix} x_{11}^* & x_{12}^* & \cdots & y_1^* \\ 0 & x_{22}^* & \cdots & y_2^* \\ \cdots & 0 & \cdots & \cdots \\ 0 & 0 & \cdots & y_n^* \end{bmatrix}$$

Derivation

To find a transformation H which can rotate the vector \mathbf{X}_1 to \mathbf{X}_1^* , one simple way is to construct an isosceles triangle where \mathbf{X}_1^* is a reflection of \mathbf{X}_1 :

$$\mathbf{X}_1^* = \mathbf{X}_1 - 2\langle \mathbf{X}_1, u \rangle u = \mathbf{X}_1 - 2uu^\top \mathbf{X}_1 = H_1 \mathbf{X}_1,$$

where

$$u = \frac{\mathbf{X}_1 - \mathbf{X}_1^*}{\|\mathbf{X}_1 - \mathbf{X}_1^*\|}, \quad H_1 = I - 2uu^\top.$$

Derivation

To find a transformation H which can rotate the vector \mathbf{X}_1 to \mathbf{X}_1^* , one simple way is to construct an isosceles triangle where \mathbf{X}_1^* is a reflection of \mathbf{X}_1 :

$$\mathbf{X}_1^* = \mathbf{X}_1 - 2\langle \mathbf{X}_1, u \rangle u = \mathbf{X}_1 - 2uu^\top \mathbf{X}_1 = H_1 \mathbf{X}_1,$$

where

$$u = \frac{\mathbf{X}_1 - \mathbf{X}_1^*}{\|\mathbf{X}_1 - \mathbf{X}_1^*\|}, \quad H_1 = I - 2uu^\top.$$

Recursion:

Let $\mathbf{X}^{(1)}$ be the sub-matrix of the resulting \mathbf{X} with the first row and the first column removed from \mathbf{X} . Apply the Householder reflection on the sub-matrix $\mathbf{X}^{(1)}$, while maintaining the first row and the first column of \mathbf{X} . This amounts to left multiplying \mathbf{X} by an orthogonal matrix H_2 .

Let $H = H_p \dots H_2 H_1$, we have $H\mathbf{X} = R$. Let $Q = H^\top$, we obtain the QR decomposition $\mathbf{X} = QR$

Python Code

```
import numpy as np
from scipy import linalg

def qr(A):
    n, m = A.shape
    R = A.copy()
    Q = np.eye(n)

    for k in range(m-1):
        x = np.zeros((n, 1))
        x[k:, 0] = R[k:, k]
        v=x
        v[k] = x[k] + np.sign(x[k,0]) * np.linalg.norm(x)
        s = np.linalg.norm(v)
        if s != 0:
            u=v/s
            R -= 2 * np.dot(u, np.dot(u.T, R))
            Q -= 2 * np.dot(u, np.dot(u.T, Q))
    Q = Q.T
    return Q, R

A = rand(100,100)
B = copy(A)
(Q,R) = householder(A)
B = Q*R
```


Gram-Schmidt Vs Householder

- ① Column vs row transformation
- ② Gram-Schmidt suffers more from numerical instability