

Note on Fermionizing Ising CFT

Yu Liu

July 01, 2026

Abstract

Ising and Majorana CFT duality

Contents

1	Ising CFT and Majorana CFT Dual	1
1.1	Background Gauge Field Language	1
1.1.1	$\mathbb{Z}_2$ Gauge Field	1
1.1.2	Spin Structure	2
1.2	Ising/Majorana CFT Duality	4
1.2.1	Partition Function Data	4
1.2.2	Ising = Ising/ $\mathbb{Z}_2$ (Kramers-Wannier Duality)	4
1.2.3	Majorana = (Ising $\otimes$ Arf/ $\mathbb{Z}_2$ (Fermionization)	5
1.2.4	Ising = Majorana/ $\mathbb{Z}_2$ (Bosonization)	6
1.2.5	Majorana' = Arf $\otimes$ Majorana	6
1.2.6	Majorana' and Ising/ $\mathbb{Z}_2$	6
1.2.7	Final Comments	7
	Bibliography	8

1 Ising CFT and Majorana CFT Dual

1.1 Background Gauge Field Language

The bosonization of Ising and Majorana CFT can be understood through gauging discrete symmetry. Here we give a brief, non rigorous introduction.

1.1.1 $\mathbb{Z}_2$ Gauge Field

Boundary conditions and defect insertion can be understood in a unified way using the language of coupling to a $\mathbb{Z}_2$ gauge theory. We introduce a background gauge field

$$A \in H^1(\Sigma_g, \mathbb{Z}_2). \quad (1)$$

This object admits several equivalent interpretations:

- as a cohomology group,
- as a flat $\mathbb{Z}_2$ connection,
- as a homomorphism from the fundamental group, $A \in \text{Hom}(\pi_1(\Sigma_g), \mathbb{Z}_2)$.

Concretely, A assigns a $\mathbb{Z}_2$ value to each non-contractible cycle. Choosing a canonical homology basis $\{a_1, b_1, \dots, a_g, b_g\}$, we write

$$A(a_i) = \alpha_i \in \mathbb{Z}_2, \quad A(b_i) = \beta_i \in \mathbb{Z}_2, \quad (2)$$

so the gauge field is completely specified by

$$A = (\alpha_1, \beta_1, \dots, \alpha_g, \beta_g) \in (\mathbb{Z}_2)^{2g}. \quad (3)$$

Given two $\mathbb{Z}_2$ gauge fields $A, B \in H^1(\Sigma_g, \mathbb{Z}_2)$, their cup product is an element

$$A \cup B \in H^2(\Sigma_g, \mathbb{Z}_2) \simeq \mathbb{Z}_2. \quad (4)$$

On a closed Riemann surface this evaluates to a number:

$$\int_{\Sigma_g} A \cup B \in \mathbb{Z}_2. \quad (5)$$

In terms of the cycle basis, writing $A = (\alpha_i, \beta_i)$ and $B = (\alpha'_i, \beta'_i)$,

$$\int_{\Sigma_g} A \cup B = \sum_{i=1}^g (\alpha_i \beta'_i - \beta_i \alpha'_i) \mod 2. \quad (6)$$

Since $-1 \equiv 1$ in $\mathbb{Z}_2$, this simplifies to

$$\int_{\Sigma_g} A \cup B = \sum_{i=1}^g (\alpha_i \beta'_i + \beta_i \alpha'_i) \mod 2. \quad (7)$$

For the torus T^2 the cup product table is:

$\int_{\mathbb{T}^2} A \cup B$	$A_0 = (0, 0)$	$A_1 = (1, 0)$	$A_2 = (0, 1)$	$A_3 = (1, 1)$
$A_0 = (0, 0)$	0	0	0	0
$A_1 = (1, 0)$	0	0	1	1
$A_2 = (0, 1)$	0	1	0	1
$A_3 = (1, 1)$	0	1	1	0

Table 1: Cup product integral on $\mathbb{T}^2$ for the four $\mathbb{Z}_2$ gauge fields, computed modulo 2.

Partition functions with different boundary conditions of a bosonic theory (or a $\mathbb{Z}_2$ defect-twisted theory) can be understood as coupling to different background $\mathbb{Z}_2$ gauge fields:

$$Z[A], \quad A \in H^1(\Sigma_g, \mathbb{Z}_2), \quad (8)$$

where A specifies the twisting of fields along non-contractible cycles. The conventions are:

- 0 for Periodic Boundary Condition (did not cross a defect),
- 1 for Anti-Periodic Boundary Condition (crossed a defect).

For example, on the torus $\Sigma_1 = T^2$ with cycles (a, b) we write $A = (\alpha, \beta) \in \mathbb{Z}_2 \times \mathbb{Z}_2$, and the four boundary conditions are

$$Z_{P,P} = Z[(0, 0)], \quad Z_{P,AP} = Z[(0, 1)], \quad Z_{AP,P} = Z[(1, 0)], \quad Z_{AP,AP} = Z[(1, 1)]. \quad (9)$$

1.1.2 Spin Structure

Definition: Mathematically, a spin structure can be defined via the quadratic refinement of a $\mathbb{Z}_2$ gauge field,

$$\rho : H^1(\Sigma_g, \mathbb{Z}_2) \rightarrow \mathbb{Z}_2. \quad (10)$$

Physically, we simply understand it as specifying the boundary conditions of fermions along the non-contractible cycles. As with the $\mathbb{Z}_2$ gauge field, we assign a label to each cycle:

- 1 for Periodic Boundary Condition (Ramond, R),
- 0 for Anti-Periodic Boundary Condition (Neveu-Schwarz, NS).

Remark

This convention is **opposite** to the bosonic case: there, 0 denotes periodic.

On the torus T^2 , a spin structure determines whether the fermion is periodic or antiperiodic along each cycle:

$$\rho = (\text{a-cycle}, \text{b-cycle}) \in \{\text{AP}, \text{P}\}^2. \quad (11)$$

Two spin structures are related by a background $\mathbb{Z}_2$ gauge field $A \in H^1(\Sigma_g, \mathbb{Z}_2)$ via

$$\rho + A = (\rho_a + A_a \mod 2, \rho_b + A_b \mod 2), \quad (12)$$

which physically corresponds to flipping the fermion boundary condition along cycles where A is nonzero.

Arf Invariant: The Arf invariant is a $\mathbb{Z}_2$ -valued number depending only on the spin structure; it distinguishes “even” from “odd” spin structures. Mathematically it is defined using quadratic refinements acting on basis elements.

On T^2 with standard cycles (a, b) we label spin structures with 0 (AP) and 1 (P). The four spin structures and their Arf invariants are:

Spin structure (ρ_a, ρ_b)	a -cycle	b -cycle	Arf(ρ)
(0, 0)	0	0	0
(1, 0)	1	0	0
(0, 1)	0	1	0
(1, 1)	1	1	1

Table 2: Arf invariants of the four spin structures on T^2 .

Useful Computational Result: Now list out some useful calculational result of Background $\mathbb{Z}_2$ gauge field and spin structure:

Cup product calculation, For all $\mathbb{Z}_2$ gauge fields $s \in H^1(\Sigma_g, \mathbb{Z}_2)$:

$$\frac{1}{2^g} \sum_s (-1)^{\int s \cup t} = \begin{cases} 2^g & \text{if } t = 0 \\ 0 & \text{otherwise.} \end{cases} \quad (13)$$

Arf Invariant of spin structure and background gauge field interacting:

$$\text{Arf}[a + b + \rho] = \text{Arf}[a + \rho] + \text{Arf}[b + \rho] + \text{Arf}[\rho] + \int a \cup b. \quad (14)$$

Arf invariant and spin structure sum:

$$\frac{1}{2^g} \sum_s (-1)^{\text{Arf}(s+\rho) + \text{Arf}[\rho] + \int s \cup t} = (-1)^{\text{Arf}[t+\rho]}. \quad (15)$$

1.2 Ising/Majorana CFT Duality

The Ising CFT and Majorana CFT are related by a web of dualities. We use the torus partition function as an illustrative example; the dualities are claimed to hold on general closed Riemann surfaces. Here is a diagram illustrating the dualities:




Figure 1: Duality web relating Ising CFT and Majorana CFT.

1.2.1 Partition Function Data

We collect the partition functions with explicit spin structure (fermionic theory) or background gauge field (bosonic theory) together with their character expansions.

Partition Function	Spin Structure	Characters
$Z_{\text{AP},\text{AP}}$	$Z[(0,0)]$	$ \chi_0 ^2 + \chi_{1/2} ^2 + \bar{\chi}_0\chi_{1/2} + \bar{\chi}_{1/2}\chi_0$
$Z_{\text{AP},P}$	$Z[(0,1)]$	$ \chi_0 ^2 + \chi_{1/2} ^2 - \bar{\chi}_0\chi_{1/2} - \bar{\chi}_{1/2}\chi_0$
$Z_{P,\text{AP}}$	$Z[(1,0)]$	$2 \chi_{1/16} ^2$
$Z_{P,P}$	$Z[(1,1)]$	0

Table 3: Partition functions and characters for Majorana CFT.

Partition Function	BG Gauge Field	Characters
$Z_{P,P}$	$Z[(0,0)]$	$ \chi_0 ^2 + \chi_{1/2} ^2 + \chi_{1/16} ^2$
$Z_{P,\text{AP}}$	$Z[(0,1)]$	$ \chi_0 ^2 + \chi_{1/2} ^2 - \chi_{1/16} ^2$
$Z_{\text{AP},P}$	$Z[(1,0)]$	$\bar{\chi}_0\chi_{1/2} + \bar{\chi}_{1/2}\chi_0 + \chi_{1/16} ^2$
$Z_{\text{AP},\text{AP}}$	$Z[(1,1)]$	$-\bar{\chi}_0\chi_{1/2} - \bar{\chi}_{1/2}\chi_0 + \chi_{1/16} ^2$

Table 4: Partition functions and characters for Ising CFT.

This calculation data is given by many old school CFT textbooks. For example, the yellow book and Ginsparg's lecture notes. Now let's use the above data to verify the dualities in the web.

1.2.2 Ising = Ising/ $\mathbb{Z}_2$ (Kramers-Wannier Duality)

This duality is a $\mathbb{Z}_2$ orbifold exchanging the A-series and D-series in the minimal model classification. For the Ising CFT it corresponds to the Kramers-Wannier duality, which relates the

partition function of the low-temperature phase to that of the high-temperature phase. Since the A-series and D-series are isomorphic for the Ising model, the orbifold gives a transformation between two isomorphic theories.

The traditional perspective proceeds in two steps:

1. **Project** the theory onto the $\mathbb{Z}_2$ -even Hilbert space:

$$Z \rightarrow \frac{1}{2}(Z_{P,P} + Z_{P,AP}). \quad (16)$$

2. **Restore** modular invariance by adding the $\mathbb{Z}_2$ -twisted sector Hilbert space:

$$+ \frac{1}{2}(Z_{AP,P} + Z_{AP,AP}). \quad (17)$$

This gives the same partition function as the Ising CFT with PP boundary conditions:

$$Z_{\text{orb}} = \frac{1}{2}(Z_{P,P} + Z_{P,AP} + Z_{AP,P} + Z_{AP,AP}) = Z_{P,P}. \quad (18)$$

Now we use the background gauge field language to give a more general discussion of this gauging process.

Theorem 1.1

$$\text{Ising} = \text{Ising}/\mathbb{Z}_2$$

Coupling to a background gauge field A , the partition functions of the two theories are related by

$$Z_{\text{Ising}/\mathbb{Z}_2}[A] = \sum_{a \in H^1(\Sigma_g, \mathbb{Z}_2)} Z_{\text{Ising}}[a] (-1)^{\int a \cup A}. \quad (19)$$

Example. For $A = (0, 0)$:

$$Z_{\text{Ising}/\mathbb{Z}_2}[(0, 0)] = \frac{1}{2} \sum_{a \in H^1(\Sigma_g, \mathbb{Z}_2)} Z_{\text{Ising}}[a] = Z_{\text{Ising}}[(0, 0)]. \quad (20)$$

For $A = (0, 1)$:

$$\begin{aligned} Z_{\text{Ising}/\mathbb{Z}_2}[(0, 1)] &= \frac{1}{2}(Z_{\text{Ising}}[(0, 0)] + Z_{\text{Ising}}[(0, 1)] - Z_{\text{Ising}}[(1, 0)] - Z_{\text{Ising}}[(1, 1)]) \\ &= Z_{\text{Ising}}[(0, 1)]. \end{aligned} \quad (21)$$

One can verify from these calculations that the two theories are isomorphic.

◇ YL: In fact, this story have more to tell, the dual model have a SM realization and it can be seen in SM theory, moreover, we can also see this and inverse process in CFTs, by gauging a dual symmetry. But I'm not familiar with these thus not talking about them here.

1.2.3 Majorana = (Ising $\otimes$ Arf/ $\mathbb{Z}_2$ (Fermionization)

Theorem 1.2

$$\text{Majorana} = \text{Ising} \otimes \text{Arf}/\mathbb{Z}_2$$

$$Z_{\text{Majorana}}[A + \rho] = \frac{1}{2^g} (-1)^{\text{Arf}[\rho]} \sum_{a \in H^1(\Sigma_g, \mathbb{Z}_2)} Z_{\text{Ising}}[a] (-1)^{\text{Arf}[a + \rho] + \int a \cup A}. \quad (22)$$

Example. Take $A = 0$ and fix the spin structure $\rho = (0, 0)$ (two NS boundary conditions). Since $\text{Arf}[(0, 0)] = 0$, the sign $(-1)^{\text{Arf}[a + \rho]}$ equals -1 only for $a = (1, 1)$. Therefore:

$$\begin{aligned}
Z_M[(0,0)] &= \frac{1}{2}(Z[(0,0)] + Z[(1,0)] + Z[(0,1)] - Z[(1,1)]) \\
&= |\chi_0|^2 + |\chi_{1/2}|^2 + \bar{\chi}_0 \chi_{1/2} + \bar{\chi}_{1/2} \chi_0.
\end{aligned} \tag{23}$$

This matches the expected result from the Majorana partition function table exactly.

1.2.4 Ising = Majorana/ $\mathbb{Z}_2$ (Bosonization)

Theorem 1.3

Ising = Majorana/ $\mathbb{Z}_2$

$$Z_{\text{Ising}}[A] = \frac{1}{2^g} (-1)^{\text{Arf}[A+\rho] + \text{Arf}[\rho]} \sum_{a \in H^1(\Sigma_g, \mathbb{Z}_2)} Z_{\text{Majorana}}[a + \rho] (-1)^{\int a \cup A}. \tag{24}$$

Example. Take $A = (0,0)$:

$$\begin{aligned}
Z_I[(0,0)] &= \frac{1}{2}(Z_M[(0,0)] + Z_M[(1,0)] + Z_M[(0,1)] + Z_M[(1,1)]) \\
&= |\chi_0|^2 + |\chi_{1/2}|^2 + |\chi_{1/16}|^2.
\end{aligned} \tag{25}$$

1.2.5 Majorana' = Arf $\otimes$ Majorana

One can define a dual theory when considering the Ramond sector of Majorana CFT. When computing the partition function in the Ramond sector, there is a freedom in choosing the fermion parity assignment for the two degenerate ground states $|\sigma_{\perp}\rangle$ and $|\mu_{\perp}\rangle$:

- $(-1)^F |\sigma_{\perp}\rangle = +|\sigma_{\perp}\rangle$ and $(-1)^F |\mu_{\perp}\rangle = -|\mu_{\perp}\rangle$,
- $(-1)^F |\sigma_{\perp}\rangle = -|\sigma_{\perp}\rangle$ and $(-1)^F |\mu_{\perp}\rangle = +|\mu_{\perp}\rangle$.

In principle these two choices define two different theories, yet they share the same torus partition function since it does not matter which state is labeled even or odd.

The two theories are related by the Arf invariant partition function (also called the Kitaev chain [1]).

Theorem 1.4

Majorana' = Arf $\otimes$ Majorana

$$Z_{\text{Majorana}'}[A + \rho] = (-1)^{\text{Arf}[A+\rho]} Z_{\text{Majorana}}[A + \rho]. \tag{26}$$

This relation is trivially consistent: changing the fermion parity does not change the partition function of Majorana CFT. Indeed, the only nonzero Arf invariant on T^2 occurs for spin structure $(1,1)$, where the partition function is already zero.

One can show that the two theories are related by a chiral transformation. Although chiral symmetry is a classical symmetry, it carries an anomaly that produces exactly:

$$Z[\rho] \rightarrow Z[\rho] \times (-1)^{\text{Arf}[\rho]}. \tag{27}$$

See [2] for a detailed explanation.

1.2.6 Majorana' and Ising/ $\mathbb{Z}_2$

From the relations above together with mathematical properties of the Arf invariant and the cup product integral, one can prove that Majorana' and Ising/ $\mathbb{Z}_2$ satisfy the same duality of partition functions as Ising CFT and Majorana CFT. One can see this using the above dualities and the properties of the Arf invariant and cup product integral.

There is an important result from this duality, is that the Majorana' theory in fact is related to the original Majorana CFT by a chiral transformation. The partition function is different due to the chiral anomaly.

And this means that after gauging the $\mathbb{Z}_2$ symmetry in the Majorana CFT and the Majorana' theory, we get Ising CFT and the orbifolded theory, this means that the Kramers-Wannier duality is in fact a consequence of the chiral anomaly of Majorana CFT.

This is further developed in the language of non-invertible symmetries. The Kramers-Wannier duality is a non-invertible symmetry of the Ising CFT, and it can be seen as a consequence of the chiral anomaly of the Majorana CFT [3].

1.2.7 Final Comments

All the results above are established in [2]. They follow from a “seed duality” which is not proved in that paper but appears to be a long-known result; the proof seems to appear in a mathematical literature.

◇ YL: I have no idea why the Arf invariant is identical to the Kitaev chain. Moreover, I don't quite understand the dual symmetry generated after gauging a symmetry, nor how gauging the dual symmetry returns one to the original theory. These are all very interesting topics but I'm not talking about them here.

Bibliography

- [1] A. Y. Kitaev, “Unpaired Majorana Fermions in Quantum Wires,” *Physics-Uspekhi*, vol. 44, no. 10S, pp. 131–136, 2001, doi: [10.1070/1063-7869/44/10S/S29](https://doi.org/10.1070/1063-7869/44/10S/S29).
- [2] A. Karch, D. Tong, and C. Turner, “A Web of 2d Dualities: Z2 Gauge Fields and Arf Invariants,” *SciPost Physics*, vol. 7, no. 1, p. 7, 2019, doi: [10.21468/SciPostPhys.7.1.007](https://doi.org/10.21468/SciPostPhys.7.1.007).
- [3] S.-H. Shao, “What's Done Cannot Be Undone: TASI Lectures on Non-Invertible Symmetries.” 2024.