

A Minimal Introduction to CFT

Yu Liu

March 8, 2026

Abstract

I recently came to review some basic concepts in Conformal Field Theory (CFT) for some semester projects. When going through the renowned textbook "Conformal Field Theory" by Di Francesco, Mathieu, and Sénéchal, I found it structures poorly in chapter 4-6, without a clear logical flow and indication of the core concepts. Thus, I write this note for myself to extract the essential ideas from the book and emplasize the core concepts in CFT. This note is by no means complete or rigorous, but I hope it can serve as a concise reference for myself and others who want to get a quick overview of CFT.

Contents

1	Fundamental QFT	3
1.1	Symmetry and Conserved Current	3
1.1.1	A more standard definition of Symmetry	3
1.1.2	Noether's Theorem and Conserved Current	4
1.1.3	Series of Conserved Currents	5
1.2	Energy-Momentum Tensor	5
1.2.1	Canonical Energy-Momentum Tensor	5
1.2.2	Symmetric Traceless Energy-Momentum Tensor	5
1.3	Euclidean Formalism of Path Integral	6
1.3.1	Correlation Functions in QFT	6
1.3.2	Euclidean Path Integral as vaccum Expectation Value	7
1.3.3	Wick Theorem	7
1.4	Symmetry Transformation of Correlation Functions	7
1.4.1	Transformation of Correlation Functions under Symmetry	7
1.4.2	Ward Identity	8
1.4.3	Conserved Charge as Generator of Symmetry	9
2	CFT Path-Integral Formalism	10
2.1	2D Conformal Transformation	10
2.1.1	Holomorphic function as conformal transformation	10
2.1.2	Holomorphic Coordinate	10
2.2	Primary Fields	10
2.2.1	Definition of Primary Fields	10
2.2.2	Correlation Function of Primary Fields	11
2.3	Ward Identity of Primary Fields	11
2.3.1	Ward Identity in Normal Coordinate	11
2.3.2	Ward Identity in Holomorphic Coordinate	12
2.3.3	Remarks on 2 Conventions of E-M Tensor	15
2.4	Conformal Ward Identity	15
2.4.1	Conformal Ward Identity for General Fields	15
2.4.2	Remark on the convention of infinistesimal variation	16

2.5	OPE Structure	16
2.5.1	Definition of OPE	16
2.5.2	OPE in CFT	16
2.6	Complementary Remarks	18
2.6.1	Transformation Rules of E-M Tensor	18
3	CFT Operator Formalism	19
3.1	Radial Quantization	19
3.1.1	Radial Quantization Setup	19
3.1.2	Hermite Conjugation in Radial Quantization	20
3.1.3	Mode Expansion in Radial Quantization	20
3.1.4	OPE in Radial Quantization	20
3.1.5	Commutator from OPE	21
3.2	Symmetry Operator of CFT	22
3.2.1	Symmetry Charges and Algebra	22
3.2.2	Symmetry Generators in Radial Quantization	23
3.3	Hilbert Space of CFT	24
3.3.1	Hilbert Space Structure	24
3.3.2	Hilbert Space Realization	25
3.3.3	Inner Product	27
3.4	Primary and Descendent Fields	28
3.4.1	State-Operator Correspondence	28
3.4.2	Descendent Operators	28
3.4.3	Operator Data of CFT	29
3.4.4	Correlation Functions with Descendant Operators	29
3.4.5	OPE with Descendant Operators	30
3.5	Complementary Remarks	30
3.5.1	Radial Quantization on Cylinder	30
4	Summary: General Structure of CFT	31
4.1	Hilbert Space and Operator Contents	31
4.2	Correlation Functions	31
5	Axiomatic Formalism	32
5.1	BPZ Axioms of CFT	32
5.2		32
6	Introduction to Conformal Bootstrap	33
6.1	Introduction to Conformal Bootstrap	33
6.2	Unitary Theory	33
6.3	Crossing Symmetry	33

1 Fundamental QFT

{sec:Fundam

1.1 Symmetry and Conserved Current

1.1.1 A more standard definition of Symmetry

In Prof. Rattazzi's lecture the symmetry of a field theory is defined as:

Definition 1. *Symmetry(QFT EPFL)*

Consider a field theory with a series of dynamical fields $\phi_a(x)$ and a dynamical Lagrangian $\mathcal{L}[\phi]$. A symmetry of the action is a transformation with parameters α :

$$x'^\mu = f^\mu(x, \alpha) \quad (1)$$

$$\phi'_a(x') = F_a(\phi(x), \alpha) \quad (2)$$

so that satisfy that:

$$d^4x [\mathcal{L}(\phi_a(x), \partial\phi_a(x))] = d^4x' [\mathcal{L}(\phi'_a(x'), \partial'\phi'_a(x')) + \partial'_\mu K^\mu(\phi')] \quad (3)$$

Remark:

We haven't say anything about what **transformation** physically means. Because it has 2 understandings with exactly the same mathematical expression:

- **Active Transformation:** The coordinates are fixed, but the fields are transformed.
- **Passive Transformation:** The coordinates are transformed, and the field transform covariantly.

This definition can be rewrite a bit more elegantly when we define the change of Action under this coordinate transformtaion:

Definition 2. *Induced Change of Field and Acton under Transformation*

Consider a field theory with dynamical fields ϕ_a and a action S that takes the form:

$$S = \int d^d x \mathcal{L}(\phi(x), \partial_\mu \phi(x)) \quad (4)$$

Form a transformation:

$$x'^\mu = f^\mu(x, \omega) \quad (5)$$

$$\phi'(x') = F(\phi(x), \omega) \quad (6)$$

We can induce a **change of the field configuration**:

$$\phi(x) \rightarrow \phi'(x) = F(\phi(f^{-1}(x)), \omega) \quad (7)$$

We define the action after the as:

$$S' = \int d^d x \mathcal{L}(\phi'(x), \partial_\mu \phi'(x)) \quad (8)$$

We define Symmetry as:

Definition 3. Symmetry(QFT standard)

A transformation is a symmetry of the action if the action is invariant up to a total derivative:

$$S' = S + \int d^d x \partial_\mu K^\mu \quad (9) \quad \{\text{eq:Symmetry}\}$$

Plugging in the definition of action we have:

$$\delta S[\phi] = \text{Boundary Term} \quad (10)$$

For we can see that $S' - S$ is the standard definition of variation of action functional induced by the transformation. Thus we can rephrase as: **A transformation is a symmetry if the induced variation of the action functional is a boundary term.**

We can Prove that these two definitions are equivalent. We consider eq. (9) and do a variable substitution $x \rightarrow x'$, we have:

$$S' = \int d^d x' \mathcal{L}(\phi'(x'), \partial'_\mu \phi'(x')) + \int d^d x' \partial'_\mu K^\mu = S = \int d^d x \mathcal{L}(\phi(x), \partial_\mu \phi(x)) \quad (11)$$

Thus we compare the integrands and get the first definition.

Remark:

Why do we use the Second definition? Because in the path integral formalism this is more natural and make the derivation more clean. However, the first definition is more intuitive easier to understand.

1.1.2 Noether's Theorem and Conserved Current

Consider a transformation that is a **Symmetry of Rigid Parameters**, we can have a Coserved Current and the form of the current is related to the variation of action functional.

Theorem 1. Noether's Theorem(Field Theory Standard)

Consider a field theory with dynamical fields ϕ_a and a action S that takes the standard form. Assume that we have a transformation that is a symmetry:

$$x'^\mu = x^\mu + \omega_a \frac{\delta x^\mu}{\delta \omega_a} \quad (12)$$

$$\phi'(x') = \phi(x) + \omega_a \frac{\delta \mathcal{F}}{\delta \omega_a}(x) \quad (13)$$

where ω_a are **rigid parameters**. Then there exists a conserved current J^μ_a that satisfies, if we **lift the rigid parameters to local parameters** $\omega_a(x)$:

$$\delta S = - \int dx J^\mu_a \partial_\mu \omega_a + \text{Boundary Terms} \quad (14)$$

We only have the derivative term $\partial_\mu \omega_a$ this is because the transformation is a symmetry for rigid parameters. One explicit form of J^μ_a is (**Canonical Form**):

$$J^\mu_a = \left\{ \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial_\nu \phi - \delta^\mu_\nu \mathcal{L} \right\} \frac{\delta x^\nu}{\delta \omega_a} - \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \frac{\delta \mathcal{F}}{\delta \omega_a} \quad (15)$$

If the EoM is satisfied, we have the conservation equation:

$$\partial_\mu J^\mu_a = 0 \quad (16)$$

Proof:

[leave it for later. Its real standard]

1.1.3 Series of Conserved Currents

The Conserved Current is not Unique. We can always add a term of the form:

$$J^\mu_a \rightarrow J^\mu_a + \partial_\nu B^{\mu\nu}_a \quad \text{where} \quad B^{\mu\nu}_a = -B^{\nu\mu}_a \quad (17)$$

Then it is also a conserved current and Satisfy the relation with the variation of action functional. This is because:

$$\int d^d x \partial_\nu B^{\mu\nu}_a \partial_\mu \omega_a = \int d^d x \partial_\mu (B^{\mu\nu}_a \partial_\nu \omega_a) - \int d^d x B^{\mu\nu}_a \partial_\mu \partial_\nu \omega_a \quad (18)$$

The first term is a boundary term and the second term is symmetric in $\mu\nu$ while $B^{\mu\nu}_a$ is antisymmetric in $\mu\nu$ thus it vanishes.

1.2 Energy-Momentum Tensor

1.2.1 Canonical Energy-Momentum Tensor

Consider the spacetime translation symmetry:

$$x'^\mu = x^\mu + \epsilon^\mu \quad (19)$$

$$\phi'(x') = \phi(x) \quad (20)$$

We can construct its Canonical Conserved Current which is defined as the canonical

Definition 4. Canonical Energy-Momentum Tensor

The Canonical Energy-Momentum Tensor is defined as the conserved current of spacetime translation symmetry:

$$T^\mu_c{}_\nu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial_\nu \phi - \delta^\mu_\nu \mathcal{L} \quad (21)$$

According to the Noether's Theorem, we can see that if a theory has translation symmetry, we lift the rigid translation parameters ϵ^μ to local parameters $\epsilon^\mu(x)$, we have:

$$\delta S = - \int d^d x T^\mu_c{}_\nu \partial_\mu \epsilon^\nu + \text{Boundary Terms} \quad (22)$$

and if the EoM is satisfied, we have the conservation equation:

$$\partial_\mu T^\mu_c{}_\nu = 0 \quad (23)$$

This is generally, the canonical E-M tensor is **not symmetric** in its indices and has a **non-vanishing trace**.

1.2.2 Symmetric Traceless Energy-Momentum Tensor

If the theory has Lorentz Symmetry and Scale Symmetry, we can always construct a symmetric traceless energy-momentum tensor from the canonical one by adding an improvement term.

Theorem 2. Symmetric Traceless Energy-Momentum Tensor

Consider a field theory with Lorentz and Scale Symmetry. We can construct a **Symmetric and Traceless Energy-Momentum Tensor**:

$$T^{\mu\nu} = T_c^{\mu\nu} + \partial_\rho B^{\rho\mu\nu} + \frac{1}{2} \partial_\lambda \partial_\rho X^{\lambda\rho\mu\nu} \quad (24)$$

where B and X are constructed tensors. Moreover, the Lorentz and Scale Conserved Currents can be constructed from this symmetric traceless energy-momentum tensor as:

$$J_{\text{Lorentz}}^{\mu\rho\sigma} = T^{\mu\rho} x^\sigma - T^{\mu\sigma} x^\rho \quad (25)$$

$$J_{\text{Scale}}^\mu = T^{\mu\nu} x_\nu \quad (26)$$

Remark:

The above statement is a bit rough.

- If the system has **Lorentz Symmetry**, we can always construct a symmetric energy-momentum tensor by adding an improvement term $\partial_\rho B^{\rho\mu\nu}$ where $B^{\rho\mu\nu} = -B^{\mu\rho\nu}$. This is called **Belinfante-Rosenfeld procedure**.
- If the system has **Scale Symmetry**, we can only construct a traceless energy-momentum tensor when the **Virial Current** is a total derivative.

$$V^\mu = \partial_\alpha \sigma^{\alpha\mu} \quad (27)$$

In this case we can add the second improvement term $\frac{1}{2} \partial_\lambda \partial_\rho X^{\lambda\rho\mu\nu}$ to make the energy-momentum tensor traceless.

Proof: See yellow book chap 4.2.2 and chap 2.5.1.

- From now on we will always use the symmetric traceless energy-momentum tensor when we talk about energy-momentum tensor.

1.3 Euclidean Formalism of Path Integral**1.3.1 Correlation Functions in QFT**

In QFT, we focus on the vacuum expectation value of operators as observables. Instead of working canonically we can use Euclidean path integral formalism to compute these vacuum expectation values.

Definition 5. Correlation Function in Canonical Formalism

In canonical formalism, we define the correlation function of operators $\{\text{quantized classical observables}\}$ $\mathcal{O}_1(x_1), \mathcal{O}_2(x_2), \dots, \mathcal{O}_n(x_n)$ as the vacuum expectation value:

$$\langle \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \cdots \mathcal{O}_n(x_n) \rangle = \langle 0 | T \{ \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \cdots \mathcal{O}_n(x_n) \} | 0 \rangle \quad (28)$$

Note it is important to have the time-ordering operator $T\{\dots\}$ here, without it we cannot have a well-defined interpretation of correlation function.

1.3.2 Euclidean Path Integral as vacuum Expectation Value

We can obtain the correlation function in Euclidean path integral formalism as:

Theorem 3. Correlation Function in Euclidean Path Integral Formalism

In Euclidean path integral formalism, we can compute the correlation function from classical observables $\mathcal{O}_1(x_1), \mathcal{O}_2(x_2), \dots, \mathcal{O}_n(x_n)$ as:

$$\langle \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \dots \mathcal{O}_n(x_n) \rangle = \frac{\int D\phi \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \dots \mathcal{O}_n(x_n) e^{-S_E[\phi]}}{\int D\phi e^{-S_E[\phi]}} \quad (29)$$

where $S_E[\phi]$ is the Euclidean action functional and $\mathcal{O}(x)$ on the right hand side are classical functionals of fields ϕ instead of operators.

From now on we will always use the Euclidean path integral formalism to compute correlation functions and we will use S which should denote the Euclidean action functional.

1.3.3 Wick Theorem

In canonical formalism, we define the correlation function with a time ordering of operators. However, it is difficult to calculate the time ordering operators vacuum expectation values. However, we can define another ordering of operators called **Normal Ordering**

Definition 6. Normal Ordering

Consider a set of creation and annihilation operators $a_i^\dagger, a_i$ as the canonical value of the theory.

The normal ordering of a operator product $\mathcal{O}$ denoted as $:\mathcal{O}:$ is defined as the operator product with all the creation operators moved to the left of all the annihilation operators.

- Vacuum expectation value of a normal ordered operator product is zero.

The Wick Theorem states that :

Theorem 4. Wick Theorem

We first define the notation of contraction:

$$:\overline{\phi_1 \phi_2 \phi_3 \phi_4} := \phi_1 \phi_3 : \langle \phi_2 \phi_4 \rangle \quad (30)$$

The relation between time ordering and normal ordering of operator products is given by:

$$T\{\phi_1 \phi_2 \dots \phi_n\} = :\phi_1 \phi_2 \dots \phi_n : + \text{all possible contractions} \quad (31)$$

1.4 Symmetry Transformation of Correlation Functions

1.4.1 Transformation of Correlation Functions under Symmetry

Consider a correlation function of fields of the following form:

$$\langle \Phi(x_1) \dots \Phi(x_n) \rangle = \frac{1}{Z} \int [d\Phi] \Phi(x_1) \dots \Phi(x_n) \exp -S[\Phi] \quad (32)$$

We have a theorem of its transformation under a symmetry transformation:

Theorem 5. Correlation Functions under Symmetry

{thm:correla

If we have a symmetry transformation of classical fields $\Phi(x) \rightarrow \Phi'(x') = F(\Phi(x), \alpha)$ with $x'^\mu = f^\mu(x, \alpha)$, then the correlation function have the following identity:

$$\langle \Phi(x'_1) \cdots \Phi(x'_n) \rangle = \langle F(\Phi(x_1)) \cdots F(\Phi(x_n)) \rangle \quad (33)$$

If we rewrite $x_i = f^{-1}x'_i$, then it means that:

$$\langle \Phi(x_1) \cdots \Phi(x_n) \rangle = \langle F(\Phi(f^{-1}(x_1))) \cdots F(\Phi(f^{-1}(x_n))) \rangle \quad (34)$$

In the infinitesimal form, we can define the quantum variation operator as:

Definition 7. Quantum Variation Operator

Consider a symmetry transformation of classical fields $\Phi(x) \rightarrow \Phi'(x') = F(\Phi(x), \omega)$ with $x'^\mu = f^\mu(x, \omega)$. We define the generator of the transformation as G_a where:

$$\Phi'(x) - \Phi(x) = -i\omega_a G_a \Phi(x) \quad (35)$$

Then we define the quantum variation of the correlation function as:

$$\delta_\omega \langle \Phi(x_1) \cdots \Phi(x_n) \rangle \equiv -i\omega_a \sum_{i=1}^n \langle \Phi(x_1) G_a \Phi(x_i) \cdots \Phi(x_n) \rangle \quad (36)$$

Then we can rewrite theorem 8 in the infinitesimal form as:

$$\delta_\omega \langle \Phi(x_1) \cdots \Phi(x_n) \rangle = 0 \quad (37) \quad \{\text{eq:preward}$$

Remark:

Note that here we **only vary the field configuration like the classical case. We do not vary the action functional** because the transformation is a symmetry of the action, the variation of action functional is a boundary term which does not contribute to the path integral.

1.4.2 Ward Identity

The above identity eq. (37) can be rewrite in another form when we use the trick of lifting the rigid parameters to local parameters. We have:

Theorem 6. Ward Identity

{eq:Ward Id

Consider a symmetry transformation of classical fields $\Phi(x) \rightarrow \Phi'(x') = F(\Phi(x), \omega)$ with $x'^\mu = f^\mu(x, \omega)$. If we lift the rigid parameters ω_a to local parameters $\omega_a(x)$, then the variation of action functional is given by:

$$\delta_\omega S[\Phi] = - \int d^d x J^\mu_a \partial_\mu \omega_a(x) \quad (38)$$

where J^μ_a is the conserved current associated to this symmetry. Then the Ward Identity states that:

$$\partial_\mu \langle J^\mu_a(x) \Phi(x_1) \cdots \Phi(x_n) \rangle = i \sum_{i=1}^n \delta(x - x_i) \langle \Phi(x_1) \cdots G_a \Phi(x_i) \cdots \Phi(x_n) \rangle \quad (39)$$

Proof: See yellow book chap 2.4.4. we use the fact that:

$$\langle X \rangle = \frac{1}{Z} \int [d\Phi'] (X + \delta X) \exp - \left\{ S[\Phi] + \int dx \partial_\mu j_a^\mu \omega_a(x) \right\} \quad (40)$$

1.4.3 Conserved Charge as Generator of Symmetry

From the Canonical Formalism of QFT, we know that the conserved charge is the generator of symmetry transformation. We can also see this from the Ward Identity.

To match the canonical formalism, we first have to define "time", and then we define the conserved charge as:

$$Q_a = \int d^{d-1}x J_a^0(t, \vec{x}) \quad (41)$$

Then we integrate the Ward Identity over the box between $t - \epsilon$ and $t + \epsilon$ and take the limit $\epsilon \rightarrow 0^+$, where $t = x_1^0$ we have:

$$\langle Q_a(t_+) \Phi(x_1) Y \rangle - \langle Q_a(t_-) \Phi(x_1) Y \rangle = -i \langle G_a \Phi(x_1) Y \rangle \quad (42)$$

This is exactly the canonical formalism statement if we use theorem 3:

$$[Q_a, \Phi] = -i G_a \Phi \quad (43)$$

Where we come back to the standard result of canonical formalism.

2 CFT Path-Integral Formalism

{sec:Fundam

Follow chapter 5 of the yellow book.

2.1 2D Conformal Transformation

2.1.1 Holomorphic function as conformal transformation

We consider a general transformation on 2D metric:

$$g^{\mu\nu} \rightarrow \left(\frac{\partial w^\mu}{\partial z^\alpha} \right) \left(\frac{\partial w^\nu}{\partial z^\beta} \right) g^{\alpha\beta} \quad (44)$$

Then we say this transformation is conformal if:

$$g_{\mu\nu} \rightarrow \Lambda(z) g_{\mu\nu} \quad (45)$$

Theorem 7. *In 2D, any transformation that is holomorphic (antiholomorphic) is conformal.*

2.1.2 Holomorphic Coordinate

We use the holomorphic coordinate to express everything in 2D:

$$z = z^0 + iz^1, \quad \bar{z} = z^0 - iz^1, \quad z^0 = \frac{z + \bar{z}}{2}, \quad z^1 = \frac{z - \bar{z}}{2i}; \quad (46)$$

$$\partial = \frac{1}{2}(\partial_0 - i\partial_1), \quad \bar{\partial} = \frac{1}{2}(\partial_0 + i\partial_1), \quad \partial_0 = \partial + \bar{\partial}, \quad \partial_1 = i(\partial - \bar{\partial}). \quad (47)$$

2.2 Primary Fields

In the context of conformal field theory, we call every observable a field, not just the dynamical field in normal QFT.

2.2.1 Definition of Primary Fields

There are fields transform covariantly under all conformal transformation, which are called **Primary Fields**. They should be a generalization of fields within lorentz representation with a dimension.

Definition 8. *Primary Field (normal coordinate)*

A field $\phi(x)$ is called Primary field if under conformal transformation $x \rightarrow x'$, it transforms as:

$$\phi'(x') = \Omega(x)^{-\Delta} e^{-is\Theta(x)} \phi(x). \quad (48)$$

with Δ the scaling dimension and s the conformal spin. And:

$$J^\mu{}_\nu(x) = \frac{\partial x'^\mu}{\partial x^\nu}. \quad \Omega(x) \equiv \sqrt{\det J(x)} \quad , \quad (49)$$

Θ can be get by solving $\text{Tr } J = 2\Omega \cos \Theta \quad \varepsilon^{\alpha\beta} J_{\alpha\beta} = 2\Omega \sin \Theta$

If a field has scaling dimension Δ and conformal spin s , we can define its holomorphic and antiholomorphic dimensions as:

$$h = \frac{1}{2}(\Delta + s) \quad \bar{h} = \frac{1}{2}(\Delta - s) \quad (50)$$

Definition 9. Primary Field (holomorphic coordinate)

A field $\phi(z, \bar{z})$ is called Primary field if under conformal transformation $z \rightarrow w(z)$, it transforms as:

$$\phi'(w, \bar{w}) = \left(\frac{dw}{dz}\right)^{-h} \left(\frac{d\bar{w}}{d\bar{z}}\right)^{-\bar{h}} \phi(z, \bar{z}) \quad (51)$$

We can write the infinitesimal version of the transformation as:

$$\delta_{\epsilon, \bar{\epsilon}} \phi \equiv \phi'(z, \bar{z}) - \phi(z, \bar{z}) = -(h\phi\partial\epsilon + \epsilon\partial\phi) - (\bar{h}\phi\bar{\partial}\bar{\epsilon} + \bar{\epsilon}\bar{\partial}\phi) \quad (52)$$

If we write it in the normal coordinate (z^0, z^1) , we have:

$$\delta\phi = \epsilon^\mu \partial_\mu \phi - \frac{\Delta}{2} \phi \partial_\mu \epsilon^\mu - \frac{is}{2} \phi \epsilon^{\alpha\beta} \partial_\alpha \epsilon_\beta. \quad (53)$$

Which is exactly a field with scaling dimension Δ and spin s under general coordinate transformation. Thus we can see that the two definitions are equivalent.

2.2.2 Correlation Function of Primary Fields

If the theory has conformal symmetry then we know that the correlation function of Primary Fields should satisfy the condition theorem 8 or the infinitesimal version eq. (37). These forms a constraint on the correlation functions of Primary fields.

Theorem 8. Correlation Function of Primary Fields

If a theory has conformal symmetry, then the correlation function of Primary fields should satisfy:

$$\langle \phi_1(z_1, \bar{z}_1) \phi_2(z_2, \bar{z}_2) \rangle = \frac{C_{12}}{(z_1 - z_2)^{2h} (\bar{z}_1 - \bar{z}_2)^{2\bar{h}}} \quad \text{if} \quad \begin{cases} h_1 = h_2 = h \\ \bar{h}_1 = \bar{h}_2 = \bar{h} \end{cases} \quad (54)$$

$$\langle \phi_1(x_1) \phi_2(x_2) \phi_3(x_3) \rangle = C_{123} \frac{1}{z_{12}^{h_1+h_2-h_3} z_{23}^{h_2+h_3-h_1} z_{13}^{h_3+h_1-h_2}} \times \frac{1}{\bar{z}_{12}^{\bar{h}_1+\bar{h}_2-\bar{h}_3} \bar{z}_{23}^{\bar{h}_2+\bar{h}_3-\bar{h}_1} \bar{z}_{13}^{\bar{h}_3+\bar{h}_1-\bar{h}_2}} \quad (55)$$

These are standard results. We note that the function are fully fixed up to a factor C_{12} and C_{123} . In fact these two factor are not independent, if we make ϕ_3 a identity operator and put it to infinity then we can find that C_{123} reduces to C_{12} .

These constant C_{123} are called the **Structure Constants** of the theory.

2.3 Ward Identity of Primary Fields

2.3.1 Ward Identity in Normal Coordinate

If a theory contains Primary field has conformal symmetry, then we can write down the Ward identity for the correlation function of Primary fields from theorem 6, where $X = \prod_{i=1}^n \phi_i(x_i)$

is a series of Primary fields with scaling dimension Δ_i and conformal spin s_i :

$$\frac{\partial}{\partial x^\mu} \langle T_\nu^\mu(x) X \rangle = - \sum_{i=1}^n \delta(x - x_i) \frac{\partial}{\partial x_i^\nu} \langle X \rangle \quad (56) \quad \{\text{eq:Ward Id}\}$$

$$\varepsilon_{\mu\nu} \langle T^{\mu\nu}(x) X \rangle = -i \sum_{i=1}^n \delta(x - x_i) s_i \langle X \rangle \quad (57) \quad \{\text{eq:Ward Id}\}$$

$$\langle T_\mu^\mu(\mathbf{x}) X \rangle = - \sum_{i=1}^n \delta(x - x_i) \Delta_i \langle X \rangle \quad (58)$$

Remember that we use the symmetric traceless energy-momentum tensor here. This is a standard result which can be found in yellow book 5.2.1

2.3.2 Ward Identity in Holomorphic Coordinate

[here I make it more concise than the yellow book]

We can also rewrite them in holomorphic coordinate. First we rewrite the energy-momentum tensor in holomorphic coordinate, we take the following definition:

$$T(z) = -2\pi T_{zz}(z) = -\pi/2(T_{00} - T_{11} - 2iT_{01}) \quad (59) \quad \{\text{eq:holomorphy}\}$$

$$\bar{T}(\bar{z}) = -2\pi T_{\bar{z}\bar{z}}(\bar{z}) = -\pi/2(T_{00} - T_{11} + 2iT_{01}) \quad (60) \quad \{\text{eq:holomorphy}\}$$

$$T_{z\bar{z}} = \frac{1}{4} T^\mu{}_\mu \quad (61)$$

[I copy this from the fudan lecture note, I haven't check this carefully yet.] The Traceless condition $T^\mu{}_\mu = 0$ means $T_{z\bar{z}} = 0$. And the conservation condition $\partial_\mu T^{\mu\nu} = 0$ means:

$$\bar{\partial} T(z, \bar{z}) = 0, \quad \partial \bar{T}(z, \bar{z}) = 0 \quad (62)$$

Thus we see that $T(z)$ is holomorphic and $\bar{T}(\bar{z})$ is antiholomorphic.

Remark:

These are all classical results, till now we don't know that in coorelation functions these properties still hold. But we will see that they only need little modification.

Then we wish to put the ward identity in holomorphic coordinate. We need some important complex analysis results:

- Total derivative in holomorphic coordinate:

$$\int_M d^2x \partial_\mu F^\mu = \frac{1}{2} i \int_{\partial M} (-dz F^{\bar{z}} + d\bar{z} F^z) \quad (63)$$

Remark:

The integral contour is conter-clockwise for the holomorphic part and clockwise for the antiholomorphic part.(we write ∂M but it have different orientation for holomorphic and antiholomorphic part)

- Detla function times a function from residue theorem:

$$\frac{1}{2\pi i} \oint_{C_i} dw \frac{\epsilon(w)}{(w - w_i)^2} = \partial \epsilon(w_i), \quad \frac{1}{2\pi i} \oint_{C_i} dw \frac{\epsilon(w)}{w - w_i} = \epsilon(w_i) \quad (64)$$

With these in mind we notice that the LHS of the above three ward identities can be combined into a total derivative according to the properties of conformal transoformation:

$$\partial_\mu (\epsilon_\nu T^{\mu\nu}) = \epsilon_\nu \partial_\mu T^{\mu\nu} + \frac{1}{2} (\partial_\rho \epsilon^\rho) \eta_{\mu\nu} T^{\mu\nu} + \frac{1}{2} \varepsilon^{\alpha\beta} \partial_\alpha \epsilon_\beta \varepsilon_{\mu\nu} T^{\mu\nu} \quad (65) \quad \{\text{eq:derivati}\}$$

Remark:

Here we use the fact that in 2D conformal transformation we have:

$$\partial_\mu \epsilon_\nu + \partial_\nu \epsilon_\mu = \eta_{\mu\nu} \partial_\rho \epsilon^\rho \quad (66)$$

Remark:

We notice that classically $T^\mu_\mu = 0$ and $\varepsilon_{\mu\nu} T^{\mu\nu} = 0$, thus the last two terms vanish. However, in quantum case these two terms may not vanish thus we never should ignore them.

Then we integrate the LHS of eq. (65) over a region M the left hand side becomes:

$$LHS = \int_M d^2x \partial_\mu \langle T^{\mu\nu}(x) \epsilon_\nu(x) X \rangle \quad (67)$$

we use the above trick to rewrite it as a contour integral:

$$LHS = \frac{1}{2}i \int_{\partial M} (-dz \langle T^{\bar{z}\bar{z}} \epsilon_{\bar{z}} X \rangle + d\bar{z} \langle T^{zz} \epsilon_z X \rangle) \quad (68)$$

We note that according to eq. (56) the $\langle T^\mu_\mu X \rangle$ vanishes on the contour, thus we don't have the $T_{z\bar{z}}$ term here. With a rewrite we can have:

$$LHS = -\frac{1}{2\pi i} \oint_{\partial M} dz \epsilon(z) \langle T(z) X \rangle + \frac{1}{2\pi i} \oint_{\partial M} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z}) X \rangle \quad (69)$$

$$= -\frac{1}{2\pi i} \oint_C dz \epsilon(z) \langle T(z) X \rangle - \frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z}) X \rangle \quad (70)$$

$$(71)$$

Remark:

Here we use that the contour C is counter-clockwise for the holomorphic part and correspondingly clockwise for the antiholomorphic part. Similarly the contour $\bar{C}$ is counter-clockwise for the antiholomorphic part and clockwise for the holomorphic part.




Figure 1: Explanation of the Contours C and $\bar{C}$

{fig:contour}

Then we Tackle the RHS of eq. (65), we plug in the ward identities eq. (56) and get:

$$RHS = \sum_{i=1}^n \int_M d^2x \delta(x - x_i) \left[-\epsilon^\nu(x) \frac{\partial}{\partial x_i^\nu} - \frac{\Delta_i}{2} \partial_\mu \epsilon^\mu(x) - \frac{is_i}{2} \varepsilon^{\alpha\beta} \partial_\alpha \epsilon_\beta(x) \right] \langle X \rangle \quad (72)$$

$$= \sum_{i=1}^n \left[-\epsilon^\nu(x_i) \frac{\partial}{\partial x_i^\nu} - \frac{\Delta_i}{2} \partial_\mu \epsilon^\mu(x_i) - \frac{is_i}{2} \varepsilon^{\alpha\beta} \partial_\alpha \epsilon_\beta(x_i) \right] \langle X \rangle \quad (73)$$

$$= \sum_{i=1}^n \left[-\epsilon(z_i) \partial - h_i \partial \epsilon(z_i) - \bar{\epsilon}(\bar{z}_i) \bar{\partial} - \bar{h}_i \bar{\partial} \bar{\epsilon}(\bar{z}_i) \right] \langle X \rangle \quad (74)$$

If we use the second mathematical trick we mentioned above, we can rewrite it in a contour integral form:

$$RHS = -\frac{1}{2\pi i} \oint_C dw \epsilon(w) \sum_{i=1}^n \left[\frac{1}{w - z_i} \partial_{z_i} + \frac{h_i}{(w - z_i)^2} \right] \langle X \rangle \quad (75)$$

$$-\frac{1}{2\pi i} \oint_{\bar{C}} d\bar{w} \bar{\epsilon}(\bar{w}) \sum_{i=1}^n \left[\frac{1}{\bar{w} - \bar{z}_i} \partial_{\bar{z}_i} + \frac{\bar{h}_i}{(\bar{w} - \bar{z}_i)^2} \right] \langle X \rangle \quad (76)$$

Remark:

Note the the Residue theorem take the + sign when integrating conter-clockwise and - sign when integrating clockwise.

$$\oint_C dz f(z) = 2\pi i \sum_i \text{Res}(f, z_i), \quad \oint_{\bar{C}} d\bar{z} f(\bar{z}) = 2\pi i \sum_i \text{Res}(f, \bar{z}_i) = -\oint_C d\bar{z} f(\bar{z}) \quad (77)$$

thus we integrate both conter-clockwise here (C contour for the holomorphic part and $\bar{C}$ contour for the antiholomorphic part) to get the correct signs.

Thus we combine the LHS and RHS and get the Conformal Ward Identity for Primary fields:

Theorem 9. Conformal Ward Identity for Primary Fields

{thm:conform

For a series of primary fields $X = \prod_{i=1}^n \phi_i(z_i, \bar{z}_i)$ with holomorphic and antiholomorphic dimensions $(h_i, \bar{h}_i)$, we have:

$$-\frac{1}{2\pi i} \oint_C dz \epsilon(z) \langle T(z) X \rangle - \frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z}) X \rangle = \sum_{i=1}^n \left[-\epsilon(z_i) \partial - h_i \partial \epsilon(z_i) - \bar{\epsilon}(\bar{z}_i) \bar{\partial} - \bar{h}_i \bar{\partial} \bar{\epsilon}(\bar{z}_i) \right] \langle X \rangle \quad (78)$$

$$= \delta_{\epsilon, \bar{\epsilon}} \langle X \rangle \quad (79)$$

note the integral contour is different for the holomorphic and antiholomorphic parts as explained above and $\epsilon(z)^* = \bar{\epsilon}(\bar{z})$

• **Decouple of Holomorphic and Antiholomorphic Part**

We know that the holomorphic and antiholomorphic parts are dependent, and we have $\epsilon(z)^* = \bar{\epsilon}(\bar{z})$. However, we manually decouple them and keep in mind that the theory is meaningful only when we combine them back.

Thus we have the Conformal Ward Identity for Primary fields decoupled into holomorphic and antiholomorphic parts by taking $\epsilon(z) = 0$ or $\bar{\epsilon}(\bar{z}) = 0$:

Theorem 10. Conformal Ward Identity for Primary Fields (decoupled form)

For a series of primary fields $X = \prod_{i=1}^n \phi_i(z_i, \bar{z}_i)$ with holomorphic and antiholomorphic dimensions $(h_i, \bar{h}_i)$, we have:

$$-\frac{1}{2\pi i} \oint_C dz \epsilon(z) \langle T(z) X \rangle = -\frac{1}{2\pi i} \oint_C dw \epsilon(w) \sum_{i=1}^n \left[\frac{1}{w - z_i} \partial_{z_i} + \frac{h_i}{(w - z_i)^2} \right] \langle X \rangle \quad (80)$$

$$-\frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z}) X \rangle = -\frac{1}{2\pi i} \oint_{\bar{C}} d\bar{w} \bar{\epsilon}(\bar{w}) \sum_{i=1}^n \left[\frac{1}{\bar{w} - \bar{z}_i} \partial_{\bar{z}_i} + \frac{\bar{h}_i}{(\bar{w} - \bar{z}_i)^2} \right] \langle X \rangle \quad (81)$$

More conveniently, we can write them in the following local form:

$$\langle T(z)X \rangle = \sum_{i=1}^n \left[\frac{h_i}{(z - z_i)^2} + \frac{1}{z - z_i} \partial_{z_i} \right] \langle X \rangle \quad (82)$$

$$\langle \bar{T}(\bar{z})X \rangle = \sum_{i=1}^n \left[\frac{\bar{h}_i}{(\bar{z} - \bar{z}_i)^2} + \frac{1}{\bar{z} - \bar{z}_i} \partial_{\bar{z}_i} \right] \langle X \rangle \quad (83)$$

However, we should keep in mind that this local form is only valid within contour integrals of arbitrary holomorphic (antiholomorphic) functions.

2.3.3 Remarks on 2 Conventions of E-M Tensor

There are two conventions of the relation between the E-M tensor and the holomorphic one.

- **Convention of the Yellow Book:** $T(z) = -2\pi T_{zz}$
- **Other Convention:** $T(z) = T_{zz}$

As far as I know and I think people are able to track, $T(z)$ in both conventions are exactly the same. They all satisfy the same properties and given the same ward identity. **Convention are only different because the different definition of the lagrangian and $T_{\mu\nu}$ in different textbooks.**

What is different is the definition of $T_{\mu\nu}$. However, textbooks never tell us how they defines $T_{\mu\nu}$ is none unique. In fact we don't need to worry about this because, we will always use $T(z)$ as a standard.

2.4 Conformal Ward Identity

2.4.1 Conformal Ward Identity for General Fields

Our above discussion is limited to Primary fields. However, we can extend it to general fields we take the following as a axiom or assumption. We define the infinitesimal variation of the correlation containing general fields:

Axiom 1. Conformal Ward Identity for General Fields

{axm: conform

For a series of general fields $X = \prod_{i=1}^n \phi_i(z_i, \bar{z}_i)$ with holomorphic and antiholomorphic dimensions $(h_i, \bar{h}_i)$, we have:

$$\delta_{\epsilon, \bar{\epsilon}} \langle X \rangle = -\frac{1}{2\pi i} \oint_C dz \epsilon(z) \langle T(z)X \rangle - \frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z})X \rangle \quad (84)$$

or if we decouple the holomorphic and antiholomorphic parts:

$$\delta_{\epsilon} \langle X \rangle = -\frac{1}{2\pi i} \oint_C dz \epsilon(z) \langle T(z)X \rangle \quad (85)$$

$$\delta_{\bar{\epsilon}} \langle X \rangle = -\frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \langle \bar{T}(\bar{z})X \rangle \quad (86)$$

We will see that in a CFT, we may have other fields that we only know their correlation function but have no idea of their transformation properties, thus we will turn to this definition.

2.4.2 Remark on the convention of infinitesimal variation

The infinitesimal variation $\delta_{\epsilon, \bar{\epsilon}}$ has two mainly used conventions:

- **Yellow book convention:** $\delta_{\epsilon, \bar{\epsilon}} = -h\phi\partial\epsilon - \epsilon\partial\phi - \bar{h}\phi\bar{\partial}\bar{\epsilon} - \bar{\epsilon}\bar{\partial}\phi$
- **Other convention:** $\delta_{\epsilon, \bar{\epsilon}} = h\phi\partial\epsilon + \epsilon\partial\phi + \bar{h}\phi\bar{\partial}\bar{\epsilon} + \bar{\epsilon}\bar{\partial}\phi$

Using different conventions will lead to a minus sign difference in the conformal ward identity. Here we follow the yellow book convention.

2.5 OPE Structure

2.5.1 Definition of OPE

Above we notice from the conformal Ward Identity of Primary fields that the correlation function of primary fields and E-M tensor has a divergence structure when they approach each other. This is a general feature of QFT, which can be captured by the Operator Product Expansion (OPE).

Remark:

In fact, we will finally notice that this structure can both give us the data of the correlation functions and the operator commutation relation between fields.

Definition 10. Operator Product Expansion (OPE)

In a QFT, when two fields $A(z)$ and $B(w)$ in a correlation function approach each other, their product can be expanded as:

$$A(z)B(w) = \sum_{n=-\infty}^N \frac{\{AB\}_n(w)}{(z-w)^n} \quad (87)$$

We understand this equation in the sense of correlation functions. Normally, we neglect the regular part $n \leq 0$ and only keep the singular part $n > 0$. and we write $\sim$ instead of $=$ to emphasize this point.

2.5.2 OPE in CFT

From the above discussions of E-M tensor and Primary fields, we can write down the EM-Primary OPE and Primary-Primary OPE in CFT.

Theorem 11. E-M Tensor - Primary Field OPE

For a CFT the OPE between E-M tensor and Primary field $\phi(z, \bar{z})$ with holomorphic and antiholomorphic dimensions $(h, \bar{h})$ is:

$$T(z)\phi(w, \bar{w}) \sim \frac{h}{(z-w)^2}\phi(w, \bar{w}) + \frac{1}{z-w}\partial_w\phi(w, \bar{w}) \quad (88)$$

$$\bar{T}(\bar{z})\phi(w, \bar{w}) \sim \frac{\bar{h}}{(\bar{z}-\bar{w})^2}\phi(w, \bar{w}) + \frac{1}{\bar{z}-\bar{w}}\partial_{\bar{w}}\phi(w, \bar{w}) \quad (89)$$

This is a direct result of the conformal Ward identity for Primary fields. Moreover we can write down the Primary-Primary OPE. We observe the correlation function between Primary fields theorem 8 and we notice that we have a natural normalization for Primary fields:

Definition 11. Normalization of Primary Fields

We often take the normalization of Primary fields such that:

$$\langle \phi_j(z_j, \bar{z}_j) \phi_k(z_k, \bar{z}_k) \rangle = \delta_{jk} / (z_j - z_k)^{2h_j} (\bar{z}_j - \bar{z}_k)^{2\bar{h}_j}. \quad (90)$$

Thus, under this normalization, if we want theorem 8 to hold, we have to take the Primary-Primary OPE as:

Theorem 12. Primary Field - Primary Field OPE

For a CFT the OPE between two Primary fields $\phi_i(z, \bar{z})$ and $\phi_j(z, \bar{z})$ with holomorphic and antiholomorphic dimensions $(h_i, \bar{h}_i)$ and $(h_j, \bar{h}_j)$ is:

$$\phi_i(z_1, \bar{z}_1) \cdot \phi_j(z_2, \bar{z}_2) \sim \sum_k C_{ijk} (z_1 - z_2)^{-h_i-h_j+h_k} (\bar{z}_1 - \bar{z}_2)^{-\bar{h}_i-\bar{h}_j+\bar{h}_k} \phi_k(z_1, \bar{z}_1) + \dots, \quad (91)$$

Proof: I notice that in many textbooks this proof is missing, so I write it down here for completeness.

Consider the three-Point function, we expand it by assuming z_{12} is very small:

$$G_{123}(z_1, z_2, z_3) = C_{123} z_{12}^{h_3-h_1-h_2} z_{23}^{h_1-h_2-h_3} z_{13}^{h_2-h_1-h_3}, \quad (92)$$

we make a change of variable $z_{13} = z_1 - z_3 = (z_2 - z_3) + z_{12} = z_{23} + z_{12}$. Then we can expand it as:

$$z_{13}^{h_2-h_1-h_3} = (z_{23} + z_{12})^{h_2-h_1-h_3} = z_{23}^{h_2-h_1-h_3} \left(1 + \frac{z_{12}}{z_{23}} \right)^{h_2-h_1-h_3} \quad (93)$$

We plug it back to the three-point function and consider the leading order term:

$$G_{123} = C_{123} z_{12}^{h_3-h_1-h_2} z_{23}^{h_1-h_2-h_3} z_{23}^{h_2-h_1-h_3} \quad (94)$$

then through an observation that we can see $(h_1 - h_2 - h_3) + (h_2 - h_1 - h_3) = -2h_3$. So we have in the leading order:

$$G_{123} = C_{123} z_{12}^{h_3-h_1-h_2} z_{23}^{-2h_3} \quad (95)$$

There is another way to calculate this three-point function. We can first use the OPE between $\phi_1(z_1)$ and $\phi_2(z_2)$ and calculate the two-point function between the result and $\phi_3(z_3)$.

The OPE of ϕ_1, ϕ_2 to produce ϕ_3 must have the position dependence $z_{12}^{-h_1-h_2+h_3}$ to make the three-point function conformally invariant. So we take an ansatz for the OPE:

$$\phi_1(z_1) \cdot \phi_2(z_2) \sim \sum_k K_{12k} z_{12}^{-h_1-h_2+h_k} \phi_k(z_2) + \dots \quad (96)$$

We plug the ansatz into the three-point function and get:

$$G_{123} = \langle \phi_1(z_1) \phi_2(z_2) \phi_3(z_3) \rangle \sim \sum_k K_{12k} z_{12}^{-h_1-h_2+h_k} \langle \phi_k(z_2) \phi_3(z_3) \rangle \quad (97)$$

$$= \sum_k K_{12k} z_{12}^{-h_1-h_2+h_k} \delta_{k3} z_{23}^{-2h_3} \quad (98)$$

$$= K_{123} z_{12}^{-h_1-h_2+h_3} z_{23}^{-2h_3} \quad (99)$$

Thus we can conclude that $K_{123} = C_{123}$.

□

Remark:

The Above proof I only use the holomorphic part, but the antiholomorphic part is completely analogous.

Remark:

Sadly, not all paper take this normalization, so we have to be careful to distinguish the difference between OPE coefficients and correlation function structure constants. So sometimes we see:

$$\phi_i \cdot \phi_j \sim \frac{C_{ijk}}{C_{kk}} \phi_k + \dots \quad (100)$$

There is another straightforward observation. OPE of the weight field should transform as the same style under scale transformation. Thus the OPE of fields should take the following general form:

Theorem 13. General Form of OPE in CFT

For a CFT the OPE between two general fields $A(z, \bar{z})$ and $B(z, \bar{z})$ with holomorphic and antiholomorphic dimensions $(h_A, \bar{h}_A)$ and $(h_B, \bar{h}_B)$ is:

$$A(z_1, \bar{z}_1) \cdot B(z_2, \bar{z}_2) \sim \sum_k C_{AB}^k (z_1 - z_2)^{-h_A-h_B+h_k} (\bar{z}_1 - \bar{z}_2)^{-\bar{h}_A-\bar{h}_B+\bar{h}_k} O_k(z_2, \bar{z}_2) + \dots, \quad (101)$$

Where C_{AB}^k is a constant and $O_k(z, \bar{z})$ is a general field with holomorphic and antiholomorphic dimensions $(h_k, \bar{h}_k)$.

2.6 Complementary Remarks**2.6.1 Transformation Rules of E-M Tensor**

From the definition of OPE between E-M tensors and the definition of general conformal Ward identity, we can deduce the transformation rules of E-M tensor under conformal transformation. We take an infinitesimal conformal transformation $z \rightarrow z + \epsilon(z)$, then we have:

Theorem 14. Transformation Rules of E-M Tensor

Under an infinitesimal conformal transformation $z \rightarrow w(z)$, the E-M tensor transforms as:

$$T'(w) = \left(\frac{dw}{dz} \right)^{-2} \left[T(z) - \frac{c}{12} \{w; z\} \right] \quad (102)$$

$$= \left(\frac{dw}{dz} \right)^{-2} T(z) + \frac{c}{12} \{z; w\} \quad (103)$$

where $\{w; z\}$ is the Schwarzian derivative defined as:

$$\{w; z\} = \frac{(d^3w/dz^3)}{(dw/dz)} - \frac{3}{2} \left(\frac{d^2w/dz^2}{dw/dz} \right)^2 \quad (104)$$

We notice that the schwarzian term vanishes for global conformal transformation, thus the E-M tensor transforms as a quasi-primary field with holomorphic dimension $h = 2$.

3 CFT Operator Formalism

{sec:CFT Op

3.1 Radial Quantization

The above discussion is purely from the path-integral formalism. However, we can also understand CFT from the canonical quantization approach. This approach is called Radial Quantization and the result is called Operator Formalism.

3.1.1 Radial Quantization Setup

Axiom 2. Radial Quantization

In 2D CFT we use radial direction as the time direction. The Hilbert Space is defined on a circle with fixed radius.

If you want to ask why we do this, well, because this is the only way that gives us amazing results in CFT... Now we can make everything look like normal canonical quantization:

- We make all field an operator on the Hilbert Space! though knowing nothing about their commutation relation or even the Hilbert Space structure yet.
- We make the Diation Operator (though we doesn't know what the hell it is) the Hamiltonian in radial quantization.
- We define time ordering as radial ordering, which orders operators according to their radius (the smaller radius operator is on the right).
- ...

In the radial quantization, the time direction is the radial direction and the equal time slice is a circle with fixed radius. If we write the coordinate into the holomorphic coordinate and anti-holomorphic then it is shown by the following diagram:




Figure 2: Radial Quantization in Holomorphic Coordinate

{fig:holoco

And just like normal QFT, we often define operators at a radius by a contour integral of operators on the radius (for example the mode operators in QFT is defined by a integral on a time slice). For example, we can define an operator A_1 from an operator $a(z)$ as

$$A_1 = \oint a(z) dz \quad (105)$$

3.1.2 Hermite Conjugation in Radial Quantization

{sec:Hermite

In normal canonical quantization, we define the Hermite Conjugation of operators. In radial quantization, we can define the Hermite Conjugation as well but a natural definition is a bit different:

Definition 12. *Hermite Conjugation of Primary Field in Radial Quantization*

For Primary Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$, its Hermite Conjugation is defined as:

$$[\phi(z, \bar{z})]^\dagger = \bar{z}^{-2h} z^{-2\bar{h}} \phi(1/\bar{z}, 1/z) \quad (106)$$

This is a strange definition, but it works in many places, so I won't claim more on it. [\[I don't think I fully understand this\]](#)

3.1.3 Mode Expansion in Radial Quantization

In normal QFT, we often expand field operators in terms of mode operators. In radial quantization, we can do the same thing. For example, for a Primary Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$, we can make a Laurent expansion:

Definition 13. *Mode Expansion of Primary Field in Radial Quantization*

{defi:Mode E

The Mode Expansion of a Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$ is given by:

$$\phi(z, \bar{z}) = \sum_{m \in \mathbf{Z}} \sum_{n \in \mathbf{Z}} z^{-m-h} \bar{z}^{-n-\bar{h}} \phi_{m,n} \quad (107)$$

$$\phi_{m,n} = \frac{1}{2\pi i} \oint dz z^{m+h-1} \frac{1}{2\pi i} \oint d\bar{z} \bar{z}^{n+\bar{h}-1} \phi(z, \bar{z}) \quad (108)$$

- Attention the integral contour is counter-clockwise for the z integral but clockwise for the $\bar{z}$ integral. This is adopted to keep the same orientation as the normal integral on $x - y$ plane.

We use this definition for convention, we will then see its convenience through the Hermite Conjugation of Mode Operators and commutation calculations. We then calculate the Hermite Conjugation of Mode Operators:

Theorem 15. *Hermite Conjugation of Mode Operators*

{thm:Hermite

The Hermite Conjugation of the Mode Operators $\phi_{m,n}$ of a Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$ is given by:

$$\phi_{m,n}^\dagger = \phi_{-m,-n} \quad (109)$$

3.1.4 OPE in Radial Quantization

In path-integral formalism we define OPE within the correlation functions and say nothing about operators (because we don't have operators in path-integral formalism). However, in radial quantization we have operators, thus we can define OPE as an operator relation:

Theorem 16. OPE as Operator Relation

Consider two operators $A(z)$ and $B(w)$, their OPE is defined as the operator relation

$$\mathcal{R}A(z)B(w) = \sum_{n=-\infty}^N \frac{\{AB\}_n(w)}{(z-w)^n} \quad (110)$$

where $\mathcal{R}$ is the radial ordering operator, which orders operators according to their radius (the smaller radius operator is on the right).

Proof: due to the relation between correlation functions in path integral formalism and canonical formalism, we can see that this result is straightforward.

The Radial Ordering property of OPE is very important for it naturally generates a commutator structure, which leads to a pronound result.

3.1.5 Commutator from OPE

The first amazing result of radial quantization is that we can get the commutator of two operators from their OPE. We first consider an equality:

Theorem 17. Commutators of local operators from OPE

Consider two local operators $a(z)$ and $b(w)$, and an operator A defined as:

$$A = \oint dz a(z) \quad (111)$$

Then we have:

$$\oint_w dz \mathcal{R}a(z)b(w) = \oint_{C_1} dz a(z)b(w) - \oint_{C_2} dz b(w)a(z) = [A, b(w)] \quad (112) \quad \{\text{eq:Commutator}\}$$

The Proof is quite straightforward. We note that in the middow part we have to change the order of the operators because of the definition of radial ordering.

Remark:

Note that the definition of redial ordering is different for fermions, we have:

$$\mathcal{R}a(z)b(w) = \begin{cases} a(z)b(w) & |z| > |w| \\ -b(w)a(z) & |z| < |w| \end{cases} \quad (113)$$

Thus, for fermionic operators the above theorem should be modified to **changing the commutator to anti-commutator**

We can generalize this to get the commutator of two operators from their OPE:

Theorem 18. Commutator from OPE

Consider two operators on a radial quantization Hilbert Space $A(z)$ and $B(w)$, their commutator is given by:

$$[A, B] = \oint_0 dw \oint_w dz \mathcal{R}a(z)b(w) \quad (114)$$

Where $\mathcal{R}a(z)b(w)$ is exatly the OPE of these two local operators.

3.2 Symmetry Operator of CFT

3.2.1 Symmetry Charges and Algebra

In canonical quantization, we define the symmetry charge as the quantization of the noether charge which is also the generator of symmetry transformation by commutators. The property of OPE between general fields axiom 1 leads to that:

$$\delta_\epsilon \Phi(w) = -[Q_\epsilon, \Phi(w)] \quad (115)$$

where Q_ϵ is:

$$Q_\epsilon = \frac{1}{2\pi i} \oint_C dz \epsilon(z) T(z) \quad (116)$$

The integral contour C is a circle around the origin with w inside it.

Proof: we just plug in the conformal ward identity to eq. (112).

Remark:

In fact this claim is not so accurate. In fact, if we only want to act on the field at w , the contour should be a circle around the point w . But now we just assume that the operator act on every point inside the contour C , which is a equal time slice.

Then we get the theorem:

Theorem 19. *Symmetry Chages in Radial Quantized CFT*

The Symmetry Charge associated to a conformal transformation $z \rightarrow z + \epsilon(z)$ is given by:

$$Q_\epsilon = \frac{1}{2\pi i} \oint dz \epsilon(z) T(z) \quad (117)$$

We can as well analogue the following as sort of "Symmetry Current":

$$J_\epsilon(z) = \epsilon(z) T(z) \quad (118)$$

We notice that this is a superposition of infinite many symmetry charges. Thus, to make its mathematical structure clear, we can make a Laurent expansion of $T(z)$, canonically we use the lorant expansion defined above definition 13:

$$T(z) = \sum_{n \in \mathbb{Z}} z^{-n-2} L_n \quad L_n = \frac{1}{2\pi i} \oint dz z^{n+1} T(z) \quad (119)$$

$$\bar{T}(\bar{z}) = \sum_{n \in \mathbb{Z}} \bar{z}^{-n-2} \bar{L}_n \quad \bar{L}_n = \frac{1}{2\pi i} \oint d\bar{z} \bar{z}^{n+1} \bar{T}(\bar{z}) \quad (120)$$

- Again we note that the integral contour is conter-clockwise for the z integral but clockwise for the $\bar{z}$ integral. This is adopted to keep the same orientation as the normal integral on $x - y$ plane.

Thus the Symmetry Charge can be written as:

$$Q_\epsilon = \sum_{n \in \mathbb{Z}} \epsilon_n L_n \quad \epsilon(z) = \sum_{n \in \mathbb{Z}} \epsilon_n z^{n+1} \quad (121)$$

We thus can see that the true internal structure of the conformal symmetry is given by the operators L_n and $\bar{L}_n$.

Theorem 20. Genuine Symmetry Charges and Symmetry Algebra

The Symmetry Charge of Radial Quantized CFT is generated by the operators L_n and $\bar{L}_n$, which satisfy the Virasoro Algebra:

$$[L_n, L_m] = (n - m)L_{n+m} + \frac{c}{12}n(n^2 - 1)\delta_{n+m,0} \quad (122)$$

$$[L_n, \bar{L}_m] = 0 \quad (123)$$

$$[\bar{L}_n, \bar{L}_m] = (n - m)\bar{L}_{n+m} + \frac{c}{12}n(n^2 - 1)\delta_{n+m,0} \quad (124)$$

Proof, we just use the OPE trick of EM Tensor to calculate the commutators.

Another important statement is the Hermite Conjugation of Symmetry Charges. From theorem 15 we can directly get:

Theorem 21. Hermite Conjugation of Symmetry Charges

The Hermite Conjugation of the Symmetry Charges L_n and $\bar{L}_n$ are given by:

$$L_n^\dagger = L_{-n} \quad \bar{L}_n^\dagger = \bar{L}_{-n} \quad (125)$$

3.2.2 Symmetry Generators in Radial Quantization**Remark:**

In a ordinary QFT, we can define the Hamiltonian Equivalently in many ways:

- The legendre transformation of Lagrangian
- The integral of the time-time component of E-M Tensor on equal time slice
- The conserve charge of time translation symmetry
- Generator of time translation

We may find that the first three definition has subtlety, for example the lagrangian is equivalent if we add some constant term or times some constants which depends on convention. However, with a concrete definition of the infinitesimal transformation, we have a concrete generator. This definition is the most robust one.

Thus, in CFT, we **define** the Hamiltonian as the generator of time translation in radial quantization.

In a QFT the Hamiltonian is the Time Direction Generator. Thus in the content of radial quantization the Hamiltonian is the Dilation Generator. If we use the result in the holomorphic coordinate, we can see that:

Theorem 22. Hamiltonian (Dilation Operator) in Radial Quantization

The Hamiltonian (Dilation Operator) in Radial Quantization is given by:

$$H = L_0 + \bar{L}_0 \quad (126)$$

In terms of EM Tensor, it is given by:

$$H = \frac{1}{2\pi i} \oint_C dz \, z T(z) + \frac{1}{2\pi i} \oint_C d\bar{z} \, \bar{z} \bar{T}(\bar{z}) \quad (127)$$

$$= L_0 + \bar{L}_0 \quad (128)$$

Proof: This is a **Very Important** Claculation, I'll complete it how.

We define the Hamiltonian as the dilation transformation generator, the dilation transformation is defined as:

$$z \rightarrow z' = (1 + \epsilon)z \quad \bar{z} \rightarrow \bar{z}' = (1 + \bar{\epsilon})\bar{z} \quad (129)$$

Thus, the generator of this transformation is given by:

$$Q_{\epsilon\bar{\epsilon}} = \frac{1}{2\pi i} \oint_C dz \epsilon(z) T(z) + \frac{1}{2\pi i} \oint_{\bar{C}} d\bar{z} \bar{\epsilon}(\bar{z}) \bar{T}(\bar{z}) \quad (130)$$

Moreover, we can see that the space-time translation operators in radial quantization are given by:

Theorem 23. Translation Operators in Radial Quantization

The Translation Operators in Radial Quantization are given by:

$$P_z = L_{-1} \quad P_{\bar{z}} = \bar{L}_{-1} \quad (131)$$

Which corresponds to translation in z and $\bar{z}$ direction respectively.

These operators can be understand as a combination of radial translation and rotation. and in the holomorphic coordinate they are exactly the translation in z and $\bar{z}$ direction.

3.3 Hilbert Space of CFT

3.3.1 Hilbert Space Structure

From usual QM we know that the Hilbert Space shall form a representation of the symmetry algebra of the theory. This is sort of a empirical fact, but it works very well in QFT. Thus, we can make it a axiom in CFT:

Axiom 3. Hilbert Space as Virasoro Algebra Representation

The Hilbert Space of a 2D CFT forms a representation of the Virasoro Algebra generated by L_n and $\bar{L}_n$. Which may be constructed by the highest weight representation of the Virasoro Algebra.

The representation of 2 virasoro algebras can be constructed by the tensor product of two single Virasoro Algebra representations. Thus, we only need to consider the representation of a single Virasoro Algebra.

We further introduce the concept of Heighest Weight Representation to build the representation of the Virasoro Algebra:

- **Highest Weight State:** assume there exists a state $|h, \bar{h}\rangle$ in the Hilbert Space such that:

$$L_0|h\rangle = h|h\rangle \quad L_n|h\rangle = 0 \quad n > 0 \quad (132)$$

- **Highest Weight Representation:** we can build a representation of the Virasoro Algebra by acting the lowering operators L_{-n} ($n > 0$) on the highest weight state $|h\rangle$. The states in this representation are called Descendant States.
- **General Representation Space:** The representation space of the Virasoro Algebra is given by the direct sum of all highest weight representations:

$$\mathcal{R} = \bigoplus_{h, \bar{h}} \mathcal{V}_h \otimes \bar{\mathcal{V}}_{\bar{h}} \quad (133)$$

Remark:

This subsection what we discuss is pure math and L_n is just abstract lie algebra. By "state" we just mean an abstract vector in a vector space. We haven't yet connect this math structure to the physical operators and states in CFT. This will be done in the next subsection.

Then we want to find a physical realization of this math structure in CFT.

3.3.2 Hilbert Space Realization

From normal QFT, we know that we can assume a well defined vacuum state $|0\rangle$ in the Hilbert Space. And then we act all possible operators on the vacuum to get all kinds of states in the Hilbert Space. In this general CFT construction, we know that the only operators we have now is:

- The Symmetry Charges L_n and $\bar{L}_n$
- The Primary Fields $\phi(z, \bar{z})$

The question is that can we get the structure of direct sum over highest weight representations (Virasoro Algebra Representation) by acting these operators on some vacuum? The answer is yes, we can.

- **Step 1: Define the Vacuum State**

we first define the vacuum state $|0\rangle$ as the state invariant under all global conformal transformations:

Definition 14. Vacuum State in CFT

The Vacuum State $|0\rangle$ in CFT is defined as the state invariant under all global conformal transformations and well defined when acted on by all local operators. In terms of Symmetry Charges, it satisfies:

$$L_{-1}|0\rangle = L_0|0\rangle = L_1|0\rangle = 0 \quad (134)$$

$$\bar{L}_{-1}|0\rangle = \bar{L}_0|0\rangle = \bar{L}_1|0\rangle = 0 \quad (135)$$

Now if we also assume that the state $T(0)|0\rangle$ is well defined, then we can further get:

$$L_n|0\rangle = \bar{L}_n|0\rangle = 0 \quad n \geq -1 \quad (136)$$

This also means that the vacuum energy is zero:

$$\langle 0|T(z)|0\rangle = \langle 0|\bar{T}(\bar{z})|0\rangle = 0 \quad (137)$$

- **Step 2: Define Primary States**

We then try to act Primary Fields on the vacuum to get some states. We define:

Definition 15. Primary State in CFT

The Primary State $|h, \bar{h}\rangle$ associated to a Primary Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$ is defined as:

$$|h, \bar{h}\rangle = \phi(0, 0)|0\rangle \quad (138)$$

This state has real unique properties. To show this we first calculate the operator commutator between L_n and $\phi(z, \bar{z})$:

$$[L_n, \phi(z, \bar{z})] = z^n ((n+1)h + z\partial_z) \phi(z, \bar{z}) \quad n \geq -1 \quad (139)$$

$$[\bar{L}_n, \phi(z, \bar{z})] = \bar{z}^n ((n+1)\bar{h} + \bar{z}\partial_{\bar{z}}) \phi(z, \bar{z}) \quad n \geq -1 \quad (140)$$

With these commutators we can calculate the action of L_n and $\bar{L}_n$ on the Primary State:

$$L_0|h, \bar{h}\rangle = h|h, \bar{h}\rangle \quad \bar{L}_0|h, \bar{h}\rangle = \bar{h}|h, \bar{h}\rangle \quad (141)$$

$$L_n|h, \bar{h}\rangle = \bar{L}_n|h, \bar{h}\rangle = 0 \quad n > 0 \quad (142)$$

This means that:

Theorem 24. Primary State as Highest Weight State

The Primary State $|h, \bar{h}\rangle$ associated to a Primary Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$ behaves like a Highest Weight State of the Virasoro Algebra with highest weight $(h, \bar{h})$.

• **Step 3: Build the Hilbert Space**

Naturally, we then find all the primary states and act the lowering operators onto them to build a representation of the Virasoro Algebra. States corresponds to Virasoro Generators acting on Primary States are called Descendant States:

Theorem 25. Descendant State in CFT

The Descendant States in CFT are defined as the states obtained by acting the lowering operators L_{-n} ($n > 0$) and $\bar{L}_{-n}$ ($n > 0$) on Primary States.

$$L_{-k_1} L_{-k_2} \cdots L_{-k_n} |h\rangle \quad (1 \leq k_1 \leq \cdots \leq k_n) \quad (143)$$

$$\bar{L}_{-l_1} \bar{L}_{-l_2} \cdots \bar{L}_{-l_m} |\bar{h}\rangle \quad (1 \leq l_1 \leq \cdots \leq l_m) \quad (144)$$

We then take the direct sum of all these representations to build the Hilbert Space.

Definition 16. Hilbert Space of CFT

The Hilbert Space $\mathcal{H}$ of a CFT is given by the direct sum of all highest weight representations of the Virasoro Algebra built from all Primary States:

$$\mathcal{H} = \bigoplus_{h, \bar{h}} \mathcal{V}_h \otimes \bar{\mathcal{V}}_{\bar{h}} = \bigoplus_{h, \bar{h}} \{L_{-k_1} L_{-k_2} \cdots L_{-k_n} |h, \bar{h}\rangle \quad (1 \leq k_1 \leq \cdots \leq k_n)\} \quad (145)$$

• **Final Step: Test? is it done?**

We finally show that all states generated by primary fields and E-M tensor acting randomly on the vacuum can be expressed as linear combinations of the above defined states in the Hilbert Space. I want to discuss a few cases:

1. **Case 0: Single Primary and many Virasoro Generators**

We only have to commute them to the right order to get the canonical form.

2. **Case 1: Many Primaries** consider a state generated by many primary fields acting on the vacuum:

$$|\psi\rangle = \phi_1(z_1, \bar{z}_1)\phi_2(z_2, \bar{z}_2)\cdots\phi_n(z_n, \bar{z}_n)|0\rangle \quad (146)$$

We can use the OPE of primary fields (Though we haven't yet discussed it) but consistently make it a sum of single primary field and descendants acting on the vacuum.

3. **Case 2: Only E-M Tensor** consider a state generated only by E-M Tensor acting on the vacuum. In fact by definition:

$$T(0)|0\rangle = L_{-2}|0\rangle \quad (147)$$

In fact we can view the vacuum as a primary field with conformal dimension $(0,0)$, thus E-M tensor is the 2nd level descendent of the vacuum primary field. Thus it has nothing different from the zero and first case.

4. **Case 3: Field on Points other than 0** consider a state generated by a field acting on a point other than 0:

$$|\psi\rangle = \phi(z, \bar{z})|0\rangle \quad z \neq 0 \quad (148)$$

We can use the translation symmetry generated by L_{-1} and $\bar{L}_{-1}$ to translate the point to 0:

$$\phi(z+w, \bar{z}+\bar{w}) = e^{zL_{-1}+\bar{z}\bar{L}_{-1}}\phi(w, \bar{w})e^{-zL_{-1}-\bar{z}\bar{L}_{-1}} \quad (149)$$

Then due to the vacuum is invariant under translation, we have:

$$\phi(z, \bar{z})|0\rangle = e^{zL_{-1}+\bar{z}\bar{L}_{-1}}\phi(0,0)|0\rangle \quad (150)$$

which is in the Hilbert Space.

Thus we can argue that all states generated by primary fields and E-M tensor acting randomly on the vacuum can be expressed as linear combinations of the states in the Hilbert Space defined above.

[Nearly all textbook skips this part, but it is really important]

Remark:

Though we write the Hilbert Space as a theorem. Its in fact more like something we only know because we only have the knowledge of primary field and E-M tensor. So in fact it is more like an axiom than a theorem.

3.3.3 Inner Product

As a valid Hilbert Space, we also need to define an inner product. We find that the above definition of Hermite Conjugation in radial quantization section 3.1.2 gives us a natural inner product:

Definition 17. Inner Product in Radial Quantization

The Inner Product between two states $|\psi_1\rangle$ and $|\psi_2\rangle$ in Radial Quantization is defined as:

$$\langle\psi_1|\psi_2\rangle = \langle 0|[\psi_1]^\dagger\psi_2|0\rangle \quad (151)$$

Pratically we can calculate the inner product by commuting all the Virasoro Generators and make the Lowering Operators to act on Primary States. Then we compute the inner product between Primary States:

Theorem 26. Inner Product between Primary States

The inner product between two primary states $|h_1, \bar{h}_1\rangle = \phi_1(0,0)|0\rangle$ and $|h_2, \bar{h}_2\rangle = \phi_2(0,0)|0\rangle$ is determined by the two-point function of their corresponding primary fields:

$$\langle h_1, \bar{h}_1 | h_2, \bar{h}_2 \rangle = \lim_{z, \bar{z}, w, \bar{w} \rightarrow 0} \langle 0 | \phi_1(z, \bar{z})^\dagger \phi_2(w, \bar{w}) | 0 \rangle \quad (152)$$

$$= \lim_{z, \bar{z}, w, \bar{w} \rightarrow 0} z^{-2h_1} \bar{z}^{-2\bar{h}_1} \left\langle 0 \left| \phi_1\left(\frac{1}{\bar{z}}, \frac{1}{z}\right) \phi_2(w, \bar{w}) \right| 0 \right\rangle \quad (153)$$

$$= \lim_{\xi, \bar{\xi} \rightarrow \infty} \bar{\xi}^{2h_1} \xi^{2\bar{h}_1} \langle 0 | \phi_1(\bar{\xi}, \xi) \phi_2(0, 0) | 0 \rangle, \quad (\xi = 1/\bar{z}, \bar{\xi} = 1/z) \quad (154)$$

$$= C_{12}, \quad (155)$$

where C_{12} is the coefficient of the two-point function $\langle \phi_1(z, \bar{z}) \phi_2(0, 0) \rangle = C_{12} z^{-2h_1} \bar{z}^{-2\bar{h}_1}$. In particular, primary states are orthogonal unless they share the same conformal dimensions:

$$\langle h_1, \bar{h}_1 | h_2, \bar{h}_2 \rangle = C_{12} \delta_{h_1, h_2} \delta_{\bar{h}_1, \bar{h}_2}. \quad (156)$$

3.4 Primary and Descendent Fields

We can see that the Hilbert space is made by Primaries and Descendents. In fact, the operators in CFT also have the same structure. We then further this in this subsection.

3.4.1 State-Operator Correspondence

We now notice that we assume the spectrum is exactly given by states at 0 point acting on the vacuum. This leads to a amazing result called State-Operator Correspondence:

Theorem 27. State-Operator Correspondence in CFT

There is a one-to-one correspondence between local operators $\mathcal{O}(z, \bar{z})$ in CFT and states $|\mathcal{O}\rangle$ in the Hilbert Space given by the following relation:

$$|\mathcal{O}\rangle = \mathcal{O}(0, 0)|0\rangle \quad (157)$$

3.4.2 Descendent Operators

We then notice that we can define an operator corresponding to the descendant states as well:

Definition 18. Descendant Operator in CFT

The Descendant Field of a Primary Field $\phi(z, \bar{z})$ with conformal dimension $(h, \bar{h})$ is defined as the operator obtained by acting the Virasoro Generators L_{-n} ($n > 0$) and $\bar{L}_{-n}$ ($n > 0$) on the Primary Field:

$$\phi^{(-n)}(w) \equiv (L_{-n}\phi)(w) = \frac{1}{2\pi i} \oint_w dz \frac{1}{(z-w)^{n-1}} T(z) \phi(w) \quad (158)$$

Generally we can act many Virasoro Generators to get higher level descendant fields:

$$\phi^{(-k, -n)}(w) \equiv (L_{-k} L_{-n} \phi)(w) = \frac{1}{2\pi i} \oint_w dz (z-w)^{1-k} T(z) (L_{-n} \phi)(w) \quad (159)$$

In particular, we have:

$$\phi^{(0)}(w) = h\phi(w) \quad \text{and} \quad \phi^{(-1)}(w) = \partial\phi(w) \quad (160)$$

3.4.3 Operator Data of CFT

Thus more as a assumption, but with good reason (just like the Hilbert Space is), we define the Operator content of CFT as:

Theorem 28. Operator Content of CFT

The Operator Content of a CFT is given by all Primary Fields and their Descendant Fields.

3.4.4 Correlation Functions with Descendant Operators

Using the conformal ward identity of primary fields we can construct a relation between correlation functions with descendant operators and correlation functions with only primary fields:

Theorem 29. Correlation Functions with Descendant Operators(1)

The Correlation Functions with Descendant Operators can be expressed as differential operators acting on correlation functions with only primary fields.

For a descendant operator $\phi^{(-n)}(z)$ of a primary field $\phi(z)$ with conformal dimension h , and a series of Pirmary Fields X we have:

$$\langle \phi^{(-n)}(w) X \rangle = \mathcal{L}_{-n} \langle \phi(w) X \rangle, \quad (n \geq 1) \quad (161)$$

Where:

$$\mathcal{L}_{-n} = \sum_i \left\{ \frac{(n-1)h_i}{(w_i - w)^n} - \frac{1}{(w_i - w)^{n-1}} \partial_{w_i} \right\} \quad (162)$$

Proof: see the following calculation:

$$\langle \phi^{(-n)}(w) X \rangle = \frac{1}{2\pi i} \oint_w dz (z - w)^{1-n} \langle T(z) \phi(w) X \rangle \quad (163)$$

$$= -\frac{1}{2\pi i} \oint_{\{w_i\}} dz (z - w)^{1-n} \sum_i \left[\frac{1}{z - w_i} \partial_{w_i} \langle \phi(w) X \rangle + \frac{h_i}{(z - w_i)^2} \langle \phi(w) X \rangle \right] \quad (164)$$

$$= \mathcal{L}_{-n} \langle \phi(w) X \rangle, \quad (n \geq 1) \quad (165)$$

We can generalize to higher level descendant operators and even to the case X containing descendant operators as well:

Theorem 30. Correlation Functions with Descendant Operators(2)

The Correlation Functions with Descendant Operators can be expressed as differential operators acting on correlation functions with only primary fields.

For a descendant operator $\phi^{(-k, -n)}(z)$ of a primary field $\phi(z)$ with conformal dimension h , and a series of Pirmary Fields X we have:

$$\langle \phi^{(-k, -n)}(w) X \rangle = \mathcal{L}_{-k} \mathcal{L}_{-n} \langle \phi(w) X \rangle, \quad (k, n \geq 1) \quad (166)$$

Where $\mathcal{L}_{-n}$ is defined as above. More generally, we have:

$$\langle \phi^{(-k_1, \dots, -k_n)}(w) X \rangle = \mathcal{L}_{-k_1} \cdots \mathcal{L}_{-k_n} \langle \phi(w) X \rangle \quad (167)$$

In general, a correlation function with many descendant operators can be expressed as differential operators acting on correlation functions with only primary fields, but may take more detailed consideration.

3.4.5 OPE with Descendant Operators

Similar to the Primaries, we can calculate the OPE of Descendant fields and the E-M Tensor. Please refer to yellow book 6.6.2 for details.

We can also calculate the full OPE between two Primary Fields including all descendant fields. Please refer to yellow book 6.6.3 for details.

3.5 Complementary Remarks

3.5.1 Radial Quantization on Cylinder

We use the holomorphic coordinate to do the radial quantization. In fact another more physical and meaningful way is to do the radial quantization on a cylinder coordinate system. Which is related to the holomorphic coordinate by the conformal transformation:

4 Summary: General Structure of CFT

{sec:Summary

4.1 Hilbert Space and Operator Contents

In a General CFT, the Hilbert Space and all Operator Contents can be classified into the Primaries and Their Descendent.

Important: Hilbert Space

The Hilbert Space $\mathcal{H}$ of a CFT is spanned by the highest weight representations of the Virasoro Algebra. Each highest weight representation is generated by a Primary State $|h, \bar{h}\rangle$ and its Descendent States $L_{-n_1}L_{-n_2}\dots|h, \bar{h}\rangle$ ($n_i > 0$).

$$\mathcal{H} = \bigoplus_{h, \bar{h}} \mathcal{V}_h \otimes \bar{\mathcal{V}}_{\bar{h}}, \quad (168)$$

And the Operators are classified similarly

Important: Operator Content

The Operator Content of a CFT is classified by the Primary Operators $\phi_{h, \bar{h}}(z, \bar{z})$ and their Descendent Operators $\phi_{h, \bar{h}}^{(-k_1, \dots, -k_n)}(z, \bar{z})$ ($k_i > 0$).

4.2 Correlation Functions

{sec:CFT Cor

To know all correlation functions of a CFT is equivalent to knowing all physical observables of the CFT which is the ultimate goal of studying a CFT. In fact the correlation function of a CFT is fully determined by the OPEs which we also called the **Operator Algebra** of the CFT.

Definition 19. Operator Algebra

The Operator Algebra is the collection of all OPEs between the operators in the CFT.

If we know the Operator Algebra of a CFT, we can in principle reduce any N-point Correlation Functions to 2-Point Correlation Functions (which is fully determined by conformal symmetry). Thus the full information of a CFT correlation function is encoded in its Operator Algebra.

In fact, the operator algebra of a CFT can be determined if we know a series of data, we write this as the following theorem

Theorem 31. Determination of Operator Algebra

The Operator Algebra of a CFT is fully determined if we know the following data:

- Central Charge c
- Hilbert Space or the Operator Content
- Structural Constants C_{abc}

Proof: See section 6.6.3 of the yellow book for reference. It just calculate the OPE and find that it only depends on these data.

Then the question is, which combination of these data can give a self-consistent CFT? This is the core idea of Conformal Bootstrap which we will discuss in the next section.

5 Axiomatic Formalism

{sec:Bootstr

[please fill in here a short discussion about the BPZ formalism of CFT] [follow Mnev's lecture notes on CFT]

The above three sections we explicitly and implicitly made some assumptions apart from the assumption of conformal symmetry :

- A CFT should contain Primary Fields and their descendants only
- General CFT should contain a E-M Tensor with a suitable OPE
- Conformal Ward identity holds beyond primary fields
- A global conformal invariant vacuum $|0\rangle$
- Hilbert Space is spanned by the highest weight representations of the Virasoro Algebra
- ...

These assumptions are really messy. However, they made the final structure of CFT beautiful and clear. Thus, we generally won't use these explicit and implicit assumptions in the above sections to define a CFT (or we will fall into mess again), but we will make the clear results (OPE Structure, Virasoro Algebra Representation Spectrum , state operator correspondence...) as a defining feature of a CFT. These feature gives a abstract but clear definition of a CFT, this work is first done by Belavin, Polyakov and Zamolodchikov in their famous BPZ paper thus I'll here give a brief introduction to BPZ formalism of CFT.

5.1 BPZ Axioms of CFT

5.2

6 Introduction to Conformal Bootstrap

{sec:Intro

6.1 Introduction to Conformal Bootstrap

The core idea of conformal bootstrap is to ask what kinds of combination of the three data (central charge, Hilbert space/operator content, structural constants) can give a self-consistent CFT. The self-consistency conditions include:

- Unitarity of the Hilbert Space
- Crossing Symmetry of the 4-Point Correlation Functions
- Modular Invariance of the Torus Partition Function

6.2 Unitary Theory

6.3 Crossing Symmetry