


Figure 1 — AT2 regeneration arrests during bleomycin-induced fibrosis; Fgfr2 is retained on stalled cells


A Epithelial cell-state landscape

Mouse bleomycin model (Kobayashi PATS). AT2 progenitors bifurcate: completing or arrested.




B Arrest accumulates during fibrotic consolidation

Fraction of Krt8⁺ ADI cells per time point. n=9 at day 28 excluded.




C Krt8 (arrest marker)

5205 / 17324 cells express (30%)




Cldn4 (arrest marker)

1195 / 17324 cells express (7%)




Fgfr2 (target receptor)

1883 / 17324 cells express (11%)




Kobayashi PATS (GSE141634) + Choi DATPs (GSE145031). Fgfr2 IIIb/IIIc isoform caveat: AT2 cells express predominantly IIIb; total Fgfr2 used as surrogate.

A Analysis pipeline




B Epithelial cell states




C Top 30 sender ligands (mouse NicheNet)

Completing vs arrested fate (geneset_o). Highlighted: FGFR2



A Receptor candidates: composite score



B Per-method scores





Figure 3 — FGF10 sender-side: supply is episodic in mouse and low in human IPF

A Mouse bleomycin time course: Fgf10⁺ Plin2-low fibroblasts

Strunz GSE141259. Pseudobulk Wilcoxon vs PBS. $p = 0.029$ for fibrotic early (d10+d14 pooled).



B Human fibroblasts: FGF10 expression

Adams GSE136831 + Habermann GSE135893. Pseudobulk Wilcoxon. n.s. = $p \geq 0.05$.




Figure 4 — FGFR2 receiver: expression maintained; co-expressed with arrest markers

