



Memo version 2.0

BESIII Analysis Memo

BAM-xxx

BESIII Analysis Memo (January 27, 2026)

Cross section measurement of process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ from threshold to 2.0 GeV via ISR technique

Guang-Yan Xiao ^{a,c1} Ya-Qian Wang ^b Lei Zhang ^a Shen-Jian Chen ^a Chang-Zheng Yuan ^c

^aNanjing University

^bHebei University

^cInstitute of High Energy Physics, CAS

Referee Committee

XXX (Chair)^d, XXX^e, XXX^f

^dXXX University

^eXXX University

^fXXX University

Pubcomm Editors

Editor A^g, Editor B^h

^gUni. G

^hUni. H

HN : <http://hnb3.ihep.ac.cn/HyperNews/get/paper835.html>

Abstract

XXX

¹E-mail: xiaogy@smail.nju.edu.cn

20 **List of changes**

21 For an easy reading of the document, major changes in various versions with respect to the previous
22 one are listed below.

23 **Version 1.0**

- 24 • First version of memo.

25	Contents	
26	1 Introduction	3
27	2 BEPCII and BESIII detector	4
28	3 Data and MC samples	5
29	4 Event selections	7
30	5 Background Study	9
31	5.1 estimate of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$	9
32	5.2 estimate of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$	11
33	6 Unfolding	14
34	7 Efficiency correction	15
35	8 Systematic uncertainties	16
36	9 Summary	17

1 Introduction

to fill

2 BEPCII and BESIII detector

The Beijing Electron-Positron Collider II (BEPCII) —a double-ring multi-bunch collider—is operating in the τ -charm energy region (2 to 4.9 GeV) with the design luminosity of $1 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$ [1]. The Beijing Spectrometer III (BESIII) operating at BEPCII is a high-precision, general-purpose detector that records symmetric e^+e^- collisions provided by the BEPCII storage ring. BESIII is approximately cylindrically symmetric with 93% coverage of the solid angle around the e^+e^- interaction point (IP). The components of the apparatus ordered by distance from the IP, are a 43-layer small-cell main drift chamber (MDC), a time-of-flight (TOF) system based on plastic scintillators with two layers in the barrel region and one layer in the end-cap region (the end-cap TOF system has been updated with the multi-gap resistive plate chamber in 2014 [2]), a 6240-cell CsI(Tl) crystal electromagnetic calorimeter (EMC), a superconducting solenoid magnet providing a 1.0 T magnetic field aligned with the beam axis, and resistive plate muon-counter layers interleaved with steel. The momentum resolution for charged tracks in the MDC is 0.5% for transverse momenta with 1 GeV/c. The energy resolution in the EMC is 2.5% in the barrel region and 5.0% in the end-cap region for 1 GeV photons. For charged tracks, particle identification (PID) for charged tracks combines measurements of the energy deposit dE/dx in MDC and flight time in TOF and forms likelihoods $\mathcal{L}(h)(h = K, \pi)$ for a hadron h hypothesis. More details about the BESIII detector are provided in Ref [3].

Table 1: Beam energies and integrated luminosities of the data samples used in this work, quoted from Ref. [4, 5].

$E_{\text{beam}}(\text{GeV})$	Integrated luminosity (pb^{-1})	Data taking time	RunNo
3.773	$2931.8 \pm 0.2 \pm 13.8$ [4]	2010(round03) 2011(round04)	11414 – 13988, 14395 – 14604 20448 – 23454
3.773	4995.0 ± 19.0 [5]	2022(round15)	70522 – 73929
3.773	8157.0 ± 31.0 [5]	2023(round16)	74031 – 78536
3.773	4191.0 ± 16.0 [5]	2024(round17)	78615 – 81094

Table 2: Processes and luminosity scales of the inclusive MC samples used in this analysis. Separate samples are available for each data-taking round. Most processes use samples with an integrated luminosity equivalent to ten times that of the data.

Process	Luminosity scale
$e^+e^- \rightarrow \psi(3770) \rightarrow D^0\bar{D}^0$	$10 \times$
$e^+e^- \rightarrow \psi(3770) \rightarrow D^+D^-$	$10 \times$
$e^+e^- \rightarrow \tau^+\tau^-$	$10 \times$
$e^+e^- \rightarrow \psi(3770) \rightarrow \text{non-}D\bar{D}$	$10 \times$
$e^+e^- \rightarrow q\bar{q}$	$10 \times$
$e^+e^- \rightarrow \gamma_{\text{ISR}}\psi(2S)$	$10 \times$
$e^+e^- \rightarrow \gamma_{\text{ISR}}J/\psi$	$10 \times$
$e^+e^- \rightarrow e^+e^-$	$0.25 \times$
$e^+e^- \rightarrow \gamma\gamma$	$3 \times$
$e^+e^- \rightarrow \mu^+\mu^-$	$15 \times$

3 Data and MC samples

We use an e^+e^- collision sample corresponding to an integrated luminosity of 20 fb^{-1} , collected by BESIII detector at center-of-mass energy of 3.773 GeV, under BOSS-7.1.2 version. The details are listed in Table 1, quoted from Ref. [4, 5].

Simulated Monte Carlo (MC) samples are employed to study backgrounds, optimize event selection, and estimate detection efficiencies. Two categories of MC samples are utilized in this analysis:

- **The inclusive MC samples** comprehensively model known physics processes at $\sqrt{s} = 3.773 \text{ GeV}$, including $D\bar{D}$ production, $\tau^+\tau^-$ pair production, initial-state-radiation (ISR) production of vector charmonium states (J/ψ and $\psi(2S)$), QED processes, and continuum light hadron production. The charm working group provides separate samples for the different data-taking rounds(round03-04, round15, round16, and round17) [7]. In this analysis, we use inclusive MC samples with an integrated luminosity equivalent to ten times that of the data for most processes, except for the QED processes $e^+e^- \rightarrow e^+e^-$, $\gamma\gamma$, and $\mu^+\mu^-$, where we use the samples with the scaling factors provided by the charm working group. The processes and the corresponding luminosity scales of

the samples used in this analysis are summarized in Table 2.

- **Exclusive signal and background MC samples** generated specifically for this analysis using the PHOKHARA event generator [8, 9, 10] (version 9.1). These samples are produced independently for each data-taking round (round03, round04, round15, round16, round17) to account for variations in detector conditions. Sufficient statistics are generated for each process to ensure reliable efficiency determination and background studies. The generated processes include:

$$- e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma_{\text{ISR}} \text{ (signal process)}$$

$$- e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$$

$$- e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$$

These exclusive samples are used for signal efficiency studies, background characterization, and validation of the analysis methodology.

4 Event selections

The process under investigation is $e^+e^- \rightarrow \pi^+\pi^-\pi^0\gamma_{\text{ISR}}$, whose final states include two charged tracks, one neutral π^0 , and one ISR photon. The following selection criteria is applied to all the data samples. Here the figures are shown by taking $\sqrt{s} = 3.773\text{GeV}$ round16 data as an example. The distributions related to the event selection of other data sets are shown in Appendix. XXX. Situations are similar to each other.

- Good charged track: A charged track is taken as a good MDC track if it originates from the beam vertex and lies within the MDC polar angle region:
 - $|\cos\theta| < 0.93$, where θ is the polar angle with respect to the beam axis.
 - $|\Delta r| < 1\text{ cm}$, $|\Delta z| < 10\text{ cm}$, where $|\Delta r|$ and $|\Delta z|$ is distance of closest approach to the interaction point (IP) perpendicular to the beam axis and along the beam axis, respectively.
 - the ratio of the calorimeter-deposited energy to the track momentum E_{EMC}/p is required to be less than 0.8 for both of the charged tracks.
 - no PID requirements for π^+ , π^- .
- γ selection:
 - $E > 0.025\text{ GeV}$ (barrel: $|\cos\theta| < 0.8$), $E > 0.050\text{ GeV}$ (endcap: $0.86 < |\cos\theta| < 0.92$).
 - $0 \leq t_{\text{TDC}} \leq 14(\times 50\text{ns})$ to suppress fake photons due to electronic noise or beam background.
 - the closest angle with all charged tracks $\theta > 10^\circ$ to suppress background from radiation of charged tracks.
- $\pi^0 \rightarrow \gamma\gamma$ selection:
 - two photon invariant mass $M_{\gamma\gamma}$ within $(0.115, 0.150)\text{ GeV}/c^2$.
 - To improve the momentum resolution, 1C kinematic fit is performed. Here, the 1C means π^0 mass constraint, and the updated 4-momentum will be used in the further analysis. $\chi_{\text{KF}}^2 < 10$

Candidate events are selected by requiring exactly two good charged tracks, accompanied by at least one reconstructed π^0 and a minimum of three good photon showers. This ensures the presence of at least one π^0 and one isolated photon from Initial-State Radiation (γ_{ISR}). A four-constraint (4C) kinematic fit is then applied to the event, and the candidate with the smallest $\chi_{4\text{C}}^2$ value is chosen for further analysis.

After identifying an ISR photon from the isolated photon showers, the invariant mass of the remaining $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ system is reconstructed. As expected for ISR processes, the resulting center-of-mass energy $\sqrt{s'}$ of the $\pi^+\pi^-\pi^0$ system shows a continuous spectrum, characteristic of the radiative return mechanism. This continuum behavior is illustrated in Fig. 1 across the $\omega(782)$, $\phi(1020)$, and ω'/ω'' mass regions.




Figure 1: Invariant mass spectra of MC in the (a) $\omega(782)$, (b) $\phi(1020)$, and (c) ω'/ω'' mass regions. The distributions correspond to $M(pK_S^0)$, $M(p\pi^0)$, and $M(K_S^0\pi^0)$, respectively. The "Truth-Matched" distribution (green dashed line) corresponds to reconstructed events where the momentum directions of the π^0 and γ_{ISR} successfully match their truth-level counterparts. A loose requirement of $\chi_{\text{KF}}^2 < 100$ is applied to the kinematic fit. The continuous spectrum of $\sqrt{s'}$ for the $\pi^+\pi^-\pi^0$ system demonstrates the radiative return mechanism via ISR.

5 Background Study

In the study of background processes, we categorize the main backgrounds into $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$, $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$, QED processes, and others. The primary backgrounds originate from processes with additional π^0 s, such as $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ and $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$. In the inclusive analysis, the cross sections for these processes were not correctly configured. The $\mathcal{L} \times \sigma_{\text{hadron}}$ method (where $\mathcal{L}$ is the luminosity and σ_{hadron} is the hadronic cross section) was employed to verify that these processes constitute the dominant backgrounds. (cross section calculation fill here). Precise yield estimates for these two components will subsequently be obtained using data-driven methods.

Based on the rough background estimate from $\mathcal{L} \times \sigma_{\text{hadron}}$, the signal yield was approximated by subtracting the background from the data, allowing for a preliminary significance calculation. Using this approach, a figure of merit (FOM) defined as $S/\sqrt{S+B}$ was constructed to evaluate the signal significance. To optimize the event selection, a scan over the kinematic fit χ^2 was performed, with the result shown in Fig. 2. The Round 16 optimization yields an optimal cut of $\chi^2_{\text{KF}} < 24$.




Figure 2: (a) Distribution of the kinematic fit χ^2 for the four-constraint (4C) fit. (b) The corresponding figure of merit ($S/\sqrt{S+B}$) as a function of the applied χ^2 cut value. The vertical dashed line indicates the chosen optimal cut of $\chi^2_{\text{KF}} < 24$ from Round 16 optimization.

The 3π invariant mass distribution obtained with the $\chi^2_{\text{KF}} < 24$ selection is presented in Fig. 3. The plot displays the data in comparison with the stacked Monte Carlo predictions for the signal and background components.

5.1 estimate of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

In estimating the $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ background, the following strategy was adopted:

It is assumed that the total yield of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ events produced in the data is a fixed quantity. This process enters our signal region primarily when a photon from one of the π^0 decays is misidentified as an initial-state radiation photon (γ_{ISR}). Consequently, two distinct selection efficiencies can be calculated: ϵ_{sig} (the efficiency for $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ events to be selected as signal candidates) and ϵ_{bkg} (the efficiency for the same process to enter the signal region as background).




Figure 3: The 3π invariant mass distribution after applying the $\chi^2_{\text{KF}} < 24$ selection. The stacked histogram shows the contributions from the signal process $e^+e^- \rightarrow \pi^+\pi^-\pi^0$ and various background processes, including the dominant $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ and $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$ backgrounds.

When $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ events are selected as signal, the same analysis procedure as for the signal is applied to construct the 3π invariant mass spectrum, thereby obtaining the distribution of this process in data. Using the ratio of these two efficiencies, $R = \epsilon_{\text{bkg}}/\epsilon_{\text{sig}}$, the observed $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ signal distribution can be converted into an estimate of its background distribution in the signal region. Specifically:

$$N_{\text{bkg}} = R \times N_{\text{sig}}^{\text{obs}} = \frac{\epsilon_{\text{bkg}}}{\epsilon_{\text{sig}}} \times N_{\text{sig}}^{\text{obs}},$$

where $N_{\text{sig}}^{\text{obs}}$ is the number of observed $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ signal events in data, and N_{bkg} is the estimated number of background events from this process in the signal region.

This method provides a data-driven conversion from the observed signal distribution to the background distribution through efficiency correction.

When treating $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ as the signal, the selection criteria are designed such that the same event sample also satisfies a loose selection for $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0$. This ensures that the invariant mass of the three-pion system is automatically defined within the latter process.

The combined selection proceeds as follows: events are required to meet the criteria for both processes. The first step applies the full $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ selection:

• **$\pi^+\pi^-\pi^0\pi^0$ Candidate Selection:**

- Exactly two good charged tracks.
- $n_{\text{Pi0}} \geq 2$ & $n_{\text{GoodShowers}} \geq 4$ (ensuring at least two π^0 s).
- Application of a total 4-momentum constraint to the initial e^+e^- collision (4C kinematic fit).
- The candidate with the minimum $\chi^2_{4\text{C}}$ is chosen.
- A requirement of $\chi^2_{\text{KF}} < 10$ on the kinematic fit.

Subsequently, within the same event, a candidate for $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ is formed by reinterpreting one of the π^0 s as the γ_{ISR} :

• $\gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ **Candidate Selection:**

- The same two good charged tracks.
- $n\text{Pi0} \geq 1$ & $n\text{GoodShowers} \geq 3$ (ensuring at least one π^0 and one photon assigned as γ_{ISR}).
- Application of the same total 4-momentum constraint to the initial e^+e^- collision (4C kinematic fit).
- The candidate with the minimum χ^2_{4C} is chosen.
- No additional kinematic fit χ^2 requirement is applied at this stage.

This dual-tag strategy guarantees that the reconstructed 3π system from the $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ hypothesis is kinematically well-defined, and its invariant mass can be reliably calculated for the subsequent analysis.

The selection described above results in a highly pure sample of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ events. The invariant mass spectrum of the corresponding three-pion system from these events, which directly enters the background estimation, is shown in Fig. 4. This observed distribution in data will be scaled by the efficiency ratio $R = \epsilon_{\text{bkg}}/\epsilon_{\text{sig}}$ to estimate the final background contribution in the signal region. All other backgrounds will be subtracted directly.




Figure 4: The 3π invariant mass spectrum from selected $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ events (red points), overlaid with the scaled Monte Carlo simulation (blue histogram). The distribution is obtained after applying the dual-tag selection criteria and represents the pure $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$ component in data.

5.2 estimate of $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0\gamma_{\text{ISR}}$

The second dominant background, $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0\pi^0$, is estimated using an analogous data-driven approach. The selection follows a similar dual-tag strategy to obtain a sample. The selection

criteria are outlined below:

Simultaneously, events must satisfy both the $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0\pi^0$ selection, where the γ_{ISR} is treated as an untagged missing momentum, and the tagged $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ loose selection criteria.

• $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0\pi^0$ **Candidate Selection (Untagged γ_{ISR}):**

- Exactly two good charged tracks.
- $n\text{Pi0} \geq 2$ & $n\text{GoodShowers} \geq 4$ (requiring at least two π^0 s).
- The γ_{ISR} is not explicitly reconstructed from a shower. Instead, its 4-momentum is inferred as a *missing* component within the kinematic fit.
- A total 4-momentum constraint to the initial e^+e^- collision is applied (4C fit).
- The candidate with the minimum $\chi^2_{4\text{C}}$ is chosen.
- A requirement of $|\cos \theta_{\text{Miss}}| > 0.99$ is applied to the direction of the missing 4-momentum to ensure it is consistent with an ISR photon along the beam axis.
- A kinematic fit quality cut of $\chi^2_{\text{KF}} < 10$ is applied.

• $\gamma_{\text{ISR}}\pi^+\pi^-\pi^0$ **Candidate Selection (within the same event):**

- The same two good charged tracks.
- $n\text{Pi0} \geq 1$ & $n\text{GoodShowers} \geq 3$ (ensuring at least one π^0 and one photon tagged as γ_{ISR}).
- A total 4-momentum constraint to the initial e^+e^- collision is applied (4C fit).
- The candidate with the minimum $\chi^2_{4\text{C}}$ is chosen.
- No additional kinematic fit χ^2 requirement is applied.




Figure 5: The 3π invariant mass spectrum from selected $e^+e^- \rightarrow \gamma_{\text{ISR}}\pi^+\pi^-\pi^0\pi^0$ events (green points). The distribution is obtained after applying the dual-tag selection and serves as a clean template for background estimation.

195 The application of this dual-tag selection strategy resulted in a sample with minimal contamina-
196 tion from other processes. The high purity observed is a consequence of the stringent criteria. The
197 corresponding 3π invariant mass distribution from this sample, which will be used for the background
198 estimation, is shown in Fig. 5.

199 The observed distribution will be similarly scaled by the appropriate efficiency ratio $R' = \epsilon'_{\text{bkg}}/\epsilon'_{\text{sig}}$
200 to estimate its contribution in the signal region.

201 **6 Unfolding**

202 to fill

203 **7 Efficiency correction**

204 to fill

205 **8 Systematic uncertainties**

206 to fill

207 **9 Summary**

208 to fill

References

- [1] M. Ablikim *et al.* (BESIII Collaboration), *Future Physics Programme of BESIII*, *Chinese Phys. C* **44** (2020).
- [2] R. X. Yang *et al.*, *MRPC detector for the BESIII E-TOF upgrade*, *Journal of Instrumentation*, **Volume 9**, September 2014. DOI: [10.1088/1748-0221/9/09/C09032](https://doi.org/10.1088/1748-0221/9/09/C09032).
- [3] M. Ablikim *et al.* [BESIII], DOI: [10.1016/j.nima.2009.12.050](https://doi.org/10.1016/j.nima.2009.12.050).
- [4] M. Ablikim *et al.* [BESIII], DOI: [10.1016/j.physletb.2015.11.043](https://doi.org/10.1016/j.physletb.2015.11.043).
- [5] M. Ablikim *et al.* [BESIII Collaboration], DOI: [10.1088/1674-1137/ad70a0](https://doi.org/10.1088/1674-1137/ad70a0).
- [6] S. Agostinelli *et al.* [GEANT4], *Nucl. Instrum. Meth. A* **506**, 250-303 (2003).
- [7] BESIII Collaboration, DocDB: 1483.
- [8] H. Czyż, A. Grzełińska, J. H. Kühn, and G. Rodrigo, DOI: [10.1140/epjc/s2006-02614-7](https://doi.org/10.1140/epjc/s2006-02614-7).
- [9] G. Rodrigo, A. Gehrmann-De Ridder, M. Guillaume, and J. H. Kühn, DOI: [10.1007/s100520100784](https://doi.org/10.1007/s100520100784).
- [10] G. Rodrigo, H. Czyż, J. H. Kühn, and M. Szopa, DOI: [10.1007/s100520200912](https://doi.org/10.1007/s100520200912).

223 **APPENDIX**