

# Lecture 2

Xiaoqian Liu

## 1 Overview

In this lecture, we introduce the concept of convexity, including convex sets and some convexity-preserving operations.

## 2 Convex sets

A subset  $C$  of  $\mathcal{H}$  is **convex** if

- $\forall \alpha \in (0, 1), \alpha C + (1 - \alpha)C \subset C$ .
- That is,  $\forall \alpha \in (0, 1), \forall x \in C, \forall y \in C, \alpha x + (1 - \alpha)y \in C$ .
- Or equivalently,  $\forall x \in C, \forall y \in C, (x, y) \subset C$ , where  $(x, y)$  denotes the segment between  $x$  and  $y$ .

Examples/counterexamples:

- A ball  $B(x; \rho)$  is convex.
- Half space and affine space are convex.
- The union of two disjoint balls are not convex.

## 3 Convexity-preserving operations

- Let  $(C_i)_{i \in I}$  be a family of convex subsets of  $\mathcal{H}$ , then  $\cap_{i \in I} C_i$  is convex.
  - union, complement are not in general
- Let  $(C_i)_{i \in I}$  be a finite family of convex subsets of  $\mathcal{H}$ , and let  $(\alpha_i)_{i \in I} \in \mathbb{R}$ , then  $\sum_{i \in I} \alpha_i C_i$  is convex.
- Let  $\mathcal{G}$  be a Euclidean space, and  $L : \mathcal{H} \rightarrow \mathcal{G}$  be a linear operator. Let  $C \in \mathcal{H}$  and  $D \in \mathcal{G}$  be convex subsets. Then,

- $L(C) = \{Lx \mid x \in C\}$  is convex
- $L^{-1}(D) = \{x \in \mathcal{H} \mid Lx \in D\}$  is convex