
MI window (1.5 s)
reference window (main)
alternative reference / control window
A Acute (Liu 2024): n = 50
cue
MI
break
MI / reference windows
alternative / control
0
1
2
3
4
5
6
7
8
screen: cue: visual cue for left/right hand (2 s) · MI: 4 s · break: 2 s (screen content as in the dataset description)
main reference = cue period 0.25 s; alternatives = break 6.25 / 6.5 s
B Chronic (Lu 2026): n = 14
prep
MI
feedback
MI / reference windows
alternative / control
0
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
screen: prep: relax; tactile brush input ends at 7.5 s (onset not stated) · MI cue at 9 s · feedback 13–17 s
main reference = preparation 0.5 s; split-half candidates 3.0 / 6.0 s
C Lower-limb (Liu 2025): n = 27 released, 25 analysed at the first assessment
fixation
text
task
break
MI / reference windows
alternative / control
0
1
2
3
4
5
6
7
8
9
10
11
12
screen: task: MI trial shows gait-phase digit in orange circle; idle trial shows black circle
reference = same windows of idle trials; control = same-trial fixation 0.5 s
D Paretic-hand / sixth-finger (Wang 2026): n = 24 released, 22 analysed at the first assessment
rest
prep circle
MI
MI / reference windows
alternative / control
0
1
2
3
4
5
6
7
8
9
10
11
12
13
14
screen: rest: green fixation cross · prep: grey circle · MI: picture of hand / sixth finger
main reference = rest 4.0 + 5.5 s; sensitivity = 1.0 + 4.0 s
time within trial (s)