

# Predicting Covid-19 EMS Incidents from Daily Hospitalization Trends

Ngoc Mai Tran

*Department of Mathematics, University of Texas at Austin, TX 78712, USA.*

Evdokia Nikolova

*Department of Electrical and Computer Engineering, University of Texas at Austin, TX 78712, USA.*

David Kulpanowski

*Department of Emergency Medical Services, City of Austin*

Yangxinyu Xie

*Department of Mathematics, University of Texas at Austin, TX 78712, USA.*

Joshua Ong

*Department of Electrical and Computer Engineering, University of Texas at Austin, TX 78712, USA.*

---

## Abstract

*Introduction:* The aim of our retrospective study was to quantify the impact of Covid-19 on the temporal distribution of Emergency Medical Services (EMS) demand in Travis County, Austin, Texas and propose a robust model to forecast Covid-19 EMS incidents.

*Methods:* We analyzed the temporal distribution of EMS calls in the Austin-Travis County area between January 1st, 2019 and December 31st, 2020. Change point detection was performed to identify critical dates marking changes in EMS call distributions and time series regression was applied for forecasting Covid-19 EMS incidents.

*Results:* Two critical dates marked the impact of Covid-19 on the distribution of EMS calls: March 17th, when the daily number of non-pandemic EMS incidents dropped significantly, and May 13th, by which the daily number of EMS calls climbed back to 75% of the number in pre-Covid-19 time. New daily count of the hospitalization of Covid-19 patients alone proves a powerful predictor of the number of pandemic EMS calls, with an  $r^2$  value equal to 0.85. In particular, for every 2.5 cases where EMS takes a Covid-19 patient to a hospital, 1 person is admitted.

*Conclusion:* The mean daily number of non-pandemic EMS demand was significantly less than the period prior to Covid-19 pandemic. The number of EMS calls for Covid-19 symptoms can be predicted from the daily new hospitalization of Covid-19 patients. These findings may be of interest to EMS departments as they plan for future pandemics, including the ability to predict pandemic-related calls in an effort to adjust a targeted response.

*Keywords:* Emergency Medical Services, Pandemics, Covid-19


---

*Email addresses:* ntran@math.utexas.edu (Ngoc Mai Tran), nikolova@austin.utexas.edu (Evdokia Nikolova), David.kulpanowski@AustinTexas.gov (David Kulpanowski), yx4247@utexas.edu (Yangxinyu Xie), joshong@utexas.edu (Joshua Ong)

## 1. Introduction

The ongoing outbreak of the coronavirus disease 2019 (Covid-19) has caused overwhelming disruptions to healthcare systems around the globe (1, 2, 3, 4), especially the emergency medical services (EMS). Researchers have conducted extensive studies on different aspects of the EMS since the outbreak of Covid-19, including the temporal distribution of EMS demand and emergency department visits (5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17), pre-hospital patient assessment (18, 19), medical resource availability and allocation (20, 21, 22, 23), personnel protective equipment (24, 25, 26), EMS response practices and strategies (27, 28, 29, 14) and ethical considerations (30, 31).

Figure 1: Spatial Distribution of EMS Incidents in 2019-2020<sup>1</sup>. Each light pink dot represents an incident during the 2019-2020 period. The total number of EMS calls in was 246,809. Noticeably, EMS incidents were prevalent along I-35 highway in Austin and were most frequent at downtown Austin.



Globally, the EMS utilization rates varied during the early stage of the Covid-19 outbreak. The number of EMS calls increased in some parts of Europe (France (32), Ankara, Turkey (12), Copenhagen, Denmark (14)) and Saudi Arabia (11). However, other observational studies suggested a significant decrease in the number of EMS calls across the United States (5, 7, 8, 33), Canada (34), and other parts of Europe (Italy (9, 35), England (10), Finland (36)). Nonetheless, there is very limited information available after the beginning of summer 2020, when states began to lift travel restrictions and people became better informed of the nature of the virus.

There is a long history of forecasting that has explored many factors to model EMS demand. The study of forecasting models dates back to the 1970s (37, 38). Earlier studies in cities across the United States

<sup>1</sup>This plot is extracted from the City of Austin Open Data Portal: <https://data.austintexas.gov/dataset/City-of-Austin-Incidents-Map/m4s3-4qdj>

incorporated socio-demographic and socioeconomic variables to forecast daily demand (38, 39, 40, 41). Other studies also found that weather factors and seasonality (season, day of the week, and holidays) could also be used to predict the daily ambulance demand (42, 43, 44, 45).

Nevertheless, these EMS models were developed under normal conditions. It is unclear whether they are able to adapt to large-scale disasters such as the Covid-19 pandemic. Studies in the 2016 Melbourne thunderstorm asthma epidemic (46) discovered a positive correlation between thunderstorm asthma cases and increased EMS demand. To our best knowledge, no models specifically targeting Covid-19 related EMS demand have been proposed. However, earlier studies in France (47, 48) and Israel (49) hinted correlations between Covid-19 case hospitalization and Covid-19 EMS demand.


The objective of this study is twofold. First, we seek to examine the long-term impact of Covid-19 on the temporal distribution of emergency calls and the average response time of ambulance assignment, dispatch and arrival. Second, we seek to design a predictive model to forecast the daily number of Covid-19 pandemic EMS calls.

## 2. Methods

### 2.1. Data Description

Our retrospective study is based on two datasets from the City of Austin Open Data Portal. The first dataset contains all records of EMS incidents in Austin Travis County from January 1st, 2019 to December 31st, 2020 (50). These EMS incident records do not contain any identifiable information. For each incident, this dataset included its problem type, call disposition, and the date of the incident. Additionally, it included EMS response times including the time elapsed, in minutes, to assign the call, dispatch the ambulance, and for the ambulance to arrive. For a comprehensive list of descriptions of the dataset, see Table 1.

Figure 2: 6 Major Hospitals in Austin<sup>2</sup>. The center of each circle represents the geolocation of each hospital. The size of the circle is proportional to the percentage of EMS incidents disposed to the hospital (see Figure 3). The two biggest hospitals are Dell Seton Medical Center and South Austin Hospital.



<sup>2</sup>This plot is extracted from the City of Austin Open Data Portal: <https://data.austintexas.gov/dataset/City-of-Austin-Hospitals-Map/df29-4hn7>

Raw names	Names	Description
Problem	Problem Type	The problem type of the emergency.
Time_PhonePickUp_Date	Date	The date of the EMS call.
Time_First_Unit_Assigned	Assignment Time	The length of time, in minutes, after the EMS call was picked up and before the first ambulance received notification of the emergency and was assigned.
Time_First_Unit_Enroute	Dispatch Time	The length of time, in minutes, after the first ambulance was assigned and before the ambulance set off toward the emergency.
Time_First_Unit_Arrived	Arrival Time	The length of time, in minutes, after the first ambulance wheels began to roll and before the ambulance arrived at the emergency and the wheels stopped.
Call_Disposition	Call Disposition	The final disposition of the event, such as cancelled call, transported to hospital, etc. If the ambulance transported the patient to a hospital, the name of the hospital would be specified. If the emergency call was from another government agency, for example, the Austin Fire Department, then the call disposition would be labelled as "referred". Other types of call dispositions fall into the category of "defunct calls", whose subcategories are defined in Table 3; in this case, the call disposition of an incident would be labelled as its respective subcategory.

Table 1: ATCEMS database: incident records

The second dataset consists of the daily frequencies of all incidents from January 1st, 2019 to December 31st, 2020 (51). This aggregated dataset distinguishes Covid-19 pandemic incidents from incidents of other problem types, as well as the defunct incidents from incidents of other types of call dispositions. Defunct calls are defined as the EMS incidents where neither the call was from another government agency, nor did the ambulance transport the patient to a hospital. For a complete list of descriptions of the subcategories of defunct calls, see Table 3. The Covid-19 hospitalization data consists of the daily count of the hospital admission of Covid-19 patients in the Austin-Round Rock metropolitan statistical area. This hospitalization data was requested from the University of Texas Covid-19 Modeling Consortium (52). For a comprehensive list of descriptions of each series of daily frequencies in the dataset, see Table 2.

## 2.2. Study Setting and Background

The total number of EMS calls during the 2019 to 2020 period was 246,809. On average, the 37 ambulances in Austin Travis County responded to  $338 \pm 34$  calls per day. The spatial distribution of EMS incidents is shown in Figure 1. Noticeably, EMS incidents were prevalent along I-35 highway in Austin and were most frequent in downtown Austin. There are 6 major hospitals in Travis County (for the locations of these hospitals, see Figure 2; for the percentages of EMS incidents classified by call dispositions, see Figure 3), among which Dell Seton Medical Center and South Austin Hospital are the two biggest ones, covering 26% of all EMS incidents. In summary, the 6 major hospitals took care of 47% of all EMS incidents. Meanwhile, refusal (20%), cancellation (7%), missing patient (5%) are also among the highest percentages.

There were 22 types of problems with the highest number of calls, which in sum comprise 88% of the total number of incidents. For a complete list of problem types and their frequencies in the period of 2019 to 2020, see Figure 4. Notably, there were 12,926 emergency calls related to Covid-19 pandemic.

The distribution of EMS response times in 2019-2020 is shown in Figure 5. On average, after an EMS call was picked up, the mean time for an ambulance assignment was 1.21 minutes and the median was 1.03

Raw names	Names	Description
total_ts	Total	The daily frequency of all EMS incidents.
defunct_calls_pandemic_removed.ts	Non-pandemic Defunct	The daily frequency of non-pandemic defunct EMS incidents.
Pandemic_defunct_calls_ts	Pandemic Defunct	The daily frequency of pandemic defunct EMS incidents.
Incidents_pandemic_removed.ts	Non-pandemic Effect	The daily frequency of non-pandemic, non-defunct EMS incidents.
Pandemic_effect_ts	Pandemic Effect	The daily frequency of pandemic, non-defunct EMS incidents.
hospitalisation_ts	Hospitalization	The daily frequency of the newly admitted Covid-19 patients to hospitals in the Austin-Round Rock metropolitan statistical area.

Table 2: ATCEMS database: incident daily frequencies

Call Disposition	Description
Refusal	A refusal means the situation when the ambulance arrives on scene and speaks with the patient, the patient refuses medical help.
No Patient	The ambulance responds to the address provided but does not find a patient.
Other	Call disposition is unspecified.
Call Cancelled	
Duplicate Call	
False Alarm Call	
Information Call Only	

Table 3: The subcategories of defunct calls

minutes. The mean and median time for an assigned ambulance to dispatch were both 1.05 minutes. The average time it took for a dispatched ambulance to arrive on the scene was 7.14 minutes and the median was 6.1 minutes.<sup>3</sup>

### 2.3. Statistical Analysis


The statistical analysis was performed using R statistical computing and graphics environment. The R code is available on GitHub.<sup>4</sup> To evaluate the impact of Covid-19 on non-pandemic incidents, we applied change point detection from the R software package "changepoint.np". We identified changes in mean and variance (the function "cpt.meanvar") with binary segmentation method (53) and Bayesian information criterion (BIC) penalty assuming underlying normal distribution. To restrict our attention to the impact of the pandemic only, we chose the maximal number of change points as 2.

After identifying the change point dates, Student's t-tests were applied to make various comparisons between the first period (pre-pandemic) and the third period (post-pandemic). To identify which of the 22 major problems (see figure 4 for a complete list) were impacted by the outbreak of Covid-19 pandemic, we use Student's t-test with Bonferroni correction to compare the mean number of daily calls for each

<sup>3</sup>The average response time varies among the 6 major hospitals. For a detailed comparison, see the appendix section.

<sup>4</sup><https://github.com/Xieyangxinyu/AustinEMS2020>

Figure 3: Call Disposition of EMS Incidents in 2019-2020. The percentage was calculated as the number of incidents with a type of disposition against the total number of incidents, which was 246,809. The 6 major hospitals in Travis County are Dell Seton Medical Center, South Austin Hospital, Saint Davids Medical Center, North Austin Hospital, Seton Medical Center and Seton Northwest Hospital. Noticeably, some types of defunct calls had high percentages, including refusal (20%), cancellation (7%) and missing patient (5%).




problem type between the first period and the third period. In other words, we set the significance cut-off at  $0.05/22 = 0.002273$ . Another comparison of the mean time of EMS responses (assignment, dispatch and arrival) was made using one-sided Student's t-test.

To predict the daily frequency of pandemic EMS incidents, time series regression with change point detection was applied in the modeling process. The regressor is the daily frequency of the Covid-19 hospitalization (the "Hospitalization" time series in Table 2). First, we identified multiple change points in the daily Covid-19 hospitalization data. Specifically, we applied exact change point detection, the pruned exact linear time (PELT, (54)) method, on variance with modified Bayesian information criterion (MBIC) penalty assuming underlying normal distribution. Second, to reduce noise and thus improve the predictive power of our model, we smoothed the time series of daily Covid-19 hospitalization by an average of a period of 7 days. Smoothing the daily hospitalization is reasonable because in practical settings epidemiological models commonly output hospitalization forecasts as "expected values", which would then be fed into our model as inputs. Next, we parsed the daily hospitalization data into different periods in accordance with detected change point dates by adding dummy variables. Lastly, after splitting the training (80%) and testing set (20%), we applied exhaustive (non-stepwise) selection of autoregressive integrated moving average (ARIMA) models and binomial thinning (55). The function used was "auto.arima" from the R software package "forecast". To evaluate the robustness of our model, we estimated the  $r^2$  value using smoothed daily pandemic EMS incident data to remove unnecessary random noise. To evaluate the predictive power of our model, we estimated the mean squared error and standard error of prediction residual using the original daily pandemic EMS incident data.

### 3. Results

This section consists of two major parts. The first subsection identifies two critical dates, March 17th and May 13th, indicating the change in mean and variance of the temporal distribution of non-pandemic EMS calls. (Unless otherwise specified, in the result section, we use non-pandemic EMS calls to refer to non-pandemic effect EMS calls and pandemic EMS calls to refer to pandemic effect EMS calls.) To further investigate the lasting impact of the outbreak of Covid-19, we separated the timeline into three periods:

Figure 4: Major Problem of EMS Incidents in 2019-2020. There were 22 types of problems with the highest number of calls, which in sum comprise 88% of the total number of incidents. The frequency of each major problem was above 3000 in the period of 2019-2020. The red color highlights the Covid-19 pandemic. The purple color signifies the problem types with a significant drop in period 3 in comparison to period 1, and the gray color matches with the problem types without a significant drop (see Table 5 in the result section.)



period 1, from January 1st, 2019 to March 17th, 2020 with  $n = 442$ , represents the pre-pandemic norm; period 2, from March 18th to May 12th in 2020, represents the sudden outbreak of Covid-19 in Austin; and period 3, after May 13th in 2020, represents the new normal. Comparing period 1 and period 3, we found that the average daily number of non-pandemic EMS incidents dropped and the average EMS response time became slower.

In the second section, we propose a time series regression model predicting the daily frequency of pandemic EMS incidents. The regressor, the daily frequency of the Covid-19 hospitalization, is parsed into 4 stages since March 2020. The regression model obtained an  $r^2$  value equal to 0.85 and indicated that for every 2.5 cases where EMS took a Covid-19 patient to a hospital, 1 person was admitted.

### 3.1. Impact of the pandemic


Via change point detection in mean and variance of the temporal distribution of non-pandemic EMS calls, with binary segmentation method and BIC penalty, we found two critical dates marking the impact of Covid-19 on non-pandemic EMS calls (shown as purple vertical lines in Figure 6). The first date was March 17th, around the time of the Austin shelter-in-place order. The average daily number of non-pandemic EMS incidents dropped significantly, from 225.69 in period 1 to 155.84 in period 2.

The second critical date was May 13th, or the beginning of summer, by which the daily number of EMS calls climbed back to a new plateau, which is about 75% of that before March 17th (169.53 in period 3 versus 225.69 in period 1). In sum, the total number of non-pandemic EMS calls decreased during Covid-19 times.

Noticeably, despite the disruption of Covid-19, non-pandemic defunct calls did not witness the same level of decrease in daily frequency (shown by the orange line in Figure 6). Nevertheless, the proportion of defunct calls rose from 35% of the overall number of non-pandemic EMS calls in period 1 to 39% in period 3. For a complete comparison of the number of non-pandemic EMS incidents per day among periods 1, 2, and 3, see Table 4.

As for pandemic-related incidents, the red line represents pandemic EMS calls disposed to hospitals, and the brown line represents pandemic related defunct calls in Figure 6. Note that pandemic EMS calls are closely correlated with the daily frequency of the newly admitted Covid-19 patients to hospitals in the

Figure 5: Performance of EMS Responses in 2019-2020



Austin-Round Rock metropolitan statistical area, shown by the green line. Our time-series model in the next subsection found that this hospitalization trend alone is a powerful predictor for pandemic EMS calls.

A closer comparison between period 1 and period 3 showed that the frequencies of only certain types of problems dropped, while others remained unaffected by Covid-19. The complete list of comparisons of the number of non-pandemic EMS incidents of each problem type can be found in Table 5.

The performance of EMS response was also impacted since the outbreak of Covid-19. The mean response time for each action, assignment, dispatch and arrival, was slower during period 2 and period 3 than that in period 1 (one-sided t-test,  $p$ -value  $< 0.01$ ). In particular, the arrival time during period 3 is 0.73 minutes slower than during period 1 for each EMS incident. A complete list of comparisons of the mean and median of response time is in Table 6. The reader can also see from Figure 7 that the mean (solid line) and median (dashed line) of the distributions of the response time has slightly shifted to the right from period 1 to period 3.


### 3.2. Covid-19 hospitalization predicts pandemic EMS calls

The exact change point detection on variance with MBIC penalty yielded 4 Covid-19 hospitalization stages since March 2020. The four change point dates are shown by the purple vertical lines in Figure 8. In particular, we see rises in Covid-19 hospitalizations during summer 2020 (June 8th - August 18th) and winter 2020 (November 5th - December 31st). This resonated with the global pandemic second and third waves. To achieve optimal modeling performance, we avoided the uncertainties at the earliest stage of the pandemic by only fitting and testing the model using data from April 9th to December 31st in 2020. Thus, we set the dummy variable "1st\_period" to 1 during the period from April 9th - June 8th, the dummy variable "2nd\_period" to 1 during the period from June 8th - August 17th and the dummy variable "3rd\_period" to 1 during the period from August 17th - December 31st.

Comparison of number of non-pandemic EMS incidents							
	period 1		period 2		period 3		
	n = 442		n = 56		n = 233		
	Mean	SD	Mean	SD	Mean	SD	
Admitted	225.69	19.43	155.84	20.23	169.53	14.76	
Defunct	122.35	21.71	113.59	19.30	109.34	12.73	

Table 4: Comparison of number of non-pandemic EMS incidents per day among period 1 (before March 17th, pre-pandemic), period 2 (March 18th - May 12th), period 3 (after May 13th, post-pandemic). SD stands for standard deviation. Overall, the total number of non-pandemic EMS calls decreased during Covid-19 times.

Figure 6: Daily Frequency of EMS Incidents in 2019-2020. The two purple lines mark the two critical change point dates, March 17th and May 13th. Each colored line indicates the temporal distribution of a time series defined in Table 2. The darker color represents the smoothed, 7-day average, trend, while the lighter color represents the raw frequencies. In summary, the total number of non-pandemic EMS calls decreased during Covid-19 times.



The exhaustive selection of ARIMA models output 0 autoregressive order, 0 degree of differencing, and 0 moving average order. The coefficients of the linear regression model, with the pandemic EMS Calls regressed on the smoothed (7-day average) hospitalization data, is specified in Table 7. This model obtained an  $r^2$  value equal to 0.85. Moreover, the residual standard error of the training and test set are 6.62 and 5.44, respectively, which mainly characterize for the random fluctuations on daily counts. The reader can also see from Figure 9 that most of the actual daily frequencies of pandemic EMS incidents fall within the 95% prediction interval. Therefore, daily new hospitalization of Covid-19 cases alone proves a powerful and robust indicator of pandemic-related EMS demand. The 0.40 estimate of slope suggests that on average, for every 2.5 cases where EMS took a Covid-19 patient to a hospital, 1 person was admitted. Moreover, moving from the 1st period to 3rd period, there was a decreasing offset between pandemic EMS incidents and hospitalization numbers.

#### 4. Discussion

The daily number of non-pandemic EMS incidents dropped precipitously since the beginning of Covid-19 pandemic and the declaration of the local state of emergency. This resonates with earlier studies in the United States (5, 6, 8, 33), Italy (9, 35), England (10), Canada (34) and Finland (36). Even though the number of non-pandemic EMS incidents increased at the beginning of summer, it remained consistently

Comparison of number of non-pandemic EMS incidents by problem type			
	period 1 <i>n</i> = 442	period 3 <i>n</i> = 233	
	Mean	Mean	p-value
Cardiac Arrest	4.03	4.54	0.999
Psychiatric	16.52	17.49	0.996
Assault	17.86	18.45	0.904
Stroke	6.51	6.55	0.561
Abdominal Pain	7.65	7.51	0.271
Alarm Activation	8.52	8.23	0.128
Injury	10.67	10.24	0.061
Overdose	12.16	11.42	0.008
Diabetic	4.91	4.45	0.007
Altered Mentation	20.90	14.15	< 0.0001
Attended Patient	10.43	6.63	< 0.0001
Chest Pain	19.17	11.10	< 0.0001
Community Health Assist	11.56	8.06	< 0.0001
Fall	30.47	27.27	< 0.0001
Hemorrhage	8.51	6.48	< 0.0001
Respiratory	24.83	16.19	< 0.0001
Seizure	15.18	12.91	< 0.0001
Sick	31.78	14.79	< 0.0001
Syncopal Episode	8.66	6.62	< 0.0001
Traffic Injury	27.39	21.37	< 0.0001
Unconscious	9.46	7.68	< 0.0001

Table 5: One-sided (greater) student-t comparison of number of non-pandemic EMS incidents per day among period 1 (before March 17th), period 2 (March 18th - May 12th), period 3 (after May 13th); via Bonferroni correction,  $\alpha = 0.00227$

lower than pre-pandemic era. More interestingly, the proportion of non-pandemic defunct calls rose after the outbreak of Covid-19 pandemic. The rise of refusals to transportation to hospitals has also been observed in Israel (56). This corroborates with earlier studies in the United States suggesting that people showed reluctance to receive hospitalization out of fear of contagion (13, 57).

This significant decrease in non-pandemic EMS incidents may also be partially explained by travel restrictions (attended patient, community health assist, fall, hemorrhage, traffic injury) and misclassification into Covid-19 cases (chest pain, respiratory, and sick). An overall decrease in traffic-related incidents were also observed in Israel (56) and Ontario, Canada (34) during period 2. People may have been traveling less frequently due to the shelter-in-place order issued on March, 2020, and possibly were less exposed to injury-prone locations. A study in Connecticut found that the mean daily vehicle miles traveled and the mean daily counts of crashes significantly decreased in the post stay-at-home period in 2020 (58). Our comparison between period 1 and 3 in Table 5 showed that the frequencies of traffic related incidents remained lower in the long run than those of the pre-pandemic time, despite the lifting of Texas stay-at-home order.


The decrease in emergency calls regarding chest pain, sickness and respiratory problems were also found in Netherlands (59) and Ontario, Canada (34). One possible explanation is that calls previously classified as "chest pain", "respiratory" and "sick", due to their similarity to Covid-19 symptoms, became categorized as "pandemic" (34).

Prior studies have found increase in EMS calls regarding cardiac arrest (56, 34, 60), stroke (34) mental health (56) and overdose (34), during the early stages of the Covid-19 pandemic. Our study confirmed that, in the long term, the EMS demand for these problems did not decrease. Interestingly, although earlier finding in Ontario, Canada (61) showed a decrease in emergency calls related to assault during the period

	Assignment Time		Dispatch Time		Arrival Time	
	Mean	Median	Mean	Median	Mean	Median
period 1	1.18	1.02	1.01	1.00	6.88	5.87
period 2	1.33	1.12	1.08	1.10	7.52	6.40
period 3	1.25	1.05	1.11	1.10	7.61	6.55

Table 6: Comparison of the average EMS response time (in minutes) among period 1 (before March 17th), period 2 (March 18th - May 12th), period 3 (after May 13th). Assignment time, dispatch time and arrival time were all slower during period 2 and period 3 than those in period 1.

Figure 7: Distribution of EMS response times (in minutes) among period 1 (before March 17th), period 2 (March 18th - May 12th), period 3 (after May 13th). The mean is represented by the solid line and median by the dashed line. Both mean and median of the three types of response time has slightly shifted to the right from period 1 to period 3.



Coefficients	intercept	hospitalization	1st_period	2nd_period	3rd_period
Estimate	15.09774	0.40327	13.87507	7.90718	6.72668
Standard Error	2.15766	0.04341	1.99988	1.34397	1.76075
train	$r^2 = 0.85$				
test	Residual standard error: 6.619 on 209 degrees of freedom				
	$r^2 = 0.84$				
	Mean squared error: 29.012				
	Standard error of prediction residual: 5.435				


Table 7: Coefficients of Time Series Regression with Change Point detection

March - May 2020, our study found that the daily average number of emergency calls related to assault in Austin was higher during post-pandemic time (period 3) than pre-pandemic time (period 1).

Despite the overall decrease in emergency call volume, we observed statistically significant prolongation of the EMS response time. Local news reports in Austin suggested that the EMS department did not receive adequate funding to maintain sufficient personnel and ambulances during the pandemic, leading to delays in response time (62, 63). However, the prolongation of EMS response time after the beginning of Covid-19 pandemic was also found in Finland (36) and Western Pennsylvania (33). It is unclear what other factors might have contributed to the increase in EMS response time.

Previous studies reporting the patterns of EMS demand under the impact of Covid-19 in France (47, 48)

Figure 8: Covid-19 daily hospitalization. The four change points on variance in the temporal distribution of Covid-19 hospitalization are shown by the purple vertical lines.



and Israel (49) suggested that the increase of EMS calls for Covid-19 symptoms followed the same shape as for confirmed Covid-19 patients. Our study confirms this intuition by quantifying the correlation between the daily new hospitalization of Covid-19 patients and pandemic-related EMS demand. Covid-19 hospitalization projection models have been investigated extensively (52, 64, 65, 66, 67). Our regression model, which can use forecasts of the Covid-19 hospitalization count as input, can thus serve as a simple and convenient tool for EMS departments to quickly forecast pandemic-related EMS demand. Interestingly, as time progressed, the offset between pandemic EMS and hospitalization numbers became smaller. The reason remains unclear. We hypothesize that people became better at making informed decisions about Covid-19-related EMS Calls. At the beginning of the outbreak, due to a lack of knowledge of the epidemic, people might have made haste to raise red flags even though they were not contaminated by Covid-19. However, we did not find supporting evidence for our hypothesis.

## 5. Limitations and Future Directions


Our analysis and the proposed time series model was restricted to the data observed in Austin. It will be interesting to see whether a similar simple model can be fitted to forecast pandemic-related EMS demand in other cities. Moreover, the decrease in the offset between pandemic EMS and hospitalization numbers stood out very interesting. However, we are unable to find a satisfying reason and we hope for future studies to test our hypothesis.

Another limitation was that our regression model forecasts the overall number of Covid-19 EMS cases, without differentiating the priority levels or the geolocation of each incident. The ability to identify and forecast urgent incidents may improve EMS further. Additionally, a more fine-grained model incorporating spatial information may help identify incident hotspots within the city.

## 6. Conclusions

This study analyzed the impact of the Covid-19 pandemic on EMS call distributions and response time. We have confirmed that, overall, non-pandemic EMS calls in Austin have significantly decreased since the beginning of the Covid-19 pandemic, yet the proportion of non-pandemic defunct calls increased. In terms of problem types, we observed that despite an overall decrease in call volumes, some types of problems seemed

Figure 9: Time series regression with change point detection. The darker blue line represents the predicted value, the lighter blue area represents the 95% prediction interval. The red line represents the actual daily frequencies of pandemic EMS incidents, consistent with Figure 6.



to remain unaffected. Specifically, EMS calls related to traffic and Covid-19 symptoms decreased while calls related to cardiac arrest, stroke, mental health, overdose and assault did not. These results resonated with those in other geographical areas around the globe. Meanwhile, the response time of EMS was significantly prolonged since the outbreak of the Covid-19 pandemic. This information can provide an opportunity for dialogue among EMS agencies, government officials and health care partners on adjusting changes in EMS demand and shortening response time during future pandemics.

Moreover, we proposed a time series regression model with change point detection to predict the daily frequency of pandemic EMS incidents. The daily new hospitalization of Covid-19 patients proved a powerful predictor for the number of Covid-19-related EMS calls. In particular, this model suggested that for every 2.5 cases where EMS took a Covid-19 patient to a hospital, 1 person was admitted. This model may be of interest to other EMS departments as they plan for future pandemics, including the ability to predict pandemic-related calls in an effort to adjust a targeted response.

### Acknowledgement

This project is funded by the Good Systems – UT Austin Bridging Barriers Initiative (<https://bridgingbarriers.utexas.edu/good-systems/>). We would like to thank Prof. Junfeng Jiao for suggesting the CDC SVI data and Mauricio Tec for providing data on the hospital admission of Covid-19 patients in the Austin-Round Rock metropolitan statistical area. We also thank Natasha Stewart for the insightful comments on earlier drafts.

### References

- [1] W. H. Organization, et al., Coronavirus disease 2019 (COVID-19): situation report, 86 (2020).
- [2] S. M. Moghadas, A. Shoukat, M. C. Fitzpatrick, C. R. Wells, P. Sah, A. Pandey, J. D. Sachs, Z. Wang, L. A. Meyers, B. H. Singer, A. P. Galvani, Projecting hospital utilization during the COVID-19 outbreaks in the United States, *Proceedings of the National Academy of Sciences* 117 (2020) 9122–9126.
- [3] J. Willan, A. J. King, K. Jeffery, N. Bienz, Challenges for NHS hospitals during covid-19 epidemic, *BMJ* 368 (2020).
- [4] R. Guharoy, E. Krenzelok, J. Noviasky, Race Against the Pandemic: The United States and Global Health, *Journal of Emergency Medicine* 60 (2021) 402–406.
- [5] E. B. Lerner, C. D. Newgard, N. C. Mann, Effect of the Coronavirus Disease 2019 (COVID-19) Pandemic on the U.S. Emergency Medical Services System: A Preliminary Report, *Academic Emergency Medicine* 27 (2020) 693–699.

- [6] B. Boserup, M. McKenney, A. Elkbuli, The impact of the COVID-19 pandemic on emergency department visits and patient safety in the United States, *The American Journal of Emergency Medicine* 38 (2020) 1732–1736.
- [7] K. P. Hartnett, A. Kite-Powell, J. DeVies, M. A. Coletta, T. K. Boehmer, J. Adjemian, A. V. Gundlapalli, et al., Impact of the COVID-19 pandemic on emergency department visits—United States, January 1, 2019–May 30, 2020, *Morbidity and Mortality Weekly Report* 69 (2020) 699.
- [8] M. M. Jeffery, G. D'Onofrio, H. Paek, T. F. Platts-Mills, W. E. Soares, J. A. Hoppe, N. Genes, B. Nath, E. R. Melnick, Trends in emergency department visits and hospital admissions in health care systems in 5 states in the first months of the COVID-19 pandemic in the US, *JAMA internal medicine* 180 (2020) 1328–1333.
- [9] M. Lazzerini, E. Barbi, A. Apicella, F. Marchetti, F. Cardinale, G. Trobia, Delayed access or provision of care in Italy resulting from fear of COVID-19, *The Lancet Child & Adolescent Health* 4 (2020) e10–e11.
- [10] J. Thornton, Covid-19: A&E visits in England fall by 25% in week after lockdown, 2020.
- [11] A. Al-Wathinani, A. J. Hertelendy, S. Alhuishi, A. Mobrad, R. Alhazmi, M. Altuwaijri, M. Alanazi, R. Alotaibi, K. Goñiewicz, Increased Emergency Calls during the COVID-19 Pandemic in Saudi Arabia: A National Retrospective Study, *Healthcare* 9 (2021).
- [12] İ. San, E. Usul, B. Bekgöz, S. Korkut, Effects of COVID-19 pandemic on emergency medical services, *International Journal of Clinical Practice* (2020) e13885.
- [13] E. WongLaura, E. HawkinsJessica, L. MurrellKaren, et al., Where are all the patients? Addressing Covid-19 fear to encourage sick patients to seek emergency care, *NEJM Catalyst Innovations in Care Delivery* (2020).
- [14] T. Jensen, M. G. Holgersen, M. S. Jespersen, S. N. Blomberg, F. Folke, F. Lippert, H. C. Christensen, Strategies to handle increased demand in the COVID-19 crisis: A coronavirus EMS support track and a web-based self-triage system, *Prehospital Emergency Care* 25 (2020) 28–38.
- [15] A. D. Lucero, A. Lee, J. Hyun, C. Lee, C. Kahwaji, G. Miller, M. Neeki, J. Tamayo-Sarver, L. Pan, Underutilization of the Emergency Department During the COVID-19 Pandemic, *Western Journal of Emergency Medicine* 21 (2020) 15.
- [16] A. A. Butt, A. M. Azad, A. B. Kartha, N. A. Masoodi, R. Bertolini, A.-B. Abou-Samra, Volume and acuity of emergency department visits prior to and after COVID-19, *The Journal of emergency medicine* 59 (2020) 730–734.
- [17] R. Montagnon, L. Rouffilange, G. Agard, P. Benner, N. Cazes, A. Renard, Impact of the COVID-19 pandemic on emergency department use: focus on patients requiring urgent revascularization, *The Journal of Emergency Medicine* 60 (2021) 229–236.
- [18] A. R. Fernandez, R. P. Crowe, S. Bourn, S. E. Matt, A. L. Brown, A. B. Hawthorn, J. Brent Myers, COVID-19 preliminary case series: characteristics of EMS encounters with linked hospital diagnoses, *Prehospital Emergency Care* 25 (2020) 16–27.
- [19] B. Y. Yang, L. M. Barnard, J. M. Emert, C. Drucker, L. Schwarcz, C. R. Counts, D. L. Murphy, S. Guan, K. Kume, K. Rodriguez, et al., Clinical characteristics of patients with coronavirus disease 2019 (COVID-19) receiving emergency medical services in King County, Washington, *JAMA network open* 3 (2020) e2014549–e2014549.
- [20] C. Gibson, C. Ventura, G. D. Collier, Emergency Medical Services resource capacity and competency amid COVID-19 in the United States: preliminary findings from a national survey, *Heliyon* 6 (2020) e03900.
- [21] J. L. Hick, D. Hanfling, M. K. Wynia, A. T. Pavia, Duty to plan: health care, crisis standards of care, and novel coronavirus SARS-CoV-2, *NAM Perspectives* (2020).
- [22] A. Devereaux, H. Yang, G. Seda, V. Sankar, R. C. Maves, N. Karanjia, J. S. Parrish, C. Rosenberg, P. Goodman-Crews, L. Cederquist, et al., Optimizing Scarce Resource Allocation During COVID-19: Rapid Creation of a Regional Health-Care Coalition and Triage Teams in San Diego County, California, *Disaster Medicine and Public Health Preparedness* (2020) 1–7.
- [23] K. E. Schreyer, A. Daniel, L. L. King, A. Blome, M. DeAngelis, K. Stauffer, K. Desrochers, W. Donahue, N. Politarhos, C. Raab, et al., Emergency department management of the Covid-19 pandemic, *The Journal of emergency medicine* 59 (2020) 946–951.
- [24] D. L. Murphy, L. M. Barnard, C. J. Drucker, B. Y. Yang, J. M. Emert, L. Schwarcz, C. R. Counts, T. Y. Jacinto, A. M. McCoy, T. A. Morgan, J. E. Whitney, J. V. Bodenman, J. S. Duchin, M. R. Sayre, T. D. Rea, Occupational exposures and programmatic response to COVID-19 pandemic: an emergency medical services experience, *Emergency Medicine Journal* 37 (2020) 707–713.
- [25] M. Jalili, How should emergency medical services personnel protect themselves and the patients during COVID-19 pandemic?, *Frontiers in Emergency Medicine* 4 (2020) e37–e37.
- [26] H. Ehrlich, M. McKenney, A. Elkbuli, Defending the front lines during the COVID-19 pandemic: Protecting our first responders and emergency medical service personnel, *The American Journal of Emergency Medicine* (2020).
- [27] S. Spina, F. Marrazzo, M. Migliari, R. Stucchi, A. Sforza, R. Fumagalli, The response of Milan's Emergency Medical System to the COVID-19 outbreak in Italy, *The Lancet* 395 (2020) e49–e50.
- [28] J. G. Cabañas, J. G. Williams, J. M. Gallagher, J. H. Brice, COVID-19 Pandemic: The Role of EMS Physicians in a Community Response Effort, *Prehospital Emergency Care* (2020) 1–8.
- [29] D. A. Ghazali, A. Ouersighni, M. Gay, V. Audebault, T. Pavlovsky, E. Casalino, Feedback to Prepare EMS Teams to Manage Infected Patients with COVID-19: A Case Series, *Prehospital and Disaster Medicine* 35 (2020) 451–453.
- [30] B. J. Maguire, K. Shearer, J. McKeown, S. Phelps, D. R. Gerard, K. A. Handal, B. O'Neill, The ethics of PPE and EMS in the COVID-19 era, *JEMS* (2020).
- [31] A. H. Shadyab, E. M. Castillo, J. J. Brennan, T. C. Chan, V. M. Tolka, Ethnic Disparities in COVID-19 among Older Adults Presenting to the Geriatric Emergency Department, *The Journal of Emergency Medicine* (2021).
- [32] Y. Penverne, J. Jenvrin, E. Montassier, EMS dispatch center activity during the COVID-19 containment, *The American Journal of Emergency Medicine* (2020).
- [33] T. Satty, S. Ramgopal, J. Elmer, V. N. Mosesso, C. Martin-Gill, EMS responses and non-transports during the COVID-19

- pandemic, *The American Journal of Emergency Medicine* (2020).
- [34] R. Ferron, G. Agarwal, R. Cooper, D. Munkley, The effect of COVID-19 on emergency medical service call volumes and patient acuity: a cross-sectional study in Niagara, Ontario, *BMC Emergency Medicine* 21 (2021) 39.
  - [35] G. Mantica, N. Riccardi, C. Terrone, A. Gratarola, Non-COVID-19 visits to emergency departments during the pandemic: the impact of fear, *Public Health* 183 (2020) 40.
  - [36] L. Laukkonen, S. Lahtinen, J. Liisanantti, T. Kaakinen, A. Ehrola, L. Raatineni, Early impact of the COVID-19 pandemic and social restrictions on ambulance missions, *European Journal of Public Health* (2021). Ckab065.
  - [37] W. K. Hall, Management science approaches to the determination of urban ambulance requirements, *Socio-economic Planning Sciences* 5 (1971) 491–499.
  - [38] C. A. Aldrich, J. C. Hisserich, L. B. Lave, An analysis of the demand for emergency ambulance service in an urban area., *American Journal of Public Health* 61 (1971) 1156–1169.
  - [39] K. F. Siler, Predicting demand for publicly dispatched ambulances in a metropolitan area., *Health Services Research* 10 (1975) 254.
  - [40] T. O. Kvålsseth, J. M. Deems, Statistical models of the demand for emergency medical services in an urban area., *American Journal of Public Health* 69 (1979) 250–255.
  - [41] C. E. McConnel, R. W. Wilson, The demand for prehospital emergency services in an aging society, *Social Science & Medicine* 46 (1998) 1027–1031.
  - [42] H. T. Wong, P. C. Lai, Weather inference and daily demand for emergency ambulance services, *Emergency Medicine Journal* 29 (2012) 60–64.
  - [43] M. L. McCarthy, S. L. Zeger, R. Ding, D. Aronsky, N. R. Hoot, G. D. Kelen, The challenge of predicting demand for emergency department services, *Academic Emergency Medicine* 15 (2008) 337–346.
  - [44] N. Channouf, P. L'Ecuyer, A. Ingolfsson, A. N. Avramidis, The application of forecasting techniques to modeling emergency medical system calls in Calgary, Alberta, *Health care management science* 10 (2007) 25–45.
  - [45] H. Setzler, C. Saydam, S. Park, EMS call volume predictions: A comparative study, *Computers & Operations Research* 36 (2009) 1843–1851.
  - [46] E. Andrew, Z. Nehme, S. Bernard, M. J. Abramson, E. Newbigin, B. Piper, J. Dunlop, P. Holman, K. Smith, Stormy weather: a retrospective analysis of demand for emergency medical services during epidemic thunderstorm asthma, *bmj* 359 (2017).
  - [47] C.-. A.-U.-I.-I. Group, Early indicators of intensive care unit bed requirement during the COVID-19 epidemic: A retrospective study in Ile-de-France region, France, *PloS one* 15 (2020) e0241406.
  - [48] B. Riou, et al., Emergency calls are early indicators of ICU bed requirement during the COVID-19 epidemic, *medRxiv* (2020).
  - [49] E. Jaffe, R. Sonkin, R. Strugo, E. Zerath, Evolution of emergency medical calls during a pandemic - An emergency medical service during the COVID-19 outbreak, *The American journal of emergency medicine* (2020) S0735-6757(20)30527-1.
  - [50] D. Kulpanowski, University of Texas Good Systems and ATCEMS, 2021. URL: <https://data.austintexas.gov/dataset/University-of-Texas-Good-Systems-and-ATCEMS/szrz-ausx>.
  - [51] D. Kulpanowski, University of Texas Good Systems and ATCEMS 1, 2021. URL: <https://data.austintexas.gov/dataset/University-of-Texas-Good-Systems-and-ATCEMS-1/2srm-55iz>.
  - [52] M. Tec, M. Lachmann, S. J. Fox, R. Pasco, S. Woody, J. Starling, M. Dahan, K. Gaither, J. Scott, L. A. Meyers, Austin COVID-19 transmission estimates and healthcare projections (2020).
  - [53] A. J. Scott, M. Knott, A cluster analysis method for grouping means in the analysis of variance, *Biometrics* (1974) 507–512.
  - [54] R. Killick, P. Fearnhead, I. A. Eckley, Optimal detection of changepoints with a linear computational cost, *Journal of the American Statistical Association* 107 (2012) 1590–1598.
  - [55] C. H. Weiβ, Thinning operations for modeling time series of counts—a survey, *AStA Advances in Statistical Analysis* 92 (2008) 319.
  - [56] M. Siman-Tov, R. Strugo, T. Podolsky, O. Blushtein, An assessment of treatment, transport, and refusal incidence in a National EMS's routine work during COVID-19, *The American Journal of Emergency Medicine* (2021).
  - [57] F. Moroni, M. Gramigna, S. Ajello, A. Beneduce, L. Baldetti, L. M. Vilca, A. Cappelletti, A. M. Scandroglio, L. Azzalini, Collateral damage: medical care avoidance behavior among patients with myocardial infarction during the COVID-19 pandemic, *Case Reports* 2 (2020) 1620–1624.
  - [58] M. L. Doucette, A. Tucker, M. E. Auguste, A. Watkins, C. Green, F. E. Pereira, K. T. Borrup, D. Shapiro, G. Lapidus, Initial impact of COVID-19's stay-at-home order on motor vehicle traffic and crash patterns in Connecticut: an interrupted time series analysis, *Injury prevention* 27 (2021) 3–9.
  - [59] E. R. de Koning, M. J. Boogers, J. Bosch, M. de Visser, M. J. Schalij, S. L. M. A. Beeres, Emergency medical services evaluations for chest pain during first COVID-19 lockdown in Hollands-Midden, the Netherlands, *Netherlands Heart Journal* 29 (2021) 224–229.
  - [60] S. A. Goldberg, R. E. Cash, G. Peters, S. G. Weiner, P. G. Greenough, R. Seethala, The impact of COVID-19 on statewide EMS use for cardiac emergencies and stroke in Massachusetts, *Journal of the American College of Emergency Physicians Open* 2 (2021) e12351.
  - [61] K. A. Muldoon, K. M. Denize, R. Talarico, D. B. Fell, A. Sobiesiak, M. Heimerl, K. Sampsel, COVID-19 pandemic and violence: rising risks and decreasing urgent care-seeking for sexual assault and domestic violence survivors, *BMC Medicine* 19 (2021) 20.
  - [62] A. Weber, Austin EMS Medics Are Overworked And Overloaded During COVID-19. It's Taking A 'Huge Mental Toll.', 2020/07/23. URL: <https://www.kut.org/austin/2020-07-23/>

- austin-ems-medics-are-overworked-and-overloaded-during-covid-19-its-taking-a-huge-mental-toll, accessed: 2021/05/21.
- [63] T. Balagtas, Austin EMS Association says pandemic has impacted emergency response times, 2020/07/22. URL: <https://cbsaustin.com/news/local/austin-ems-association-says-pandemic-has-impacted-emergency-response-times>, accessed: 2021/05/21.
- [64] S. M. Moghadas, A. Shoukat, M. C. Fitzpatrick, C. R. Wells, P. Sah, A. Pandey, J. D. Sachs, Z. Wang, L. A. Meyers, B. H. Singer, et al., Projecting hospital utilization during the COVID-19 outbreaks in the United States, *Proceedings of the National Academy of Sciences* 117 (2020) 9122–9126.
- [65] Z. Du, X. Wang, R. Pasco, M. Petty, S. J. Fox, L. A. Meyers, Covid-19 healthcare demand projections: 22 texas cities, UT COVID-19 Consortium (2020).
- [66] E. S. Gel, M. Jahn, T. Lant, A. R. Muldoon, T. Nelson, H. M. Ross, COVID-19 healthcare demand projections: Arizona, *PloS one* 15 (2020) e0242588.
- [67] J. O. Ferstad, A. J. Gu, R. Y. Lee, I. Thapa, A. Y. Shin, J. A. Salomon, P. Glynn, N. H. Shah, A. Milstein, K. Schulman, et al., A model to forecast regional demand for COVID-19 related hospital beds, *medRxiv* (2020).
- [68] J. Chen, A. K. Gupta, Testing and locating variance changepoints with application to stock prices, *Journal of the American Statistical association* 92 (1997) 739–747.

## Appendix

### *Comparison of mean response time across hospitals*

One-way analysis of variance (ANOVA) suggested that the mean response time varied for the 6 major hospitals (F-test, p-value < 0.01 for assignment, dispatch and arrival). This is also shown in the mean plots in Figure 10. In particular, Dell Seton Medical Center, located in downtown Austin, had a lower response time than other hospitals. However, the slight difference in time may not have practical significance. A complete list of the mean and median response time for each of the 6 hospitals is given in Table 8.

	Assignment Time		Dispatch Time		Arrival Time	
	Mean	Median	Mean	Median	Mean	Median
Dell Seton Med Ctr	1.16	1.03	0.98	1.00	6.34	5.57
North Austin Hospital	1.22	1.08	1.07	1.08	6.97	6.20
Saint Davids Med Ctr	1.22	1.08	1.01	1.03	6.61	5.88
Seton Med Ctr	1.23	1.08	1.04	1.07	7.02	6.27
Seton Northwest	1.26	1.12	1.11	1.13	6.79	6.07
South Austin Hospital	1.17	1.05	1.04	1.07	7.17	6.30

Table 8: Comparison of mean response time across hospitals

### *Comparison of change point locations with respect to various choices of penalty*

To evaluate the impact of pandemic on non-pandemic incidents, we identified changes in mean and variance with binary segmentation method and BIC penalty assuming underlying normal distribution. To restrict our attention to the impact of the pandemic only, we chose the maximal number of change points as 2. When we allowed a greater number of change points, binary segmentation with both BIC penalty and Schwarz information criterion (SIC (68)) penalty also identified the date April 8th, around which the non-pandemic incidents began to increase from the lowest point. We have also tested the PELT method, which failed to produce meaningful results. To identify changes in mean and variance, the PELT method was sensitive to random noise and produced too many dates. When we restricted the PELT method to variance only, it failed to identify the date March 17th.

To identify multiple change points in the daily hospitalization data, we applied PELT method on variance with MBIC penalty assuming underlying normal distribution. We chose PELT on variance with MBIC penalty because it produced a relatively small number of change points, which could help avoid overfitting. The PELT method with other types of penalties (BIC, Akaike information criterion (AIC), SIC), as well as the Binary Segmentation method, however, produced too many change points ( $\geq 6$ ).

*Time series regression model without change point detection*

Table 9 gives the output of a single variable time series regression model. The mean squared error of this model is 40.501, which is worse than that of our proposed model (table 7).

Coefficients	intercept	hosp
Estimate	26.10542	0.27774
Standard Error	0.94579	0.02735
train		$r^2 = 0.59$
		Residual standard error: 7.665 on 212 degrees of freedom
test		$r^2 = 0.54$
		Mean squared error: 40.501
		Standard error of prediction residual: 6.420

Table 9: Coefficients of Simple Time Series Regression

Figure 10: Comparison of response time across hospitals - mean plot

