

Combinatorial Hypothesis Testing

April 13, 2023

0 Contents



1 Introduction

1

2 Moment Methods

1

3 Extension

1

1 Introduction



Suppose we observe an n -dimensional vector $\mathbf{X} = (X_1, \dots, X_n)$. The null hypothesis H_0 is that the components of \mathbf{X} are independent and identically distributed (i.i.d.) standard normal random variables. We denote the probability measure and expectation under H_0 by \mathbb{P}_0 and \mathbb{E}_0 , respectively.

Combinatorics kicks in as we consider the alternative hypotheses: consider a class $\mathcal{C} = \{S_1, \dots, S_N\}$ of N sets of indices such that $S_k \subset \{1, \dots, n\}$ for all $k = 1, \dots, N$. Under H_1 , there exists an $S \in \mathcal{C}$ such that X_i has a distribution determined by whether i is in S :

1. In its simplest form, as discussed in [1–3], we consider

$$X_i \text{ has distribution } \begin{cases} \mathcal{N}(0, 1), & \text{if } i \notin S \\ \mathcal{N}(\mu, 1), & \text{if } i \in S \end{cases}$$

where $\mu > 0$ is a positive parameter and components of \mathbf{X} are independent.

- 2.

2 Moment Methods



3 Extension



3 References



- [1] Louigi Addario-Berry, Nicolas Broutin, Luc Devroye, and Gábor Lugosi. On combinatorial testing problems. 2010.

Last updated April 13, 2023

-
- [2] Ery Arias-Castro, Emmanuel J Candes, and Arnaud Durand. Detection of an anomalous cluster in a network. *The Annals of Statistics*, pages 278–304, 2011.
 - [3] Ery Arias-Castro, Emmanuel J Candes, Hannes Helgason, and Ofer Zeitouni. Searching for a trail of evidence in a maze. 2008.