


21 forecasters on the late windows (origins after each set's training cutoff). Five baselines are given: ANKYRA's information set; the six test sets were scored once, after the model had been fixed. NH: non-household; HH: households. b: 95% unit-and-month bootstrap interval of the unit-equal log RMS ratio.

a Mean per-unit rank on the six test sets (lower is better)

	BDG2	Cambridge	HEEW	EWELD	GoiEner NH	GoiEner HH	Mean
ANKYRA	4.6	4.6	4.8	6.6	5.2	7.2	5.5
Ridge (per unit)	6.7	7.3	7.3	9.0	7.0	6.6	7.3
Chronos-2-X	7.1	6.8	6.7	9.2	7.7	9.5	7.8
iTransformer-X	10.8	5.6	7.9	9.1	7.4	8.2	8.2
TimesFM	8.5	8.9	7.1	7.9	8.7	9.6	8.4
TimesFM-X	8.5	9.1	7.5	9.2	8.4	8.5	8.6
Chronos-2	9.7	9.5	8.5	8.1	8.2	9.2	8.9
iTransformer	10.4	7.8	9.6	10.0	6.8	8.8	8.9
PatchTST	10.5	8.0	8.4	10.2	10.1	7.9	9.2
DLinear	8.4	10.7	11.3	11.6	9.0	9.3	10.0
GBT-T	8.0	7.0	7.6	13.8	14.7	9.6	10.1
TiDE	8.6	12.0	11.6	11.9	10.0	9.3	10.5
4-week profile	12.9	10.4	12.4	10.2	13.8	12.8	12.1
LSTM	12.4	15.1	15.7	14.4	11.2	9.2	13.0
Previous-month profile	14.7	14.2	14.7	10.2	13.3	11.2	13.1
Holt-Winters	14.5	12.9	13.3	12.3	12.9	12.6	13.1
MSTL	14.0	15.0	12.4	12.1	15.2	16.9	14.2
Seasonal naive (week)	14.5	15.6	13.9	12.3	16.0	16.3	14.8
Seasonal naive (day)	16.6	18.8	15.4	11.5	14.8	15.2	15.4
GBT (zero-shot)	17.2	14.9	17.3	15.1	15.8	14.9	15.9
Last-year profile	12.3	16.7	17.5	16.5	14.7	18.1	16.0

b Head-to-head on the test sets



Same information · trained
 Same information · zero-shot foundation model
 Load only · zero-shot foundation model
 Load only · statistical
 Load only · trained
 Per-unit regression (calendar, temperature)
 Profiles, naive, zero-shot GBT