

The origin-continuity layer of ankyra-lite 0.2.0 on against off, on the same carrier and anchoring layer (the seasonal naive of the state rule on state windows in both, where the layer is bypassed), on the test windows of every population, by lead time. Improvement 100[1 − exp(r)] of the hourly error with the layer, r the unit-equal log RMS ratio over the hours of the segment (hour 0; hours 0–2; forecast day 1; days 2–7; days 8–31; the whole window); 95% unit-and-month bootstrap intervals; positive = lower error with the layer. Energy: the monthly energy error of the delivered trajectory, which the layer changes; the release's energy readout is read before the layer (readout rule A) and does not change. a: the units of all populations pooled (resampled together). b: each population; filled cells and bold numbers: the interval excludes zero; the colour scale is compressed. Figure 9 shows the hourly error by forecast day against the baselines. ankyra-lite 0.2.0: carrier recipe Rw (trained from scratch with early stopping on the 93-day validation set, then 8 more epochs with the validation span returned and each training window's loss weighted by the inverse-noise rule; six seeds averaged; figure 1b), the anchoring layer, the state rule and the origin-continuity layer; hourly errors score the delivered trajectory after that layer, energy, level and peak are read before it (readout rule A).

a Pooled: 1,891 units



b By population: hourly error by lead time, and energy

	hour 0	hours 0–2	day 1	days 2–7	days 8–31	whole window	energy (deliv.)
BDG2	**+39.2**	**+32.2**	**+19.2**	**+3.7**	+0.21	**+0.79**	+0.66
Cambridge	**+22.6**	**+17.7**	**+7.6**	+0.44	−0.11	+0.20	−1.2
HEEW	**+31.9**	**+27.5**	**+14.3**	**+3.1**	**+0.28**	**+0.86**	+2.6
EWELD	**+37.0**	**+28.4**	**+7.5**	+1.1	+0.30	**+0.50**	−1.5
GoiEner NH	**+32.4**	**+29.8**	**+12.0**	**+2.4**	**+0.27**	**+0.51**	+3.3
GoiEner HH	**+18.4**	**+13.5**	**+8.4**	**+1.7**	**+0.15**	**+0.36**	+2.8
Oslo	**+35.1**	**+30.0**	**+12.6**	−0.97	**−0.53**	−0.33	−1.5
Drammen	**+42.8**	**+39.5**	**+25.1**	+2.3	−0.07	+1.0	−0.20
CINELDI	**+24.8**	**+23.9**	**+6.4**	+0.16	−0.13	+0.01	−2.8
pooled	**+28.0**	**+23.4**	**+11.1**	**+1.9**	**+0.17**	**+0.46**	+1.9



Windows per population (state windows, bypassed by the layer): BDG2 474 (33); Cambridge 730 (0); HEEW 658 (0); EWELD 523 (245); GoiEner NH 890 (13); GoiEner HH 696 (35); Oslo 581 (0); Drammen 689 (0); CINELDI 475 (0). Layer record (results/continuation_record.csv): no forecast day 1–7 of the last observed day's workday class, so forecast day 7 is the reference day, on 33 windows (EWELD 33); spans whose mean is at or below zero (x = 0, flagged; a release rule): 0. Mean of the delivered trajectory against the energy readout (rule A): median 0.77%, 99th percentile 7.8%, maximum 310% (relative to the readout).