
ankyra-lite against the zero-shot foundation-model variants on the test windows of every population, by the release's primary estimators. a: hourly error, the unit-equal RMS ratio over the 744 hours (ankyra-lite's delivered trajectory, after the continuation layer); b: monthly energy error, 744 × the error of the window mean (ankyra-lite: its trajectory before the continuation layer, readout rule A; each variant: its delivered trajectory); c: monthly peak, ankyra-lite's peak readout (read before the continuation layer) against the same readout applied to each variant's trajectory (the release's rule). The layer's own effect by lead time is in figure 9c. Improvement 100[1 − exp(r)] with 95% unit-and-month bootstrap intervals; filled = resolved in ankyra-lite's favour, red = resolved against it, open = not resolved; shaded: ankyra-lite better. Above each panel: the count of its contrasts resolved better, not resolved and resolved worse. TimesFM is TimesFM 2.5; the -X variants are given the release's shared covariate set (no observed future weather), the others read the load only. The other 16 baselines are in figures 2, 3, 5 and 11 (21 forecasters in all). ankyra-lite 0.2.0: carrier recipe Rw (trained from scratch with early stopping on the 93-day validation set, then 8 more epochs with the validation span returned and each training window's loss weighted by the inverse-noise rule; six seeds averaged; figure 1b), the anchoring layer, the state rule and the origin-continuity layer; hourly errors score the delivered trajectory after that layer, energy, level and peak are read before it (readout rule A).




A unit whose error is exactly zero drops out of a unit-equal contrast (on BDG2, the three near-zero meters, where the state rule makes ankyra-lite's hourly error zero). GoiEner NH / HH: non-household / households.