




a–c: the training-recipe study behind recipe R (the span return; the released recipe Rw adds the inverse-noise loss weights to it), on the test windows (read before: a description, not a confirmation). All three recipes start from the same checkpoints (three seeds) trained from scratch with early stopping on the 93-day validation set. F and R then train exactly 8 more epochs: F on the unchanged training set (control); R with the validation span returned to the training set. V instead re-splits with a proportional validation set (10% of the population's own record, 38–93 days) and stops early on it (5 to 30 epochs; the start weights count as epoch 0). Improvement of the first recipe over the second (unit-equal RMS ratio of the three-seed ensembles; positive = the first is better); 95% intervals combine the unit-and-month bootstrap with the spread over seeds (Welch); filled = interval excludes zero. Switch layer: short-record test windows whose season had no same-season month in the training set before the return and has one after it. On the four long-record populations V's validation set is the same 93 days. d–e: the 0.2.0 study's test read (once; the test windows had been read before, so directions only), the configurations named by recipe (R, Rw), seeds (3: seeds 0–2; 6) and the continuation layer: 0.2.0 (Rw6 with the layer) against the 0.1.0 configuration (R3) and the single changes (the layer on the same Rw6 carrier, the loss weighting Rw6 vs R6, six vs three seeds R6 vs R3); positive = the first is better; 95% intervals as recorded (z intervals that combine the unit-and-month bootstrap with the spread of the two three-seed groups where the record has them, else unit-and-month only). Energy and peak are read before the layer (readout rule A), so the layer contrast is zero by construction on the energy rows (the readout and the sum of the trajectory before the layer) and on the peak; the energy of the delivered trajectory, which the layer does change, is in figure 9c.











Resolved against R: Six development populations, days 2–7, R vs F -1.35%. Lowest population estimate of R vs F (hourly): Drammen -2.9%.