
21 forecasters (20 baselines and ankyra-lite) on the same test windows of nine populations (origins after each population's training cutoff of the trained baselines, where every forecaster has a forecast). A re-evaluation: these windows had been read before. Mean: over the populations shown. NH/HH: non-/households. b: 95% unit-and-month bootstrap interval of the log RMS ratio. ankyra-lite 0.2.0: carrier recipe Rw (trained from scratch with early stopping on the 93-day validation set, then 8 more epochs with the validation span returned and each training window's loss weighted by the inverse-noise rule; six seeds averaged; figure 1b), the anchoring layer, the state rule and the origin-continuity layer; hourly errors score the delivered trajectory after that layer, energy, level and peak are read before it (readout rule A). Linear-context: the TiDE implementation of the release, whose forecast is a linear map of the load context.

**a Mean per-unit rank on the nine test sets (lower is better)**

| | BDG2 | Cambridge | HEEW | EWELD | GoiEner NH | GoiEner HH | Oslo | Drammen | CINELDI | Mean |
|---|---|---|---|---|---|---|---|---|---|---|
| **ankyra-lite** | **4.6** | **3.0** | **5.0** | **6.9** | **4.8** | **5.1** | **1.8** | **1.7** | **2.8** | **4.0** |
| Ridge (per unit) | 6.7 | 7.4 | 7.3 | 9.0 | 7.1 | 6.7 | 9.6 | 4.7 | 5.6 | **7.1** |
| Chronos-2-X | 7.1 | 7.0 | 6.7 | 9.2 | 7.8 | 9.6 | 5.5 | 3.6 | 8.4 | **7.2** |
| iTransformer-X | 10.8 | 5.9 | 7.9 | 9.0 | 7.5 | 8.4 | 3.9 | 6.0 | 6.8 | **7.4** |
| iTransformer | 10.4 | 8.0 | 9.5 | 10.0 | 6.8 | 9.0 | 6.2 | 5.8 | 7.3 | **8.1** |
| PatchTST | 10.5 | 8.2 | 8.5 | 10.2 | 10.2 | 8.1 | 6.3 | 6.5 | 7.2 | **8.4** |
| GBT-T | 8.0 | 7.3 | 7.7 | 13.8 | 14.7 | 9.9 | 2.6 | 5.9 | 9.4 | **8.8** |
| TimesFM-X | 8.5 | 9.2 | 7.5 | 9.2 | 8.4 | 8.6 | 10.1 | 9.2 | 8.5 | **8.8** |
| TimesFM | 8.4 | 8.9 | 7.0 | 7.9 | 8.6 | 9.6 | 9.7 | 11.4 | 9.6 | **9.0** |
| Chronos-2 | 9.7 | 9.5 | 8.5 | 8.1 | 8.2 | 9.2 | 9.8 | 9.7 | 8.4 | **9.0** |
| DLinear | 8.4 | 10.8 | 11.3 | 11.5 | 9.1 | 9.5 | 9.7 | 7.2 | 7.6 | **9.5** |
| Linear-context | 8.7 | 12.1 | 11.6 | 11.9 | 10.0 | 9.5 | 9.7 | 8.2 | 7.0 | **9.8** |
| 4-week profile | 12.8 | 10.5 | 12.3 | 10.1 | 13.8 | 12.9 | 13.2 | 16.4 | 11.6 | **12.6** |
| Holt–Winters | 14.5 | 13.0 | 13.3 | 12.3 | 12.9 | 12.8 | 14.6 | 15.0 | 13.2 | **13.5** |
| LSTM | 12.4 | 15.1 | 15.8 | 14.5 | 11.3 | 9.4 | 14.9 | 15.2 | 16.2 | **13.9** |
| Previous-month profile | 14.8 | 14.2 | 14.7 | 10.2 | 13.4 | 11.3 | 17.3 | 18.8 | 14.0 | **14.3** |
| MSTL | 14.0 | 14.9 | 12.3 | 12.0 | 15.1 | 16.8 | 16.8 | 15.6 | 17.3 | **15.0** |
| Seasonal naive (week) | 14.5 | 15.6 | 13.9 | 12.2 | 16.0 | 16.3 | 16.6 | 15.9 | 17.0 | **15.3** |
| Last-year profile | 12.3 | 16.7 | 17.6 | 16.5 | 14.7 | 18.1 | 13.7 | 15.8 | 17.5 | **15.9** |
| GBT (zero-shot) | 17.2 | 14.9 | 17.2 | 15.1 | 15.8 | 15.0 | 18.6 | 18.3 | 16.1 | **16.5** |
| Seasonal naive (day) | 16.6 | 18.8 | 15.4 | 11.4 | 14.8 | 15.2 | 20.5 | 20.0 | 19.6 | **16.9** |

**b Head-to-head on the test sets**




Same information · trained
Same information · zero-shot foundation model
Load only · zero-shot foundation model
Load only · statistical
Load only · trained
Per-unit regression (calendar, temperature)
Profiles, naive, zero-shot GBT